2001-04

Primary structure and sexual stage-specific expression of a LAMMER protein kinase of *Plasmodium falciparum*.

Li, JL

<http://hdl.handle.net/10026.1/10323>

10.1016/s0020-7519(01)00126-6

Int J Parasitol

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Research note

Primary structure and sexual stage-specific expression of a LAMMER protein kinase of *Plasmodium falciparum*[☆]Ji-Liang Li^{a,b,*}, Geoffrey A.T. Targett^a, David A. Baker^a^aDepartment of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK^bInstitute of Tropical Medicine, The First Medical University, Nanfang Hospital, Guangzhou GD510515, China

Received 9 November 2000; received in revised form 2 January 2001; accepted 2 January 2001

Abstract

We have isolated a LAMMER-like gene from *Plasmodium falciparum* by vectorette technique. The gene consists of 3316 bp encoding a protein 881 amino acids with a predicted molecular mass of approximately 106.7 kDa. The encoded protein, termed PflAMMER, is composed of two distinct domains. The N-terminal domain is not related to any previously described protein kinases and has several interesting features including multiple consensus phosphorylation sites for a range of protein kinases, a number of RS/SR dipeptides, a large proportion of charged amino acids, two putative nuclear localisation signals and 14 copies of a tetramer DKYD repeats. The C-terminal domain is characteristic of a kinase in the LAMMER family with the highest homology to the *Arabidopsis thaliana* AFC3 kinase. Genomic restriction analysis showed that PflAMMER is encoded by a single copy gene in the parasite genome. A single transcript of approximately 3800 nucleotides is expressed specifically in the sexual stage, indicating that PflAMMER may be important in regulating the processes of sexual differentiation of the parasite. © 2001 Australian Society for Parasitology Inc. Published by Elsevier Science Ltd. All rights reserved.

Keywords: Malaria; Protein kinase; LAMMER; Gametocyte; *Plasmodium falciparum*

The resistance of *Plasmodium falciparum* to drugs and the resistance of mosquitoes to insecticides have resulted in a resurgence of malaria in the tropics and subtropics. Therefore, development of effective vaccines and new anti-malarial drugs has become a top priority. A better understanding of the various cellular processes at different stages of the parasite will help to identify new targets for vaccine and drug development. Gametocytogenesis within the human erythrocytes and gametogenesis in the mosquito midgut are two key processes in the parasite life cycle. However, the molecular mechanisms involved in control of the sexual development are unclear (Lobo and Kumar, 1998). We are interested in signal transduction pathways involved in the sexual differentiation of *P. falciparum*. The first step toward this goal has been to identify the components, mainly protein kinases and phosphatases, of the signal transduction pathways. Recently, we have reported several sexual stage-specific genes encoding either protein serine/threonine

phosphatases (Li and Baker, 1997, 1998) or protein kinases (Li et al., 1996, 2000). In this paper, we describe the molecular cloning and characterisation of a novel gene encoding a *P. falciparum* LAMMER protein kinase (PflAMMER). PflAMMER possesses a large N-terminal extension and is expressed specifically in the sexual stage of the parasite life cycle, indicating that it may be important in regulating the processes of sexual stage development.

Protein kinases possess a catalytic domain of approximately 30 kDa within which some sequences are highly conserved among all members of the family (Hanks et al., 1988). Initially, although our interests lay in the *P. falciparum* protein kinases in general, cyclin-dependent kinases were targeted. Therefore, two primers, cdcI (5'-GAAA-AA/GITIGGIGAA/GGGIAC-3', 1798–1817) and cdcII (5'-TCCAA/TAAGTCCATACATCACT-3', 2395–2415), were constructed based on the conserved sequences of subdomains I (EKIGEGT) and IX (SDVWSF/YG) and used in PCR. A band of 617 bp was amplified from *P. falciparum* 3D7A genomic DNA (data not shown) and inserted into the pGEM-T vector (Promega). The sequence data revealed that the PCR clone (cdcI-II) encodes 205 amino acids of a kinase-like protein. The A + T content and codon usage are typical of the coding region of *P. falciparum* genes (Weber, 1988). Southern blot analysis

[☆] Nucleotide sequence data reported in this paper are available in the GenBank™, EMBL and DDJB databases under the accession number AF104915.

* Corresponding author. Present address: Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Oxford OX3 9D5, UK. Tel.: +44-1865-222419; fax: +44-1865-222431.

E-mail address: lij@icrf.icnet.uk (J.-L. Li).

Fig. 1. A schematic representation of a partial restriction map of the *Pflammer* gene and the overlapping fragments used to determine its nucleotide sequence. A, *AccI*; B, *Bam*HI; E, *Eco*RV; H, *Hinc*II; R, *Rsa*I. The open box represents the coding region of *Pflammer*.

confirmed that the *cdcI-II* clone is derived from parasite DNA (data not shown). To isolate the full-length gene, two specific primers, PSK1 (5'-GTTTCGAACAACTTT-TACAGCGTAG-3', 1863–1887) (to obtain further sequence in the 5' direction) and PSK2 (5'-CTCCAGAAAGTTATATTAATTTGGG-3', 2357–2381) (to obtain further sequence in the 3' direction), were constructed on the basis of known sequences and used in PCR to screen vectorette libraries (Li et al., 1996). Four fragments (PSK1-*Rsa*I, PSK1-*Sau*3AI, PSK2-*Rsa*I and PSK2-*Dra*I) (Fig. 1) were obtained and sequenced. As expected, PSK1-*Sau*3AI contains all sequence of PSK1-*Rsa*I while PSK2-*Dra*I covers a full-length of PSK2-*Rsa*I. The sequence data of PSK1-*Sau*3AI permitted construction of the PSK5 primer (5'-CATACTTGTTGTCA-TATTTGTCCTC-3', 1048–1072) and subsequent screening of vectorette libraries. Based on the sequence of PSK5-*Taq*I, the PSK27 primer (5'-CATCTTGTTA-TAACGCTTCTGTTCTC-3', 518–543) was designed to produce the PSK27-*Dra*I fragment. Database searches revealed that in the *P. falciparum* tag database there was a PNABQ25TR fragment containing all sequence of PSK27-*Dra*I. Nucleotide sequence analysis revealed a putative start codon in PNABQ25TR and a stop codon in PSK2-*Dra*I. In order to confirm the sequence obtained from the overlapping fragments, a pair of primers, PSKB (5'-AGATCTCTCGA-GATGGGTTATTCATCGAATTTGTA-3', 234–257) and PSKE (5'-TTAATAGTACTCATAATT-TTCTTC-3', 2755–2778), were used to amplify the gene from genomic

DNA and the PCR product was sequenced in both strands (see Fig. 1). The sequence derived from overlapping fragments consists of 3316 bp and contains an open reading frame starting with an ATG codon at nucleotide 133 and terminating with a TAA codon at nucleotide 2776. The sequence and codon usage in the coding region are typical for a *P. falciparum* gene (Weber, 1988). The A + T contents of both flanking regions (91.9 and 84.4%, respectively) are characteristically higher than that of the coding region (75.5%). Two putative polyadenylation signals (AATAAA) are found at nucleotides +242 and +321, downstream of the TAA termination codon. A diffuse G + T-rich sequence occurs at nucleotide +24, downstream of the first polyadenylation signal. The existence of introns in the coding sequence was excluded by reverse transcription-PCRs (Li and Baker, 1997; Li et al., 2000; Li and Cox, 2000) using a number of primer pairs that cover the whole coding region.

The open reading frame encodes a protein of 881 amino acids with a predicted Mr of approximately 106.7 kDa. Database searches revealed that the amino acid sequence of PflAMMER shares 56–60% similarity and 37–41% identity with kinases in the LAMMER family across the catalytic domain. PflAMMER has the highest homology to the *Arabidopsis thaliana* AFC3 kinase (41% identity, 60% similarity) (Bender and Fink, 1994). Fig. 2 shows a sequence alignment of LAMMER kinases from diverse species. PflAMMER is composed of two distinct domains. The N-terminal domain, consisting of 547 amino acid residues, is not related to any previously described protein

Fig. 2. Alignment of the predicted amino acid sequence of PflAMMER with other members in the LAMMER family. The GenBank™/EMBL/DDJB database accession numbers are as follows: *P. falciparum* PflAMMER, AF104915; human hCLK1, P49759; hCLK2, P49760; hCLK3, P49761; mouse mCLK1, P22518; mCLK2, O35491; mCLK3, O35492; mCLK4, O35493; rat rCLK3, Q63117; *D. melanogaster* DOA, P49762; *S. cerevisiae* KNS1, P32350; *S. pombe* KATB, Q10156; *A. thaliana* AFC1, P51566; AFC2, P51567; AFC3, P51568; and *N. tabacum* PK12, U73937. Sequences were aligned with the CLUSTAL W (1.60) multiple sequence alignment programme. The amino acid residues are numbered to the left of the sequence. Identical residues are indicated with asterisks and conservative changes indicated with dots. The eleven canonical subdomains of protein kinases (Hanks et al., 1988) are indicated by roman numerals. The residues conserved in the catalytic domain of the protein kinase family are highlighted with bold and the LAMMER kinase signature motifs are shaded with black. The boundary of the N-terminal and catalytic domains is indicated by arrows.

kinases and has several interesting features. Firstly, it is rich in serine/threonine [14% (74/547)] and tyrosine [14% (73/547)], forming a number of potential phosphorylation sites for a range of known protein kinases (Kennelly and Krebs,

1991). Thus, there is a strong possibility for regulation of the PfLAMMER activity by reversible phosphorylation of the N-terminal segment. Secondly, the N-terminal domain contains many serine and arginine residues [15% (92/

```

PFLAMMER 1 MTELNIIDNNEKHSYKLYKYCRNYINDPKNMYMGYSNLYKNSPKYIYDTRNNSNN
PFLAMMER 61 NNGYHHNYNGYQIRNKKRNSFETVSNKSKHKIKSKYKYNNEPESFRDYDYKSPSY
PFLAMMER 121 SSVKKIYDREQKYNKMFVESRNRNENHNKYNINYNLSRKNYYSYKKIYIDRRNFDT
PFLAMMER 181 FGQGNIIYNNEYLLINDGSSNLKRLHYAKGFSRSRSNKYNNTYLEKSKMKMYSSRHN
PFLAMMER 241 IYRYHNGNKNKSISRWDENYTKDIYQFSSSLHYKRRSNFTRDDNSIKSKMYENKTD
PFLAMMER 301 HLYSYEDKYNKYDDKYDDKYDDKYDDKYDDKYDDKYDDKYDDKYDDKYDDKYDDKYD
PFLAMMER 361 EEEHREQEYKKKKISFNNSTNNKSIINDYDKRELKRRKKKYSNESKVDYSLKRDENH
PFLAMMER 421 HTNNNSYLNENKNSNLNNYVAVNRKRDKEYISDSNKSFGSNKGSYMYQKKKLHYDK

```

```

N-region catalytic domain
AFC2 87 IPELGGDLTP--RYKIYSKMGEGTGGVLECDWRER--KEMVAIVKVRGKYREAAIMEI
PK12 85 MFELGENTLT--RYKILKIGEGTGGVLECDWRQ--KGFVAIKIRISIKYREAAIMEV
AFC1 104 VFVVGDLTP--RYQILSKMGEGTGGVLECDWRD--KEYVAIKIRISIKYREAAIMEI
AFC3 60 VFSLRDLTP--RYKILSKMGEGTGGVLECDWRD--KEYVAIKIRISIKYREAAIMEI
hCLK1 150 IQSGDVLSA--RYEIVDTLGEAGFGKVVCEIDHKGAGRHVAVKIVKNVDYRECAARSEI
hCLK1 149 IQSGDVLSA--RYEIVDTLGEAGFGKVVCEIDHKGAGRHVAVKIVKNVDYRECAARSEI
hCLK4 148 IQSGDVLRA--RYEIVDTLGEAGFGKVVCEIDHKGAGRHVAVKIVKNVDYRECAARSEI
hCLK2 152 IYHVGDLQE--RYEIVDTLGEAGFGKVVCEIDHKGAGRHVAVKIVKNVDYRECAARSEI
hCLK2 151 IYHVGDLQE--RYEIVDTLGEAGFGKVVCEIDHKGAGRHVAVKIVKNVDYRECAARSEI
hCLK3 145 VCRIGDLQE--RYEIVDTLGEAGFGKVVCEIDHKGAGRHVAVKIVKNVDYRECAARSEI
hCLK3 145 VCRIGDLQE--RYEIVDTLGEAGFGKVVCEIDHKGAGRHVAVKIVKNVDYRECAARSEI
DOA 159 IYHVGDLQE--RYEIVDTLGEAGFGKVVCEIDHKGAGRHVAVKIVKNVDYRECAARSEI
KATB 351 IYHVGDLQE--RYEIVDTLGEAGFGKVVCEIDHKGAGRHVAVKIVKNVDYRECAARSEI
KNS1 300 IYHVGDLQE--RYEIVDTLGEAGFGKVVCEIDHKGAGRHVAVKIVKNVDYRECAARSEI
PFLAMMER 541 SWKKGMLNN--AFVIRKMGDGTGGVLECDWRD--KEYVAIKIRISIKYREAAIMEI

```

```

AFC2 144 EMLQQLGKHDKGNC--RCVOIRNPFYDNRHICIVFEKLGSSLYDFLRKNNSYRFPDLYRE
PK12 142 DVQLQLGKHDKGNC--RCVOIRNPFYDNRHICIVFEKLGSSLYDFLRKNNSYRFPDLYRE
AFC1 161 DVQLQLGKHDKGNC--RCVOIRNPFYDNRHICIVFEKLGSSLYDFLRKNNSYRFPDLYRE
AFC3 117 DVQLQLGKHDKGNC--RCVOIRNPFYDNRHICIVFEKLGSSLYDFLRKNNSYRFPDLYRE
hCLK1 208 QVLEHLNTDPNSTFRVCQMLEWFEHGHICIVFELGLSTYDFIKENSLFPFMDHHRK
hCLK1 207 QVLEHLNTDPNSTFRVCQMLEWFEHGHICIVFELGLSTYDFIKENSLFPFMDHHRK
hCLK4 206 QVLEHLNTDPNSTFRVCQMLEWFEHGHICIVFELGLSTYDFIKENSLFPFMDHHRK
hCLK2 210 NVLEKIKEDKPNKLCVQMFDFWYDGHGMCIFELGLSTYDFIKENSLFPFMDHHRK
hCLK2 209 NVLEKIKEDKPNKLCVQMFDFWYDGHGMCIFELGLSTYDFIKENSLFPFMDHHRK
hCLK3 203 NVLEKIKEDKPNKLCVQMFDFWYDGHGMCIFELGLSTYDFIKENSLFPFMDHHRK
hCLK3 203 NVLEKIKEDKPNKLCVQMFDFWYDGHGMCIFELGLSTYDFIKENSLFPFMDHHRK
hCLK3 203 NVLEKIKEDKPNKLCVQMFDFWYDGHGMCIFELGLSTYDFIKENSLFPFMDHHRK
DOA 216 NALEKIAQDKPHCDLHCKMIDWFDYDGHGMCIFELGLSTYDFIKENSLFPFMDHHRK
KATB 408 RVLTQIAHSDPTENKCIQLRDFYDGHGMCIFELGLSTYDFIKENSLFPFMDHHRK
KNS1 360 RITQTLINNDPQGGQCLLRECFDYKNCICVLDLYGRSITDFMCSNGIAGS--ALISG
PFLAMMER 598 DILKIKQNDINN--NVKYHGKFMHYDGHGMCIFELGLSTYDFIKENSLFPFMDHHRK

```

```

AFC2 203 IGWQLLECVAFMDRLHIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
PK12 201 IGWQLLECVAFMDRLHIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
AFC1 220 LGQQLLECVAFMDRLHIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
AFC3 176 LGQQLLECVAFMDRLHIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
hCLK1 268 MAYQICKSVNPLHNNKLTIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
hCLK1 267 MAYQICKSVNPLHNNKLTIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
hCLK4 266 MAYQICKSVNPLHNNKLTIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
hCLK2 270 MAYQICKSVNPLHNNKLTIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
hCLK2 269 MAYQICKSVNPLHNNKLTIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
hCLK3 263 MAYQICKSVNPLHNNKLTIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
hCLK3 263 MAYQICKSVNPLHNNKLTIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
DOA 276 MAYQICKSVNPLHNNKLTIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
KATB 468 LSQQLFVSFVAFHSLGLVHIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
KNS1 419 HCRQLRSVCFDLHGLIITIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-
PFLAMMER 657 YCIEILKALNLYKMSLTIDLKPNIDLVSSDYVVKIPEYKGS--RLQ----RDVCY--KR-

```

```

AFC2 256 VPSSAIAKVIDFGSTAYERPDHNYIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
PK12 256 VPSSAIAKVIDFGSTAYERPDHNYIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
AFC1 275 VPSSAIAKVIDFGSTAYERPDHNYIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
AFC3 228 VPSSAIAKVIDFGSTAYERPDHNYIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK1 316 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK1 315 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK4 314 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK2 318 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK2 317 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK3 311 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK3 311 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
DOA 324 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
KATB 516 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
KNS1 479 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
PFLAMMER 710 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT

```

```

AFC2 316 GEALFQTHENIHLAMMERVIG-PPFQOMLKVVDRH-----SEKYVRRG-
PK12 316 GEALFQTHENIHLAMMERVIG-PPFQOMLKVVDRH-----SEKYVRRG-
AFC1 288 GEALFQTHENIHLAMMERVIG-PPFQOMLKVVDRH-----SEKYVRRG-
AFC3 288 GEALFQTHENIHLAMMERVIG-PPFQOMLKVVDRH-----SEKYVRRG-
hCLK1 375 GFTVFPTHDSKHLAMMERVIG-PLPKHMIQKTR-----KRYFPHHD-
hCLK1 374 GFTVFPTHDSKHLAMMERVIG-PLPKHMIQKTR-----KRYFPHHD-
hCLK4 373 GFTVFPTHDSKHLAMMERVIG-PIPAHMIQKTR-----KRYFPHHD-
hCLK2 377 GFTVFPTHDSKHLAMMERVIG-PIPSRMIRKTR-----KRYFPHHD-
hCLK2 376 GFTVFPTHDSKHLAMMERVIG-PIPSRMIRKTR-----KRYFPHHD-
hCLK3 370 GFTVFPTHDSKHLAMMERVIG-PIPSRMIRKTR-----KRYFPHHD-
hCLK3 370 GFTVFPTHDSKHLAMMERVIG-PIPSRMIRKTR-----KRYFPHHD-
DOA 383 GFTVFPTHDSKHLAMMERVIG-PIPSRMIRKTR-----KRYFPHHD-
KATB 575 GFTVFPTHDSKHLAMMERVIG-PIPSRMIRKTR-----KRYFPHHD-
KNS1 539 GFTVFPTHDSKHLAMMERVIG-PIPSRMIRKTR-----KRYFPHHD-
PFLAMMER 770 GFTVFPTHDSKHLAMMERVIG-PIPSRMIRKTR-----KRYFPHHD-

```

```

AFC2 359 -RLDWPGGATSRD-----SLKAVLKL-----PRLQNL
PK12 359 -RLDWPGGATSRD-----SLKAVLKL-----PRLQNL
AFC1 379 -RLDWPGGATSRD-----SLKAVLKL-----PRLQNL
AFC3 332 -RLDWPGGATSRD-----SLKAVLKL-----PRLQNL
hCLK1 416 -RLDWDEHSSAGR-----VYRRCCK-----PLKEFM
hCLK1 415 -RLDWDEHSSAGR-----VYRRCCK-----PLKEFM
hCLK4 414 -RLDWDEHSSAGR-----VYRRCCK-----PLKEFM
hCLK2 418 -RLDWDEHSSAGR-----VYRRCCK-----PLKEFM
hCLK2 417 -RLDWDEHSSAGR-----VYRRCCK-----PLKEFM
hCLK3 411 -GLVDENSSDGR-----VYRRCCK-----PLKEFM
hCLK3 411 -GLVDENSSDGR-----VYRRCCK-----PLKEFM
DOA 423 -GLVDENSSDGR-----VYRRCCK-----PLKEFM
KATB 619 -GLVDENSSDGR-----VYRRCCK-----PLKEFM
KNS1 599 -GLVDENSSDGR-----VYRRCCK-----PLKEFM
PFLAMMER 815 -GLVDENSSDGR-----VYRRCCK-----PLKEFM

```

```

AFC2 385 IM-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
PK12 385 VM-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
AFC1 405 IM-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
AFC3 358 VS-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
hCLK1 441 LS-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
hCLK1 440 LS-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
hCLK2 443 TS-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
hCLK2 442 TS-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
hCLK3 436 LQ-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
hCLK3 436 LQ-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
DOA 436 LQ-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
KATB 645 FA-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
KNS1 659 LQ-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF
PFLAMMER 838 LY-----QHVDSHAGE-----LINMVGQLRFPDPERITAREALRHFF

```

```

AFC2 423 FARRR-----XII-----
PK12 423 FTKDQYRRL-----XII-----
AFC1 443 FTRSRREQSIIPFPNPNPHFLYNQKN-----XII-----
AFC3 396 FKSSS-----XII-----
hCLK1 477 FDLLKKSI-----XII-----
hCLK1 476 FYPLKKHT-----XII-----
hCLK4 475 FDLLKKR-----XII-----
hCLK2 479 FARLRAEPPN-KL-----WDSSRDISR-----
hCLK2 478 FARLRAEPPN-KL-----WDSSRDISR-----
hCLK3 472 FAGLTPEERS-----FHSSRNPSR-----
hCLK3 472 FAGLTPEERS-----FHSSRNPSR-----
hCLK3 472 FAGLTPEERS-----FHSSRNPSR-----
DOA 484 FDRLLPHHRVGEVSNKQPLSSGSSSRERSHSLSR
KATB 682 FTQPISSNL-----
KNS1 719 FNLGLDDGIATYNNYNTQG-----
PFLAMMER 874 LEENEYEEY-----

```

```

AFC2 256 VPSSAIAKVIDFGSTAYERPDHNYIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
PK12 256 VPSSAIAKVIDFGSTAYERPDHNYIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
AFC1 275 VPSSAIAKVIDFGSTAYERPDHNYIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
AFC3 228 VPSSAIAKVIDFGSTAYERPDHNYIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK1 316 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK1 315 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK4 314 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK2 318 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK2 317 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK3 311 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
hCLK3 311 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
DOA 324 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
KATB 516 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
KNS1 479 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT
PFLAMMER 710 LNPDIKVVDFGSATYDDEHHTIVSTRHRYAPEVILGLGWSYPCDWSVSGCIIIVELCT

```

547)], indicating that PflAMMER is an SR-like protein (Nayler et al., 1997). Close inspection of the sequence revealed 11 RS/SR dipeptides and two RSRS motifs in this region, implying that the N-terminal segment may interact with other SR proteins (Wu and Maniatis, 1993; Kohtz et al., 1994). Thirdly, a large proportion of charged amino acids [37% (203/547) K, R, E and D] exists in the N-terminal segment, suggesting further that this region may be involved in the protein-protein interactions. Fourthly, the N-terminal segment possesses at least two putative nuclear localisation signals (Dingwall and Laskey, 1991), implying that PflAMMER may be a nuclear protein. Fifthly, there are 14 copies of a tetramer DKYD repeats at residue positions 307 to 482. Finally, in contrast to those of the known kinases in the LAMMER family, the N-terminal segment is the largest extension of LAMMERs yet described, placing PflAMMER in a distinct category within the LAMMER family. The kinase catalytic domain of PflAMMER is composed of 334 residues and contains all 11 conserved subdomains of the protein kinase family (Hanks et al., 1988). It has almost all of the characteristic features of a kinase including 15 invariant residues, the ATP-binding motif and the catalytic motif (Hanks et al., 1988; Knighton et al., 1991; DeBondt et al., 1993). The conserved sequences (HTDLKPENIL) in subdomain VI and (EHLAMMESII) in subdomain X are characteristic for LAMMER kinases (Yun et al., 1994). However, a highly conserved R residue in the LAMMER motif is replaced by an S residue in PflAMMER. A similar change was also found in mammalian CLK3 kinases (Hanes et al., 1994; Becker et al., 1996; Nayler et al., 1997). By analogy with the crystallised struc-

ture of two protein kinases (Knighton et al., 1991; DeBondt et al., 1993), the LAMMER motif lies in an α -helix below the substrate-binding cleft, potentially allowing it to contact with substrates. Therefore, the R to S change in the LAMMER motif might affect substrate specificity of PflAMMER. The sequences (DLKPEN) in subdomain VI indicate that PflAMMER is a serine/threonine kinase (Hanks et al., 1988); however, the sequences (TRQYRAPE) in subdomain VIII suggest that PflAMMER could also be a tyrosine kinase (Hanks et al., 1988; Howell et al., 1991). Indeed, many kinases in the LAMMER family have been demonstrated to exhibit a dual-specificity, phosphorylating not only serine and threonine but also tyrosine (Ben-David et al., 1991; Howell et al., 1991; Yun et al., 1994; Duncan et al., 1995; Lee et al., 1996; Sessa et al., 1996). Therefore, PflAMMER may be a parasite dual-specificity protein kinase. It is worth noting that, up to date, no tyrosine kinase has been reported in *P. falciparum*.

To investigate the structural organisation of the *Pflammer* gene in the *P. falciparum* genome, clone 3D7A genomic DNA was digested with a number of restriction enzymes and analysed by Southern blotting. Hybridisation of the *cdcI-II* (see Fig. 1) probe revealed a single band in digests with *Bam*HI, *Bcl*II, *Eco*RI, *Eco*RV, *Hinc*II and *Rsa*I, and two bands in digest with *Acc*I, consistent with the restriction map (Fig. 3a), suggesting strongly that *Pflammer* is encoded by a single copy gene in the parasite genome. However, hybridisation of the same blot with the *PSK1-Sau*3AI fragment detected two bands (one predominant band and the other faint) in digests with *Acc*I, *Bcl*II, *Eco*RI, *Eco*RV and *Hinc*II, and three bands in digest with *Rsa*I, contradictory to

Fig. 3. Southern blot analysis of the *Pflammer* gene. Four micrograms of genomic DNA from *P. falciparum* clone 3D7A were digested with restriction enzymes, electrophoresed on a 1.0% agarose gel, transferred onto a nylon membrane, and probed with the *cdcI-II* fragment (a) and *PSK1-Sau*3AI fragment (b and c), respectively. The blot was washed either in a normal condition (at 56°C) (a and b) (Li et al., 1996) or in a higher condition (at 65°C) (c). Lanes 1–7 correspond to digests with *Acc*I, *Bam*HI, *Bcl*II, *Eco*RI, *Eco*RV, *Hinc*II and *Rsa*I. The sizes of 1–Kb DNA markers (M) are given in kilobase pairs to the left.

Fig. 4. Northern blot analysis of the *Pflammer* gene. Ten micrograms of total RNA extracted from the asexual erythrocytic stages (A) and the sexual erythrocytic stage (S) of *P. falciparum* (3D7A) were fractionated in a denaturing formaldehyde gel, blotted onto a nylon membrane and hybridised to radiolabeled probes. The positions of *P. falciparum* rRNA subunits (18 and 28S) are indicated by arrows. a, b, c and d are autoradiographs of the membrane probed with the *Pflammer* gene (*cdcI-II* and *PSK1-Sau3AI*), the *Pfppkac* gene (Li and Cox, 2000) and the *Pfpp-α* gene (Li and Baker, 1998), and exposed for 44, 1 and 24 h, respectively. *Pflammer* detected a transcript of approximately 3800 nucleotides in the sexual stage, *Pfppkac* hybridised with a band of approximately 1800 nucleotides in the asexual stage and *Pfpp-α* hybridised with a band of approximately 3900 nucleotides in the sexual stage.

the restriction map (Fig. 3b). The faint bands disappeared gradually with higher stringency washing conditions (Fig. 3c), indicating the presence of a *PSK1-Sau3AI*-related gene in the *P. falciparum* genome. To date, at least four LAMMER kinases have been identified in mouse (Ben-David et al., 1991; Howell et al., 1991; Nayler et al., 1997), three in human (Johnson and Smith, 1991; Hanes et al., 1994), three in *A. thaliana* (Bender and Fink, 1994), one in rat (Becker et al., 1996), one in *Drosophila melanogaster* (Yun et al., 1994), one in *Saccharomyces cerevisiae* (Padmanabha et al., 1991), one in *Schizosaccharomyces pombe* (the database accession number Q10156) and one in *Nicotiana tabacum* (Sessa et al., 1996). Our southern results suggest that only one LAMMER kinase exists in *P. falciparum*.

In order to obtain some information on how *Pflammer* mRNA levels are regulated during parasite development and differentiation, a northern blot containing equal quantities of total RNA prepared from cultures enriched in stage III to stage V gametocytes and from mixed asexual erythrocytic stages was probed with the *cdcI-II* clone. A single transcript of approximately 3800 nucleotides in size was detected only in the lane containing the sexual stage RNA, migrating just behind of the 28S rRNA bands (Fig. 4a). This was also confirmed by hybridising the same blot with the *PSK1-Sau3AI* fragment (Fig. 4b). The result is in sharp contrast to the *lammer* genes of humans, mice and insects, for which multiple transcripts were detected due

to the alternative and incomplete splicings (Howell et al., 1991; Johnson and Smith, 1991; Hanes et al., 1994; Yun et al., 1994; Duncan et al., 1995; Nayler et al., 1997). As internal controls for hybridisation to the asexual and sexual stage mRNA, the same blot was hybridised with *Pfppkac*, an asexual stage-specific gene (Li and Cox, 2000), and *Pfpp-α*, a sexual stage-specific gene (Li and Baker, 1998) (Fig. 4c,d). The result suggests that *PfLAMMER* is involved in sexual stage-specific events. In rodent, it has been shown that *mCLK2*, *mCLK3*, *mCLK4* and *rCLK3* are predominantly expressed in testis, indicating that these kinases regulate a testicular function (Becker et al., 1996; Nayler et al., 1997). In *Arabidopsis*, the *AFC1* kinase gene is able to complement yeast signal transduction mutants via activation of the transcription factor *STE12* (Bender and Fink, 1994). In *Drosophila*, *DOA* is critical to the development of the fly embryo and affects eye differentiation and sex determination by phosphorylation of the SR proteins and consequently regulation of alternative pre-mRNA splicing (Yun et al., 1994; Du et al., 1998). In mammalian cells, over expression of the *CLK1* kinase in PC12 cells seems to initiate their differentiation into neural derivatives possibly through activation of a mitogen-activated protein kinase cascade (Myers et al., 1994). The mouse CLKs have been demonstrated to bind to and phosphorylate SR-rich mRNA splicing factors (Colwill et al., 1996; Nayler et al., 1997). Overexpression of *CLK1* in COS cells leads to the subcellular redistribution of the SR-proteins, and to alternations in mRNA splicing in vivo (Colwill et al., 1996; Duncan et al., 1997). Recently, it has been shown that both *mCLK1* and *hCLK2* can phosphorylate and activate the tyrosine phosphatases *PTP-1B* and *YPTP1* in vitro (Moeslein et al., 1999). By analogy, therefore, we speculate that *PfLAMMER* may play an important role in signal transduction pathway(s) involved in the sexual differentiation of the parasite perhaps by phosphorylation of the sexual stage-specific mRNA splicing factors and/or protein phosphatases.

Acknowledgements

This work was supported in part by funds from the Wellcome Trust. J.-L. Li was supported by the Sino-British Friendship Scholarship Scheme (SBFSS), the Great Britain-China Educational Trust and Henry Lester Trust.

References

- Becker, W., Kentrup, H., Heukelbach, J., Joost, H.G., 1996. cDNA cloning and characterisation of rat *Clk3*, a LAMMER kinase predominately expressed in testis. *Biochim. Biophys. Acta* 1312, 63–67.
- Ben-David, Y., Letwin, K., Tannock, L., Bernstein, A., Pawson, T., 1991. A mammalian protein kinase with potential for serine/threonine and tyrosine phosphorylation is related to cell cycle regulators. *EMBO J.* 10, 317–25.
- Bender, J., Fink, G.R., 1994. *AFC1*, a LAMMER kinase from *Arabidopsis*

- thaliana*, activates STE12-dependent processes in yeast. Proc. Natl. Acad. Sci. USA 91, 12105–9.
- Colwill, K., Pawson, T., Andrews, B., Prasad, J., Manley, J.L., Bell, J.C., Duncan, I.P., 1996. The Clk/Sty protein kinase phosphorylates SR splicing factors and regulates their intranuclear distribution. EMBO J. 15, 265–75.
- DeBondt, H.L., Rosenblatt, J., Jancarik, J., Jones, H.D., Morgan, D., Kim, S.H., 1993. Crystal structure of cyclin-dependent kinase 2. Nature 363, 595–602.
- Dingwall, C., Laskey, R.A., 1991. Nuclear targeting sequences- a consensus? Trends Biochem. Sci. 16, 478–81.
- Du, C., McGuffin, M.E., Dauwalder, B., Rabinow, L., Mattox, W., 1998. Protein phosphorylation plays an essential role in the regulation of alternative splicing and sex determination in *Drosophila*. Mol. Cell 2, 741–50.
- Duncan, P.I., Howell, B.W., Marius, R.M., Drmanic, S., Douville, E.M.J., Bell, J.C., 1995. Alternative splicing of STY, a nuclear dual specificity kinase. J. Biol. Chem. 270, 21524–31.
- Duncan, P.I., Stojdl, D.F., Marius, R.M., Bell, J.C., 1997. In vivo regulation of alternative pre-mRNA splicing by the Clk1 protein kinase. Mol. Cell Biol. 17, 5996–6001.
- Hanes, J., von der Kammer, H., Klaudiny, J., Scheit, K.H., 1994. Characterisation by cDNA cloning of two new human protein kinases. J. Mol. Biol. 244, 665–72.
- Hanks, S.K., Quinn, A.M., Hunter, T., 1988. The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. Science 241, 42–52.
- Howell, B.W., Afar, D.E.H., Lew, J., Douville, E.M., Icely, P.L., Gray, D.A., Bell, J.C., 1991. STY, a tyrosine-phosphorylating enzyme with sequence homology to serine/threonine kinases. Mol. Cell Biol. 11, 568–72.
- Johnson, K.W., Smith, K.A., 1991. Molecular cloning of a novel human cdc2/CDC28-like protein kinase. J. Biol. Chem. 266, 3402–7.
- Kennelly, P.J., Krebs, E.G., 1991. Consensus sequences as substrate specificity determinants for protein kinases and protein phosphatases. J. Biol. Chem. 266, 15555–8.
- Knighton, D.R., Zheng, J., TenEyck, L.F., Ashford, V.A., Xuong, N.H., Taylor, S.S., Sowadski, J.M., 1991. Crystal structure of the catalytic subunit of cyclic adenosine monophosphate-dependent protein kinase. Science 253, 407–14.
- Kohtz, J.D., Jamison, S.F., Will, C.L., Zuo, P., Luhrmann, R., Garcia-Blanco, M.A., Manley, J.L., 1994. Protein-protein interactions and 5' splice-site recognition in mammalian mRNA precursors. Nature 368, 119–24.
- Lee, K., Du, C., Horn, M., Rabinow, L., 1996. Activity and autophosphorylation of LAMMER protein kinases. J. Biol. Chem. 271, 27299–303.
- Li, J.L., Baker, D.A., 1997. Protein phosphatase β , a putative type-2A protein phosphatase from the human malaria parasite *Plasmodium falciparum*. Eur. J. Biochem. 249, 98–106.
- Li, J.L., Baker, D.A., 1998. A putative protein serine/threonine phosphatase from *Plasmodium falciparum* contains a large N-terminal extension and five unique inserts in the catalytic domain. Mol. Biochem. Parasitol. 95, 287–95.
- Li, J.L., Cox, L.S., 2000. Isolation and characterisation of a cAMP-dependent protein kinase catalytic subunit gene from *Plasmodium falciparum*. Mol. Biochem. Parasitol. 109, 157–63.
- Li, J.L., Robson, K.J.H., Chen, J.L., Targett, G.A.T., Baker, D.A., 1996. Pfmkr, a MO15-related protein kinase from *Plasmodium falciparum*: gene cloning, sequence, stage-specific expression and chromosome localisation. Eur. J. Biochem. 241, 805–13.
- Li, J.L., Baker, D.A., Cox, L.S., 2000. Sexual stage-specific expression of a third calcium-dependent protein kinase from *Plasmodium falciparum*. Biochim. Biophys. Acta 1491, 341–9.
- Lobo, C.A., Kumar, N., 1998. Sexual differentiation and development in the malaria parasite. Parasitol. Today 14, 146–50.
- Moeslein, F.M., Myers, M.P., Landreth, G.E., 1999. The CLK family kinases, CLK1 and CLK2, phosphorylate and activate the tyrosine phosphatase, PTP-1B. J. Biol. Chem. 274, 26697–704.
- Myers, M.P., Murphy, M.B., Landreth, G., 1994. The dual-specificity CLK kinase induces neuronal differentiation of PC12 cells. Mol. Cell Biol. 14, 6954–61.
- Nayler, O., Stamm, S., Ullrich, A., 1997. Characterisation and comparison of four serine- and arginine-rich (SR) protein kinases. Biochem. J. 326, 693–700.
- Padmanabha, R., Gehrung, S., Snyder, M., 1991. The KNS1 gene of *Saccharomyces cerevisiae* encodes a non-essential protein kinase homologue that is distantly related to members of the CDC28/cdc2 gene family. Mol. Gen. Genet. 229, 1–9.
- Sessa, G., Raz, V., Savaldi, S., Fluhr, R., 1996. PK12, a plant dual-specificity protein kinase of the LAMMER family, is regulated by the hormone ethylene. Plant Cell 8, 2223–34.
- Weber, J.L., 1988. Molecular biology of malaria parasites. Exp. Parasitol. 66, 143–70.
- Wu, J.Y., Maniatis, T., 1993. Specific interactions between proteins implicated in splice site selection and regulated alternative splicing. Cell 75, 1061–70.
- Yun, B., Farkas, R., Lee, K., Rabinow, L., 1994. The *Doa* locus encodes a member of a new protein kinase family and is essential for eye and embryonic development in *Drosophila melanogaster*. Genes Dev. 8, 1160–73.