

2018-03-02

Haloferax volcanii Proteome Response to Deletion of a Rhomboid Protease Gene

Costa, MI

<http://hdl.handle.net/10026.1/11853>

10.1021/acs.jproteome.7b00530

Journal of Proteome Research

American Chemical Society

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

This is an accepted manuscript of an article published by ACS in Journal of Proteome Research (accepted January 5 2018) available at:

<https://pubs.acs.org/doi/10.1021/acs.jproteome.7b00530>

=====

1
2
3 *Haloferax volcanii* proteome response to deletion of a rhomboid protease
4
5
6 gene
7

8
9
10 Mariana I. Costa¹✉, Micaela Cerletti¹✉, Roberto A. Paggi¹, Christian Trötschel², Rosana E.
11
12 De Castro¹, Ansgar Poetsch^{2,3*} and María I. Giménez^{1*}.
13
14

15
16 ¹Instituto de Investigaciones Biológicas, Universidad Nacional de Mar del Plata (UNMDP). Consejo
17
18 Nacional de Investigaciones Científicas y Técnicas (CONICET), Funes 3250 4^{to} nivel, (7600) Mar
19
20 del Plata, Provincia de Buenos Aires, Argentina. ²Plant Biochemistry, Faculty of Biology &
21
22 Biotechnology, Ruhr University Bochum, 44801 Bochum, Germany. ³ School of Biomedical and
23
24 Healthcare Sciences, Plymouth University, Plymouth PL4 8AA, United Kingdom
25
26
27
28
29

30
31 Key words: *Archaea*, *Haloferax volcanii*, rhomboid protease, shotgun proteomics, protease
32
33 substrate.
34
35
36
37
38

39 **Abstract**
40
41

42 Rhomboids are conserved intramembrane serine proteases involved in cell signaling processes.
43
44 Their role in prokaryotes is scarcely known and remains to be investigated in *Archaea*. We
45
46 previously constructed a rhomboid homolog deletion mutant ($\Delta\rho II$) in *Haloferax volcanii*, which
47
48 showed reduced motility, increased novobiocin sensitivity and an N- glycosylation defect. To
49
50 address the impact of *rhoII* deletion on *H. volcanii* physiology, the proteomes of mutant and
51
52 parental strains were compared by shotgun proteomics. A total of 1847 proteins were identified
53
54 (45.8 % of *H. volcanii* predicted proteome) from which 103 differed in amount. Additionally, the
55
56
57
58
59
60

1
2
3 mutant strain evidenced 99 proteins with altered electrophoretic migration, suggesting differential
4 post-translational processing/modification. Integral membrane proteins that evidenced variations in
5 concentration, electrophoretic migration and/or semi-tryptic cleavage in the mutant were considered
6 as potential RhoII targets. These included a PrsW protease homolog (which was less stable in the
7 mutant strain), a predicted halocyanin and six integral membrane proteins potentially related to the
8 mutant glycosylation (S-layer glycoprotein, Agl15) and cell adhesion/motility (flagellin1,
9 HVO_1153, PilA1 and PibD) defects. This study investigated for the first time the impact of a
10 rhomboid protease on the whole proteome of an organism.
11
12
13
14
15
16
17
18
19
20
21
22
23

24 **Introduction**

25
26
27 Proteases cleave their protein/peptide substrates in an irreversible reaction and through this activity
28 they catalyze the fate and activity of other proteins. They control appropriate protein folding,
29 localization, activation (or inactivation) and shedding from cell surfaces. Therefore, proteases
30 control a wide range of important cellular functions including DNA replication, cell cycle
31 progression, cell proliferation, cell-cell communication, differentiation, morphogenesis, neuronal
32 outgrowth, homeostasis, wound healing, immunity, angiogenesis and apoptosis^{1,2}.
33
34
35
36
37
38
39
40

41 Intramembrane proteolysis, that is, the cleavage of peptide bonds within the plane of the cell
42 membrane, is a mechanism conserved in the three Domains of Life. The genomes of every
43 sequenced organism encode for intramembrane proteases (IMPs). These enzymes exert diverse
44 biological functions and are considered as promising future drug targets³. Among IMPs, three major
45 classes have been described, the GxGD-aspartyl proteases (eukaryal signal peptide peptidase SPP
46 and presenilin families), rhomboids and site 2 proteases (S2P)².
47
48
49
50
51
52
53
54

55 Rhomboid proteases (Rho) are widespread serine proteases and to date they are the best understood
56
57
58
59
60

1
2
3 IMPs⁴. In eukaryotic organisms they have been implicated in a variety of processes. In *Drosophila*
4 *melanogaster* a Rho activates epidermal growth factor receptors^{5,6}. In mitochondria the mammalian
5 PARL rhomboid protease regulates cell homeostasis and has been implicated in the biology of
6
7 diabetes and Parkinson's disease⁷. Rhomboids in parasites regulate adhesion to the host through
8 processing of adhesins⁸. Recently a role in fungal hypoxia adaptation through sterol regulatory
9 element-binding protein activation has been reported⁹.

10
11
12
13
14
15
16
17 In prokaryotes, the role of Rho has been explored to a lesser extent. Up to date, the function of only
18 one prokaryotic Rho, AarA from *Providencia stuartii*, has been elucidated. AarA cleaves an N-
19 terminal extension of the twin arginine translocation (Tat) complex component TatA, allowing for
20 the assembly of a functional Tat translocon and, hence, for the export of a still unidentified *quorum*
21 *sensing* signal¹⁰. In addition, some bacterial rhomboid deletion mutants have been characterized,
22 even though no specific protease substrates were reported. A null mutant in the *Escherichia coli*
23 rhomboid protease GlpG showed no evident phenotype under laboratory conditions, except for a
24 mild increase in resistance to cefotaxime. However, GlpG was shown to cleave heterologous
25 substrates. Specific targets of other Rho, such as Spitz, Gurken and Keren of *D. melanogaster*, were
26 proteolytically processed by GlpG¹¹. A chimeric protein, consisting of the N-terminal and
27 periplasmically localized β -lactamase domain, a LacY-derived transmembrane segment (TMS) and
28 a cytoplasmic maltose binding protein domain, could be cleaved *in vitro* by this Rho¹² and, in
29 addition, GlpG could cleave truncated forms of the transporter protein MdfA¹³. In *Bacillus subtilis* a
30 rhomboid protease gene (*yqgP*) null mutant displayed a cell division defect and decreased glucose
31 uptake¹⁴. *Mycobacterium smegmatis* rhomboid null mutants exhibited impaired biofilm formation
32 and augmented antibiotic sensitivity¹⁵.

33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51 The Rho family is widely represented through archaeal genomes¹⁶. Particularly, the halophilic
52 model archaeon *Haloferax volcanii* encodes two Rho homologs, HVO_1474 (RhoI) and HVO_0727
53 (RhoII). We have previously obtained and characterized a Rho mutant deleted in the *rhoII* gene in
54
55
56
57
58
59
60

1
2
3 *H. volcanii*. The mutant strain (MIG1) showed reduced motility on soft agar plates, increased
4
5 novobiocin sensitivity and a truncation in a novel oligosaccharide bound to position N732 of the S-
6
7 layer glycoprotein (SLG)¹⁷. However, global changes on the proteome that may explain these
8
9 phenotypes still need to be determined.
10

11
12 Comparative proteomics is a valuable tool to analyze the impact of the absence/overexpression of a
13
14 particular protease on the proteome of a certain organism and allows for identification of putative
15
16 endogenous substrates. In the case of processive proteases, accumulation of a particular protein in a
17
18 protease defective strain with respect to the control strain would be an evidence of that protein
19
20 being a specific substrate. This method was applied by our group to identify candidate targets of the
21
22 LonB protease in *H. volcanii*¹⁸. On the other hand, the situation for regulatory enzymes, such as
23
24 Rho, is different. These proteases cut at specific sites releasing or exposing fragments of the
25
26 substrate, and therefore, protease activity may be evidenced by changes in electrophoretic migration
27
28 and/or stability of the (unprocessed) substrate protein.
29
30
31

32
33 In this work, a quantitative high-throughput comparative proteomics approach was applied in *H.*
34
35 *volcanii* to understand the importance of Rho in archaeal physiology and to identify putative natural
36
37 targets of RhoII.
38
39
40
41
42

43 **Experimental Section**

44 **Strains and growth conditions**

45
46
47
48
49
50 *H. volcanii* H26 (DS70 Δ *pyrE2*)¹⁹ and MIG1¹⁷ were grown in MGM 18% or Hv-Min medium²⁰ at
51
52 42 °C and 150 rpm. When needed, uracil or tryptophan (50 μ g/ml) was added to Hv-Min.
53
54

55
56 *E. coli* DH10 β (Invitrogen) and GM33 (Marinus, 1973) were grown in Luria-Bertani (LB) medium
57
58
59
60

1
2
3 at 37 °C and 150 rpm. Kanamycin (50 µg/ml) or ampicillin (100 µg/ml) was added to the medium
4
5 when indicated. *E. coli* cells were transformed by the CaCl₂ method²¹.
6
7

8 **Cell fractionation**

9

10
11 *H. volcanii* cultures were grown up to the indicated optical density at 600 nm (OD₆₀₀) and cells were
12 harvested by centrifugation (10,000 xg 10 min, 4 °C). Culture medium was re-centrifuged under the
13 same conditions and the supernatant was passed through a 0.22 µm nitrocellulose filter and
14 precipitated with TCA 10% (v/v). Precipitated proteins were washed twice with 80% (v/v) acetone
15 and left to dry 30 min at room temperature. Cell pellets were resuspended in 50 mM HCl-Tris, 2 M
16 NaCl (pH 7.5) and disrupted by ultrasound (3 x 30 s, 80 W). Lysates were clarified (17,000 xg for
17 20 min at 4 °C) and membranes were pelleted by centrifugation (100,000 xg for 1 h at 4 °C). The
18 membranes were washed with the same buffer and both, supernatant (cytoplasmic fraction) and
19 membranes, were re-centrifuged for 30 min in the same conditions. To remove remaining salts,
20 cytoplasm and membrane proteins were precipitated overnight with 100% acetone at 4 °C followed
21 by centrifugation (17,000 xg for 20 min at 4 °C). The precipitated proteins were washed twice with
22 80% (v/v) and once with 100% acetone and left to dry for 30 min at room temperature. All samples
23 were resuspended in 1x SDS-PAGE loading buffer²² containing 0.1% (w/v) SDS and 0.1 M DTT,
24 incubated for 10 min at 70 °C and conserved at -20 °C until used.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 **SDS-PAGE**

43
44

45 Samples (30 µg per lane) were loaded onto 10% (w/v) polyacrylamide gels containing 0.1% SDS.
46
47 The gels were run at room temperature (20 mA) until samples passed the stacking gel and all
48 proteins were concentrated into one protein band in the separation gel. For the PROTOMAP
49 assay²³, samples were allowed to run 2 cm beyond the stacking gel (Fig. S1). Proteins were
50 visualized with a coomassie brilliant blue (CBB-G250) stain²⁴.
51
52
53
54
55
56
57
58
59
60

In-gel tryptic digestion

Protein bands were excised from the gels and cut into small cubes (ca. 1 × 1 mm) which were completely destained according to Schluesener and colleagues²⁵. In the case of PROTOMAP assay, each lane was divided in four 0.5 cm sections which were treated separately in the same manner (Figure 1). Gel pieces were dried by incubation with 100% acetonitrile for 10 min at room temperature and then incubated with 50 mM DTT in 25 mM NH₄HCO₃ (30 min at 60 °C) to reduce proteins disulfide bonds. The gel pieces were dried again with acetonitrile and proteins were alkylated treating the gel pieces with 50 mM iodoacetamide in 25 mM NH₄HCO₃ (1 h at room temperature in darkness). After washing (with destaining solution) and drying the gel pieces in a SpeedVac, trypsin (porcine, sequencing grade, Promega) solution (12.5 ng ml⁻¹ in 25 mM ammonium bicarbonate, pH 8.6) was added. Protein digestion and peptide elution was performed according to Cerletti et al¹⁸. The extracted peptides were dried using a SpeedVac and stored at -20 °C. Before MS-analysis peptides were resuspended in 20 µl of buffer A (0.1% formic acid in water, ULC/MS, Biosolve, Netherlands) with 2% (v/v) acetonitrile by sonication for 10 min and transferred to LC-MS grade glass vials (12 × 32 mm glass screw neck vial, Waters, USA). Each measurement was performed with 8 µl of sample.

One-dimensional nLC-ESI-MS/MS

An UPLC HSS T3 column (1.8 µm, 75 µm × 150 mm, Waters, Milford, MA, USA) and an UPLC Symmetry C₁₈ trapping column (5 µm, 180 µm × 20 mm, Waters, Milford, MA, USA) for LC as well as a PicoTip Emitter (SilicaTip, 10 µm i.d., New Objective, Woburn, MA, USA) were used in combination with the nanoACQUITY gradient UPLC pump system (Waters, Milford, MA, USA) coupled to a LTQ Orbitrap Elite mass spectrometer (Thermo Fisher Scientific Inc., Waltham, MA, USA). For elution of the peptides a linear gradient with increasing concentration of buffer B (0.1% formic acid in acetonitrile, ULC/MS, Biosolve, Netherlands) from 1% to 40% within 165 min was

1
2
3 applied, followed by a linear gradient from 40% to 99% acetonitrile concentration within 15 min
4
5 (0–5 min: 1% buffer B; 5–10 min: 5% buffer B; 10–165 min: 40% buffer B; 165–180 min: 99%
6
7 buffer B; 180–195 min: 1% buffer B) at a flow rate of 400 nL min⁻⁵ and a spray voltage of 1.5–
8
9 1.8 kV. The column was re-equilibrated at 1% buffer B within 15 min. The analytical column oven
10
11 was set to 55 °C and the heated desolvation capillary was set to 275 °C. The LTQ Orbitrap Elite via
12
13 instrument method files of Xcalibur (Rev. 2.1.0) in positive ion mode. The linear ion trap and
14
15 Orbitrap were operated in parallel, i.e. during a full MS scan on the Orbitrap in the range of 150–
16
17 2,000 *m/z* at a resolution of 240,000 MS/MS spectra of the 20 most intense precursors, from most
18
19 intense to least intense, were detected in the ion trap. The relative collision energy for collision-
20
21 induced dissociation (CID) was set to 35%. Dynamic exclusion was enabled with a repeat count of
22
23 1 and 60 seconds exclusion duration window. Singly charged and ions of unknown charge state
24
25 were rejected from MS/MS.
26
27
28

29 **Protein identification and quantification**

30
31
32 Protein identification was performed by SEQUEST²⁶ and MS Amanda²⁷ algorithms embedded in
33
34 Proteome Discoverer 1.4 (Thermo Electron © 2008–2012) searching against the complete proteome
35
36 database of *H. volcanii* DS2 containing 4035 entries exported from the Halolex database²⁸ on
37
38 9/24/2013. The mass tolerance for precursor ions and fragment ions was set to 6 ppm and 0.4 Da,
39
40 respectively. Only tryptic peptides with up to two missed cleavages were accepted. As static peptide
41
42 modification the carbamidomethylation of cysteine and as variable peptide modification the
43
44 oxidation of methionine as well as the conversion of glutamine (Gln) to pyro-glutamate (pyro-glu)
45
46 at any N-terminus were admitted. The false discovery rate (FDR) was determined with the
47
48 percolator validation in Proteome Discoverer 1.4 and the q-value was set to 1%²⁹. For protein
49
50 identification the mass spec format-(msf)-files were filtered with peptide confidence “high” and two
51
52 different peptides per protein. Additionally protein grouping options were enabled as default, which
53
54 means consider only PSMs (peptide scoring mass spectrum matches), with confidence at least
55
56
57
58
59
60

1
2
3 “medium” and consider only PSMs with delta CN better than 0.15. Proteins were quantified by
4 spectral counting^{30, 31} and a protein was considered as significantly regulated at a p value ≤ 0.05
5 (Student’s t-test) and fold ≥ 1.5 . Only proteins that had on average at least 5 PSMs were taken into
6 consideration.
7
8
9
10

11 **Generation of an expression vector of His tagged PrsW**

12
13
14
15
16 The predicted PrsW protease gene (HVO_0408) was PCR-amplified from *H. volcanii* H26 genomic
17 DNA. The amplicon (1038 bp) was synthesized using primers FWPrsWNcoI (5’
18 ATATCCATGGACCAGCAGGACCCA 3’) and RvPrsWBamHI (5’
19 ATATGGATCCTCAGTCCTCGAACTCG 3’) in a reaction mixture containing Phusion GC buffer
20 (1x), 0.2 mM dNTPs, 0.6 mM of each primer 4% (v/v) DMSO and 0.5 units of Phusion DNA
21 polymerase (New England Biolabs) in a final volume of 25 μ l. The program used was 98 °C (30
22 sec), 30 cycles of 98 °C (10 sec)-60 °C (30 sec)-72 °C (30 sec) and a final extension of 72 °C (1
23 min). The PCR product was cloned in the ZeroBlunt® TOPO cloning vector (ThermoFisher, Inc.).
24 The identity of the amplification product was confirmed by sequencing (Macrogen, Korea). The
25 DNA fragment corresponding to PrsW was excised from the ZeroBlunt® TOPO by digestion with
26 NcoI and BamHI and subcloned with an N- terminal 6x his tag in the *H. volcanii* expression vector
27 pTA963³². The ligation product was transformed into *E. coli* DH10 β and then passed through *E.*
28 *coli* GM33 in order to obtain non-methylated DNA and finally, introduced in *H. volcanii* H26 or
29 MIG1 by the standard PEG method²⁰.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 **Protein stability determination**

47
48
49
50 *H. volcanii* H26 and MIG1 harboring the pTA963-PrsW his tagged construction were grown until
51 they reached a OD₆₀₀ of 0.3-0.5 and then 1 mM tryptophan was added and left O/N in order to
52 induce the *ptnaA* promoter. After incubation with tryptophan, 100 μ g/ml puromycin was added to
53 the cultures. Samples (5 ml) were taken at the indicated times and membrane fractions were
54
55
56
57
58
59
60

1
2
3 obtained as detailed above.
4
5

6 **Western blotting**

7

8
9 Polyacrylamide gels (12% w/v) were transferred to PVDF membranes (80 mA, 2h) and blocked
10 with TBST buffer [25mM Tris-HCl, pH 7.4, 0.01% (v/v) Tween 20)] with 5% (w/v) skimmed milk.
11
12 Membranes were incubated with 1:5000 anti-His antibody (Biolegend) in blocking buffer, washed
13 with TBST (3 x 10 min), and incubated with 1:10,000 alkaline phosphatase-conjugated anti-mouse
14 IgG (Sigma-Aldrich) for 2 h. After washing (TBST, 3 x 10 min), blots were developed with 0.33
15 mg/ml nitro blue tetrazolium and 0.01 mg/ml 5-bromo-4-chloro-3-indolyl-phosphate.
16
17
18
19
20
21
22

23 **Cell adhesion assay**

24

25
26 Surface adhesion was assessed based on the method described by Tripepi et al³³. *H. volcanii*
27 cultures (3 ml) at an OD₆₀₀ of ~0.3 were incubated in 50 ml sterile screw cap plastic tubes. Glass
28 coverslips were inserted into each tube at an angle of 90° and were incubated without shaking. After
29 the indicated time intervals, coverslips were removed with forceps, submerged for 2 min in 2%
30 (v/v) acetic acid and allowed to air dry. Dry coverslips were stained in 0.1% (w/v) crystal violet for
31 10 min, washed (x3) with distilled water, air dried and then observed by light microscopy using a
32 Nikon Eclipse Ti microscope.
33
34
35
36
37
38
39
40
41
42
43
44

45 **Results and Discussion**

46

47 **Global analysis of total proteins identified in *H. volcanii* wt and $\Delta\rho$ II (MIG1)**

48
49
50
51 **strains**

52
53
54 In order to investigate the relevance of Rho in *H. volcanii* physiology, the differential proteome
55
56
57
58
59
60

1
2
3 map between the parent (H26) and a mutant strain (MIG1) containing a chromosomal deletion in
4 one of the two Rho homolog genes, *rhoII* (HVO_0727), was assessed. Culture media supernatant,
5 membrane and cytoplasmic proteins obtained from triplicate cultures of both strains harvested in
6 exponential (exp) (OD₆₀₀ 0.5-0.6) and stationary (st) (OD₆₀₀ 1.8-1.9) growth phases were subjected
7 to SDS-PAGE, digested with trypsin and then analyzed by nano-LC-MS/MS as described in the
8 Experimental Section. As Rho exert their proteolytic activity within the membrane environment,
9 their natural substrates are expected to be integral membrane proteins. Thus, in order to analyze the
10 membrane fraction in more detail in search of putative RhoII direct targets, the same assay was
11 repeated only with membrane fractions (triplicates) from *H. volcanii* H26 and MIG1 at exp growth
12 phase.
13
14
15
16
17
18
19
20
21
22
23
24

25 Out of 4035 total proteins from the predicted *H. volcanii* proteome, 1847 (45.8 %) were identified,
26 including all strains, growth phases and cell fractions of both experiments. The complete list of
27 detected proteins is included in Table S1. From these, 276 (14.9 %) are predicted to be integral
28 membrane proteins, 133 (7.2 %) membrane-associated either by a predicted signal peptide and/or
29 lipobox, and the remaining 1438 (77.8 %) as soluble proteins (Table S1). Figure 1 shows the
30 number of identified proteins distributed according to strain and growth phase (Fig. 1A) and
31 functional category (Fig. 1B). The protease RhoII (HVO_0727) was not detected in membrane
32 fractions of *H. volcanii* H26 (exp and st phase) in none of the two assays. This outcome was not
33 unexpected since integral membrane proteins containing several TMS and/or hydrophobic stretches,
34 such as RhoII are hardly detected by MS technology, as they are underrepresented in trypsin
35 digestion products³⁴. However, when these samples were used in a PROTOMAP assay, RhoII was
36 detected in the parental strain but not in the mutant confirming the absence of this protease in MIG1
37 (see below).
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Total identified protein distribution in *H. volcanii* H26 and MIG1 strains. Protein detection was based on a minimum of two different peptides (in at least one biological replicate) with a minimal false discovery rate of $q\text{-value} \leq 1\%$. **A.** Venn diagrams³⁵ showing the distribution of identified proteins in both strains and growth stages (center) and for each growth stage, the distribution of proteins according to cellular fraction (upper diagrams). **B.** Functional categories corresponding to identified proteins, according to HaloLex Database. **MET** (metabolism, blue): AA amino acid metabolism, CHM carbohydrate metabolism, CIM central intermediary metabolism, COM coenzyme metabolism, EM energy metabolism, LIP lipid metabolism, NUM nucleotide metabolism. **GIP** (genetic information processing, green): CHP chaperones, RMT RNA maturation, RRR replication, repair, recombination, TC transcription, TL translation. **TP_CP** (transport and cellular processes, red): CP cellular processes, SEC protein secretion, MOT motility, TP small molecule transport. **ENV** (environmental information processing, grey): REG gene regulation, SIG signal transduction. **MIS** (miscellaneous functions, purple): GEN general enzymatic function: ISH transposases and ISH-encoded proteins, MIS miscellaneous. **UNASS** (unassigned function, yellow): CHY conserved hypothetical protein, HY hypothetical protein.

Proteins differentially represented in the *H. volcanii* RhoII deficient strain

Using the criteria described in M&M, a total of 103 proteins were differentially represented between the *H. volcanii* H26 parent and MIG1 strains. Table 1 shows the proteins which evidenced significant variation in amount, organized by growth phase and cellular fraction. Thirty-three proteins (4 from culture medium supernatants, 9 from membrane and 20 from cytoplasm) were enriched while 70 proteins (23 from culture medium supernatant, 32 from membrane and 15 from cytoplasm) decreased their concentration. Most of the proteins that showed variation in their concentration were detected in membrane fractions, in accordance with RhoII exerting its role within the membrane.

The proteins differing between strains in the exp and st growth phases were not coincident, suggesting a growth phase-dependent effect of the rhoII knock-out (Table 1). The impact of the RhoII protease deficiency in cytoplasm and membrane protein samples was more evident in vegetative cells (exp phase) than in non-actively growing cells (st phase), since 60 vs. 16 proteins varied in abundance, respectively. The opposite situation was observed for the secretome fraction in which only 3 proteins were differential (accumulated) in vegetative cells, while 24 varied in concentration (1 increased and 23 decreased) in st phase cultures. It is likely that changes in the cell proteome, produced in absence of rhoII during the exp growth phase, may be reflected on the secretome of the mutant at a later stage of growth. A volcano plot (Fig. 2) was generated in order to visualize the dataset changes obtained in these experiments, by plotting $-\log_{10}$ (p-value) vs. \log_2 (fold-change) for proteins differentially detected in MIG1 samples taken in exp (Fig. 2A) and st (Fig. 2B) phase of growth. Of those proteins meeting statistically significant criteria (p value ≤ 0.05 ; fold ≥ 1.5), 63 were detected in exp phase (27 overrepresented, 36 underrepresented) and 40 in st phase (6 enriched, 34 underrepresented).

The differential proteins belonged to several functional classes (Figure 3, Table 1). Even though the

1
2
3 majority of these proteins corresponded to the miscellaneous and general metabolic pathways,
4
5 proteins involved in translation and solute transport were the most represented, suggesting that these
6
7 processes are affected by RhoII. However, the growth performance of the mutant was not affected
8
9 at standard laboratory conditions¹⁷, probably due to a functional overlap between the two Rho
10
11 homologs present in *H. volcanii*.
12

13
14 In cytoplasm fractions, 35 proteins were differentially represented in the mutant cells. As all Rho
15
16 substrates identified so far are integral membrane proteins, these differential proteins are probably a
17
18 consequence of yet unidentified proteolytic and/or gene expression processes regulated by RhoII,
19
20 representing an indirect/secondary effect of the knock-out mutation.
21
22

23
24 Differential proteins with predicted TMS that may have been released to the cytoplasm as a
25
26 consequence of RhoII activity were not identified in the cytoplasm. This observation suggests that,
27
28 if present, they were either degraded or not detected under the experimental conditions used in this
29
30 work.
31
32

33
34 The proteins which showed altered levels in the cytoplasm fractions were very diverse and they
35
36 could not be associated to a defined route or metabolic pathway.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Graphical representation of quantitative proteomics data. Total detected proteins from exp (A) and st (B) growth phases are ranked in a volcano plot according to their statistical p -value (\log_{10}) and their relative abundance ratio (\log_2 fold change) between MIG1 and H26 strains (x-axis). Off-centered spots are those that vary the most between both groups. Each data point represents a single unique protein identified in our proteomic analysis, color coded as follows: gray indicates no significant change and colored dots indicate significant negative (red) and positive (green) changes.




Figure 3. Distribution of the MIG1 differentially expressed proteins into functional classes, according to the Halolex database. For abbreviations, see Fig. 1.

Proteins differentially represented in the membrane fraction (Potential direct targets of RhoII)

A total of 41 proteins were affected by the *rhoII* mutation in membrane fractions (Table 1) comprising 40.2% of the total affected proteins in MIG1. Fourteen proteins detected as differential are predicted to have at least 1 TMS and three (without TMS) contain signal peptides that likely target them to the cell surface. The remaining proteins affected in the mutant may be associated to the membrane via interactions with other molecules (i.e. lipids, other proteins) and likely represent indirect targets of RhoII. For instance, uracil DNA glycosylase (accumulated 28.7 fold in membranes of exponentially growing cells) which participates in DNA base excision repair has been reported to be associated to the mitochondrial inner membrane through electrostatic interactions³⁶. Nonetheless, it cannot be discarded that some of these proteins may represent contamination from cytoplasmic origin.

1
2
3 Rho are not processive proteases, therefore, their activity may expose or release a portion of the
4 substrate, leading to changes in size, stability and/or location of the cleaved substrate protein.
5
6 Consequently, integral membrane proteins which evidenced variations in amount (accumulated or
7 decreased) will be discussed as putative targets of RhoII. Proteins with altered electrophoretic
8 migration were also taken into consideration (see next section).
9
10
11
12

13
14 Rho cleave target proteins within or near a TMS. Most characterized Rho substrates are single
15 TMS, type I or III proteins (N-terminus facing the extracellular side with or without a signal
16 peptide, respectively), even though exceptions are known such as *Drosophila melanogaster* Star,
17 which is a type II membrane protein (N-terminus facing the intracellular side of the cell)³⁷. It has
18 been proposed that Rho so far characterized from eukaryotes and bacteria recognize two elements in
19 their natural substrates: part of the TMS and a sequence motif³⁸.
20
21
22
23
24
25
26

27
28 Among the integral membrane proteins which evidenced differential levels between MIG1 and the
29 parental strain, only HVO_2150, encoding a halocyanin homolog which accumulated in MIG1 (7.8
30 fold), fulfills the requirements to be a Rho substrate following the criteria proposed by Strisovsky
31 et al.³⁹ (1TMS, type I or III membrane protein and presence of a 7 amino acid conserved motif).
32
33
34
35
36

37
38 Halocyanins are blue copper membrane associated proteins from halophilic archaea that are thought
39 to act as mobile electron carriers in a way similar to plant plastocyanins⁴⁰. Cu²⁺ regulates
40 plastocyanin abundance in cyanobacteria, algae and higher plants, which show a decrease in
41 plastocyanin abundance when subjected to Cu deficiency. However, the mechanism of down-
42 regulation varies, as some organisms show Cu-induced variations in mRNA levels while others
43 evidence Cu deficiency-induced plastocyanin proteolysis⁴¹⁻⁴³. In *Chlamydomonas*, Cu deficiency-
44 induced proteolysis was proposed to be carried out by the IMP RSEP1 based in gene expression
45 studies⁴². Halocyanin was purified from the haloalkaliphilic archaeon *Natronobacterium pharaonis*
46 and the gene was cloned and sequenced⁴⁴, however regulation of halocyanin levels by Cu has not
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 been confirmed in haloarchaea.
4
5

6 In *Arabidopsis sp.*, mutations in two P-type ATPases (PAA1 and PAA2) that deliver Cu to the
7 chloroplast and thylakoid lumen caused 80–90% reduction of plastocyanin abundance^{45, 46}. While
8 plant chloroplast P-type ATPases import Cu to the chloroplast lumen, prokaryotic P-type ATPases
9 contribute to Cu detoxification by copper efflux out of the cell⁴⁷. In the thermophilic and
10 thermoacidophilic archaea *Archaeoglobus fulgidus* and *Sulfolobus solfataricus* CopA ATPases
11 are active Cu export pumps that confer Cu and Ag tolerance^{48, 49}. In this study, a P-type ATPase
12 (*copA*, HVO_1751) showed a 2.1 fold reduction in abundance in the MIG1 strain (Table 1). A
13 functional link between halocyanin and the CopA homolog in *H. volcanii* could be hypothesized.
14 Whether RhoII regulates directly halocyanin concentration through processing or indirectly (i.e.
15 by processing of a P-type ATPase and therefore regulating Cu and halocyanin levels) remains to
16 be determined and is focus of future research.
17
18
19
20
21
22
23
24
25
26
27
28

29
30
31 *Escherichia coli* rhomboid GlpG was shown to cleave heterologous multi pass membrane
32 proteins¹³ and the rhomboid protease 1 (RHBDD1) was reported to cleave the 6 TMS TSAP6
33 protein in human cell lines, allowing for a probably TSAP6-dependent regulation of exosome
34 trafficking⁵⁰. Therefore, in addition to halocyanin, the polytopic integral membrane proteins
35 which evidenced variations in their abundance in MIG1 were also considered as putative RhoII
36 substrates.
37
38
39
40
41
42
43

44 Changes in the concentration of proteins involved in regulation of gene expression may explain
45 differences in amount of the protein products of the genes they control. Two polytopic membrane
46 sensor histidine kinases (HVO_0570 and HVO_B0273) were less abundant in MIG1 exp phase
47 samples (Table 1) with negative folds of 1.5 and 13.9, respectively and may represent RhoII
48 direct targets. It has been reported that the Rho PARL mediates cleavage of the histidine kinase
49 PINK1 in the mitochondrial inner membrane, which regulates the activity of respiratory complex
50
51
52
53
54
55
56
57
58
59
60

1
2
3 I and the clearance of damaged mitochondria by mitophagy⁵¹.
4
5

6 Four integral membrane metalloproteases showed lower protein levels in MIG1 than in the parental
7 strain (Table 1): peptidase T homolog (HVO_0477, 13.1 fold), Abi/CAAX protease (HVO_1997,
8 10.7 fold), HtpX- like protease (HVO_A0045, 1.6 fold) and PrsW protease homolog (HVO_0408,
9 28.2 fold), suggesting that metal-dependent proteolysis at the membrane level is, by some means,
10 regulated by RhoII. This pattern may reflect a specific effect of RhoII on individual proteases but
11 may also be a consequence of altered metal availability caused by variations in metal transporters
12 (Table 1). It is likely that some proteins observed as stabilized in the mutant (Table 1) may
13 accumulate as a consequence of the level reduction of these proteases. To the best of our
14 knowledge, these enzymes have not been characterized in *Achaea*, and the fact that in this study
15 they were detected at the protein level suggests that they are functional enzymes in *H. volcanii*.
16
17
18
19
20
21
22
23
24
25
26
27

28 In *Bacillus subtilis* the PrsW peptidase inactivates the anti-sigma factor RsiW by
29 performing the first of several proteolytic cleavages, resulting in the transcription of genes
30 controlled by the extra cytoplasmic function sigma factor σ^W ⁵². To validate the results
31 obtained in the proteomic approach, the *in vivo* stability of the *H. volcanii* PrsW protease
32 was compared in the parental and MIG1 strains. Expression of the *prsW* gene, cloned with
33 an N-terminal His tag downstream the *H. volcanii* tryptophanase promoter (*ptnA*), was
34 induced in both strains by addition of tryptophan (1mM) to the growth medium. Cells were
35 treated with puromycin to block protein translation and the stability of PrsW over time was
36 assessed by Western blotting. While PrsW electrophoretic mobility was not affected in the
37 mutant, the levels of PrsW were significantly reduced after 16 h (Fig. 4), compared to the
38 parental strain. This result correlates with the reduced PrsW levels observed in the mutant
39 in the proteomic approach and suggest that PrsW is affected by RhoII, likely through a
40 proteolytic signaling cascade at the membrane in *H. volcanii* cells. Alternatively, as the
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

recombinant protein bears an N-terminal 6His tag, processing by RhoII of a small peptide from the C-terminus (not detected as a shift in migration in the SDS-PAGE) may increase the stability of PrsW in the parental strain. As the biological function of PrsW in archaea is not known, at present it is not possible to correlate/evaluate to what extent the deficiency in PrsW affects the physiology of the *H. volcanii* MIG1 mutant.




Figure 4. Stability of PrsW homolog in MIG1 strain. *H. volcanii* H26 and MIG1 cells harboring plasmid pTA963 containing the *prsW* gene with a 6x histidine tag under the *ptnA* promoter were grown to exp phase in presence of tryptophan 1 mM. At that point puromycin (100 $\mu\text{g}/\text{ml}$) was added and samples were taken at the indicated times and subjected to SDS-PAGE followed by western blotting with anti his tag antibody (A). Position of Molecular weight markers (kDa) is

1
2
3 indicated at the left. The arrow points the band corresponding to PrsW and the asterisk a
4 degradation product. The band corresponding to PrsW was quantified with ImageJ (B). The plotted
5 data correspond to two independent experiments. Error bars, S.D.
6
7
8
9

10 11 12 13 **Proteins differentially represented in the secretome of *H. volcanii***

14
15 The growth medium of *H. volcanii* MIG1 (exp plus st phases) showed 27 proteins with variations in
16 their level, compared to the wt (Table 1). Out of these, 8 are predicted secretory proteins, 4 are
17 expected to be membrane associated via at least 1 TMS, and the remaining are likely soluble
18 proteins. The presence of proteins without a predicted signal peptide in culture medium supernatant
19 is not uncommon; actually, evidence from proteomics studies suggests that there are significant
20 differences between predicted and experimental archaeal secretomes^{53, 54}. These differences may be
21 due to unconventional secretion such as vesicle transport, or a consequence of cell lysis. In any
22 case, soluble proteins (with or without a signal peptide) which displayed differences in their levels
23 in MIG1 supernatants may represent an indirect effect of the *rhoII* deletion.
24
25
26
27
28
29
30
31
32
33

34
35 Four integral membrane proteins were underrepresented in MIG1 culture media supernatants (Table
36 1). Out of these, SecD (HVO_1976), flagellin 1 (HVO_1210) and signal peptidase I (HVO_2603)
37 were detected in both strains while the conserved hypothetical protein (HVO_1153)] was not
38 detected in the supernatants of MIG1 cultures. The absence of peptides from HVO_1153 in the
39 culture medium supernatant of MIG1 could be a consequence of the lack of processing by RhoII.
40
41
42
43
44
45
46

47 SecD and Sec1a are proteins involved in the Sec protein translocation pathway^{55, 56}. Even though
48 they were both affected in the mutant, the proteomic data did not provide additional evidence
49 suggesting that this pathway is altered in MIG1.
50
51
52
53

54
55 HVO_1153 is a conserved hypothetical protein with 1 TMS. BLAST searches indicated that this
56
57
58
59
60

1
2
3 protein has homology with archaeal flagellin domain-containing proteins and also with bacterial
4 adhesins, and thus the lack of processing of HVO_1153 in MIG1 may contribute to the observed
5 defects in motility and cell adhesion (see below) previously evidenced in the mutant. Amino acid
6 sequence analysis with Pattrinprot (https://npsa-prabi.ibcp.fr/cgi-bin/pattern_pattrinprot.pl) evidenced
7 the presence of a Rho consensus motif within the predicted TMS of this hypothetical protein. Thus
8 it can be hypothesized that the presence of peptides from HVO_1153 in the culture medium of the
9 wt strain may be due to RhoII proteolytic activity.

10
11
12
13
14
15
16
17
18
19 Flagellin 1 is the major component of the *H. volcanii* flagellum and deletion of the corresponding
20 gene, *flagA1*, renders non motile cells³³. Flagella are released from the cell surface through
21 centrifugation, therefore, the reduced level of flagellin peptides in the culture supernatant of MIG1
22 suggests that flagella are either less abundant and/or not properly assembled which may contribute
23 with the reduced motility reported previously for this mutant¹⁷. Maturation of flagellin involves N-
24 glycosylation and cleavage by PibD signal peptidase III prior to flagellum assembly³³. Whether the
25 role of RhoII regulating the level of flagellin in the culture medium is direct or indirect (i. e
26 regulating flagellin glycosylation and flagellum assembly) needs to be experimentally determined.
27 This also applies to Sec11a and HVO_1153, as both have predicted glycosylation sites, according to
28 NetNGlyc server⁵⁷.

44 **Identification of proteins with differential migration in the *H. volcanii***

47 ***rhoII* KO mutant**

49
50 Rho cleave their substrates at specific sites releasing peptides which often play a regulatory role⁵⁸.
51 Therefore, to detect variations in the electrophoretic migration of individual proteins that may arise
52 from differential processing in the mutant vs. wt strain, a simplified PROTOMAP assay was
53
54
55
56
57
58
59
60

1
2
3 performed²³. The results are shown in Table 2.
4
5

6 In this assay RhoII (HVO_0727) was detected in membranes from the parental strain while it was
7 not evidenced in MIG1, confirming the absence of the RhoII protease in this mutant. This finding
8 agreed with the observation that a higher percentage of total proteins with predicted TMS were
9 detected in this experiment compared to the previous analysis (21 vs. 12%, respectively) (Tables S1
10 and S3).
11
12
13
14
15
16
17

18 A total of 99 proteins showed differences in amount (fold ≥ 1.5) in at least one of the gel sections
19 (Table 2). Twenty-one (21) integral membrane proteins displayed differences comparing the mutant
20 vs. wt strain and may represent a direct consequence of RhoII proteolytic activity
21 (absence/presence). In addition, 78 predicted soluble proteins showed differential migration in the
22 PROTOMAP experiment. These probably represent proteins that were secondarily/indirectly
23 affected by the *rhoII* deletion, as a consequence of absence of processing by other protease/s or by
24 another altered posttranslational modification process.
25
26
27
28
29
30
31
32
33

34 Several proteins involved in solute transport across cell membranes showed altered electrophoretic
35 migration including 16 ABC transporter components for lipoproteins, peptides, Zn^{2+} , cobalamin,
36 phosphate and sugars (Table 2). In addition, three integral membrane metal transporters (probable
37 Zn/Cd transport protein-HVO_2602 and P-type ATPase-HVO_1751, Mg transport protein CorA-
38 HVO_1675) displayed altered migration with negative folds of 8.4 (section 3), 1.7 (section 3) and
39 5.2 (section 2), respectively. P-type ATPase also showed reduced levels in quantitative proteomics
40 experiment (see above). It is possible that differences in metal ion availability would in turn cause
41 to some extent variations in other metalloprotein levels (i.e. metalloproteases, see above). Metal
42 ions have a key role in the physiology of cells, as they act as the redox centers for metalloproteins
43 such as cytochromes, blue copper proteins, and iron-sulfur proteins which play a vital role in
44 electron transport. On the other hand, any disturbance in metal ion homeostasis could produce toxic
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 effects on cell viability. Among *Archaea*, thermophiles, hyperthermophiles and methanogens
4 utilize P-type ATPases and ABC transporters for metal transport and balancing. However, metal
5 homeostasis in haloarchaea has not been extensively studied⁵⁹. Independently if the observed
6 difference in concentration of metal transporter proteins is a direct or indirect effect of RhoII
7 absence, these data, together with the semi quantitative proteomics results (Table 1) suggest that
8 metal homeostasis is affected in MIG1. However, under standard laboratory conditions, this effect
9 was not reflected in cell proliferation and growth.
10
11
12
13
14
15
16
17
18
19
20
21

22 **Proteins with differential migration related to the phenotypes of the MIG1** 23 **strain**

24 ***Soluble proteins***

25 DNA gyrase is an essential bacterial enzyme that catalyzes the ATP-dependent negative super-
26 coiling of double-stranded closed circular DNA. Gyrase is a topoisomerase involved in the control
27 of topological transitions of DNA⁶⁰. The active form of this enzyme is a tetramer of two A and two
28 B subunits⁶¹. Novobiocin is a DNA gyrase inhibitor that binds to the active enzyme⁶². In this work
29 we observed that the DNA gyrase subunit A was reduced by about 2 fold in section 2 of the
30 PROTOMAP gel (Table 2). The abnormal migration of subunit A in the *rhoII* mutant suggests that
31 migration of this polypeptide is altered (i.e. affected by post translational modification) generating a
32 DNA gyrase more susceptible to novobiocin inhibition that could account for the enhanced
33 sensitivity to novobiocin displayed by the MIG1 strain¹⁷.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 In *H. volcanii*, the *rhoII* gene is part of an operon with the sequence encoding EndV (HVO_0726),
54 an endonuclease involved in DNA repair in *E. coli*⁶³. We have shown that *endV* mRNA is expressed
55 in *H. volcanii* H26 and MIG1 at similar levels¹⁷. MIG1 cells irradiated with UV light in liquid
56
57
58
59
60

1
2
3 minimal medium cultures, showed a delay in resuming growth when compared to the wt (Figure
4 S2), evidencing that the mutant has difficulty in coping with UV stress. In this study, the soluble
5 protein RadA (HVO_0104) which is involved in DNA repair after UV irradiation⁶⁴ was reduced by
6 1.9 fold in section 2 in the mutant (Table 2). Even though the functional link between RhoII, EndV
7 and RadA remains to be addressed, our data suggest that RhoII may participate in the regulation of
8 DNA repair processes in *H. volcanii*. A link between UV irradiation and Rho has been reported for
9 RHBDL2 which showed upregulation upon UV treatment in human keratinocytes⁶⁵.

22 ***Integral membrane proteins***

23
24
25 *H. volcanii* is surrounded by a hexagonally packed surface S-layer, formed by the SLG. This
26 glycoprotein contains 7 predicted N-glycosylation sites. Out of these, N83, N13, N274 and N279
27 are decorated with a pentasaccharide which also occurs in *H. volcanii* flagellins and pilins⁶⁵⁻⁶⁷. In
28 addition, position N498 is decorated with a tetrasaccharide synthesized only in low salt growth
29 conditions⁶⁸. Our previous work showed that N-732 in the SLG harbors a novel sulfoquinovose
30 containing oligosaccharide (GlcNAc-GlcNAc(Hex)₂-(SQ-Hex)₆) which is truncated in the MIG1
31 strain¹⁷.

32
33
34 Several enzymes potentially involved in protein glycosylation were detected as differential by
35 shotgun proteomics (Table 1) and in the PROTOMAP experiment (Table 2). A list along with their
36 proposed/known roles is provided in Table S2. Agl15 (HVO_2050), a flippase which participates in
37 the synthesis of low salt tetrasaccharide at position N498 of SLG⁶⁷, was the only affected
38 glycosylase predicted to be an integral membrane protein (14 TMS) with an 8.1 fold reduction at
39 section 3 of the PROTOMAP. Whether this protein is involved in the N-732 glycosylation of SLG
40 is not known. However, if this were the case, Agl15 processing/activation by RhoII may account at
41 least in part for the differences in SLG glycosylation observed in the mutant¹⁷. Alternatively, it is
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 also possible that RhoII cleaves an unknown substrate that regulates the level of one or more of
4 these glycosylation enzyme/s (Table S2) leading to an abnormal glycosylation pattern of the SLG.
5
6

7
8 SLG is initially synthesized with a C-terminal transmembrane domain and following proteolytic
9 cleavage upstream of the TMS, the truncated protein is transferred to a lipid anchor⁶⁹. The
10 involvement of the ArtA archaeosortase in the proteolytic processing of the C-terminus of this
11 glycoprotein was reported by Abdul Halim et al⁷⁰. In this study SLG was enriched 1.5 fold in
12 section 2 and 1.7 fold in section 3 of the PROTOMAP (Table 2). The differential electrophoretic
13 migration of this protein may be due to the previously observed truncation of the oligosaccharide
14 bound to N732 in the mutant strain. However, it cannot be ruled out that RhoII may also contribute
15 to proteolytic processing of SLG.
16
17
18
19
20
21
22
23
24
25

26 The preflagellin peptidase PibD was reduced 7.7 fold in section 3 and evidenced a 2 fold reduction
27 in level when calculating the total fold (Table 2), indicating that its electrophoretic migration was
28 altered but also that the protein concentration was lower in MIG1 cells. PibD is known to process
29 preflagellins and prepilins prior to flagella and pili assembly in archaea⁷¹. It was shown that a
30 knock-out mutation of *pibD* abolished swimming motility and adhesion to glass surfaces in *H.*
31 *volcanii*³³. Therefore, decreased amounts of PibD in MIG1 cells may explain the reduced motility of
32 this strain in soft agar plates¹⁷. In order to determine whether cell adhesion was also affected in the
33 MIG1 mutant, we examined the ability of these cells to adhere to glass slides. As observed in Fig. 5,
34 the Rho mutant showed reduced adhesion to glass plates compared to the parental strain. In
35 addition, PilA1 (HVO_0972), a substrate of PibD⁷¹, was accumulated in section 4 of the gel. It
36 could be proposed that RhoII is involved in proteolytic maturation/processing of PibD affecting
37 pilin (and probably flagellin, see Table 1) processing and/or assembly and therefore *H. volcanii*
38 motility.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 5. Cells from *H. volcanii* MIG1 strain show reduced adhesion to glass surfaces.

Adhesion to glass coverslips was tested using a previously described protocol³³. Coverslips were placed in 50 ml plastic tubes, containing 6 ml of MGM 18% medium cultures of *H. volcanii* H26 and MIG1 at an OD₆₀₀ of 0.3. After 3 or 6 h incubation at 42 °C, cells were fixed with 2% acetic acid, stained with 0.1% crystal violet and observed by light microscopy. Light micrographs of the coverslips were taken at 100x magnification.

Analysis of semi-tryptic peptides in integral membrane proteins

Detection of proteins with different levels and/or electrophoretic mobility variations was performed against a trypsin digestion database. In order to identify peptides that may result from RhoII proteolytic activity, a search of semi-tryptic peptides was also performed. This analysis focused on proteins with semi-tryptic peptide/s (found exclusively in the wt) inside or near a TMS and which evidenced a Rho cleavage motif (Table S4). A total of 10 candidates were identified in all the experiments performed, including PibD, hypothetical proteins and small solute transporters. However, it has to be noted that it is not known whether archaeal rhomboid proteases recognize the same cleavage motif as their bacterial and eukaryotic counterparts. Experimental validation and cleavage site identification of *H. volcanii* RhoII predicted substrates will contribute to clarify this

1
2
3 particular issue.
4
5
6
7
8

9 **Concluding remarks**

10
11
12
13 This study shows for the first time, the effect of a rhomboid protease gene deletion on the proteome
14 of an archaeon. Previous studies have examined the influence of Rho deficiency on specific
15 subproteomes, such as *A. thaliana* chloroplasts⁷² and the secretome of *Trichomonas vaginalis*⁷³,
16 however, our study focused on a more comprehensive analysis of the relevance of Rho at the whole
17 proteome level.
18
19
20
21
22

23
24 Elimination of *rhoII* from *H. volcanii* chromosome had a global impact on the proteome, affecting
25 soluble, membrane-associated as well as secreted proteins with diverse biological functions,
26 suggesting a role for this protease in the regulation of processes such as solute transport, Cu
27 homeostasis, signal transduction, protein glycosylation, cell adhesion and motility. Interestingly,
28 some of these proteins are associated with the differential phenotypes observed in the *rhoII* mutant.
29 A schematic representation of the proteins showing differences in their levels and/or electrophoretic
30 migration and their involvement in the generation of the mutant phenotypes is provided in Fig. 6.
31 Ongoing work is focused on the validation of selected candidate substrates of RhoII which will
32 contribute to get insight on the biological function of this family of IMPs in the *Archaea* domain.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 6. Schematic representation of the effect of deleting *rhoII* in *H. volcanii* physiology.

Elimination of *rhoII* from the chromosome causes several phenotypes in *H. volcanii* cells. Proteins that showed variations in level and/or electrophoretic mobility in this work and that are potentially involved in the generation of these phenotypes are listed in the corresponding box. Blue font denotes proteins that showed differential migration in the PROTOMAP assay. The arrows indicate proteins with augmented (green) or decreased concentration (red) in the mutant. The proposed role of RhoII in the processing of the conserved hypothetical protein HVO_1153 and PibD (affecting cell adhesion and motility) is detailed on the right.

Supporting information

Figure S1. Cartoon representation of SDS gels used for comparative quantitative proteomics or for the simplified PROTOMAP assay sample preparation.

Figure S2. Effect of UV irradiation on *H. volcanii* MIG1 growth.

Table S1. Proteins detected in (membrane, cytoplasm and cell culture supernatants) of *H. volcanii* H26 and MIG1 at exp and st phase of growth

Table S2. Proteins involved in N-glycosylation with affected level and/or migration in MIG1. CE: cytoplasm of exp phase. ME: membrane fraction of exp phase.

Table S3. Proteins detected in membranes of exponentially growing H26 and MIG1 strains in the PROTOMAP assay.

Table S4. Semitryptic peptides with a rhomboid cleavage consensus detected in membrane proteins.

The authors declare no competing financial interest.

AUTHOR INFORMATION

Corresponding authors

* María I. Giménez. Email: migimen@mdp.edu.ar. Tel: 54-223-4753030 int 12.

* Ansgar Poestch. Email: ansgar.poetsch@rub.de.

Author contributions

¥ Both authors contributed equally to this work.

ACKNOWLEDGEMENTS

This work was supported by CONICET Grant (PIP-1106), UNMDP Grant EXA632/13. Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) Grant PICT1477. MIC is an ANPCyT Doctoral Fellow and MC is a CONICET Post-Doctoral Fellow.

ABBREVIATIONS

Rho, rhomboid proteases; IMP, intramembrane protease; TMS, transmembrane segment; exp, exponential; st, stationary; nLC-ESI-MS/MS, Nano-scale liquid chromatographic electrospray ionization tandem mass spectrometry; SLG, S-layer glycoprotein.

REFERENCES

1. Lopez-Otin, C.; Overall, C. M., Protease degradomics: a new challenge for proteomics. *Nature reviews. Molecular cell biology* **2002**, *3* (7), 509-19.
2. Gimenez, M. I.; Cerletti, M.; De Castro, R. E., Archaeal membrane-associated proteases: insights on *Haloferax volcanii* and other haloarchaea. *Frontiers in microbiology* **2015**, *6*, 39.
3. Verhelst, S. H., Intramembrane proteases as drug targets. *The FEBS journal* **2017**, **284**(10), **1489-1502**.

- 1
2
3 4. Freeman, M., The rhomboid-like superfamily: molecular mechanisms and biological roles.
4
5 *Annual review of cell and developmental biology* **2014**, *30*, 235-54.
6
7
- 8 5. Lee, J. R.; Urban, S.; Garvey, C. F.; Freeman, M., Regulated intracellular ligand transport
9 and proteolysis control EGF signal activation in *Drosophila*. *Cell* **2001**, *107* (2), 161-71.
10
11
- 12 6. Urban, S.; Lee, J. R.; Freeman, M., A family of Rhomboid intramembrane proteases
13 activates all *Drosophila* membrane-tethered EGF ligands. *The EMBO journal* **2002**, *21* (16), 4277-
14 86.
15
16
17
18
19
- 20 7. Spinazzi, M.; De Strooper, B., PARL: The mitochondrial rhomboid protease. *Seminars in cell*
21 & *developmental biology* **2016**, *60*, 19-28.
22
23
24
- 25 8. Sibley, L. D., The roles of intramembrane proteases in protozoan parasites. *Biochimica et*
26 *biophysica acta* **2013**, *1828* (12), 2908-15.
27
28
29
- 30 9. Dhingra, S.; Kowlaski, C. H.; Thammahong, A.; Beattie, S. R.; Bultman, K. M.; Cramer, R. A.,
31 RbdB, a Rhomboid Protease Critical for SREBP Activation and Virulence in *Aspergillus fumigatus*.
32 *mSphere* **2016**, *1* (2), e00035-16.
33
34
35
36
37
- 38 10. Stevenson, L. G.; Strisovsky, K.; Clemmer, K. M.; Bhatt, S.; Freeman, M.; Rather, P. N.,
39 Rhomboid protease AarA mediates quorum-sensing in *Providencia stuartii* by activating TatA of
40 the twin-arginine translocase. *Proceedings of the National Academy of Sciences of the United*
41 *States of America* **2007**, *104* (3), 1003-8.
42
43
44
45
46
47
- 48 11. Akiyama, Y.; Maegawa, S., Sequence features of substrates required for cleavage by GlpG,
49 an *Escherichia coli* rhomboid protease. *Molecular microbiology* **2007**, *64* (4), 1028-37.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 12. Maegawa, S.; Ito, K.; Akiyama, Y., Proteolytic action of GlpG, a rhomboid protease in the
4 Escherichia coli cytoplasmic membrane. *Biochemistry* **2005**, *44* (41), 13543-52.
5
6
7
8
9 13. Erez, E.; Bibi, E., Cleavage of a multispinning membrane protein by an intramembrane
10 serine protease. *Biochemistry* **2009**, *48* (51), 12314-22.
11
12
13
14 14. Mesak, L. R.; Mesak, F. M.; Dahl, M. K., Expression of a novel gene, gluP, is essential for
15 normal Bacillus subtilis cell division and contributes to glucose export. *BMC microbiology* **2004**, *4*,
16
17
18 13.
19
20
21 15. Kateete, D. P.; Katabazi, F. A.; Okeng, A.; Okee, M.; Musinguzi, C.; Asiimwe, B. B.; Kyobe, S.;
22 Asiimwe, J.; Boom, W. H.; Joloba, M. L., Rhomboids of Mycobacteria: characterization using an
23 aarA mutant of Providencia stuartii and gene deletion in Mycobacterium smegmatis. *PloS one*
24
25
26 **2012**, *7* (9), e45741.
27
28
29
30
31 16. Kinch, L. N.; Grishin, N. V., Bioinformatics perspective on rhomboid intramembrane
32 protease evolution and function. *Biochimica et biophysica acta* **2013**, *1828* (12), 2937-43.
33
34
35
36 17. Parente, J.; Casabuono, A.; Ferrari, M. C.; Paggi, R. A.; De Castro, R. E.; Couto, A. S.;
37 Giménez, M. I., A rhomboid protease gene deletion affects a novel oligosaccharide N-linked to the
38 S-layer glycoprotein of Haloferax volcanii. *The Journal of biological chemistry* **2014**, *289* (16),
39
40
41 11304-17.
42
43
44
45
46 18. Cerletti, M.; Paggi, R. A.; Guevara, C. R.; Poetsch, A.; De Castro, R. E., Global role of the
47 membrane protease LonB in Archaea: Potential protease targets revealed by quantitative
48 proteome analysis of a lonB mutant in Haloferax volcanii. *Journal of proteomics* **2015**, *121*, 1-14.
49
50
51
52
53 19. Nieuwlandt, D. T., Palmer, J. R., Armbruster, D. T., Kuo, Y. P., Oda, W. and Daniels, C. J., In
54 *Archaea: A Laboratory Manual*, Robb, F. T., Place, A. R., Sowers, K. R., Schreier, H. J., DasSarma, S.
55
56
57
58
59
60

1
2
3 and Fleischmann, E. M., Ed. Cold Spring Harbor Laboratory Press: Cold Spring Harbor, NY, 1995; pp
4
5 161-162.

6
7
8 20. Dyall-Smith, M., The halohandbook. *Protocols for haloarchaeal genetics*. Edited by: Dyall-
9
10 Smith M. University of Melbourne **2009**, 144 pp.
11
12 <http://www.haloarchaea.com/resources/halohandbook/>

13
14
15 21. Sambrook, J.; Russell, D. W., Molecular cloning: a laboratory manual. *Cold Spring Harbor*
16
17 *Laboratory Press, Cold Spring Harbor, NY 2001*.

18
19
20 22. Laemmli, U. K., Cleavage of structural proteins during the assembly of the head of
21
22 bacteriophage T4. *Nature* **1970**, 227 (5259), 680-5.

23
24
25 23. Dix, M. M.; Simon, G. M.; Cravatt, B. F., Global identification of caspase substrates using
26
27 PROTOMAP (protein topography and migration analysis platform). *Methods in molecular biology*
28
29 **2014**, 1133, 61-70.

30
31
32 24. Candiano, G.; Bruschi, M.; Musante, L.; Santucci, L.; Ghiggeri, G. M.; Carnemolla, B.;
33
34 Orecchia, P.; Zardi, L.; Righetti, P. G., Blue silver: a very sensitive colloidal Coomassie G-250
35
36 staining for proteome analysis. *Electrophoresis* **2004**, 25 (9), 1327-33.

37
38
39 25. Schluesener, D.; Fischer, F.; Kruip, J.; Rogner, M.; Poetsch, A., Mapping the membrane
40
41 proteome of *Corynebacterium glutamicum*. *Proteomics* **2005**, 5 (5), 1317-30.

42
43
44 26. Eng, J. K.; McCormack, A. L.; Yates, J. R., An approach to correlate tandem mass spectral
45
46 data of peptides with amino acid sequences in a protein database. *Journal of the American Society*
47
48 *for Mass Spectrometry* **1994**, 5 (11), 976-989.
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 27. Dorfer, V.; Pichler, P.; Stranzl, T.; Stadlmann, J.; Taus, T.; Winkler, S.; Mechtler, K., MS
4 Amanda, a universal identification algorithm optimized for high accuracy tandem mass spectra.
5
6 Amanda, a universal identification algorithm optimized for high accuracy tandem mass spectra.
7
8 *Journal of proteome research* **2014**, *13* (8), 3679-84.
9
- 10 28. Pfeiffer, F.; Broicher, A.; Gillich, T.; Klee, K.; Mejia, J.; Rampp, M.; Oesterhelt, D., Genome
11 information management and integrated data analysis with HaloLex. *Archives of microbiology*
12
13 **2008**, *190* (3), 281-99.
14
15
- 16 29. Kall, L.; Storey, J. D.; MacCoss, M. J.; Noble, W. S., Posterior error probabilities and false
17 discovery rates: two sides of the same coin. *Journal of proteome research* **2008**, *7* (1), 40-4.
18
19
20
21
- 22 30. Liu, H.; Sadygov, R. G.; Yates, J. R., 3rd, A model for random sampling and estimation of
23 relative protein abundance in shotgun proteomics. *Analytical chemistry* **2004**, *76* (14), 4193-201.
24
25
26
27
- 28 31. Haussmann, U.; Qi, S. W.; Wolters, D.; Rogner, M.; Liu, S. J.; Poetsch, A., Physiological
29 adaptation of *Corynebacterium glutamicum* to benzoate as alternative carbon source - a
30 membrane proteome-centric view. *Proteomics* **2009**, *9* (14), 3635-51.
31
32
33
34
35
- 36 32. Allers, T.; Barak, S.; Liddell, S.; Wardell, K.; Mevarech, M., Improved strains and plasmid
37 vectors for conditional overexpression of His-tagged proteins in *Haloferax volcanii*. *Applied and*
38
39 *environmental microbiology* **2010**, *76* (6), 1759-69.
40
41
42
43
- 44 33. Tripepi, M.; Imam, S.; Pohlschroder, M., *Haloferax volcanii* flagella are required for motility
45 but are not involved in PibD-dependent surface adhesion. *Journal of bacteriology* **2010**, *192* (12),
46
47 3093-102.
48
49
50
- 51 34. Fischer, F.; Poetsch, A., Protein cleavage strategies for an improved analysis of the
52 membrane proteome. *Proteome science* **2006**, *4*, 2.
53
54
55
56
57
58
59
60

- 1
2
3 35. Heberle, H.; Meirelles, G. V.; da Silva, F. R.; Telles, G. P.; Minghim, R., InteractiVenn: a web-
4 based tool for the analysis of sets through Venn diagrams. *BMC Bioinformatics* 2015, 16:169.
5
6
7
8 36. Stuart, J. A.; Mayard, S.; Hashiguchi, K.; Souza-Pinto, N. C.; Bohr, V. A., Localization of
9 mitochondrial DNA base excision repair to an inner membrane-associated particulate fraction.
10
11
12 *Nucleic acids research* 2005, 33 (12), 3722-32.
13
14
15
16 37. Tsruya, R.; Wojtalla, A.; Carmon, S.; Yogev, S.; Reich, A.; Bibi, E.; Merdes, G.; Schejter, E.;
17 Shilo, B. Z., Rhomboid cleaves Star to regulate the levels of secreted Spitz. *The EMBO journal* 2007,
18
19
20 26 (5), 1211-20.
21
22
23 38. Strisovsky, K., Structural and mechanistic principles of intramembrane proteolysis--lessons
24 from rhomboids. *The FEBS journal* 2013, 280 (7), 1579-603.
25
26
27
28 39. Strisovsky, K.; Sharpe, H. J.; Freeman, M., Sequence-specific intramembrane proteolysis:
29 identification of a recognition motif in rhomboid substrates. *Molecular cell* 2009, 36 (6), 1048-59.
30
31
32
33 40. Scharf, B.; Engelhard, M., Halocyanin, an archaebacterial blue copper protein (type I) from
34 Natronobacterium pharaonis. *Biochemistry* 1993, 32 (47), 12894-900.
35
36
37
38 41. Zhang, L.; McSpadden, B.; Pakrasi, H. B.; Whitmarsh, J., Copper-mediated regulation of
39 cytochrome c553 and plastocyanin in the cyanobacterium Synechocystis 6803. *The Journal of*
40
41
42 *biological chemistry* 1992, 267 (27), 19054-9.
43
44
45
46 42. Kropat, J.; Gallaher, S. D.; Urzica, E. I.; Nakamoto, S. S.; Strenkert, D.; Tottey, S.; Mason, A.
47
48
49 Z.; Merchant, S. S., Copper economy in Chlamydomonas: prioritized allocation and reallocation of
50
51
52 copper to respiration vs. photosynthesis. *Proceedings of the National Academy of Sciences of the*
53
54 *United States of America* 2015, 112 (9), 2644-51.
55
56
57
58
59
60

- 1
2
3 43. Abdel-Ghany, S. E.; Pilon, M., MicroRNA-mediated systemic down-regulation of copper
4 protein expression in response to low copper availability in Arabidopsis. *The Journal of biological*
5 *chemistry* **2008**, *283* (23), 15932-45.
6
7
8
9
10 44. Mattar, S.; Scharf, B.; Kent, S. B.; Rodewald, K.; Oesterhelt, D.; Engelhard, M., The primary
11 structure of halocyanin, an archaeal blue copper protein, predicts a lipid anchor for membrane
12 fixation. *The Journal of biological chemistry* **1994**, *269* (21), 14939-45.
13
14
15
16
17 45. Shikanai, T.; Muller-Moule, P.; Munekage, Y.; Niyogi, K. K.; Pilon, M., PAA1, a P-type
18 ATPase of Arabidopsis, functions in copper transport in chloroplasts. *The Plant cell* **2003**, *15* (6),
19 1333-46.
20
21
22
23
24 46. Abdel-Ghany, S. E.; Muller-Moule, P.; Niyogi, K. K.; Pilon, M.; Shikanai, T., Two P-type
25 ATPases are required for copper delivery in Arabidopsis thaliana chloroplasts. *The Plant cell* **2005**,
26 17 (4), 1233-51.
27
28
29
30
31
32 47. Migocka, M., Copper-transporting ATPases: The evolutionarily conserved machineries for
33 balancing copper in living systems. *IUBMB life* **2015**, *67* (10), 737-45.
34
35
36
37
38 48. Arguello, J. M., Identification of ion-selectivity determinants in heavy-metal transport P1B-
39 type ATPases. *The Journal of membrane biology* **2003**, *195* (2), 93-108.
40
41
42
43 49. Völlmecke, C.; Drees, S. L.; Reimann, J.; Albers, S.-V.; Lübben, M., The ATPases CopA and
44 CopB both contribute to copper resistance of the thermoacidophilic archaeon *Sulfolobus*
45 *solfataricus*. *Microbiology* **2012**, *158* (6), 1622-1633.
46
47
48
49
50 50. Wan, C.; Fu, J.; Wang, Y.; Miao, S.; Song, W.; Wang, L., Exosome-related multi-pass
51 transmembrane protein TSAP6 is a target of rhomboid protease RHBDD1-induced proteolysis. *PLoS*
52 *one* **2012**, *7* (5), e37452.
53
54
55
56
57
58
59
60

- 1
2
3 51. Wai, T.; Saita, S.; Nolte, H.; Muller, S.; Konig, T.; Richter-Dennerlein, R.; Sprenger, H. G.;
4
5 Madrenas, J.; Muhlmeister, M.; Brandt, U.; Kruger, M.; Langer, T., The membrane scaffold SLP2
6
7 anchors a proteolytic hub in mitochondria containing PARL and the i-AAA protease YME1L. *EMBO*
8
9 *reports* **2016**, *17* (12), 1844-1856.
10
11
12
13 52. Heinrich, J.; Hein, K.; Wiegert, T., Two proteolytic modules are involved in regulated
14
15 intramembrane proteolysis of *Bacillus subtilis* RsiW. *Molecular microbiology* **2009**, *74* (6), 1412-26.
16
17
18 53. Ellen, A. F.; Zolghadr, B.; Driessen, A. M.; Albers, S.-V., Shaping the archaeal cell
19
20 envelope. *Archaea* **2010**, 608243, 13 pp.
21
22 54. Feng, J.; Wang, J.; Zhang, Y.; Du, X.; Xu, Z.; Wu, Y.; Tang, W.; Li, M.; Tang, B.; Tang, X. F.,
23
24 Proteomic analysis of the secretome of haloarchaeon *Natrinema* sp. J7-2. *Journal of proteome*
25
26 *research* **2014**, *13* (3), 1248-58.
27
28
29 55. Hand, N. J.; Klein, R.; Laskewitz, A.; Pohlschroder, M., Archaeal and bacterial SecD and SecF
30
31 homologs exhibit striking structural and functional conservation. *Journal of bacteriology* **2006**, *188*
32
33 (4), 1251-9.
34
35
36
37 56. Fink-Lavi, E.; Eichler, J., Identification of residues essential for the catalytic activity of
38
39 Sec11b, one of the two type I signal peptidases of *Haloferax volcanii*. *FEMS microbiology letters*
40
41 **2008**, *278* (2), 257-60.
42
43
44
45
46 57. Gupta, R.; Brunak, S., Prediction of glycosylation across the human proteome and the
47
48 correlation to protein function. *Pacific Symposium on Biocomputing. Pacific Symposium on*
49
50 *Biocomputing* **2002**, 310-22.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 58. Urban, S.; Dickey, S. W., The rhomboid protease family: a decade of progress on function
4 and mechanism. *Genome biology* **2011**, *12* (10), 231.
5
6
7
8 59. Srivastava, P.; Kowshik, M., Mechanisms of metal resistance and homeostasis in
9 haloarchaea. *Archaea* **2013**, 732864.
10
11
12
13 60. Reece, R. J.; Maxwell, A., DNA gyrase: structure and function. *Critical reviews in*
14 *biochemistry and molecular biology* **1991**, *26* (3-4), 335-75.
15
16
17
18 61. Bush, N. G.; Evans-Roberts, K.; Maxwell, A., DNA Topoisomerases. *EcoSal Plus* 2015;
19 doi:10.1128/ecosalplus.ESP-0010-2014.
20
21
22
23
24 62. Heide, L., New aminocoumarin antibiotics as gyrase inhibitors. *International journal of*
25 *medical microbiology : IJMM* **2014**, *304* (1), 31-6.
26
27
28
29 63. Kuraoka, I., Diversity of Endonuclease V: From DNA Repair to RNA Editing. *Biomolecules*
30 **2015**, *5* (4), 2194-206.
31
32
33
34 64. Woods, W. G.; Dyall-Smith, M. L., Construction and analysis of a recombination-deficient
35 (radA) mutant of *Haloferax volcanii*. *Molecular microbiology* **1997**, *23* (4), 791-7.
36
37
38
39 65. Cheng, T. L.; Wu, Y. T.; Lin, H. Y.; Hsu, F. C.; Liu, S. K.; Chang, B. I.; Chen, W. S.; Lai, C. H.; Shi,
40 G. Y.; Wu, H. L., Functions of rhomboid family protease RHBDL2 and thrombomodulin in wound
41 healing. *The Journal of investigative dermatology* **2011**, *131* (12), 2486-94.
42
43
44
45 66. Kandiba, L.; Lin, C. W.; Aebi, M.; Eichler, J.; Guerardel, Y., Structural characterization of the
46 N-linked pentasaccharide decorating glycoproteins of the halophilic archaeon *Haloferax volcanii*.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 67. Tripepi, M.; You, J.; Temel, S.; Onder, O.; Brisson, D.; Pohlschroder, M., N-glycosylation of
4 Haloferax volcanii flagellins requires known Agl proteins and is essential for biosynthesis of stable
5 flagella. *Journal of bacteriology* **2012**, *194* (18), 4876-87.
6
7
8
9
10 68. Esquivel, R. N.; Schulze, S.; Xu, R.; Hippler, M.; Pohlschroder, M., Identification of Haloferax
11 volcanii Pilin N-Glycans with Diverse Roles in Pilus Biosynthesis, Adhesion, and Microcolony
12 Formation. *The Journal of biological chemistry* **2016**, *291* (20), 10602-14.
13
14
15
16
17
18 69. Kaminski, L.; Guan, Z.; Yurist-Doutsch, S.; Eichler, J., Two distinct N-glycosylation pathways
19 process the Haloferax volcanii S-layer glycoprotein upon changes in environmental salinity. *mBio*
20
21
22 **2013**, *4* (6), e00716-13.
23
24
25
26 70. Kandiba, L.; Guan, Z.; Eichler, J., Lipid modification gives rise to two distinct Haloferax
27 volcanii S-layer glycoprotein populations. *Biochimica et biophysica acta* **2013**, *1828* (3), 938-43.
28
29
30
31 71. Abdul Halim, M. F.; Pfeiffer, F.; Zou, J.; Frisch, A.; Haft, D.; Wu, S.; Tolic, N.; Brewer, H.;
32 Payne, S. H.; Pasa-Tolic, L.; Pohlschroder, M., Haloferax volcanii archaeosortase is required for
33 motility, mating, and C-terminal processing of the S-layer glycoprotein. *Molecular microbiology*
34
35
36
37 **2013**, *88* (6), 1164-75.
38
39
40
41 72. Esquivel, R. N.; Xu, R.; Pohlschroder, M., Novel archaeal adhesion pilins with a conserved N
42 terminus. *Journal of bacteriology* **2013**, *195* (17), 3808-18.
43
44
45
46 73. Knopf, R. R.; Feder, A.; Mayer, K.; Lin, A.; Rozenberg, M.; Schaller, A.; Adam, Z., Rhomboid
47 proteins in the chloroplast envelope affect the level of allene oxide synthase in Arabidopsis
48 thaliana. *The Plant journal : for cell and molecular biology* **2012**, *72* (4), 559-71.
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 74. Riestra, A. M.; Gandhi, S.; Sweredoski, M. J.; Moradian, A.; Hess, S.; Urban, S.; Johnson, P.
4
5 J., A *Trichomonas vaginalis* Rhomboid Protease and Its Substrate Modulate Parasite Attachment
6
7 and Cytolysis of Host Cells. *PLoS pathogens* **2015**, *11* (12), e1005294.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Differential subproteome of the *H volcanii* MIG1 strain. Regulation factors for three biological replicates from H26 and MIG1 strains were submitted to a Student's t-test. Proteins with p -values ≤ 0.05 were considered significant. Negative fold values indicate that the protein is less abundant in the mutant than in the wt whereas a positive fold value indicates accumulation of the protein in MIG1. For membrane fractions two independent experiments (three biological replicates each) were performed and data corresponding to both assays was included in the table. TMS: number of transmembrane segments. Sec: general translocation signal peptide. Tat: Twin arginine translocation signal peptide. FC and SC: functional class and functional sub-class, according to the Halolex database, respectively (for abbreviations see Fig. 1).

ID	Gene name	Description	TMS	Signal peptide	Lipobox	FC	SC	p value	Fold
Supernatant Exponential									
HVO_0874	-	Beta-lactamase domain protein/mRNA 3-end processing factor homolog				MIS	GEN	0.000136	17.8
HVO_2375	<i>pstS1</i>	ABC-type transport system periplasmic substrate-binding protein (probable substrate phosphate)		Tat	Yes	TP_CP	TP	0.000013	11.4
HVO_2563	<i>rpl4</i>	50S ribosomal protein L4				GIP	TL	0.047086	1.7
Membrane Exponential									

1	HVO_1038	<i>udg4</i>	Uracil-DNA glycosylase superfamily protein			MIS	GEN	0.000666	28.7
2	HVO_1344	-	SBDS family protein			MIS	GEN	0.001658	18.1
3	HVO_2150	<i>hcpG</i>	Halocyanin	1	Tat	MET	EM	0.000016	7.8
4	HVO_2464	<i>sucD</i>	Succinate--CoA ligase (ADP-forming) alpha subunit			MET	CIM	0.041494	7.1
5	HVO_1925	-	Probable GTP-binding protein			MIS	GEN	0.042352	2.6
6	HVO_2757	<i>rplI</i>	50S ribosomal protein L1			GIP	TL	0.008677	1.9
7	HVO_0316	<i>atpA</i>	A-type ATP synthase subunit A			MET	EM	0.049824	1.6
8	HVO_0570	-	Histidine kinase	3	Sec	ENV	SIG	0.043555	-1.5
9	HVO_0684	<i>gatB</i> , <i>aatB</i>	Aspartyl/glutamyl-tRNA(Asn/Gln) amidotransferase subunit B			GIP	TL	0.020533	-1.5
10	HVO_A0045	<i>htpX3</i>	HtpX-like protease	4		MIS	MIS	0.024904	-1.6
11	HVO_A0208	<i>cas5</i>	CRISPR-associated endoribonuclease Cas5, Hmari subtype			MIS	MIS	0.005354	-1.6
12	HVO_0194	<i>orc9</i>	Orc1-type DNA replication protein			MIS	MIS	0.026537	-1.8
13	HVO_0237	-	UPF0761 family protein			MIS	GEN	0.029733	-1.8
14	HVO_1357	-	receiver/bat box HTH-10 family transcription regulator			ENV	REG	0.04533	-1.8
15	HVO_B0052	-	PQQ repeat protein			MIS	GEN	0.043425	-1.9
16	HVO_0581	<i>ftsZ2</i>	cell division protein FtsZ, type II			MIS	MIS	0.024353	-2
17	HVO_1554	<i>traB</i>	TraB family protein	8		MIS	GEN	0.026824	-2
18	HVO_0874	-	beta-lactamase domain protein/mRNA 3-end processing factor homolog			MIS	GEN	0.045002	-2.6

1	HVO_0598	-	conserved hypothetical protein			UNASS	CHY	0.000252	-7.1	
2	HVO_0762	-	NUDIX family hydrolase			MIS	GEN	0.000355	-8.4	
3	HVO_1659	<i>mscS2</i>	mechanosensitive channel protein MscS	5	Sec	TP_CP	TP	0.000240	-9	
4	HVO_1997	-	Abi/CAAX family protein	7		MIS	GEN	0.026824	-10.7	
5	HVO_0255	-	Spermine/spermidine synthase family protein	7		MIS	GEN	0.000011	-10.8	
6	HVO_0477	<i>pepB3</i>	Aminopeptidase (homolog to leucyl aminopeptidase / aminopeptidase T)			MIS	GEN	0.000214	-13.1	
7	HVO_B0273	-	Sensor box histidine kinase	7		ENV	SIG	0.047159	-13.9	
8	HVO_1382	<i>nac</i>	Nascent polypeptide-associated complex protein			GIP	TL	0.003979	-14.9	
9	HVO_2060	<i>agl8</i>	NUDIX family hydrolase (low-salt glycan biosynthesis protein Agl8)			MIS	MIS	0.034081	-19.3	
10	HVO_0941	-	Conserved hypothetical protein	6		UNASS	CHY	0.000550	-24.1	
11	HVO_2318	<i>uppS2</i>	Tritrans, polycis-undecaprenyl-diphosphate synthase			MET	LIP	0.043265	-25.4	
12	HVO_1676	<i>tfb2</i>	Transcription initiation factor TFB			GIP	TC	0.006282	-27.1	
13	HVO_0408	-	PrsW family protein	9		MIS	GEN	0.000165	-28.2	
14	Cytoplasm Exponential									
15	HVO_2548	<i>rpl6</i>	50S ribosomal protein L6			GIP	TL	0.000076	9.5	
16	HVO_2772	-	rhodanese domain protein / probable metallo-beta-lactamase family hydrolase			MIS	GEN	0.000001	9.2	
17	HVO_C0074	<i>dppDF16</i>	ABC-type transport system ATP-binding protein (probable			TP_CP	TP	0.000001	9.2	

		substrate dipeptide/oligopeptide)					
HVO_2600	<i>map</i>	methionyl aminopeptidase	MIS	MIS	0.000035	8.5	
HVO_1758	<i>trxB5</i>	oxidoreductase (homolog to thioredoxin-disulfide reductase)	MIS	GEN	0.000004	7.8	
HVO_2709	-	GNAT family acetyltransferase	MIS	GEN	0.000104	6.9	
HVO_0818	<i>thrC2</i>	threonine synthase	MET	AA	0.049049	4.9	
HVO_1047	<i>qor2</i>	NADPH:quinone reductase	MIS	MIS	0.047097	4.8	
HVO_1604	<i>aspC1</i>	pyridoxal phosphate-dependent aminotransferase (probable aspartate aminotransferase)	MIS	GEN	0.044358	2	
HVO_A0406	-	conserved hypothetical protein	UNASS	CHY	0.010564	2	
HVO_2270	<i>rmeM</i>	type I restriction-modification system DNA-methyltransferase RmeM	MIS	MIS	0.008658	1.9	
HVO_0353	<i>rps12</i>	30S ribosomal protein S12	GIP	TL	0.024019	1.8	
HVO_1478	<i>tfb5</i>	transcription initiation factor TFB	GIP	TC	0.049709	1.7	
HVO_1681	-	DNA N-glycosylase	GIP	RRR	0.038567	1.6	
HVO_2345	-	flavin-dependent pyridine nucleotide oxidoreductase	MIS	GEN	0.041979	1.6	
HVO_0710	<i>pabB</i>	aminodeoxychorismate synthase component II	MET	COM	0.049446	1.5	
HVO_1249	-	conserved hypothetical protein	MIS	GEN	0.025429	1.5	
HVO_1488	<i>gnaD</i>	D-gluconate dehydratase	MET	CHM	0.002489	-1.5	
HVO_A0379	<i>hyuA2</i>	N-methylhydantoinase A(ATP-hydrolyzing)	MIS	MIS	0.013225	-1.5	
HVO_2790	<i>apbC1</i>	Fe-S cluster carrier protein ApbC	MIS	MIS	0.012758	-1.6	

HVO_2059	<i>agl12</i> , <i>rjbB</i>	dTDP-glucose 4,6-dehydratase (Agl12)				MET	CHM	0.036699	-1.7
HVO_2947	<i>pheT</i>	phenylalanine--tRNA ligase beta subunit				GIP	TL	0.0382	-1.8
HVO_0214	-	L-lactate dehydrogenase		Sec		MET	CIM	0.004094	-1.9
HVO_2695	<i>tsgA3</i>	ABC-type transport system periplasmic substrate-binding protein (probable substrate sugar)		Tat	Yes	TP_CP	TP	0.027275	-2
HVO_1401	<i>tsgA11</i>	ABC-type transport system periplasmic substrate-binding protein (probable substrate sugar)		Tat	Yes	TP_CP	TP	0.003844	-2.1
HVO_2740	<i>ndk</i>	nucleoside-diphosphate kinase				MET	NUM	0.019173	-2.2
HVO_1311	-	probable oxidoreductase (short-chain dehydrogenase family)				MIS	GEN	0.032821	-6
HVO_2684	-	UspA domain protein				MIS	GEN	0.000151	-9.1
HVO_2648	<i>serA2</i>	phosphoglycerate dehydrogenase				MIS	GEN	0.000035	-9.4
Supernatant Stationary									
HVO_0062	<i>dppA1</i>	ABC-type transport system periplasmic substrate-binding protein (probable substrate dipeptide/oligopeptide)		Tat	Yes	TP_CP	TP	0.031260	1.7
HVO_1976	<i>secD</i>	Protein-export membrane protein SecD	5	Sec		TP_CP	SEC	0.047004	-5.3
HVO_1210	<i>flgA1</i>	Flagellin 1	1			TP_CP	MOT	0.000004	-5.3
HVO_1477	-	Endonuclease domain protein		Sec	Yes	MIS	GEN	0.000004	-5.3
HVO_2607	-	PQQ repeat protein		Tat	Yes	MIS	GEN	0.000004	-5.3
HVO_0058	<i>dppF1</i>	ABC-type transport system ATP-binding protein (probable				TP_CP	TP	0.000086	-6.0

		substrate dipeptide/oligopeptide)							
HVO_A0181	-	Conserved hypothetical protein		Tat	Yes	UNASS	CHY	0.000369	-6.0
HVO_A0377	<i>hyuE</i>	Hydantoin racemase				MIS	MIS	0.000608	-6.5
HVO_2384	-	CBS domain protein				MIS	GEN	0.000010	-6.7
HVO_2603	<i>sec11a</i>	Signal peptidase I	1			TP_CP	SEC	0.000010	-6.7
HVO_1945	-	Conserved hypothetical protein		Tat	Yes	UNASS	CHY	0.000144	-6.7
HVO_2413	<i>tefla2</i>	Translation elongation factor aEF-1 alpha subunit				GIP	TL	0.035008	-7.0
HVO_2286	-	Conserved hypothetical protein		Tat	Yes	UNASS	CHY	0.027060	-7.4
HVO_1153	-	Conserved hypothetical protein	1			UNASS	CHY	0.000002	-7.5
HVO_0054	<i>glyS</i>	Glycine--tRNA ligase				GIP	TL	0.000377	-7.8
HVO_B0057	<i>cbiH2,</i> <i>cobJ2</i>	Precorrin-3B C17-methyltransferase				MET	COM	0.000124	-8.7
HVO_2815	<i>hbd1</i>	3-hydroxyacyl-CoA dehydrogenase / enoyl-CoA hydratase				MET	LIP	0.000706	-8.9
HVO_0348	<i>rpoB1</i>	DNA-directed RNA polymerase subunit B'				GIP	TC	0.000865	-9.0
HVO_2471	<i>pccB2</i>	Propionyl-CoA carboxylase carboxyltransferase component				MET	LIP	0.033685	-9.1
HVO_2554	<i>rpl14</i>	50S ribosomal protein L14				GIP	TL	0.000011	-9.5
HVO_B0184	<i>dppA14</i>	ABC-type transport system periplasmic substrate-binding protein (probable substrate dipeptide/oligopeptide)		Tat	Yes	TP_CP	TP	0.000573	-10.1
HVO_B0061	<i>cbiL</i>	Precorrin-2 C20-methyltransferase				MET	COM	0.000175	-10.2
HVO_2749	<i>rpl21e</i>	50S ribosomal protein L21e				GIP	TL	0.000001	-10.3

HVO_2452	<i>nrdJ</i>	Ribonucleoside-diphosphate reductase, adenosylcobalamin-dependent							MET	NUM	0.036975	-11.9
Membrane Stationary												
HVO_0887	<i>korB</i>	oxoglutarate--ferredoxin oxidoreductase beta subunit							MET	CIM	0.006496	1.7
HVO_2614	<i>udp2</i>	uridine phosphorylase							MET	NUM	0.023086	1.5
HVO_2989	<i>pmm4</i>	phosphohexomutase (phosphoglucomutase / phosphomannomutase)							MET	CHM	0.040726	-1.5
HVO_2069	-	RND superfamily permease	11						TP_CP	TP	0.002656	-1.6
HVO_C0054	-	hypothetical protein				Tat			UNASS	HY	0.033063	-1.6
HVO_0307	-	conserved hypothetical protein				Sec			UNASS	CHY	0.022016	-2
HVO_1751	<i>copA</i>	P-type transport ATPase (probable substrate copper/metal cation)	8						TP_CP	TP	0.011109	-2.1
HVO_0462	<i>cydA</i>	cytochrome bd ubiquinol oxidase subunit I	9						MET	EM	0.000721	-8.8
HVO_B0228	<i>tsgA9</i>	ABC-type transport system periplasmic substrate-binding protein (probable substrate sugar)				Tat	Yes		TP_CP	TP	0.000374	-8.8
HVO_1870	-	probable metalloprotease	7						MIS	GEN	0.000041	-9.9
Cytoplasm Stationary												
HVO_1804	<i>pncB</i>	nicotinate phosphoribosyltransferase							MET	COM	0.000018	8.2
HVO_2639	-	DUF1611 family protein							MIS	GEN	0.0001	6.7
HVO_2860	-	DUF181 family protein							MIS	GEN	0.0001	6.7

HVO_0087	<i>hemB</i>	porphobilinogen synthase	MET	COM	0.0376	-1.5
HVO_1575	<i>rocF</i>	Arginase	MET	AA	0.0356	-1.5
HVO_2225	-	AstE family protein	MIS	GEN	0.0232	-1.8

Table 2. Proteins with differential migration in SDS-PAGE in the MIG1 mutant. Membrane protein samples (triplicates) from exponentially growing MIG1 and wt cells were subjected to SDS-PAGE allowed to run for 2 cm and separated in four sections (top to bottom: S1-S4) which were analyzed separately by RPLC-MS/MS, as indicated in Table 1. Regulation factors were submitted to a Student's t-test and proteins with p -values ≤ 0.05 and fold ≥ 1.5 were considered significant. In grey, integral membrane proteins. Ns: not significant.

ID	Gene name	Protein name	Signal		Lipobox	FC	SC	Fold				Total Fold
			TMS	peptide				S1	S2	S3	S4	
HVO_1240	-	Conserved hypothetical protein	0	Tat	Yes	UNASS	CHY	8.8		1.8		ns
HVO_2545	<i>rpl18</i>	50S ribosomal protein L18	0			GIP	TL	8.1				ns
HVO_2302	-	Ubiquinone biosynthesis protein UbiB	3			MET	COM	1.8				ns
HVO_1847	<i>pepF</i>	Oligoendopeptidase PepF	0			MIS	MIS	1.7				ns
HVO_2495	-	DUF106 family protein	4			MIS	GEN	-1.5				ns
HVO_0541	<i>citB2. acnA</i>	aconitate hydratase	0			MET	CIM	-1.5				ns
		ABC-type transport system periplasmic substrate-binding										
HVO_C0075	<i>dppA16</i>	protein (probable substrate dipeptide/oligopeptide)	0	Tat	Yes	TP_CP	TP	-1.5				ns
		2-succinyl-5-enolpyruvyl-6-hydroxy-3-cyclohexene-1-										
HVO_1469	<i>menD</i>	carboxylate synthase	0			MET	COM	-1.6				-1.9
HVO_1242	-	Conserved hypothetical protein	0	Tat	Yes	UNASS	CHY	-7.8				ns
HVO_0893	<i>mmcA1</i>	Methylmalonyl-CoA mutase subunit A	0			MET	CIM	-10.3				ns
		ABC-type transport system ATP-binding protein (probable										
HVO_C0074	<i>dppDF16</i>	substrate dipeptide/oligopeptide)				TP_CP	TP	-14.4				ns
						ACS Paragon Plus Environment						
HVO_0519	<i>rpa2</i>	Replication protein A	0			GIP	RRR		7.1			ns

HVO_2784	<i>rps13</i>	30S ribosomal protein S13	0			GIP	TL	5.2		ns
ABC-type transport system periplasmic substrate-binding										
HVO_0447	<i>phnD1</i>	protein (probable substrate phosphate/phosphonate)	0	Tat	Yes	TP_CP	TP	2.1		2.0
HVO_2945	<i>valS</i>	Valine--tRNA ligase	0			GIP	TL	1.8		ns
HVO_0220	<i>mcm</i>	ATP-dependent DNA helicase MCM	0			GIP	RRR	1.8		ns
HVO_1305	<i>porA</i>	Pyruvate--ferredoxin oxidoreductase alpha subunit	0			MET	CIM	1.8		ns
HVO_0133	<i>ths1. cct1</i>	Thermosome subunit 1	0			GIP	CHP	1.7		ns
HVO_1590	<i>dnaK</i>	DnaK-type molecular chaperone Hsp70	0			GIP	CHP	1.6		ns
HVO_2072	<i>csg</i>	S-layer glycoprotein	1	Sec		TP_CP	CE	1.5	1.7	ns
HVO_0717	<i>ftsZ1</i>	Cell division protein FtsZ. type I	0			TP_CP	CP	1.5		ns
Probable branched-chain amino acid dehydrogenase E1										
HVO_2209	<i>oadhA4</i>	component alpha subunit	0			MET	AA	-1.5		-1.6
HVO_A0379	<i>hyuA2</i>	N-methylhydantoinase A(ATP-hydrolyzing)	0			MIS	MIS	-1.5		ns
HVO_1304	<i>porB</i>	Pyruvate--ferredoxin oxidoreductase beta subunit	0			MET	CIM	-1.5		ns
Glyceraldehyde-3-phosphate dehydrogenase (NAD)										
HVO_0481	<i>gap2</i>	(phosphorylating)	0			MET	CIM	-1.5		ns
HVO_A0271	<i>glpC2</i>	Glycerol-3-phosphate dehydrogenase subunit C	0			MET	CIM	-1.5		-1.7
HVO_3007	<i>mdh</i>	Malate dehydrogenase	0			MET	CIM	-1.6		ns
Probable oxidoreductase (aldo-keto reductase family										
HVO_1009	-	protein)	0			MIS	GEN	-1.6	1.6	ns
HVO_2617	-	TrkA-C domain protein	3			MIS	GEN	-1.7		ns
HVO_2643	<i>qor1</i>	NADPH:quinone reductase	0			MIS	MIS	-1.7		ns
HVO_0243	<i>ltaE</i>	Threonine aldolase	0			MET	AA	-1.8		-1.9

1	HVO_0832	-	Alpha/beta hydrolase fold protein	0	Sec	MIS	GEN	-1.8	-1.8	-1.6
2										
3										
4	HVO_0945	<i>cbaA</i>	ba3-type terminal oxidase subunit I	14		MET	EM	-1.8		ns
5										
6	HVO_0104	<i>radA</i>	DNA repair and recombination protein RadA	0		GIP	RRR	-1.9		ns
7										
8	HVO_1573	<i>gyrA</i>	DNA gyrase subunit A	0		GIP	RRR	-2.2		ns
9										
10			S-adenosyl-L-methionine:L-histidine 3-amino-3-							
11	HVO_1631	<i>dph2</i>	carboxypropyltransferase (GTP binding)	0		MIS	MIS	-5.2	5.8	ns
12										
13	HVO_1675	<i>corA</i>	Magnesium transport protein CorA	2		TP_CP	TP	-5.2		-1.7
14										
15			ABC-type transport system ATP-binding protein (probable							
16	HVO_B0078	<i>dppF11</i>	substrate dipeptide/oligopeptide)	0		TP_CP	TP	-5.4	13.6	ns
17										
18	HVO_1443	-	ABC-type transport system ATP-binding protein	0		TP_CP	TP	-5.5		-2.0
19										
20	HVO_2191	<i>purU</i>	Formyltetrahydrofolate deformylase	0		MET	NUM	-5.8		ns
21										
22	HVO_1006	-	ABC-type transport system ATP-binding protein	0		TP_CP	TP	-6.3		ns
23										
24	HVO_B0011	<i>soxA1</i>	Sarcosine oxidase	0		MET	AA	-6.3		-6.7
25										
26	HVO_2168	<i>moxR3</i>	AAA-type ATPase (MoxR subfamily)	0		MIS	MIS	-6.4	13.8	ns
27										
28	HVO_0951	-	Hypothetical protein	1		UNASS	HY	-6.5		-6.9
29										
30	HVO_2993	<i>pibD</i>	Prepilin peptidase PibD	6		MIS	MIS	-7.7		-2.0
31										
32	HVO_2348	<i>mptA. folE2</i>	GTP cyclohydrolase MptA	0		MET	COM	-12.4	9.0	-1.7
33										
34	HVO_1576	-	NAD-dependent epimerase/dehydratase	0		MIS	GEN		17.4	2.3
35										
36			ABC-type transport system periplasmic substrate-binding							
37	HVO_2397	<i>znuA1</i>	protein (probable substrate zinc)	0	Tat	Yes	TP_CP	TP	14.6	ns
38										
39			ABC-type transport system periplasmic substrate-binding							
40	HVO_A0611	<i>znuA2</i>	protein (probable substrate zinc)	0	Tat	Yes	TP_CP	TP	9.6	ns
41										
42	HVO_2374	<i>phoU2</i>	Transcription regulator (homolog to phosphate uptake	0		ENV	REG		9.0	ns
43										
44										
45										
46										
47										

regulator)									
HVO_0739	-	Conserved hypothetical protein	4			UNASS	CHY	9.0	ns
HVO_2943	<i>pyrD</i>	Dihydroorotate dehydrogenase (quinone)	0			MET	NUM	7.9	ns
HVO_2607	-	PQQ repeat protein	0	Tat	Yes	MIS	GEN	7.3	ns
HVO_2384	-	CBS domain protein	0			MIS	GEN	7.1	ns
HVO_2068	<i>ftsZ8</i>	FtsZ family protein, type III	0			MIS	GEN	6.1	ns
HVO_A0619	-	Conserved hypothetical protein	0	Tat	Yes	UNASS	CHY	5.8	ns
HVO_2756	<i>rpl10</i>	50S ribosomal protein L10	0			GIP	TL	5.1	ns
HVO_1506	<i>ihvC</i>	Ketol-acid reductoisomerase	0			MET	AA	2.7	ns
HVO_1570	<i>top6A</i>	DNA topoisomerase VI subunit	0			GIP	RRR	2.4	ns
HVO_0059	<i>dppD1</i>	ABC-type transport system ATP-binding protein (probable substrate dipeptide/oligopeptide)	0			TP_CP	TP	1.9	ns
HVO_2046	<i>agl7</i>	Sulfatase domain protein	0			MIS	MIS	1.8	ns
HVO_2047	-	Conserved hypothetical protein	0			UNASS	CHY	-1.5	ns
HVO_2703	<i>panB2</i>	3-methyl-2-oxobutanoate hydroxymethyltransferase	0			MET	COM	-1.6	-1.5
HVO_2544	<i>rps5</i>	30S ribosomal protein S5	0			GIP	TL	-1.6	ns
HVO_0790	<i>jba2</i>	2-amino-3,7-dideoxy-D-threo-hept-6-ulosonate synthase	0			MET	AA	-1.6	ns
HVO_1751	<i>copA</i>	P-type transport ATPase (probable substrate copper/metal cation)	8			TP_CP	TP	-1.7	-1.7
HVO_0841	<i>petD</i>	Cytochrome bc1 complex cytochrome b/c subunit	5			MET	EM	-1.7	ns
HVO_0773	-	Probable S-adenosylmethionine-dependent methyltransferase	0			MIS	GEN	-1.7	ns
HVO_2398	<i>znuC1</i>	ABC-type transport system ATP-binding protein (probable	0			TP_CP	TP	-1.7	-1.6

		substrate zinc)							
HVO_1478	<i>tfb5</i>	Transcription initiation factor TFB	0			GIP	TC	-1.8	ns
HVO_B0134	-	ABC-type transport system ATP-binding protein	0			TP_CP	TP	-1.8	-1.7
HVO_2554	<i>rpl14</i>	50S ribosomal protein L14	0			GIP	TL	-1.8	ns
HVO_2981	<i>upp</i>	Uracil phosphoribosyltransferase	0			MET	NUM	-1.9	ns
HVO_1241	<i>prcC</i>	SCO1/SenC/PrrC family protein	0	Tat	Yes	MIS	GEN	-2.0	-1.5
HVO_0536	<i>dpsA1</i>	Ferritin	0			MIS	MIS	-2.2	ns
HVO_0727	-	Rhomboid protease II	6			MIS	GEN	-6.6	-6.7
HVO_1274	-	Conserved hypothetical protein	7			UNASS	CHY	-6.6	-5.5
HVO_1532	-	DUF368 family protein	8			MIS	GEN	-6.6	ns
HVO_1914	<i>acaB3</i>	Acetyl-CoA C-acyltransferase	0			MET	LIP	-7.4	-5.1
HVO_1936	<i>cofE</i>	F420-0:gamma-glutamyl ligase	0			MET	COM	-7.4	-6.1
HVO_2318	<i>ups2</i>	Tritrans.polycis-undecaprenyl-diphosphate synthase	0			MET	LIP	-7.5	ns
HVO_2055	<i>agl15</i>	Probable low-salt glycan biosynthesis flippase Agl15	14			TP_CP	TP	-8.1	ns
HVO_2602	-	Transport protein (probable substrate zinc/cadmium)	4			TP_CP	TP	-8.4	-6.9
HVO_2952	<i>endA</i>	tRNA-splicing endonuclease	0			GIP	RMT	-8.4	-6.9
HVO_0972	<i>pilA1</i>	Pilin (PilA1)	1			MIS	MIS	11.7	2.0
HVO_2773	<i>rps2</i>	30S ribosomal protein S2 1	0			GIP	TL	10.6	ns
HVO_0531	<i>tsgB1</i>	ABC-type transport system permease protein (probable substrate sugar)	0			TP_CP	TP	8.6	ns
HVO_1110	<i>btuF</i>	ABC-type transport system periplasmic substrate-binding protein (probable substrate cobalamin)	0	Sec		TP_CP	TP	7.1	ns
HVO_2163	-	ABC-type transport system ATP-binding protein (homolog	0			TP_CP	TP	7.1	ns

to LolDCE lipoprotein release factor)									
HVO_2547	<i>rpl32e</i>	50S ribosomal protein L32e	0		GIP	TL	1.9	ns	
HVO_2564	<i>rpl3</i>	50S ribosomal protein L3	0		GIP	TL	1.7	ns	
HVO_0316	<i>atpA</i>	A-type ATP synthase subunit A	0		MET	EM	1.6	ns	
HVO_0060	<i>dppC1</i>	ABC-type transport system permease protein (probable substrate dipeptide/oligopeptide)	8		TP_CP	TP	-1.5	ns	
HVO_B0185	<i>dppB14</i>	ABC-type transport system permease protein (probable substrate dipeptide/oligopeptide)	6		TP_CP	TP	-1.6	ns	
HVO_0500	-	Conserved hypothetical protein	1		UNASS	CHY	-1.6	ns	
HVO_1198	-	UspA domain protein	0		MIS	GEN	-1.8	ns	
HVO_0146	<i>psd</i>	Phosphatidylserine decarboxylase	1		MET	LIP	-2.5	ns	
HVO_1563	-	Conserved hypothetical protein	0	Sec	UNASS	CHY	-2.6	-1.6	
HVO_1377	-	UPF0145 family protein	0		MIS	GEN	-13.4	-6.9	

For table of contents only

