2018-06

Bridging the divide: Social-ecological coherence in Marine Protected Area network design

Rees, Sian

<http://hdl.handle.net/10026.1/11996>

10.1002/aqc.2885

Aquatic Conservation: Marine and Freshwater Ecosystems

Wiley

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1 **Bridging the divide: A framework for social-ecological coherence in Marine Protected Area**
2 **network design**

3 Siân E. Rees^{1*}, Nicola Foster¹, Olivia Langmead^{1, 2}, Charly Griffiths², Steve Fletcher^{1,3}, Simon
4 J. Pittman^{1,4}, David E. Johnson^{5,6}, Martin Attrill¹.

5 ¹ Marine Institute, 3rd Floor Marine Building, Plymouth University, Plymouth PL4 8AA, UK.

6 ² Marine Biological Association of the United Kingdom.

7 ³ UNEP World Conservation Monitoring Centre.

8 ⁴ National Oceanic and Atmospheric Administration.

9 ⁵ Seascope Consultants Ltd.

10 ⁶ Global Ocean Biodiversity Initiative Programme Coordinator.

11 *Corresponding author e-mail: sian.rees@plymouth.ac.uk

12

Abstract

1. Marine Protected Areas (MPAs) and networks of MPAs are being implemented globally as a spatial management tool for achieving conservation objectives. There has been considerable progress in reaching the prescribed 10 % protected area target for 2020 outlined in the Convention on Biological Diversity Aichi Target 11 and the United Nations Sustainable Development Goal 14.
2. Application of MPA network design principles (e.g. representativity, ecological connectivity) which underpin ecological coherence are still lacking or insufficient in many regions. Poor ecological coherence hinders the ecological performance of MPA networks leading to dysfunction in the flow of ecosystem services and reduced ecosystem benefits with potentially negative consequences for human wellbeing
3. This paper presents four pivotal focus points for future progress that can bridge a gap between the ecological and the social systems. The aim is to shift the discourse of “ecological coherence” further into the social sphere and hence support the alignment of the process of designating ecologically coherent MPA networks with the ‘triple bottom line’ of economic development, environmental sustainability and social inclusion as described in the SDGs to achieve social-ecological coherence in MPA network design.

Key words: Ocean, Marine Protected Areas; Sustainable Development Goals; ecosystem services; ecological coherence.

1. Introduction

Marine Protected Areas (MPAs) are regarded as an important tool for the maintenance of marine ecosystem functionality, health, and ecosystem integrity through the conservation of significant species, habitats, or entire ecosystems (Sobel & Dahlgren, 2004). There is growing evidence that, when properly designed, resourced and managed, MPAs are an effective spatial management tool for achieving conservation objectives (Edgar et al., 2014; Lester et al., 2009; Sciberras, Jenkins, Kaiser, Hawkins & Pullin, 2013; Sheehan, Stevens, Gall, Cousens & Attrill, 2013; Stewart et al., 2009). Developments in social-ecological systems (SES) research emphasize the crucial interdependencies between the natural and the human system (Berkes, Folke & Colding, 2000; Liu et al., 2007) (Figure 1). From a SES perspective, protecting the habitats and species, which are the subject of conservation management measures within MPAs, supports ecological functions and processes (Pollnac et al., 2010; Potts et al., 2014). In turn, this delivers flows of ecosystem services that support human wellbeing (e.g. food, flood protection, opportunities for recreation) (Arkema et al., 2013; Arkema et al., 2015; McCook et al., 2010; Pollnac et al., 2010; Potts et al., 2014; Rees et al., 2014; Rees, Rodwell, Attrill, Austen & Mangi, 2010; Roberts, Bohnsack, Gell, Hawkins & Goodridge, 2001) (Figure 1). Given the high level of functional and spatial connectivity within marine ecosystems and variable and uncertain distribution of risks from ecosystem disturbances, individual MPAs are not considered to be adequate to safeguard the important ecosystem processes and services they underpin (Jones, Srinivasan & Almany, 2007; Margules & Pressey, 2000) and networks of MPAs are needed (Olsen et al., 2013).

Global MPA policy has developed to address the broader spatial requirements for marine conservation within this SES context. In 2004, Convention on Biological Diversity (CBD) Parties decided that “marine and coastal protected areas are essential tools and approaches in the conservation and sustainable use of marine and coastal biodiversity”, committing to a target of “effective conservation of at least 10% of each of the world’s ecological regions by 2010” (UNEP/CBD/COP/DEC/VII/5). In 2010, CBD Parties adopted the Strategic Plan for Biodiversity 2011-2020, including its 20 Aichi Biodiversity Targets. Among these targets, Parties reaffirmed the importance of area-based conservation measures, including MPAs, as a tool for the conservation of biological diversity and the sustainable use of its components. Specifically, Aichi Target 11 states that, ‘by 2020, at least 17% of terrestrial and inland

water, and 10% of coastal and marine areas, especially areas of particular importance for biodiversity and ecosystem services, are conserved through effectively and equitably managed, ecologically representative and well-connected systems of protected areas and other effective area-based conservation measures, and integrated into the wider landscapes and seascapes' (CBD, 2010). Aichi target 11 notably expands upon the quantitative 10% spatial target for protected areas and, through the qualitative aspects (Ecologically representative; Areas of particular importance for biodiversity and ecosystem services; Management equity and effectiveness; Well-connected and; Integration into wider landscape and seascape (Rees, Foster, Langmead, Pittman & Johnson, 2017). These qualitative aspects of Aichi reflect best practices developed in MPA network design by broadening the scope of conservation planning to be more systematic (Margules & Pressey, 2000) and also enable the wider consideration of the relationship between the protection of biodiversity and human wellbeing. The qualitative aspects of Aichi Target 11 link to the principles of "ecological coherence" (Ardron, 2008; Laffoley, Brockington & Gililand, 2006), whereby a "network of MPAs" (a collection of individual MPAs or reserves operating cooperatively and synergistically, at various spatial scales) is designed to

- Interact and support the wider environment (OSPAR, 2006, , Sects. 5.3, 6);
- Maintain the processes, functions, and structures of the intended protected features across their natural range (Laffoley, Brockington & Gililand, 2006);
- Function synergistically as a whole, such that the individual protected sites benefit from each other to achieve the above two objectives (based on OSPAR, 2006, , Sect. 5.2);
 - Additionally, an ecologically coherent network of MPAs may be designed to be resilient to changing conditions (OSPAR, 2006, , Sect. 5).

In recognition of the importance of these qualitative aspects of Aichi Target 11 the Conference of the Parties to the CBD (Decision XI/24, 2012), invited Parties to undertake major efforts to achieve all elements of Aichi Biodiversity Target 11. Progress towards this goal has been slow. The fourth edition of the Global Biodiversity Outlook (GBO 4) reported that while the quantitative elements of Aichi Target 11 (the 10% protected areas target) are on track to be achieved at the global level by 2020 for marine areas within national

jurisdiction, the other elements relating to ecological representation, coverage of areas important for biodiversity, management effectiveness, governance, and integration of protected areas into wider seascape, still need more attention in order to be achieved (Secretariat of the Convention on Biological Diversity, 2014).

Since GBO 4 there has been further sustained progress towards the 10% spatial target (Lubchenco & Grorud-Colvert, 2015). The MPAtlas reports that 2.98% of the global ocean is within an MPA, with 7.29% in national jurisdictions (Marine Conservation Institute, 2017). Proposed MPAs will add a further 3.15%, with an additional 6.33% of MPAs designated within national jurisdictions (Marine Conservation Institute, 2017). Whilst this increase represents a significant achievement in terms of increasing the spatial protection of marine habitats and species within national jurisdictions, a significant gap remains in areas beyond national jurisdiction and other parameters of ecological coherence, that are the cornerstone of Aichi target 11, are potentially lost (e.g. well-connected, ecologically representative) (Jones & De Santo, 2016). Furthermore, arguments have been put forward that suggest that the 'ease of establishment' of MPAs (in some planning areas) has overridden the need to underpin ecological coherence of marine systems (Devillers et al., 2015). Overall, it can be argued that spatial targets alone, which confer no positive or negative biodiversity outcomes, will potentially undermine efforts to halt the continued degradation and loss of marine habitats and species (D. Spalding et al., 2016; Jones & De Santo, 2016) which, in turn, then further impairs the ability of marine systems to continue to provide ecosystem services that underpin human wellbeing (Worm et al., 2006).

Since GBO 4 a number of assessments at a regional MPA network level have been undertaken to assess whether MPA networks are ecologically coherent in their current configuration. Sub-regional assessments for parameters of ecological coherence of MPA designations in North America (Jessen, Morgan & Bezaury-Creel, 2016); The Caribbean (Pittman et al., 2014); the UK (Lieberknecht, Mullier & Ardron, 2014; Ridgeway, Cornthwaite, Wright & Davies, 2014); Northern Ireland (Barnard, Burdon, Strong & Atkins, 2014); the Celtic Seas (Rees, Foster, Langmead & Griffiths, 2015b); the OSPAR region (Johnson et al., 2014; OSPAR, 2013); Chile (Tognelli, Fernández & Marquet, 2009); The English Channel (Foster et al., 2014); the Baltic (Piekäinen & Korpinen, 2008); and the NE Atlantic (Evans, Peckett & Howell, 2015), demonstrate that whilst progress is being made

towards the 10% protected area target, and some areas are moving towards ecological coherence, none of the existing MPA networks are ecologically representative of the full range of ecosystems nor are they well-connected. Additionally, whilst there has been progress towards the development of methods to assess management effectiveness (Hockings, Stolton, Leverington, Dudley & Courrau, 2006; International Union for the Conservation of Nature, 2016). Regional assessments of management effectiveness reveal a lack of progress in meeting conservation objectives and establishing management plans for networks of protected areas (OSPAR, 2013; Rodríguez-Rodríguez, Rodríguez & Abdul Malak, 2016). There is also lack of a formal process for reporting management effectiveness at a national level scale to support regional assessments (Foster et al., 2014; OSPAR, 2013).

1.1 A new policy context

Further priority is given to the relationship between oceans and human wellbeing in the United Nations (UN) Sustainable Development Goals (SDGs), designed to succeed the Millennium Development Goals (MDGs) as reference goals for the international development community for the period 2015-2030. The SDGs advocate a 'triple bottom line' approach to maintaining human wellbeing; these being economic development, environmental sustainability and social inclusion (Sachs, 2012). SDG Goal 14 to 'conserve and sustainably use the oceans, seas and marine resources for sustainable development' reaffirms the CBD 10% spatial target (SDG 14.5), but places this ecological goal firmly within the economic and social context of SES to aid global development (Figure 1). There are notable alignments between Aichi Target 11 and the SDGs not only in terms of SDG 14, but also in terms of synergies with Goal 1: End poverty in all its forms everywhere and; Goal 13: Take urgent action to combat climate change and its impacts (Diz, Morgera & Wilson, 2017; Rees, Foster, Langmead, Pittman & Johnson, 2017).

The SDGs provide an opportunity to address a more fundamental issue. Namely, global conservation policy in relation to the marine environment has seen major advances in recent years but the building blocks of conservation planning (the broader goals of achieving ecological coherence in conservation planning) are not following the same trajectory. It is noted that the discipline of conservation biology (from where ecological coherence is rooted) has grown from the 'deep green' idea that nature has an intrinsic value and that we

should protect nature ‘for nature’s sake’ (Vucetich, Bruskotter & Nelson, 2015). This idea, is shared, to some extent, by the general populous but is limited by a broad spectrum of factors such as: other competing values; institutional frameworks; social norms and; knowledge of what is or is not acceptable in terms of ‘harm’ to nature (burden of proof) (Vucetich, Bruskotter & Nelson, 2015). This ‘deep green’ idea has, in the past, pervaded in the development of conservation policy, which often appears to lack any explicit connection between the ecological and the social system. In addressing this gap there has been a movement to “restore and reemphasize the fundamental links between nature and human wellbeing” through the development and application of the ecosystem services framework to conservation policy (Armsworth et al., 2007; Millennium Ecosystem Assessment, 2005; Natural Capital Committee, 2014; TEEB, 2010).

Increasingly, it is becoming recognized that approaches rooted firmly in ecological science are not the only solution to conservation issues (Hicks et al., 2016). Here we present a set of four focus points (Figure 1) for future development that can construct a bridge between the ecological and the social systems. The aim is to shift the discourse on “ecological coherence” further into the social sphere and hence support the alignment of the process of designating ecologically coherent MPA networks with the ‘triple bottom line’ of economic development, environmental sustainability and social inclusion as described in the SDGs to achieve social-ecological coherence in MPA network design.

178 **Figure 1:** The social-ecological system. Marine ecosystems are essential to maintain human wellbeing. The links between ecosystems (marine)
 179 and human well-being (adapted from NCC 2012) **REF** The policy frameworks from the CBD Aichi Target 11 and the UN Sustainable
 180 Development Goal 14 influence how marine ecosystems are managed with the aim to underpin human wellbeing. Four focus points are
 181 presented to develop social-ecological coherent MPA network.

182

[Figure 1 here]

2. Focus points

Focus point 1: A representative network of MPAs is more likely to reduce the risk of loss of ecosystem services

Representativity refers to the inclusion of the full range of ecosystems, habitats, biotic diversity, ecological processes, and environmental gradients (e.g. depth, wave exposure) within the MPA network (HELCOM, 2010; OSPAR, 2006; Roberts et al., 2003; Rondinini, 2010; UNEP-WCMC, 2008). The objective in applying this criterion to MPA networks is to ensure representative coverage of all biodiversity and biogeographic regions within the network (Jackson, Hiscock, Evans, Seeley & Lear, 2008; Roberts et al., 2003). The key premise behind representativity is that the full range of biodiversity is protected worldwide. This includes the species, as well as evolutionary patterns, distinct communities, refugia habitats in anticipation of environmental change and a range of key ecological processes that sustain global biodiversity (Spalding et al., 2007). Also, often included in processes for MPA network design, is a criterion for ‘replication’. Replication of habitats and species within an MPA network aims to ensure natural variation and to minimize the effects of damaging events and long-term changes (resilience), adequate replication of all habitats and species is recommended within MPA networks (HELCOM, 2010; OSPAR, 2007). Replication enhances the resilience of ecosystems to change and reduces the possibility that catastrophic events may wipe out entire populations of species or habitats within the network (HELCOM, 2010; OSPAR, 2007; Roberts et al., 2003).

Representativity (and replication) of species and habitats within an MPA network aims to underpin ecological resilience and to spread risk (of permanent loss, regime shifts) across a geographically broad region. From a social perspective the notion of ‘insurance’ is familiar for material goods. The valuation of goods and benefits derived from marine ecosystems falls firmly within the social system (Figure 1). According to the insurance hypothesis from an ecological perspective, biodiversity can provide insurance for ecosystems against potential future declines in their functioning because the more species that are present provides a greater likelihood that some will maintain functioning even if others are lost (Naeem & Li, 1997). Although species diversity does not necessarily guarantee high

resilience, in theory, ecosystems with high functional redundancy (functional overlap) will be more robust to loss of species than ecosystems with low functional redundancy (Naeem 1998). In coastal marine systems, however, limited research indicates that low functional redundancy could be typical even in the highest diversity ecosystems such as coral reefs (Micheli & Halpern 2005), yet very little is known about seascape redundancy with reference to habitat patch types. Ultimately, the loss of functional processes or regime shifts can impact upon the realization or delivery of ecosystem services that support human wellbeing (Folke et al., 2004).

Several researchers have argued that the CBD target of 10% is too low to achieve the objective of protecting biodiversity underpinning ecosystem services and meeting socioeconomic priorities (O'Leary et al., 2016). At the 2016 IUCN World Conservation Congress in Hawai'i, Resolution 50 calls on the Director General and the IUCN to "designate and implement at least 30% of each marine habitat in a network of highly protected MPAs and other effective area based conservation measures (OECMs), with the ultimate aim of creating a fully sustainable ocean, at least 30% of which has no extractive activities, subject to the rights of indigenous peoples and local communities" (IUCN, 2016). Proposals to achieve this encourage IUCN State and Government Agency Members to commit to a programme of work to designate and implement 30% of national waters as MPAs and OECMs; to engage in the process of establishing MPAs in areas beyond national jurisdiction (ABNJ) and to develop a new legally-binding instrument under the United Nations Law of the Sea "for the conservation and sustainable use of marine biological diversity in areas beyond national jurisdiction, and that such a new instrument contains a robust mechanism for establishing effectively and equitably managed, ecologically representative and well connected systems of marine protected areas, including reserves. Additionally IUCN member States are urged to accelerate progress towards achieving Aichi Target 11 and all Parties to the CBD are encouraged to develop post 2020 targets to achieve spatial management measures via MPAs for 30% of marine areas.

It must be noted that IUCN resolutions do not have any legal standing and that it is governments and competent international organisations that must further these resolutions into national and international policy. From a social perspective, SDG14 aims for the conservation and sustainable use of all the oceans, seas and marine resources, underpinned

by networks of MPAs (10% spatial coverage). Processes to determine the risk of loss of ecosystem goods and benefits and identify the potential societal consequences in relation to current levels of representativity within MPA networks may resonate further with governments and civil society to increase ambition for marine conservation (and sustainable resource use). There are recent innovations in next generation risk assessments for coupled natural-human systems (Elliott et al., 2017; Holsman et al., 2017) that can be applied to marine systems. At this stage, however, without an understanding of risk, an ecologically representative network of MPAs (meeting the CBD Aichi target 11 and SGD 14 10% target) may only be considered as a minimum spatial requirement to 'insure' for human wellbeing.

Focus point 2: A connected network of MPAs requires multisectoral planning processes that integrate different forms of spatial management structures.

Functional connectivity in ecological systems describes the extent to which populations in different parts of a species' range are linked by the exchange of eggs, larvae, recruits or other propagules, juveniles or adults (Palumbi, 2003). The connectivity between two populations is dependent on: (i) the larval characteristics of the species (e.g. duration of the planktonic stage and swimming behaviour of propagules), (ii) the abundance of the source population, (iii) the availability and suitability of surrounding habitat, and (iv) the characteristics of the physical environment (e.g. speed and direction of ocean currents, temperature, salinity) (Shanks, Grantham & Carr, 2003; Treml, Halpin, Urban & Pratson, 2008). The movements of adult life stages also influences connectivity and MPA performance and therefore requires consideration in MPA network design (Green et al., 2015; IUCN-WCPA, 2008; Olds et al., 2016). Understanding larval dispersal and marine population connectivity remains a highly complex analytical challenge. Processes to model connectivity require interdisciplinary approaches that combine high-resolution biophysical modeling and empirical data on movement capabilities of individual species (Cowen, Gawarkiewicz, Pineda, Thorrold & Werner, 2007).

Progress towards MPA networks that are considered well-connected are usually based on broad structural connectivity metrics that serve as spatial proxies for actual functional connectivity e.g. distance between MPAs. Geopolitical boundaries such as territorial limits of sovereign nations (i.e. exclusive economic zone boundaries) often appear to hamper

structural connectivity between MPA networks. Recent sub-regional assessments reveal that there is virtually no connectivity of MPAs across jurisdictional boundaries (Foster et al., 2014; Jessen, Morgan & Bezaury-Creel, 2016; Rees, Foster, Langmead & Griffiths, 2015a). As stated earlier there also remain relatively few MPAs in ABNJ (Marine Conservation Institute, 2017). Whilst understanding connectivity and translating this to MPA network planning remains challenging, the gaps in connectivity point towards a compounding societal challenge of how to further marine biodiversity protection in synergy with other management structures that operate across jurisdictional boundaries and in ABNJ.

The potential for OECMs aside from statutory MPAs to contribute to ecologically representative and well-connected MPA networks is increasingly receiving attention (Borrini-Feyerabend et al., 2014; Diz et al., 2017; Dunn, Maxwell, Boustany & Halpin, 2016; Jonas, Barbuto, Jonas, Kothari & Nelson, 2014; Laffoley et al., 2017; Spalding, Meliane, Milam, Fitzgerald & Hale, 2013; Woodley et al., 2012). There is currently no formal definition of an OECM under the CBD though a IUCN Task Force on ‘Other Effective Area based Conservation Measures’ has, so far, defined OECMs as “a geographical space where de-facto conservation of nature and associated ecosystem services and cultural values is achieved and expected to be maintained in the long term regardless of specific recognition and dedication” (Borrini-Feyerabend et al., 2014).

Potential OECMs may include some of the spatial management measures of Regional Fisheries Management Organizations (RMFOs) some of which, such as Vulnerable Marine Ecosystems (VMEs), are coincidental or overlap with Ecologically or Biologically Significant Areas (EBSAs) described by CBD Regional Workshops (Johnson et al., in review). The North East Atlantic Fisheries Commission (NEAFC) has been identified as an RFMO that has implemented ‘good practice’ through enacting closures to protect deep sea ecosystems from bottom towed fishing gear (Hoydal, Johnson & Hoel, 2014; Wright, Ardrøn, Gjerde, Currie & Rochette, 2015). In Australia, spatial closures that are enacted for fisheries by the Australian Fisheries Management Authority (AFMA) exceed the spatial extent of areas that are designated as Commonwealth Marine Reserves (CMR). It is argued by Bax and Cresswell (2012) that AFMA regulated areas are more restrictive on fishing activities than the proposed zonation of fisheries activities in the CMRs. Similar arguments can be put forward for environmental protection measures implemented by the International Maritime

305 Organization (Particularly Sensitive Sea Areas, Special Areas) and the International Seabed
306 Authority (Areas of Particular Environmental Interest). Diz et al (2017) also describe how
307 locally managed marine areas (LMMAs) with targeted biodiversity conservation can support
308 fishing communities.

309 From an SES perspective and to improve social and ecological coherence of MPA networks
310 though connectivity there is a need for States and competent international organizations to
311 collaborate to join up and recognize areas of ecological significance that support the
312 delivery of ecosystem goods and benefits and to coordinate protective measures through
313 multi sectoral planning to achieve this end.

314 There remains a diverse set of sectoral and political interests involved in the establishment
315 of multilateral agreements between countries for the protection of shared resources. From
316 a national perspective, ocean policy needs to reflect the importance of connected marine
317 ecosystems with an interdepartmental structure that mandates this policy. Moving forward
318 with such structures it should be considered a priority to invest in capacity building to train
319 practitioners with interdisciplinary skills who can facilitate the inclusion of a diverse set of
320 stakeholders into new shared governance structures to develop equitable rights and
321 management with regards to a network of MPAs and OECMs that is both socially and
322 ecological coherent. For ABNJ the prospect of a future legally binding “Implementing
323 Agreement for the protection of biodiversity beyond national jurisdiction” (BBNJ), with a
324 proposed focus on four distinct topics (the so-called “package”) and one being area-based
325 management tools, provides an important opportunity to make progress on this focus point
326 (Long & Rodriguez Chaves, 2015) .

327 **Focus point 3. Increase the size and number of MPAs with higher protection levels by**
328 **formalising procedures to track the performance of MPA management measures against**
329 **socio-economic outcomes.**

330 The World Commission on Protected Areas (WCPA) states that the ecological coherence of
331 MPA networks is supported by sites with a range of protection levels that are designed to
332 meet objectives that a single reserve cannot achieve (WCPA/IUCN, 2007). Indeed those
333 areas that have proven to have the most benefits for biodiversity are ‘no-take’ marine
334 reserves (IUCN Ia Strict Nature Reserve) where extractive activities are strictly controlled

(Edgar et al., 2014; Sciberras, Jenkins, Kaiser, Hawkins & Pullin, 2013). From a perspective of habitat recovery, some no-take MPAs have been shown to support complete shifts in the structure of ecosystems and reversal of trophic cascades (Behrens & Lafferty, 2004; Guidetti, 2007; Salomon, Shears, Langlois & Babcock, 2008). The potential of these highly protected sites to enhance sustainable economic development is strongly supported with evidence in the academic literature (Aburto-Oropeza et al., 2008; Halpern, Lester & Kellner, 2009; McCook et al., 2010). However, the practical reality is that no-take MPAs, particularly in the nearshore environment, are largely viewed as being unequitable and have, in many places, been difficult to implement because of social and political opposition (Agardy et al., 2003).

Equity, the premise that there is a fair distribution of benefits and costs between individuals and groups of people, is a subject that is recognized as having a potential to influence intended conservation outcomes (Tallis, Polasky, Lozano & Wolny, 2012) and it is embedded in the Aichi Target 11 text. Some accounts emerging from the academic literature demonstrate how supporting studies that document the ecological, social and economic impacts of MPA management measures (i.e. proof of the societal benefits of protection) can convince stakeholders of the efficacy area with higher protection levels (Erisman et al., 2017; Oliver et al., 2015; Rees et al., 2016; Vandeperre et al., 2011). From a community perspective these sites are essentially ‘control sites’ that support experimental design to robustly demonstrate the impact of management measures on the ecological system (e.g. reef recovery (Sheehan, Stevens, Gall, Cousens & Attrill, 2013)) and consequently benefits to the socio-economic system (e.g. increased landings (Rees et al., 2016)).

From a perspective of social-ecological coherence and achieving higher protection levels there is a need to formalize equity in the decision making process and track MPA performance beyond biological metrics. This can be achieved through the development of protocols for the monitoring and reporting on the effectiveness of MPAs (at an individual site and network level) that include socio-economic performance indicators (monetary and non-monetary) alongside conservation objectives for an MPA to reveal which characteristics of MPAs are most beneficial and acceptable to communities and how management measures can promote these.

Lessons can be learnt from behavioral psychology in that peoples' intentions to choose pro-environmental behavior (e.g. to agree and comply with no-take zones) is predominantly influenced by self-interest and pro-social motives (Bamberg & Möser, 2007). Areas with higher protection levels that can be used as 'control sites' for scientific monitoring essentially support equity via the opportunity for interested parties to be involved with rather than excluded from MPA management. This is an alternative consensus building approach, grounded in motivations that support human well-being (what can the MPA provide for me/my community), which may, in the long-term lead to greater social acceptability and broader spatial protection at higher levels.

Focus point 4: Networks of MPAs can provide investment opportunities.

Policy appraisal tools, applied by governments, are essential to consider the wider costs and benefits to society of an intervention, e.g. an MPA or a network of MPAs. Progress towards designation of ecologically coherent MPA networks, has, in some cases, slowed down or ground to a halt when subject to a policy appraisal at the government level (McGowan & Possingham, 2015; Rees et al., 2015). For example, Fletcher et al (2015) highlight a number of weaknesses in the recent Impact Assessment (IA) undertaken by UK Government to take forward the designation of the second round of MPAs (Marine Conservation Zones) required to work towards an ecologically coherent network of MPAs in England and Wales~~the UK~~. Costs are presented as quantified monetary values calculated using conventional economic assessment methods adapted to specific marine sectors, whereas benefits are described in non-monetary qualitative terms using an ecosystem services framework not tailored to specific sectors. This difference in methodology limits the policy appraisal tool's ability to fairly assess the costs and benefits of effective conservation. Furthermore, no method is employed in the IA to take into account the cost of inaction (i.e. of doing nothing) despite evidence that demonstrates that there are opportunity costs associated with delayed conservation action or inaction (Grantham, Wilson, Moilanen, Rebelo & Possingham, 2009) and that the overall health and functionality of the marine environment is deteriorating (Jackson et al., 2001; Lotze et al., 2006; Worm et al., 2006), thus, requiring a corresponding declining economic baseline. Additionally, there is no consideration of the benefits of an ecologically coherent network of MPAs versus site-based costs. The result of this policy appraisal process is that where an MPA is contested, (considered to be inequitable though

for example, loss of income, opportunity or rights) then gaps appear in the MPA network that undermine ecological coherence. It is argued that such contention can lead to a network of 'residual' MPAs that afford no step-change in the management of activities and therefore no additional benefits for biodiversity (Devillers et al., 2015).

Undeniably the purpose of policy appraisal processes are to define social and economic equity in the decision-making processes and this can identify inequitable trade-offs between biodiversity conservation and socio-economic objectives. However, equity is more than a narrow trade-off between resource use and rights occurring directly within the boundaries of the MPA. MPAs may have both localised and broader societal benefits. A series of studies in different parts of the world are contributing to a body of evidence that supports this. For example, an economic investment by the US government (American Recovery and Reinvestment Act of 2009) to restore degraded coastal habitat (blue infrastructure) has led to job creation in the short term, further economic benefits through the rebuilding of fisheries and coastal tourism and benefits to coastal economies, such as higher property values and improved water quality (Edwards, Sutton-Grier & Coyle, 2013). McCook et al. (2010) demonstrate that the economic returns of the rezoning of the Great Barrier Reef Marine Park are estimated to be 130 times greater than the cost of management. Further protection for corals and fish could potentially have knock on benefits (opportunity costs) for the tourist industry and commercial fisheries (McCook et al., 2010). Arkema et al. (2015) showed that presence of intact reefs and coastal vegetation reduce the likelihood and magnitude of losses resulting from extreme weather events and sea-level rise. Jackson, Rees, Wilding, & Attrill (2015) demonstrate that, by providing habitat for species during essential life history stages, seagrass (*Posidonia oceanica*) meadows are worth around €78 million every year to commercial fishing and €112 million to recreational fishing in the Mediterranean. Investment in effective conservation will underpin this value and the associated employment. In the U.S. Virgin Islands, economic valuation has estimated that nearshore coral reefs are worth approximately US\$200 million annually (Van-Beukering, Brander, Zanten, Verbrugge & Lems, 2011) and in Hawai'i are estimated at US\$360 million per year (Cesar & Beukering, 2004). These are powerful socio-political arguments for investment in conservation.

New strategies are required from Governments to move decision-making beyond site-based cost benefit analysis towards broader strategies for investment in ecosystems (Natural Capital Committee, 2014). It is possible for identified centralized costs (e.g. enforcement or management) to become opportunities for investment (supporting sustainable growth) or offsetting (loss of jobs) rather than a direct trade off against site based ecosystem service benefits. Flagship projects led by national governments that integrate investment (both public and private) with the conservation of priority biodiversity areas e.g. The Brazil Blue Fund are pioneers in the development of coastal and marine conservation strategies that aim to underpin sustainable development in a new accounting model. Exactly how an 'ecology coherent' network of MPAs may support such investment strategies is yet to be fully substantiated. However, the reduction of risk (of loss of benefits) through planning for aspects of ecological coherence, such as representativity of habitats and species within an MPA network (focus point one), along with case studies that demonstrate positive social and economic outcomes (Focus point 3) would seem to support such investment strategies. Such an approach might enable MPA managers to become more influential in marine spatial planning activities that envision future scenarios of the optimal use and allocation of maritime space.

3. Conclusion

Despite the development of appropriate criteria, thresholds and policy frameworks (to develop ecologically coherent networks of MPAs, overall progress towards achieving them is slow. Whilst biodiversity considerations underpin MPA selection, the SDGs with their 17 goals to "transform our world" are broadening the focus of ecological sustainability to encompass social and economic objectives. This leads us to a wider consideration of how conservation can underpin human well-being and how the current drive towards ecologically coherent networks of MPAs can be aligned with broader policy objectives.

The key premises are that 1) MPAs are an effective spatial management tool for achieving conservation objectives; and 2) Ecologically coherent networks support the high level of functional and spatial connectivity within marine ecosystems so that the network as a whole supports those ecosystem services that underpin human wellbeing. The focus points presented here are directed at those in a position to influence MPA policy and/or MPA

469 management with the intention of shifting the discourse of “ecological coherence” into the
470 social and economic sphere. Reframing the discourse for ecological coherence in this way
471 offers opportunities for integration with other disciplines beyond conservation biology such
472 as individuals or groups that specialize, for example, in risk management, finance,
473 investment, natural capital assessments and performance management. There are also
474 opportunities for new partnerships with wider stakeholder groups who operate in the
475 marine environment but under different sectoral management strategies e.g. ocean energy.
476 Such integration may advance progress towards the aim of ecologically coherent networks
477 of MPAs and therefore support the ecological, social and economic goals (the ‘triple bottom
478 line’ of sustainable development) outlined in the SDGs.

479

4. Acknowledgements

The authors would like to thank WWF who funded the ‘Assessment of the Ecological coherence of the Celtic Seas MPA network’ which initiated ideas for this paper. Also the participants of the ‘Expert meeting’ to share experiences and lessons learned on achieving qualitative elements of Aichi Target 11 in marine and coastal areas, convened by the Secretariat of the Convention on Biological Diversity and the Secretariat of the Global Ocean Biodiversity Initiative (Berlin, 24-26 February 2016) whose ideas and discussions informed this paper. The outputs presented here reflect the views of the named authors and are not the consensus of the CBD expert group. This work was funded with ‘in-kind’ support from the authors’ institutions. DJ would like to recognize support from the Global Ocean Biodiversity Initiative International Climate Initiative supported financially by the German government, Federal Ministry for the Environment, Nature Conservation, Building and Nuclear Safety, (Bundesministerium für Umwelt, Naturschutz, Bau und Reaktorsicherheit) (BMUB)).

5. References

- Aburto-Oropeza, O., Ezcurra, E., Danemann, G., Valdez, V., Murray, J. & Sala, E. (2008) 'Mangroves in the Gulf of California increase fishery yields'. *Proceedings of the National Academy of Sciences*, 105 (30), pp. 10456-10459.
- Agardy, T., Bridgewater, P., Crosby, M. P., Day, J., Dayton, P. K., Kenchington, R., . . . Peau, L. (2003) 'Dangerous targets? Unresolved issues and ideological clashes around marine protected areas'. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 13 (4), pp. 353-367.
- Ardron, J. A. (2008) 'The challenge of assessing whether the OSPAR network of marine protected areas is ecologically coherent'. *Hydrobiologia*, 606 pp. 45-53.
- Arkema, K. K., Guannel, G., Verutes, G., Wood, S. A., Guerry, A., Ruckelshaus, M., . . . Silver, J. M. (2013) 'Coastal habitats shield people and property from sea-level rise and storms'. *Nature Clim. Change*, 3 (10), pp. 913-918.
- Arkema, K. K., Verutes, G. M., Wood, S. A., Clarke-Samuels, C., Rosado, S., Canto, M., . . . Guerry, A. D. (2015) 'Embedding ecosystem services in coastal planning leads to better outcomes for people and nature'. *Proceedings of the National Academy of Sciences*, 112 (24), pp. 7390-7395.
- Armsworth, P. R., Chan, K. M. A., Daily, G. C., Ehrlich, P. R., Kremen, C., Ricketts, T. H. & Sanjayan, M. A. (2007) 'Ecosystem-Service Science and the Way Forward for Conservation'. *Conservation Biology*, 21 (6), pp. 1383-1384.
- Bamberg, S. & Möser, G. (2007) 'Twenty years after Hines, Hungerford, and Tomera: A new meta-analysis of psycho-social determinants of pro-environmental behaviour'. *Journal of Environmental Psychology*, 27 (1), pp. 14-25.
- Barnard, S., Burdon, D., Strong, J. & Atkins, J. (2014) *The Ecological Coherence and Economic & Social Benefits of the Northern Ireland MPA Network*. Hull, UK, (Report No. YBB238-F-2014). Available.
- Bax, N. & Cresswell, I. (2012) Marine reserves not about closing fisheries, but about preserving ocean health. *The Conversation. The Conversation Trust (UK) Limited.*,
- Behrens, M. D. & Lafferty, K. D. (2004) 'Effects of marine reserves and urchin disease on southern Californian rocky reef communities'. *Marine Ecology Progress Series*, 279 pp. 129-139.

535
536 Berkes, F., Folke, C. & Colding, J. (2000) *Linking Social and Ecological Systems: Management*
537 *Practices and Social Mechanisms for Building Resilience*. Cambridge: University Press.

538
539 Borrini-Feyerabend, G., P. , Bueno, T., Hay-Edie, B., Lang, A., Rastogi, A. & Sandwith, T.
540 (2014) *A primer on governance for protected and conserved areas, Stream on Enhancing*
541 *Diversity and Quality of Governance*. Gland, Switzerland: IUCN. Available.

542
543 CBD (2010) 'Convention on Biological Diversity. COP 10. Decision X/2.Strategic Plan for
544 Biodiversity 2011-2020'.

545
546 Cesar, H. S. J. & Beukering, P. v. (2004) 'Economic Valuation of the Coral Reefs of Hawai'i'.
547 *Pacific Science*, 58 (2), pp. 231-242.

548
549 Cowen, R. K., Gawarkiewicz, G., Pineda, J., Thorrold, S. R. & Werner, F. E. (2007) 'Population
550 Connectivity in Marine Systems An Overview'. *Oceanography*, 20 (3), pp. 14-21.

551
552 D. Spalding, M., Meliane, I., J. Bennett, N., Dearden, P., G. Patil, P. & D. Brumbaugh, R.
553 (2016) 'Building towards the marine conservation end-game: consolidating the role of MPAs
554 in a future ocean'. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 26 pp. 185-
555 199.

556
557 Devillers, R., Pressey, R. L., Grech, A., Kittinger, J. N., Edgar, G. J., Ward, T. & Watson, R.
558 (2015) 'Reinventing residual reserves in the sea: are we favouring ease of establishment
559 over need for protection?'. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 25
560 (4), pp. 480-504.

561
562 Diz, D., Johnson, D., Riddell, M., Rees, S., Battle, J., Gjerde, K., . . . Roberts, J. M. (2017)
563 'Mainstreaming marine biodiversity into the SDGs: The role of other effective area-based
564 conservation measures (SDG 14.5)'. *Marine Policy*,

565
566 Diz, D., Morgera, E. & Wilson, M. (2017) 'Marine policy special issue: SDG synergies for
567 sustainable fisheries and poverty alleviation'. *Marine Policy*,

568
569 Dunn, D. C., Maxwell, S. M., Boustany, A. M. & Halpin, P. N. (2016) 'Dynamic ocean
570 management increases the efficiency and efficacy of fisheries management'. *Proceedings of*
571 *the National Academy of Sciences*, 113 (3), pp. 668-673.

572

573 Edgar, G. J., Stuart-Smith, R. D., Willis, T. J., Kininmonth, S., Baker, S. C., Banks, S., . . .
574 Thomson, R. J. (2014) 'Global conservation outcomes depend on marine protected areas
575 with five key features'. *Nature*, 506 (7487), pp. 216-220.

576
577 Edwards, P. E. T., Sutton-Grier, A. E. & Coyle, G. E. (2013) 'Investing in nature: Restoring
578 coastal habitat blue infrastructure and green job creation'. *Marine Policy*, 38 pp. 65-71.

579
580 Elliott, M., Burdon, D., Atkins, J. P., Borja, A., Cormier, R., de Jonge, V. N. & Turner, R. K.
581 (2017) "'And DPSIR begat DAPSI(W)R(M)!' - A unifying framework for marine environmental
582 management'. *Marine Pollution Bulletin*, 118 (1), pp. 27-40.

583
584 Erisman, B., Heyman, W., Kobara, S., Ezer, T., Pittman, S., Aburto-Oropeza, O. & Nemeth, R.
585 S. (2017) 'Fish spawning aggregations: where well-placed management actions can yield big
586 benefits for fisheries and conservation'. *Fish and Fisheries*, 18 (1), pp. 128-144.

587
588 Evans, J. L., Peckett, F. & Howell, K. L. (2015) 'Combined application of biophysical habitat
589 mapping and systematic conservation planning to assess efficiency and representativeness
590 of the existing High Seas MPA network in the Northeast Atlantic'. *ICES Journal of Marine
591 Science: Journal du Conseil*,

592
593 Fletcher, S. & Rees, S. (2015) *Marine Conservation Zone Impact Assessment Review. A
594 report for The Wildlife Trusts by the Centre for Marine and Coastal Policy Research,
595 Plymouth University*. 31 pp. Available.

596
597 Folke, C., Carpenter, S., Walker, B., Scheffer, M., Elmqvist, T., Gunderson, L. & Holling, C. S.
598 (2004) 'Regime Shifts, Resilience, and Biodiversity in Ecosystem Management'. *Annual
599 Review of Ecology, Evolution, and Systematics*, 35 (1), pp. 557-581.

600
601 Foster, N. L., Sciberras, M., Jackson, E., Ponge, B., Toison, V., Carrier, S., . . . Attrill, M. (2014)
602 *Assessing the Ecological Coherence of the Channel MPA Network. Report prepared by the
603 Marine Institute for the Protected Area Network Across the Channel Ecosystem (PANACHE)
604 project. INTERREG programme France (Channel) England funded project*. 156 pp. Available.

605
606 Grantham, H. S., Wilson, K. A., Moilanen, A., Rebelo, T. & Possingham, H. P. (2009) 'Delaying
607 conservation actions for improved knowledge: how long should we wait?'. *Ecology Letters*,
608 12 (4), pp. 293-301.

609
610 Green, A. L., Maypa, A. P., Almany, G. R., Rhodes, K. L., Weeks, R., Abesamis, R. A., . . .
611 White, A. T. (2015) 'Larval dispersal and movement patterns of coral reef fishes, and
612 implications for marine reserve network design'. *Biological Reviews*, 90 (4), pp. 1215-1247.

613
614 Guidetti, P. (2007) 'Potential of Marine Reserves to Cause Community-Wide Changes
615 beyond Their Boundaries

616 EL Potencial de Reservas Marinas para Provocar Cambios a Nivel de Comunidad Más Allá de
617 sus Límites'. *Conservation Biology*, 21 (2), pp. 540-545.

618
619 Halpern, B. S., Lester, S. E. & Kellner, J. B. (2009) 'Spillover from marine reserves and the
620 replenishment of fished stocks'. *Environmental Conservation*, 36 (04), pp. 268-276.

621
622 HELCOM (2010) *Towards an ecologically coherent network of well-managed Marine*
623 *Protected Areas – Implementation report on the status and ecological coherence of the*
624 *HELCOM BSPA network*. Balt. Sea Environ. Proc. No. 124B. 148 pp. Available.

625
626 Hicks, C. C., Levine, A., Agrawal, A., Basurto, X., Breslow, S. J., Carothers, C., . . . Levin, P. S.
627 (2016) 'Engage key social concepts for sustainability'. *Science*, 352 (6281), pp. 38-40.

628
629 Hockings, M., Stolton, S., Leverington, F., Dudley, N. & Courrau, J. (2006) *Evaluating*
630 *Effectiveness: A framework for assessing management effectiveness of protected areas*. 2nd
631 *Edition*. Gland, Switzerland and Cambridge, UK: IUCN. 121 pp. Available.

632
633 Holsman, K., Samhouri, J., Cook, G., Hazen, E., Olsen, E., Dillard, M., . . . Andrews, K. (2017)
634 'An ecosystem-based approach to marine risk assessment'. *Ecosystem Health and*
635 *Sustainability*, 3 (1), pp. e01256-n/a.

636
637 Hoydal, K., Johnson, D. & Hoel, A. H. (2014) *Regional governance: the case of NEAFC and*
638 *OSPAR. Chapter 16: 225-238*. eds. Garcia, S.M., Rice, J. and Charles, A., *Governance for*
639 *Fisheries and Marine Conservation: Interaction and co-evolution*. Wiley-Blackwell

640
641 International Union for the Conservation of Nature (2016) 'Green List'. International Union
642 for Conservation of Nature. (Accessed: 23 September 2016).

643
644 IUCN-WCPA (2008) *Establishing Marine Protected Area Networks—Making It Happen*. .
645 Washington, D.C: IUCN, World Commission on Protected Areas, National Oceanic and
646 Atmospheric Administration and The Nature Conservancy. 118 pp. Available.

647
648 IUCN (2016) *IUCN Resolutions, Recommendations and other Decisions*. Gland, Switzerland:
649 IUCN. 106pp pp. Available.

650
651 Jackson, E. L., Hiscock, K., Evans, J. L., Seeley, B. & Lear, D. B. (2008) *Investigating the*
652 *existing coverage and subsequent gaps in protection and providing guidance on*

653 *representativity and replication for a coherent network of Marine Protected Areas in*
654 *England's territorial waters*. Plymouth: Marine Life Information Network (MarLIN), Marine
655 Biological Association of the UK. Natural England Commissioned Reports, Number 018. 1-
656 138 pp. Available.

657

658 Jackson, E. L., Rees, S. E., Wilding, C. & Attrill, M. J. (2015) 'Use of a seagrass residency index
659 to apportion commercial fishery landing values and recreation fisheries expenditure to
660 seagrass habitat service'. *Conservation Biology*, 29 (3), pp. 899-909.

661

662 Jackson, J. B. C., Kirby, M. X., Berger, W. H., Bjorndal, K. A., Botsford, L. W., Bourque, B.
663 J., . . . Warner, R. R. (2001) 'Historical Overfishing and the Recent Collapse of Coastal
664 Ecosystems'. *Science*, 293 (5530), pp. 629-637.

665

666 Jessen, S., Morgan, L. & Bezaury-Creel, J. (2016) *Dare to be Deep: SeaStates Report on North*
667 *America's Marine Protected Areas (MPAs)*. Ottawa, Seattle, and México City: Canadian Parks
668 and Wilderness Society, Marine Conservation Institute. 52 pp. Available at:
669 <http://cpaws.org/uploads/CPAWS-Oceans-Report-2016.pdf>.

670

671 Johnson, D., Ardron, J., Billett, D., Hooper, T., Mullier, T., Chaniotis, P., . . . Corcoran, E.
672 (2014) 'When is a marine protected area network ecologically coherent? A case study from
673 the North-east Atlantic'. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24 (S2),
674 pp. 44-58.

675

676 Jonas, H. D., Barbuto, V., Jonas, H. C., Kothari, A. & Nelson, F. (2014) 'New Steps of Change:
677 Looking Beyond Protected Areas to Consider Other Effective Area-Based Conservation
678 Measures'. *PARKS*, 20 (2),

679

680 Jones, G. P., Srinivasan, M. & Almany, G. R. (2007) 'Population Connectivity and
681 Conservation of Marine Biodiversity'. *Oceanography*, 20 (3), pp. 100-111.

682

683 Jones, P. J. S. & De Santo, E. M. (2016) 'Viewpoint – Is the race for remote, very large marine
684 protected areas (VLMPAs) taking us down the wrong track?'. *Marine Policy*, 73 pp. 231-234.

685

686 Laffoley, D., Brockington, S. & Gililand, P. M. (2006) *Developing the concepts of good*
687 *environmental status and marine ecosystem objectives: some important considerations*.
688 Peterborough, UK: English Nature Research Reports, No 689. 38 pp. Available.

689

690 Laffoley, D., Dudley, N., Jonas, H., MacKinnon, D., MacKinnon, K., Hockings, M. & Woodley,
691 S. (2017) 'An introduction to 'other effective area-based conservation measures' under Aichi
692 Target 11 of the Convention on Biological Diversity: Origin, interpretation and emerging
693 ocean issues'. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 27 pp. 130-137.

694
695 Lester, S. E., Halpern, B. S., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B. I., Gaines, S.
696 D., . . . Warner, R. R. (2009) 'Biological effects within no-take marine reserves: a global
697 synthesis'. *Marine Ecology Progress Series*, 384 pp. 33-46.

698
699 Lieberknecht, L., Mullier, T. & Ardron, J. (2014) *Assessment of the ecological coherence of*
700 *the UK's marine protected area network. A report prepared for the Joint Links*. Available.

701
702 Liu, J., Dietz, T., Carpenter, S. R., Alberti, M., Folke, C., Moran, E., . . . Taylor, W. W. (2007)
703 'Complexity of Coupled Human and Natural Systems'. *Science*, 317 (5844), pp. 1513-1516.

704
705 Long, R. & Rodriguez Chaves, M. (2015) 'Anatomy of a new international instrument for
706 marine biodiversity beyond national jurisdiction: first impressions of the preparatory
707 process. '. *Environmental Liability, Law, Policy and Practice.*, 23 (6), pp. 213-229.

708
709 Lotze, H. K., Lenihan, H. S., Bourque, B. J., Bradbury, R. H., Cooke, R. G., Kay, M. C., . . .
710 Jackson, J. B. C. (2006) 'Depletion, Degradation, and Recovery Potential of Estuaries and
711 Coastal Seas'. *Science*, 312 (5781), pp. 1806-1809.

712
713 Lubchenco, J. & Grorud-Colvert, K. (2015) 'Making waves: The science and politics of ocean
714 protection'. *Science*, 350 (6259), pp. 382-383.

715
716 Margules, C. R. & Pressey, R. L. (2000) 'Systematic conservation planning'. *Nature*, 405
717 (6783), pp. 243-253.

718
719 Marine Conservation Institute (2017) 'MPAtlas [On-line]. Available at www.mpatlas.org'.
720 [Online]. Available at: www.mpatlas.org (Accessed: 13.04.2016).

721
722 McCook, L. J., Ayling, T., Cappo, M., Choat, J. H., Evans, R. D., De Freitas, D. M., . . .
723 Williamson, D. H. (2010) 'Adaptive management of the Great Barrier Reef: A globally
724 significant demonstration of the benefits of networks of marine reserves'. *Proceedings of*
725 *the National Academy of Sciences*, 107 (43), pp. 18278-18285.

726
727 McGowan, J. & Possingham, H. P. (2015) *Submission to the Commonwealth Marine Reserves*
728 *Review . A report by the ARC Centre of Excellence for Environmental Decisions The University*
729 *of Queensland*. 4 pp. Available.

730
731 Millennium Ecosystem Assessment (2005) *Ecosystems and human well-being: Synthesis*.
732 Washington, D.C.: World Resources Institute. 155 pp. Available.

733

734 Naeem, S. & Li, S. (1997) 'Biodiversity enhances ecosystem reliability'. *Nature*, 390 (6659),
735 pp. 507-509.

736

737 Natural Capital Committee (2014) *The State of Natural Capital: Restoring our Natural Assets.*
738 *Second report to the Economic Affairs Committee.* 86 pp. Available.

739

740 O'Leary, B. C., Winther-Janson, M., Bainbridge, J. M., Aitken, J., Hawkins, J. P. & Roberts, C.
741 M. (2016) 'Effective Coverage Targets for Ocean Protection'. *Conservation Letters*, 9 (6), pp.
742 398-404.

743

744 Olds, A. D., Connolly, R. M., Pitt, K. A., Pittman, S. J., Maxwell, P. S., Huijbers, C. M., . . .
745 Schlacher, T. A. (2016) 'Quantifying the conservation value of seascape connectivity: a global
746 synthesis'. *Global Ecology and Biogeography*, 25 (1), pp. 3-15.

747

748 Oliver, T. A., Oleson, K. L. L., Ratsimbazafy, H., Raberinary, D., Benbow, S. & Harris, A. (2015)
749 'Positive Catch & Economic Benefits of Periodic Octopus Fishery Closures: Do Effective,
750 Narrowly Targeted Actions 'Catalyze' Broader Management?'. *PLoS ONE*, 10 (6), pp.
751 e0129075.

752

753 Olsen, E. M., Johnson, D., Waever, P., Goni, R., Ribeiro, M. C., Rabaut, M., . . . Zaharia, T.
754 (2013) *Achieving ecologically coherent MPA networks in Europe: Science needs and*
755 *priorities.* Ostend, Belgium: European Marine Board. 1-88 pp. Available.

756

757 OSPAR (2006) *Guidance on developing an ecologically coherent network of OSPAR marine*
758 *protected areas.* London, UK: OSPAR Commission. 11 pp. Available.

759

760 OSPAR (2007) *Background document to support the assessment of whether the OSPAR*
761 *Network of Marine Protected Areas is ecologically coherent.* London, UK: OSPAR
762 Commission. Available.

763

764 OSPAR (2013) *An assessment of the ecological coherence of the OSPAR Network of Marine*
765 *Protected Areas in 2012.* London, UK: OSPAR Commission. 76 pp. Available.

766

767 Palumbi, S. R. (2003) 'Population genetics, demographic connectivity, and the design of
768 marine reserves'. *Ecological Applications*, 13 (sp1), pp. 146-158.

769

770 Piekäinen, H. & Korpinen, S. (2008) *Towards an Assessment of Ecological Coherence of the*
771 *Marine Protected Areas Network in the Baltic Sea Region.* Available.

772

773 Pittman, S. J., Monaco, M. E., Friedlander, A. M., Legare, B., Nemeth, R. S., Kendall, M. S., . . .
 774 Caldwell, C. (2014) 'Fish with Chips: Tracking Reef Fish Movements to Evaluate Size and
 775 Connectivity of Caribbean Marine Protected Areas'. *PLoS ONE*, 9 (5), pp. e96028.
 776
 777 Pollnac, R., Christie, P., Cinner, J. E., Dalton, T., Daw, T. M., Forrester, G. E., . . . McClanahan,
 778 T. R. (2010) 'Marine reserves as linked social–ecological systems'. *Proceedings of the*
 779 *National Academy of Sciences*, 107 (43), pp. 18262-18265.
 780
 781 Potts, T., Burdon, D., Jackson, E., Atkins, J., Saunders, J., Hastings, E. & Langmead, O. (2014)
 782 'Do marine protected areas deliver flows of ecosystem services to support human welfare?'.
 783 *Marine Policy*, 44 (0), pp. 139-148.
 784
 785 Rees, S., Foster, N., Langmead, O. & Griffiths, C. (2015a) *Assessment of the Ecological*
 786 *Coherence of the MPA Network in the Celtic Seas: A report for WWF-UK by the Marine*
 787 *Institute, Plymouth University and The Marine Biological Association of the United*
 788 *Kingdom*. . 167 pp. Available.
 789
 790 Rees, S., Foster, N., Langmead, O. & Griffiths, C. (2015b) 'Assessment of the Ecological
 791 Coherence of the MPA Network in the Celtic Seas: A report for WWF-UK by the Marine
 792 Institute, Plymouth University and The Marine Biological Association of the United
 793 Kingdom'. [in, 120. (Accessed: Rees, S., Foster, N., Langmead, O. & Griffiths, C.
 794
 795 Rees, S., Sheehan, E., Foster, N., Rees, A., Gall, S., Pittman, S. & Shellock, R. (2015)
 796 *Consultation on the second tranche of Marine Conservation Zones 2015. Response from the*
 797 *Marine Institute at Plymouth University*. Plymouth University. 8 pp. Available.
 798
 799 Rees, S. E., Ashley, M., Evans, L., Mangi, S., Rodwell, L., Attrill, M., . . . Rees, A. (2016) *An*
 800 *evaluation framework to determine the impact of the Lyme Bay Marine Protected Area and*
 801 *the activities of the Lyme Bay Consultative Committee on ecosystem services and human*
 802 *wellbeing. A report to the Blue Marine Foundation by research staff the Marine Institute at*
 803 *Plymouth University, Exeter University and Cefas*. Pp139. Available.
 804
 805 Rees, S. E., Fletcher, S., Gall, S. C., Friedrich, L. A., Jackson, E. L. & Rodwell, L. D. (2014)
 806 'Securing the benefits: Linking ecology with marine planning policy to examine the potential
 807 of a network of Marine Protected Areas to support human wellbeing'. *Marine Policy*, 44 (0),
 808 pp. 335-341.
 809
 810 Rees, S. E., Foster, N. L., Langmead, O., Pittman, S. & Johnson, D. E. (2017) 'Defining the
 811 qualitative elements of Aichi Biodiversity Target 11 with regard to the marine and coastal
 812 environment in order to strengthen global efforts for marine biodiversity conservation
 813 outlined in the United Nations Sustainable Development Goal 14'. *Marine Policy*,

814
815 Rees, S. E., Rodwell, L. D., Attrill, M. J., Austen, M. C. & Mangi, S. C. (2010) 'The value of
816 marine biodiversity to the leisure and recreation industry and its application to marine
817 spatial planning'. *Marine Policy*, 34 (5), pp. 868-875.

818
819 Ridgeway, A., Cornthwaite, A., Wright, H. & Davies, J. (2014) *Identifying the remaining MCZ*
820 *site options that would fill big gaps in the existing MPA network around England and*
821 *offshore waters of Wales & Northern Ireland*. JNCC. Available at:
822 <http://jncc.defra.gov.uk/page>.

823
824 Roberts, C. M., Bohnsack, J. A., Gell, F., Hawkins, J. P. & Goodridge, R. (2001) 'Effects of
825 Marine Reserves on Adjacent Fisheries'. *Science*, 294 (5548), pp. 1920-1923.

826
827 Roberts, C. M., Branch, G., Bustamante, R. H., Castilla, J. C., Dugan, J., Halpern, B. S., . . .
828 Warner, R. R. (2003) 'Application of ecological criteria in selecting marine reserves and
829 developing reserve networks'. *Ecological Applications*, 13 (1), pp. S215-S228.

830
831 Rodríguez-Rodríguez, D., Rodríguez, J. & Abdul Malak, D. (2016) 'Development and testing of
832 a new framework for rapidly assessing legal and managerial protection afforded by marine
833 protected areas: Mediterranean Sea case study'. *Journal of Environmental Management*,
834 167 pp. 29-37.

835
836 Rondinini, C. (2010) *Meeting the MPA network design principles of representation and*
837 *adequacy: developing species-area curves for habitats*. JNCC Report No. 439. 1-45 pp.
838 Available.

839
840 Sachs, J. D. (2012) 'From Millennium Development Goals to Sustainable Development
841 Goals'. *The Lancet*, 379 (9832), pp. 2206-2211.

842
843 Salomon, A. K., Shears, N. T., Langlois, T. J. & Babcock, R. C. (2008) 'CASCADING EFFECTS OF
844 FISHING CAN ALTER CARBON FLOW THROUGH A TEMPERATE COASTAL ECOSYSTEM'.
845 *Ecological Applications*, 18 (8), pp. 1874-1887.

846
847 Sciberras, M., Jenkins, S. R., Kaiser, M. J., Hawkins, S. J. & Pullin, A. S. (2013) 'Evaluating the
848 biological effectiveness of fully and partially protected marine areas'. *Environmental*
849 *Evidence*, 2 (1), pp. 1-31.

850
851 Secretariat of the Convention on Biological Diversity (2014) *Global Biodiversity Outlook 4*.
852 Montréal. 155 pp. Available.

853

854 Shanks, A. L., Grantham, B. A. & Carr, M. H. (2003) 'Propagule Dispersal Distance and the
855 Size and Spacing of Marine Reserves'. *Ecological Applications*, 13 (1), pp. S159-S169.

856

857 Sheehan, E. V., Stevens, T. F., Gall, S. C., Cousens, S. L. & Attrill, M. J. (2013) 'Recovery of a
858 Temperate Reef Assemblage in a Marine Protected Area following the Exclusion of Towed
859 Demersal Fishing'. *PLoS ONE*, 8 (12), pp. e83883.

860

861 Sobel, J. & Dahlgren, C. P. (2004) *Marine Reserves: A Guide to Science, Design and Use*.
862 Washington DC: Island Press.

863

864 Spalding, M., Meliane, I., Milam, A., Fitzgerald, C. & Hale, L. (2013) 'Protecting Marine
865 Spaces: Global Targets and Changing Approaches'. *Ocean Yearbook*, 27 pp. 213-248.

866

867 Spalding, M. D., Fox, H. E., Allen, G. R., Davidson, N., Ferdaña, Z. A., Finlayson, M., . . .
868 Robertson, J. (2007) 'Marine Ecoregions of the World: A Bioregionalization of Coastal and
869 Shelf Areas'. *BioScience*, 57 (7), pp. 573-583.

870

871 Stewart, G. B., Kaiser, M. J., Côté, I. M., Halpern, B. S., Lester, S. E., Bayliss, H. R. & Pullin, A.
872 S. (2009) 'Temperate marine reserves: global ecological effects and guidelines for future
873 networks'. *Conservation Letters*, 2 (6), pp. 243-253.

874

875 Tallis, H., Polasky, S., Lozano, J. S. & Wolny, S. (2012) *Inclusive wealth accounting for
876 regulating ecosystem services. Inclusive Wealth Report 2012. Measuring progress toward
877 sustainability* Cambridge, uk. Available.

878

879 TEEB (2010) *The Economics of Ecosystems and Biodiversity: The Ecological and Economic
880 Foundations*. London, Washington DC: UNEP. 411 pp. Available.

881

882 Tognelli, M. F., Fernández, M. & Marquet, P. A. (2009) 'Assessing the performance of the
883 existing and proposed network of marine protected areas to conserve marine biodiversity in
884 Chile'. *Biological Conservation*, 142 (12), pp. 3147-3153.

885

886 Treml, E., Halpin, P., Urban, D. & Pratson, L. (2008) 'Modeling population connectivity by
887 ocean currents, a graph-theoretic approach for marine conservation'. *Landscape Ecology*, 23
888 (1), pp. 19-36.

889

890 UNEP-WCMC (2008) *National and Regional Networks of Marine Protected Areas: A Review
891 of Progress*. Cambridge, UK: UNEP-WCMC. 156 pp. Available.

892

893 Van-Beukering, P. J. H., Brander, L., Zanten, B. v., Verbrugge, E. & Lems, K. (2011) *The*
894 *Economic Value of the Coral Reef Ecosystems of the United States Virgin Islands. Report*
895 *number R-11/06. Institute for Environmental Studies (IVM), Amsterdam.*

896

897 Vandeperre, F., Higgins, R. M., Sanchez-Meca, J., Maynou, F., Goni, R. & Martin-Sosa, P.
898 (2011) 'Effects of no-take area size and age of marine protected areas on fisheries yields: a
899 meta-analytical approach'. *Fish Fish*, 12

900

901 Vucetich, J. A., Bruskotter, J. T. & Nelson, M. P. (2015) 'Evaluating whether nature's intrinsic
902 value is an axiom of or anathema to conservation'. *Conservation Biology*, 29 (2), pp. 321-
903 332.

904

905 WCPA/IUCN (2007) *Establishing networks of marine protected areas: A guide for developing*
906 *national and regional capacity for building MPA networks. Non-technical summary report.*
907 Gland, Switzerland: IUCN. 16 pp. Available.

908

909 Woodley, S., Bertzky, B., Crawhall, N., Dudley, N., Londoño, J. M., MacKinnon, K., . . .
910 Sandwith, T. (2012) 'MEETING AICHI TARGET 11: WHAT DOES SUCCESS LOOK LIKE FOR
911 PROTECTED AREA SYSTEMS?'. *PARKS*, 18 (1), pp. 23-34.

912

913 Worm, B., Barbier, E. B., Beaumont, N., Duffy, J. E., Folke, C., Halpern, B. S., . . . Watson, R.
914 (2006) 'Impacts of Biodiversity Loss on Ocean Ecosystem Services'. *Science*, 314 (5800), pp.
915 787-790.

916

917 Wright, G., Ardron, J., Gjerde, K., Currie, D. & Rochette, J. (2015) 'Advancing marine
918 biodiversity protection through regional fisheries management: A review of bottom fisheries
919 closures in areas beyond national jurisdiction'. *Marine Policy*, 61 pp. 134-148.

920

921

922

923