2019-01

Deeper knowledge of shallow waters: reviewing the invertebrate fauna of southern African temporary wetlands

Bird, MS

<http://hdl.handle.net/10026.1/12484>

10.1007/s10750-018-3772-z.

Hydrobiologia

Springer Verlag

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Deeper knowledge of shallow waters: reviewing the invertebrate fauna of southern African temporary wetlands

Matthew S. Bird,^{1*} Musa C. Mlambo,² Ryan J. Wasserman,^{3,4} Tatenda Dalu,⁵ Alexandra J. Holland,² Jenny A. Day,⁶ Martin H. Villet,⁷ David T. Bilton,⁸ Helen M. Barber-James,^{2,7} Luc Brendonck^{9,10}

¹ Department of Zoology, University of Johannesburg, Auckland Park, Johannesburg 2006, South Africa

² Department of Freshwater Invertebrates, Albany Museum[†], Grahamstown 6140, South Africa

³ South African Institute for Aquatic Biodiversity, Grahamstown 6140, South Africa

⁴ Department of Biological Sciences and Biotechnology, Botswana International University of Science and Technology, P. Bag 16, Palapye, Botswana

⁵ Department of Ecology and Resource Management, University of Venda, Thohoyandou 0950, South Africa

⁶ Institute of Water Studies, Department of Earth Sciences, University of the Western Cape, Bellville 7535, Cape Town

⁷ Department of Zoology and Entomology, Rhodes University, Grahamstown 6140, South Africa

⁸ Marine Biology and Ecology Research Centre, School of Biological and Marine Sciences, University of Plymouth, Plymouth PL4 8AA, UK

⁹ Animal Ecology, Global Change and Sustainable Development, KU Leuven, Leuven B-3000, Belgium

¹⁰ Research Unit for Environmental Sciences and Management, Potchefstroom Campus, North-West University, Potchefstroom 2520, South Africa

* Corresponding author email: mattsbird@gmail.com

[†] A Rhodes University affiliated institution

Abstract

Temporary lentic wetlands are becoming increasingly recognised for their collective role in contributing to biodiversity at the landscape scale. In southern Africa, a region with a high density of such wetlands, information characterising the fauna of these systems is disparate and often obscurely published. Here we provide a collation and synthesis of published research on the aquatic invertebrate fauna inhabiting temporary lentic wetlands of the region. We expose the poor taxonomic knowledge of most groups, which makes it difficult to comment on patterns of richness and endemism. Only a few groups (e.g. large branchiopods, ostracods, copepods and cladocerans) appear to reach higher richness and/or endemism in temporary wetlands compared to their permanent wetland counterparts. IUCN Red List information is lacking for most taxa, thus making it difficult to comment on the conservation status of much of the invertebrate fauna. However, except for a few specialist groups, many of the taxa inhabiting these environments appear to be habitat generalists that opportunistically exploit these waterbodies and this is hypothesized as one of the reasons why endemism appears to be low for most taxa. Given that taxonomy underpins ecology, the urgent need for more foundational taxonomic work on these systems becomes glaringly apparent.

Keywords

aquatic invertebrates; wetland invertebrates; ephemeral wetlands; temporary ponds; ephemeral ponds; pans; southern Africa; African wetlands

Introduction

Invertebrates dominate the faunas of temporary wetlands worldwide, in terms of species diversity, abundance and animal biomass (Williams, 2006). Insects and crustaceans, in particular, are ubiquitous in such systems. Temporary wetlands house a suite of uniquely specialized invertebrates (Wiggins et al., 1980; Collinson et al., 1995; Brucet et al., 2005; Escalera-Vázquez & Zambrano, 2010; Strachan et al., 2015) which are particularly vulnerable to habitat and hydrological modifications (Calhoun et al., 2017; Dalu et al., 2017a). The southern African region, with its distinct climatic and geomorphic features, has one of the highest densities of temporary depression wetlands (known locally as ‘pans’) worldwide (Goudie & Wells, 1995). Reviews of the invertebrate fauna of temporary wetlands

have been conducted for a number of regions globally (e.g. Batzer et al., 1999; Batzer & Boix, 2016), but not for southern Africa; as such, this study aims to fill the knowledge gap. Southern Africa is generally defined as the area south of the Kunene River in the west and the Zambezi River in the east (including the countries of Botswana, Lesotho, Namibia, South Africa, Swaziland, Zimbabwe, and southern parts of Angola and Mozambique). These boundaries are referred to as ‘the region’ or ‘the southern African region’ throughout this review.

Providing a checklist of all the species within each invertebrate group known from temporary wetlands in the region is beyond the scope of this review and in many cases insufficient information is available. For certain groups, we summarise reported checklists and highlight gaps in taxonomic and ecological understanding. We review taxonomic and biodiversity studies pertaining to freshwater invertebrates in temporary lentic wetlands in the region, group-by-group. Our primary aim is to provide an overview of available information (taxonomic, faunistic and ecological) for each of the major invertebrate taxa found in temporary lentic wetlands in the region, highlighting major gaps in knowledge. The key adaptations that each invertebrate taxon has evolved to cope with desiccation are also briefly covered where such information exists. For general information on wetland invertebrate adaptations to drought, see Wiggins et al. (1980), Brendonck & De Meester (2003), Williams (2006), Strachan et al. (2015) and Batzer & Boix (2016).

Methods

A comprehensive review of all published material in peer-reviewed journals for the period 1960 to 2016 was conducted using a combination of the following terms: ‘temporary’, ‘seasonal’, ‘ephemeral’, ‘vernal’, ‘endorheic’, ‘rock’, ‘rain’, ‘pond’, ‘pool’, ‘pan’, ‘tarn’, ‘depression’, ‘southern Africa’, plus all the individual countries within it, and ‘invertebrates’, plus all the major taxa within. References to other important studies cited in the published material above were also referred to and in turn these were also searched thoroughly for further important cited work. In this way, the relevant literature has been comprehensively surveyed. Only those papers that tackled invertebrate populations or communities from wetland systems that temporarily dry up were included in the final list used for this review. However, in certain cases, there was no mention of whether the wetland was temporarily inundated or not. In such cases it was necessary to pay careful attention to the mention of or

inference to flagship taxa (sensu De Roeck et al., 2007) or the adaptive traits to drying conditions (Lahr, 1997) as this would confirm the temporary nature of the wetland. The review aims to focus on lentic isolated depressional wetlands that are not connected to longitudinal watercourses (i.e. non-riverine systems). However, such specific habitat information is often not available for a given taxon, and thus we include taxa that have simply been recorded as occurring in temporary wetland environments, even if information on the specific nature of these environments is not available for the taxon in question.

The material from peer-reviewed journals was further complemented by consulting key texts in the ‘grey literature’ that contain comprehensive species lists, including the Water Research Commission’s ‘Guides to the Freshwater Invertebrates of Southern Africa’ (relevant chapters are referenced separately throughout the text). For the molluscs, for example, Brown’s (1994) comprehensive account of African freshwater gastropods was used to systematically search each species’ information to establish whether or not it is known to inhabit temporary wetland environments. Similarly, for odonates, which have several species checklist publications covering the various sub-regions of southern Africa (Martens et al., 2003; Kipping, 2010; Samways & Simaika, 2016), it was assessed whether the habitat preferences for the various species have been noted or not, and this information was used to construct a list of temporary wetland-dependent species.

Major invertebrate groups

Porifera

Sponges occur in permanent and temporary freshwater systems worldwide, both lotic and lentic (Manconi & Pronzato, 2008). In southern Africa there are two known families, Spongillidae (five genera, seven species) and Potamolepidae (two genera, species information unavailable, Heeg, 2002a). There appears to be a low richness of sponge taxa in southern African freshwater environments compared to other regions (Manconi & Pronzato, 2008) but, as with many of the lesser-known phyla, additional collecting effort is likely to reveal more species (Heeg, 2002a). *Ephydatia fluviatilis* (Linnaeus, 1758) of the Spongillidae is the only known species that has often been recorded from temporary wetlands in the region, usually found on the stems of emergent macrophytes. This species appears to be distributed throughout South Africa (SA) and Namibia (Heeg, 2002a). Sponges generally survive

drought periods through formation of gemmules, which are internal buds composed of amoebocytes surrounded by an external protective coat comprising collagen layers embedded with spicules (for review, see Frost, 1991).

Cnidaria

Hydra (class Hydrozoa) is the best-known cnidarian taxon in freshwater wetlands, and is particularly abundant in temporary lentic ponds, where it is represented by solitary polyps (Williams, 2006). In the region, out of the five species of Hydrozoa recorded, only *Limnocnida tanganjicae* (Günther, 1893) is recognised as indigenous to Africa, the rest being cosmopolitan species (Rayner et al., 2002). An invasive freshwater jellyfish, *Craspedacusta sowerbyi* (Lankester, 1880), has been recorded in large numbers of man-made reservoirs (Rayner et al., 2002). Hydras are capable of both asexual and sexual reproduction and are known to survive in desiccated wetlands via the latter technique, whereby fertilized eggs are surrounded by a thecal coat that is resistant to external desiccation and extreme changes in environmental conditions in general (Williams, 2006).

Platyhelminthes (flatworms)

‘Turbellarians’ are common and widespread in southern African temporary wetlands, yet no taxonomic key exists for species or even genera in the region (Appleton, 2002a). Young (1976) recorded 80 valid species of freshwater turbellarians across Africa, but little work has been done since to describe African taxa. Artois et al. (2004) described one new genus and four new species of flatworm from ephemeral rock pools in Botswana. Intensive ecological studies of temporary rock pool invertebrate communities in south-eastern Botswana (Brendonck et al., 2002) found that turbellarians (particularly *Mesostoma* spp.) are key predators of anostracan (fairy shrimp) egg banks in these systems, seriously affecting anostracan population sizes (De Roeck et al., 2005). *Mesostoma* species have even been recorded preying on adult *Branchipodopsis* shrimp in these pools (Brendonck et al., 2000b).

Curtis (1991) reported that only one species of freshwater turbellarian, *Mesostoma brincki* Marcus 1970, had been positively identified from Namibia. In a comprehensive global collation of turbellarian species with Gondwanan affinities, Noreña et al. (2003) listed six freshwater turbellarians (*Xenostenostomum microstomoides* Reisinger, 1976; *Prorhynchus brincki* Marcus, 1955; *Bothrioplana semperi* Braun, 1881; *Mesostoma brincki*; *M. ewerum* Du Bois-Reymond Marcus 1951; *Phaenocora foliacea* (Böhmig, 1914)) from the region, but

did not indicate how many of these occur in temporary wetlands. Houben et al. (2014) reported *P. foliacea* from a temporary pond in Hluhluwe Game Reserve, SA. Tubellarians can cope with the periodic drying up of temporary wetlands by producing cysts, and these can come out of dormancy after 5-7 minutes of inundation (Van Steenkiste et al., 2010).

The other major platyhelminth groups with notable freshwater representatives are the trematodes and cestodes, both of which are parasitic. There is no existing taxonomic key for African trematodes and they are poorly known. Appleton (2002a) treats them according to groups (types) based on morphology of the cercariae. The obvious exception to an otherwise poorly-known fauna is the genus *Schistosoma*, which has seven species implicated in human or bovine bilharzia infection and that uses *Biomphalaria* or *Bulinus* gastropod snails as intermediate hosts (see ‘Molluscs’ below for further review of this topic). False tapeworms (subclass Cotyloda) constitute the group of cestodes that can be found in freshwater environments, and in the region are a common parasite of the platanna frog *Xenopus laevis* (Daudin, 1802), but their free-living larval stages are microscopic and are never seen unless specifically sought (Appleton, 2002a).

Gastrotricha

Gastrotrichs have been recorded from temporary wetlands worldwide, but very little ecological or taxonomic information is available on these organisms in freshwater environments of southern Africa (Williams, 2006). In the review of Heeg (2002b), not a single species description existed for the region. The genera that have been recorded so far in the region are cosmopolitan, being found particularly in Europe but also elsewhere (Heeg, 2002b). One recent exception is the new planktonic gastrotrich species, *Kijanebalola devastiva* Todaro, Perissinotto and Bownes, 2013, described from a shallow pond on the shores of iSimangaliso Wetland Park (KwaZulu-Natal, SA). However, despite the pond being small, it appears to be permanently inundated (Todaro et al., 2013). Nevertheless, gastrotrichs are well adapted to life in temporary wetlands through their production of dormant eggs (Ricci & Balsamo, 2000).

Bryozoa

Bryozoans have been noted as relatively common in freshwater environments in southern Africa (Heeg, 2002c; de Moor & Day, 2013), but remain largely uncollected and unstudied in the region. Heeg (2002c) comments that the bryozoan fauna of Africa appears to contain

176 mostly cosmopolitan species, but lists only four species as recorded from the region
177 (*Plumatella emarginata* (Allman, 1844), *P. repens* (Linnaeus, 1758), *Lophopodella capensis*
178 (Sollas, 1908) and *Hyalinella punctata* (Hancock, 1850)). It is highly likely that bryozoans
179 inhabit temporary wetlands across large areas of the southern African region, given that they
180 are well adapted to survive desiccation. Bryozoans germinate by producing desiccation-
181 resistant statoblasts that have hooks for clinging to large mobile animals such as waterfowl
182 and are thus able to disperse passively over wide geographical areas (Bilton et al., 2001).

183 *Nematoda*

184 Nematodes are a highly diverse group worldwide (conservatively estimated at one million
185 species, Abebe et al., 2008) and there are approximately 150 known species of freshwater
186 nematodes in southern Africa, of which most are likely to be capable of surviving dry periods
187 in moist soil (Heyns, 2002). Thirty-four of the 150 species listed by Heyns (2002) have been
188 recorded in semi-aquatic environments and are capable of surviving near-terrestrial
189 conditions, for example in moist soil during the dry phase of a temporary wetland.
190 Nematodes have commonly been recorded elsewhere as able to enter a quiescent phase
191 without water loss in response to drying in temporary ponds (Womersley & Ching, 1989;
192 Wharton, 2004; Abebe et al., 2006) and thus are not expected to be limited by extreme
193 environments in southern African temporary wetlands.

194 *Nematomorpha*

195 Information on the genera and species of the parasite phylum Nematomorpha (horsehair
196 worms) in southern African water bodies is extremely sparse, but the few records that do
197 exist suggest a wide distribution (Rayner & Appleton, 2002). Corrêa (1951) reported two
198 species of *Prostoma* as widespread in SA. In a review of known African species, Schmidt-
199 Rhaesa & Perissinotto (2016) reported a further six species from SA belonging to the genera
200 *Chordodes*, *Paragordius* and *Beatogordius*. They also described a previously unconfirmed
201 species (*Chordodes ferox* Camerano, 1897) collected from a live praying mantid floating in a
202 river in the Eastern Cape province, SA. Nine species recorded by Corrêa (1951) and Schmidt-
203 Rhaesa & Perissinotto (2016) appear to be all that is known taxonomically of the fauna from
204 the southern African region to date. Nematomorph larvae are dependent on an aquatic
205 medium, where they generally penetrate the bodies of arthropods (particularly orthopterans)
206 and spend the vast majority of their lives growing into large adult worms inside the host

207 body. Thus, nematomorphs possess traits that pre-adapt them to life in temporary wetlands
208 (Williams, 2006) and it is likely that they inhabit such environments in southern Africa.

209 *Rotifera*

210 Microscopic animals belonging to the phylum Rotifera (commonly called ‘wheel animals’)
211 are found in a wide variety of freshwater habitats from large lakes to damp moss and are well
212 adapted to life in temporary wetlands. Rotifers have two major means of dealing with
213 desiccation. Bdelloid rotifers are able to slowly dry out and then return to activity when they
214 are rehydrated, a process known as cryptobiosis (Brain, 2002). Monogonant rotifers instead
215 survive as desiccation-tolerant resting eggs (Ricci & Balsamo, 2000). Both desiccated
216 bdelloids and monogonant eggs are carried by wind and for this reason freshwater rotifers are
217 mostly cosmopolitan, being able to flourish wherever suitable conditions are encountered
218 (Brain, 2002). According to Dumont (1983), African rotifers are particularly cosmopolitan,
219 although no comprehensive species list exists. Brain (2002) provided a list and an
220 identification key for the 47 genera of freshwater rotifers known from southern Africa (8
221 digononts and 39 monogononts), but it is not known how many of these taxa are found in
222 temporary wetland environments. Anusa et al. (2012) reported eight rotifer species
223 (belonging to the genera *Asplanchna*, *Brachionus*, *Conochilus*, *Epiphanes*, *Lepadella*,
224 *Rotaria* and *Synchaeta*) as common inhabitants across a range of hydroperiods in temporary
225 rock pools in northern Zimbabwe. They found that rotifers were more abundant in pools with
226 longer hydroperiods, but were able to withstand the ephemerality of the shallow short-lived
227 pools. Dalu et al. (2016) recorded five rotifer species belonging to four genera (*Brachionus*,
228 *Euchlanis*, *Filinia*, *Pompholyx*) in an ephemeral pond in the Eastern Cape. However, more
229 species are expected to be found with more studies currently underway. Riato et al. (2014)
230 compared the zooplankton communities of temporary and permanent depression wetlands in
231 the Mpumalanga Highveld, SA, and found that *Platyias quadricornis* (Ehrenberg, 1832) only
232 occurred in the temporary wetlands. All nine species reported from the studies of Anusa et al.
233 (2012) and Riato et al. (2014) are widespread and all occur on other continents, thus
234 providing some preliminary indication that the temporary wetland rotifers in the region
235 comprise largely cosmopolitan species.

236 *Tardigrada*

237 Phylum Tardigrada ('waterbears') are tiny coelomate animals (50 µm to 1.2 mm in length)
238 represented by approximately 900 species worldwide, of which only 62 species (representing
239 13 genera) are truly aquatic (Garey et al., 2008). Very little is known about the aquatic
240 tardigrades in the region and there has been no definitive study on the group. They are not
241 commonly encountered by freshwater ecologists and are easily overlooked due to their small
242 size (Rayner, 2002). No taxonomic or ecological information on the aquatic tardigrades of
243 southern Africa is available. There are however a number of studies on the terrestrial and
244 semi-terrestrial tardigrade fauna of mosses, lichens and leaf litter habitats in the region (for
245 reviews see Meyer & Hinton, 2009; Zawierucha et al., 2012). Despite the lack of studies, it is
246 strongly suspected that tardigrades inhabit southern African temporary wetlands, given their
247 exceptional adaptations to desiccation and physico-chemical extremes (reviewed by
248 Williams, 2006; Møbjerg et al., 2011).

249 Tardigrades are able to spend long periods in a desiccated cryptobiotic state, during which
250 time they are unique among metazoans in their ability to withstand extreme physico-chemical
251 fluctuations, tolerating laboratory immersion in liquid helium at -272°C and temperatures as
252 high as 340°C (Williams, 2006). During cryptobiosis, tardigrades are also easily dispersed by
253 wind, as are their eggs. Once their environment is re-flooded with water they rehydrate and
254 become active (Williams, 2006). Incredibly, tardigrades can survive for up to 70 years in the
255 cryptobiotic state (Rayner, 2002). Life spans are more commonly less than one year however
256 (Rayner, 2002).

257 *Annelida*

258 Annelid worms found in temporary wetlands in the region include earthworms (subclass
259 Oligochaeta) and leeches (subclass Hirudinea). Although sabellid polychaetes (class
260 Polychaeta) have been recorded elsewhere in temporary wetlands (Yozzo & Diaz, 1999), they
261 have not been reported in southern African. *Ceratonereis keiskama* Day, 1953 appears to be
262 the only polychaete worm recorded in entirely freshwater environments in SA, where it has
263 been recorded from permanent freshwater lakes that are estuarine relicts, being particularly
264 abundant in Lake Sibaya in KwaZulu-Natal (Day & Day, 2002).

265 Approximately one-third of the almost 5 000 valid species of oligochaetes worldwide are
266 aquatic, of which approximately 1 100 are found in freshwater environments (Martin et al.,

2008). About 50 microdrile and 20 megadrile taxa have been recorded from southern Africa (van Hoven & Day, 2002). Whilst most of the microdrile species are cosmopolitan, the megadriles are mostly African endemics (van Hoven & Day, 2002). None of the taxonomic works specifically report on which oligochaete taxa are found in southern African temporary wetlands, but the megadrile earthworm *Lumbriculus variegatus* (Müller, 1774) (cosmopolitan species) and microdriles belonging to *Nais* (Naididae) have been reported in faunal lists from ecological studies of temporary wetlands in the region (e.g. Bird & Day, 2016). Oligochaete specimens from southern African wetland studies are most commonly reported at the ordinal level as ‘Oligochaeta’ (e.g. Anusa et al., 2012; Ferreira et al., 2012). Oligochaetes can survive desiccation in temporary wetland environments by forming dehydration-resistant cysts, empirically documented for lumbriculids (Anlauf, 1990) and naidids (Montalto & Marchese, 2005).

Leeches are found in most freshwater habitats, although they are sparse in fast-flowing water or wetlands that are highly ephemeral (Oosthuizen & Siddall, 2002). Leeches are, however, well adapted to live in temporary waterbodies, mostly due to their ability to disperse via mobile vertebrates such as waterfowl (Davies et al., 1982; Williams, 2006) or attached to large invertebrates (Oosthuizen & Siddall, 2002). Some leeches are capable of surviving droughts as adults in mucus-lined cysts (Williams, 2006). The excellent dispersal ability of leeches results in their distributions in temporary wetlands in other regions of the world being well known (Williams, 2006). There are approximately 50 species of leeches in southern Africa (Oosthuizen & Siddall, 2002), but it is unknown how many can tolerate living in temporary wetlands. Bird & Day (2016) reported two species, the African leech *Helobdella conifera* (Moore, 1933) and the globally widespread *H. stagnalis* (Linnaeus, 1758) from temporary depression wetlands in Cape Town, SA. Oosthuizen & Siddall (2002) have recorded *Placobdelloides multistriata* (Jahansson, 1909) from temporary wetlands throughout southern Africa (the species is distributed across Africa), although they remark that the leeches usually enter small temporary pools inadvertently while attached to water scorpions and other hemipterans as transport hosts. All three of the above-mentioned species occurring in temporary wetlands thus have wide geographic distributions.

Mollusca

Approximately 111 species of gastropod and 26 bivalves have been reported from southern Africa, which includes 10 alien gastropod species (Appleton, 2002b). The region has very

low freshwater molluscan endemicity at about 12.3% (Appleton, 2002b). According to Brown's (1994) habitat records, the following species are frequently found in temporary wetlands: *Pila occidentalis* (Mousson, 1887); *Lanistes ovum* Troschel, 1845; *Tomichia ventricosa* (Reeve, 1842); *Ceratophallus natalensis* (Krauss, 1848); *Lymnaea natalensis* Krauss, 1848; *L. columella* (introduced from North America, now widespread in Africa); *Physa acuta* Draparnaud, 1805 (introduced from Europe, now widespread in Africa); *Biomphalaria pfeifferi* (Krauss, 1848); *Bulinus globosus* (Morelet, 1866) *B. forskalii* (Ehrenberg, 1831); *B. reticulatus* (Mandahl-Barth, 1954); *B. tropicus* (Krauss, 1848); *B. scalaris* (Dunker, 1845); and *Aplexa marmorata* (Guilding, 1828). *Aplexa marmorata* is an invasive species from South America, which has recently been recorded from temporary wetlands as far south as Cape Town by Mlambo et al. (2011).

The genus *Ferrissia* are freshwater limpets that are found in varied habitats including seasonal pools throughout Africa (Brown, 1994). Although several studies have recorded *Ferrissia* at the generic level in South African temporary wetlands (Mlambo et al., 2011; Bird et al., 2013; Bird & Day, 2016), species level information is unavailable. Members of this genus are able to partly close the shell using a septum, which facilitates aestivation during the dry phase of a temporary wetland (Brown, 1994).

Of the African gastropods, the genus *Bulinus* excels in temporary wetland environments (Vera et al., 1995). For example, *B. depressus* and *B. scalaris* were the species most commonly found in isolated seasonal pools in Moremi Game Reserve, Botswana (Appleton et al., 2003). Similarly, *B. forskalii* was the only mollusc collected together with branchiopods in rain pools on the Makathini Flats (Hamer, 1989). The success of *Bulinus* snails can, at least in part, be put down to their ability to aestivate for up to six months and then exploit nutrient-rich temporary water bodies upon re-inundation by breeding rapidly (re-selected growth) during the short breeding season (Brown, 1994).

Bulinus and *Biomphalaria* are the two best-studied gastropod genera in southern Africa because they are the aquatic intermediate hosts of human and bovine schistosomiasis (Rubaba et al., 2016). Although species of *Biomphalaria* commonly occur in temporary habitats in the Neotropical region (Paraense, 1996), this is rare for the African counterparts. *Biomphalaria* species in the Neotropics often have lamellae in their shells' apertures that are thought to play a role in tolerating desiccation. In a southern African context, only one specimen of *B. pfeifferi* is known to have apertural lamellae (Appleton CC, *personal communication*).

Of the freshwater bivalves, members belonging to the family Sphaeriidae commonly inhabit temporary wetlands (Williams, 2006). The three genera of Sphaeriidae (*Sphaerium*, *Pisidium* and *Eupera*) present in southern Africa are in need of revision (Appleton, 2002b) and information on which sphaeriid species are known to inhabit temporary wetlands in the region appears to be lacking. Sphaeriids have been shown elsewhere to deal with desiccation by burrowing into the substrates of drying wetlands (McKee & Mackie, 1980). Most species require some residual moisture in the substrate to survive the dry season (at least 15% moisture for two Canadian sphaeriid species, McKee & Mackie, 1980) and thus are not likely to be encountered in highly ephemeral wetlands. Sphaeriids have effective dispersal mechanisms (e.g. clipping onto the limbs of mobile invertebrates and waterfowl) and are self-fertilizing hermaphrodites, adaptations that further allow them to exploit temporary environments (Williams, 2006).

Crustaceans: Malacostraca

Most freshwater-dwelling members of the class Malacostraca (i.e. Amphipoda, Isopoda and Decapoda) lack physical adaptations for diapause or dispersal. Thus, they are not commonly associated with temporary wetlands but, due mostly to behavioural adaptations, certain species are able to persist in temporary waters (Williams, 2006). Amphipods appear to be very scarce in southern African temporary wetlands and virtually no records exist of their occurrence in these systems. Mlambo et al. (2011) and Bird & Day (2016) both recorded populations of the Gondwanan relictual species *Paramelita capensis* (Barnard 1916) and *P. pinnicornis* Stewart and Griffiths 1992 in small temporary wetlands of the Kenilworth Racecourse Conservation Area in Cape Town. However, studies of how these amphipods survive the summer dry phase have not been undertaken. A study of the North American amphipod *Crangonyx pseudogracilis* Bousfield, 1958 provides evidence that this epigean species is able to inhabit parts of the water table during the dry phase of a temporary wetland and this mechanism explains both its ability to persist and its ubiquity across a network of nearby wetlands (Harris et al., 2002). The authors suggest that such subterranean explorations by epigean amphipods may be a key mechanism for amphipod survival in temporary wetlands in other parts of the world. Another interesting behavioural adaptation in amphipods is reported from a congeneric North American amphipod, *C. antennatus* Packard, 1881, which survives the desiccated phase of mud-bottom pools by digging a burrow into the mud

that is then blocked off from the outside with an air bubble (Holsinger & Dickson, 1977). The burrow itself remains hydrated and provides a temporary aquatic refuge until the pool refills.

True crabs (Brachyura) also deal with desiccation by burrowing into the substrate. At least four southern African species (namely *Potamonautes lividus* Gouws, Stewart and Reavell, 2001, *P. flavusjo* Daniels, Phiri and Bayliss, 2014, *P. isimangaliso* Peer & Gouws, 2015, and *P. calcaratus* (Gordon, 1929)) exhibit a vaulted carapace, which is an adaptation to a semi-terrestrial lifestyle away from permanent waters (Daniels et al., 2014). All of these species may appear superficially similar morphologically, but phylogenetically they are very distinct (Daniels et al., 2014; Peer et al., 2015). From north-eastern KwaZulu-Natal, *Potamonautes lividus* and *P. isimangaliso* are documented to live in close association with highly ephemeral depression wetlands (Peer et al., 2015). These crabs occupy 0.3-0.5 m deep burrows within the wetland substrate that remain moist throughout the dry season. Brachyurans associated with temporary waterbodies worldwide have often also evolved physiological air-breathing adaptations such as the ability to efficiently draw oxygen from moist air within burrows (Cumberlidge, 2009), and this appears to be the case for *P. isimangaliso* (Peer et al., 2015). Similarly, *P. calcaratus* displays the same ecological habits as *P. isimangaliso*, surviving in burrows around ephemeral pools in the KNP (Daniels et al., 2014). Another congeneric, *P. flavusjo*, burrows into peat soils in Highveld wetlands in Mpumalanga province where it remains submerged in water by following the vertical movement of the water table (Daniels et al., 2014).

The only freshwater isopods known to inhabit temporary wetlands in the region belong to the genus *Mesamphisopus* (Gouws et al., 2005). Barnard (1926) observed that *Mesamphisopus abbreviatus* (Barnard, 1927) burrows into muddy substrata in small mountain streams that seasonally run dry. It aestivates in these burrows by lowering its metabolism and breathing the moist air in the burrows, in a similar fashion to *P. isimangaliso*. *Mesamphisopus* has been reported from temporary depression wetlands in the Western Cape province (Mlambo et al., 2011; Bird et al., 2013).

To our knowledge, and based on extensive literature searching, there are no records of prawns or shrimps in the freshwater temporary wetlands of southern Africa (indigenous freshwater crayfish do not occur in Africa).

392 *Crustaceans: Large branchiopods*

393 The class Branchiopoda, as a matter of convenience rather than taxonomic distinction, are
394 usually treated as two groups: the large branchiopods (fairy shrimps, tadpole shrimps and
395 clam shrimps) and the cladocerans (water fleas). Following suit, we address the cladocerans
396 in a separate section together with their microcrustacean counterparts, the ostracods and
397 copepods.

398 The large branchiopods are the enigmatic flagship invertebrates of temporary wetlands
399 worldwide (Brendonck et al., 2008). They are specialists of lentic environments that
400 incorporate a dry phase, although in some cases they do occur in fishless semi-permanent
401 waterbodies (Ferreira et al., 2011). Their adaptations to ephemeral waterbodies have led to
402 their prevalence in arid and semi-arid regions, where such waterbodies predominate. This is
403 no exception in the largely semi-arid subcontinent of southern Africa, where they are
404 widespread and diverse. The large branchiopod fauna is well studied in the region compared
405 to most of the aquatic invertebrate groups so far covered in this review and to date
406 approximately 66 species have been recorded (Tuytens et al., 2015).

407 Large branchiopods generally exhibit a lifestyle of rapid growth and maturation, followed
408 by the production of drought-resistant eggs that can lie dormant in dry sediment under harsh
409 conditions for decades and perhaps even centuries (Brendonck et al., 2008). Generally
410 speaking, populations adopt a ‘bet-hedging’ strategy in that not all eggs hatch with the first
411 onset of inundation, but rather some eggs remain dormant and only hatch after subsequent
412 inundations so as to ensure survival of the population in a wetland that dries before a cohort
413 reaches maturity (Simovich & Hathaway, 1997; Brendonck et al., 1998; Brendonck et al.,
414 2008). Eggs are generally dispersed passively to other waterbodies via wind (Brendonck &
415 Riddoch, 1999; Vanschoenwinkel et al., 2009), ingestion by predators, which transport the
416 eggs in their gut (Rogers, 2014, 2015), or by attachment to mobile fauna (Vanschoenwinkel
417 et al., 2011). More specific aspects of the adaptations of large branchiopods to temporary
418 wetlands environments, including their biology and reproduction, are covered elsewhere (for
419 reviews see Belk et al., 1991; Brendonck, 1996; Brendonck et al., 1998; Dumont & Negrea,
420 2002; Rogers, 2009).

421 Various aspects of the ecology of southern African large branchiopods (including
422 metacommunity dynamics, competition and predation, adaptations to drying and local abiotic

conditions, human impacts, management and conservation) have received considerable research attention due largely to a series of studies of rock pools in the eastern Free State province (Vanschoenwinkel et al., 2007; Vanschoenwinkel et al., 2009; Vanschoenwinkel et al., 2010a; Vanschoenwinkel et al., 2010b; Pinceel et al., 2013; Vanschoenwinkel et al., 2013; Tuytens et al., 2014) and south-eastern Botswana (Brendonck et al., 1998; Brendonck et al., 2000a; Brendonck et al., 2000b; De Roeck et al., 2005; Jocqué et al., 2006; Jocqué et al., 2010). Recently, the large branchiopod fauna of soft-bottomed wetlands and rock pools have received ecological attention in Zimbabwe (Nhiwatiwa et al., 2011; Anusa et al., 2012; Nhiwatiwa et al., 2014; Tuytens et al., 2015; Nhiwatiwa et al., 2017a; Nhiwatiwa & Dalu, 2017; Nhiwatiwa et al., 2017b) and the Western Cape province (De Roeck et al., 2007; De Roeck et al., 2010; Mlambo et al., 2011). A scattering of studies exists from various other localities, including Namibia (Day, 1990; Curtis, 1991), Botswana (Brendonck & Riddoch, 1997), the Drakensberg (Hamer & Martens, 1998), and soft-bottomed wetlands in the Free State (Seaman et al., 1991; Meintjes, 1996), Mpumalanga (Ferreira et al., 2011, 2012), North-West (de Necker et al., 2016), Northern Cape (Hamer & Rayner, 1996) and KwaZulu-Natal (Hamer & Appleton, 1991) provinces of SA. As far as we are aware, no ecological studies on large branchiopods have been recorded from Mozambique.

The best-known of the large branchiopods is the order Anostraca (fairy shrimps). Early taxonomic descriptions in the southern African region were pioneered by G.O. Sars and E. Daday, but largest contribution to anostracans systematics was made by K.H. Barnard of the South African Museum during the 1930s-1940s. During the 1990s, various workers updated diversity and distribution data for the anostraca of the region. Hamer et al. (1994) and Brendonck & Coomans (1994) reviewed the Streptocephalidae south of the Zambezi and Kunene rivers and reported a total of 16 species. Brendonck (1995) added a new genus and species *Rhinobrachipus martensi* Brendonck, 1995 (Branchipodidae) to the southern African list of anostraca, which was collected from a shallow depression wetland in the Thomas Baines Nature Reserve in the Eastern Cape province. Hamer (1999) provided a taxonomic key for identification of the 46 southern African species of anostracans (six genera and four families) known at the time, all of which are known to occur in temporary freshwater wetlands with the exception of members of the genera *Artemia* and *Parartemia*, which are adapted to hypersaline salt lakes and salt pans. Nhiwatiwa et al. (2017b) recently described a new species, *Streptocephalus sangoensis* Nhiwatiwa, Dalu & Brendonck, 2017 in the south-eastern Lowveld of Zimbabwe, which was formerly published as *Streptocephalus*

cf. *bidentatus*, indicating that more research is still required to prove valid species status and identify more taxa.

Brendonck et al. (2000b) provided a focussed review on the genus *Branchipodopsis*, the members of which are shown to be specialists of transparent, very fresh ($<50 \mu\text{S cm}^{-1}$) and often highly ephemeral rock pools in the region. They described *Branchipodopsis* species as the ‘record holders’ of an ephemeral lifestyle in these rock pools, where populations are able to grow to maturity within the first week after inundation. Their long-distance dispersal ability by wind appears to be rare and rather they disperse to other nearby pools via overflows. This is suggested as a potential factor behind the high levels of endemism of *Branchipodopsis* species within the region. Since 2000, two new anostracan species have been described from KwaZulu-Natal province belonging to the genus *Metabbranchipus* (*M. rubra* Rogers and Hamer, 2012 and *M. prodigiosus* Rogers and Hamer, 2012) of the family Branchipodidae.

Otherwise, the only recent notable taxonomic work in southern Africa concerning the anostracans is for the genus *Artemia* (Artemiidae, commonly known as brine shrimp). *Artemia* has been in a state of taxonomic confusion for quite some time (Hamer, 1999). The genus is cosmopolitan and restricted to salt lakes and salt pans with salinity levels above approximately 40 g L^{-1} , where fish and many predatory invertebrates are absent (Kaiser et al., 2006). In southern Africa, populations seem to be a mix of bisexual and parthenogenetic forms (Kaiser et al., 2006). The sexually reproducing species cannot be distinguished by gross morphological characters and require genetic analyses to separate species. Since such work had not been conducted in the region until recently, members of the genus have been referred to at generic level. However, Baxevanis et al. (2014) attempted the genetic identification of *Artemia* species in SA. Phylogenetic and network analyses revealed that three out of the seven studied populations belong to the invasive *A. franciscana* Kellog, 1906, while four belong to *A. salina* (Linnaeus, 1758). This is the first confirmed report of the invasive *A. franciscana* in the region and highlights concern regarding its invasive ability, as demonstrated by its complete replacement of the native *A. salina* population at the Veldrift Saltworks on the west coast of SA. Given that Baxevanis et al. (2014) only investigated seven populations, the taxonomic status of *Artemia* populations in the broader southern African region stills requires investigation.

Brendonck & Riddoch (1997) conducted a comprehensive survey of the anostracan fauna of Botswana, sampling from approximately 200 localities around the country, collecting both live specimens and mud samples containing eggs. They reported 14 anostracan species, with only one (*Branchipodopsis kalaharensis* Daday, 1910) being endemic to Botswana. They attribute the higher species richness (33) and endemism (64%) in SA to greater habitable surface area, topographic and environmental heterogeneity, and research effort. Hamer & Brendonck (1997) expanded on the work of Brendonck & Riddoch (1997) by establishing a species-level checklist of the anostracan fauna for the whole region, providing a detailed distribution of each species. They divided the anostracan fauna into ten biogeographic categories, shaped largely by climatic factors: (i) widespread species occurring across southern Africa; (ii) eastern escarpment species occurring at high altitude within the Drakensberg; (iii) arid south-west species occurring in the arid and semi-arid regions of Namibia and the Karoo; (iv) tropical/subtropical species occurring on the low-lying eastern margin of southern Africa; (v) southern savannah species; (vi) Highveld species occurring in summer rainfall areas on the plateau at altitudes of 1000 – 1500 m; (vii) Cape east coast species occurring in winter-rainfall areas of the southern and south-eastern Cape margins; (viii) Cape west coast species occurring from Cape Town northwards along the west coast; (ix) Eastern Cape inland species (two species), of which *Streptocephalus dregei* Sars, 1899 occurs in the more arid lowland areas towards the south, whilst *S. spinicaudatus* Hamer and Appleton, 1993 occurs more northwards at higher altitude and with higher annual rainfall; and (x) Zimbabwean species that have been collected mainly from Zimbabwe and also Zambia, but not further south.

The order Notostraca (tadpole shrimps) are branchiopod crustaceans that are known from as far back as the Triassic Period, earning them the status of ‘living fossils’ (Fryer, 1988; Vanschoenwinkel et al., 2012), although recently this concept for the group has been challenged (Mathers et al., 2013). Rayner (1999) provides an overview of the morphology, habitat preferences, feeding, reproduction and identification of southern African notostracans. Two species of *Triops* have been recorded from southern Africa, *T. granarius* (Lucas, 1864) and *T. cancriformis* (Bosc, 1801), although the latter is rare and has only been recorded from Ovamboland (northern Namibia) and the Kalahari (Hamer & Rayner, 1995). *T. granarius* is common and widespread in the drier areas of southern Africa (<500 mm mean annual rainfall), preferring muddy temporary pools seldom exceeding one hectare in size (for more detailed distribution records see Hamer & Rayner, 1995). *Triops* taxonomy in southern Africa

(and worldwide for that matter) is not without controversy. For instance, Korn & Hundsdoerfer (2006) used ribosomal DNA markers to investigate genetic distinctiveness of *T. granarius* populations from Tunisia, Namibia and Japan, and their results strongly suggest that the three populations belong to different, possibly cryptic species. Thus, further molecular phylogenetic work on African *Triops* populations is required to more clearly resolve cryptic species complexes.

The third major group of large branchiopods frequenting temporary wetlands are the clam shrimps, although the colloquial term is used here solely for descriptive convenience. The clam shrimps consist of the order Laevicaudata (smooth clam shrimp, one family with three genera and 39 species) and, in the order Diplostraca, two suborders: Spinicaudata and Cyclestherida, with approximately 116 known species (four families and 16 genera) worldwide (Brendonck et al., 2008; Rogers et al., 2012). All three groups are primitive and known as far back as the Devonian (Gueriau et al., 2016). Despite being abundant and widespread in arid-zone temporary wetlands of the region (and worldwide), the taxonomy of the clam shrimps has been poorly studied. The systematics has been problematic and complicated by morphological plasticity in the fine details, despite generally uniform gross morphology (Rogers et al., 2012). The difficulty in finding diagnostic morphological characters appears to be a prime reason why virtually no taxonomists have tackled the systematics of southern African clam shrimps since the studies of G.O. Sars, R. Gurney, G.S. Brady, E. Daday, K.H. Barnard and V. Brehm.

Brendonck (1999) provides a taxonomic key to distinguish the southern African species of clam shrimps, although only genus-level distinction is provided in some cases (e.g. *Cyzicus* Audouin, 1837), due to difficult taxonomy. Brendonck (1999) also provides distribution records for the 18 species known at the time of his assessment. The only notable taxonomic work that includes southern African clam shrimps in recent times is a revision of the worldwide genera of the Limnadiidae (Spinicaudata) by Rogers et al. (2012), who combined morphological and molecular information to show the presence of eight well-defined genera, including two new genera (*Afrolimnadia* Rogers, Rabet & Weeks, 2012 and *Calalimnadia* Rogers, Rabet & Weeks, 2012). South African material was used for the description of *Afrolimnadia alluaudi* (Daday, 1926). *Afrolimnadia* was subsequently renamed *Gondwanalimnadia* Rogers, Rabet & Weeks, 2016 because it is a junior homonym of

551 *Afrolimnadia* Tasch, 1987, an fossil genus of spinicaudatan clam shrimps (Rogers et al.
552 2016).

553 *Crustaceans: Ostracoda*

554 Southern Africa has a diverse ostracod fauna. Martens (2001) reported about 155 species (ca.
555 55 genera) belonging to the class Ostracoda from the region, but remarked that dozens of new
556 species await description. Since this publication, several new species and genera have also
557 been added to the regional list (Martens, 2003, 2007; Jocqué et al., 2010), including the re-
558 discovery of two species of giant ostracod (*Liocypris grandis* Sars, 1924 and *Afrocypris*
559 *barnardi* Sars, 1924) that were thought to be extinct (Martens, 2003; Matzke-Karasz &
560 Martens, 2007), both from temporary wetland habitats. The most striking feature of the
561 ostracod fauna of the region is the high endemism of the Western and Eastern Cape
562 provinces, which although possessing few endemic genera, harbour many endemic species
563 (de Moor & Day, 2013). The northern and central parts of Namibia and Botswana and the
564 southern part of Zimbabwe form another belt of high endemism, with generic endemism in
565 these areas being comparable to the exceptionally rich and endemic ostracod fauna of Lake
566 Tanganyika (Jocqué et al., 2010). The ostracod fauna of southern Africa exhibits its highest
567 endemism and diversity in temporary wetland environments where few cosmopolitan species
568 occur (Martens, 2001). Ostracods are mostly benthic scavengers (although some forms are
569 pelagic) and play an important role in temporary wetland ecosystems as detritivores
570 (Martens, 2001). The success of ostracods in temporary wetlands is largely attributed to their
571 production of highly desiccation-resistant eggs, but also their ability to reproduce
572 parthenogenetically and the ease with which their eggs are passively dispersed to other
573 waterbodies via wind dispersal or attached to other animals (for review see Horne & Martens,
574 1998).

575 Martens (2001) provides a taxonomic key for the genera (and species for some recently-
576 revised genera) of southern African freshwater ostracods, many of which are found in
577 temporary wetland environments. Following this reference, the genera that have been
578 unequivocally recorded from temporary wetlands are: *Cypricercus*, *Eundacypris*,
579 *Gomphocythere*, *Globocypris*, *Korannacythere*, *Leucocythere*, *Megalocypris*,
580 *Ovambocythere*, *Physocypris*, *Pseudocypris*, *Ramotha*, *Sarscypridopsis*, *Sclerocypris*,
581 *Zonocypris*. Jocqué et al. (2006) reported *Potamocypris* and *Strandesia* from temporary rock
582 pools in Botswana, while *Amphibolocypris* (*A. arida* and *A. exigua*) were recorded from

temporay rock pools in Botswana and a temporary soft-bottomed wetland in KNP,
respectively (Jocqué et al., 2010).

In the Western Cape province, Martens (2007) recorded the genus *Mnementh* from temporary wetlands; and Martens et al. (1996) reported *Zonocypris cordata* Sars, 1924 and *Physocypris capensis* Sars, 1896, which are both found in temporary and permanent wetlands. Additionally, Bird & Day (2016) recorded the following genera *Chrissia*; *Sarscypridopsis*; and *Gomphocythere* from temporary depression wetlands within the Kenilworth Racecourse Conservation Area, Cape Town. This also included the following species: *Cypretta turgida* Sars, 1896, *Cypricercus episphaena* (Müller, 1908), *Cypridopsis vidua* (Müller, 1908), *Paracyprretta acanthifera* Sars, 1924, *P. capensis*, *Ramotha capensis* (Müller, 1908), and *Zonocypris cordata* Sars, 1924.

Reviewing the ostracod fauna of KwaZulu-Natal, Martens et al. (1998) report that most of the diversity in lowland temporary wetlands of the province is in the genera *Heterocypris* and *Cypricercus*, but diversity and endemcity in these habitats is generally low. The higher altitude temporary rock pools of the Drakensberg have high diversity and endemcity in the genera *Sarscypridopsis* and *Heterocypris*. However, most notable of the Drakensberg pools is the diversity within the family Limnocytheridae that is unmatched anywhere else in the world.

Crustaceans: Copepoda

Copepods are one of the most diverse crustacean groups, including over 14,000 known species of both free-living and symbiotic forms. Most free-living copepods are marine, but approximately 2,800 species have colonized a wide variety of freshwater habitats (Boxshall & Defaye, 2008). Copepods in freshwaters represent a pivotal component of the food webs as part of the zooplankton and benthic/littoral communities worldwide. Up to 20 genera and 51 species of freshwater copepods have been recorded from the region (Rayner, 2001). Most of the information available for the region concerns the calanoid copepods (order Calanoida) and there has been virtually no recent taxonomic research on the cyclopoids or harpacticoids. Among the calanoids, only members of the exclusively freshwater family Diaptomidae have been recorded in the region (Rayner & Heeg, 1994). The subfamily Diaptominae, containing about 410 species, is represented in the region by the genera *Tropodiaptomus* and *Thermodiaptomus*, both of which occur in more permanent waters and are not considered

further here. The diaptomid subfamily Paradiaptominae currently consists of 25 species (Suárez-Morales et al., 2015), most of which are endemic to Africa and adapted to life in temporary wetlands (Rayner, 2001). This largely endemic group includes a genus, *Lovenula*, that comprises extremely large individuals (de Moor & Day, 2013). Rayner (2001) provides a taxonomic key to the species of Diaptomidae in the region, along with cursory notes on the cyclopoid and harpacticoid taxa that have been described or recorded from the region. The following species in the subfamily Paradiaptominae have thus far been recorded from temporary wetlands in the region (for distributional information, see Day, 1990; Rayner, 1998; Rayner, 2000, 2001; Suárez-Morales & Rayner, 2004; Suárez-Morales et al., 2015): *Lovenula falcifera* (Lovén 1845), *L. africana* (Daday, 1908), *L. simplex* Kiefer, 1929, *L. raynerae* Suárez-Morales, Wasserman & Dalu 2015, *Paradiaptomus lamellatus* Sars, 1895, *P. schultzei* van Douwe, 1912, *P. similis* van Douwe, 1912, *P. peninsularis* Rayner, 1999, *P. hameri* Rayner, 1999, *Metadiaptomus capensis* (Sars, 1907), *M. purcelli* (Sars, 1907), and *M. meridianus* (van Douwe, 1912).

The cyclopoids (order Cyclopoida), on the other hand, are represented in freshwaters by the widespread family Cyclopidae; in Africa this group is represented by 167 species, a figure that resembles that of the Neotropical region (174 species), but is about half the number found in the Palaearctic region (337 species) (Boxshall & Defaye, 2008). In southern Africa, records of cyclopids are scarce and mostly consist of cosmopolitan species. This is probably the result of the 19th Century species concepts and delimitations that were extensively used and applied by European researchers who named species from different geographic regions on the basis of their resemblance with European species. Revisionary studies have proved that records of some of these “cosmopolitan” nominal species (i.e. *Acanthocyclops vernalis* (Fischer, 1853) and *A. robustus* (Sars, 1863)) represent species complexes containing undescribed taxa. The South African cyclopoid copepod fauna certainly deserves further examination with recent taxonomic standards. The following cyclopoids have been recorded in temporary wetlands in the region (for distributional information, see Day, 1990; Rayner, 2001): *A. vernalis*, *Mesocyclops major* Sars, 1927, *Thermocyclops oblongatus* Sars, 1927, *T. macracanthus* Kiefer, 1929, *T. schuurmanae* Kiefer, 1928, *Microcyclops crassipes* (Sars 1927), *M. inopinatus* (Sars, 1927), *Eucyclops* (*Afroscyclops*) *gibsoni* (Brady, 1904), *E. sublaevis* (Sars, 1927), *E. serrulatus* (Fischer, 1851), *Tropocyclops prasinus* (Fischer, 1860), *Ectocyclops phaleratus* (Koch, 1838), *Paracyclops poppei* (Rehberg, 1880), and *P. fimbriatus* (Fischer, 1853).

647 Information on the African freshwater harpacticoids (order Harpacticoida) reveals a
648 moderate diversity (108 species), which is comparable to that of India and Thailand. The
649 most diverse harpacticoid family, the Canthocamptidae is represented by only 46 species in
650 Africa, whereas more than 320 occur in the Palaearctic region (Boxshall & Defaye, 2008).
651 Records of Harpacticoida in southern Africa are extremely scarce, let alone which taxa are
652 known from temporary waters. *Nitokra dubia* Sars, 1927 and *Elaphoidella bidens coronata*
653 Kiefer, 1934 have both been recorded from temporary wetlands of the Cape Flats near Cape
654 Town, SA (Rayner, 2001). Vanschoenwinkel et al. (2008) recorded *Bryocamptus* sp. from
655 temporary rock pools of the eastern Free State province. Rayner (2001) lists *Attheyella*
656 *natalis* Brady, 1904, *A. warreni* Brady, 1913, *Harpacticus meridionalis* Sars, 1927,
657 *Cletocamptus confluens meridionalis* Kiefer, 1934 and *C. trichotus* Kiefer, 1934 as other
658 harpacticoid species that occur in southern Africa, but information could not be sourced on
659 whether these species occur in temporary wetlands.

660 Of the ecological work directed specifically at copepods, most has focussed on aspects
661 related to diapause. Diapause is a stop in the ontogenetic development induced by the need of
662 physiological adjustments in the individual; in copepods this process affects one stage only,
663 occurs regularly and is only partially related to environmental factors. Dormancy is a
664 reversible cessation of development clearly related to environmental changes and involves
665 different stages (depending on the species and environmental scenario) including eggs,
666 nauplii, copepodites and adults (Dahms, 1995). Eggs are the most common dormant stage
667 among diaptomids, and in cyclopoids dormancy is frequently related to copepodites C4 or C5
668 (Suárez-Morales et al., 2015). Diapause occurs under drying or unfavourable environmental
669 conditions and is a dominant feature of the life cycle of many copepod species and a major
670 factor in their successful colonisation of temporary wetland habitats worldwide (Dahms,
671 1995; Williams, 2006). In temporary ponds, cyclopine cyclopoids can emerge from diapause
672 within a single day of the appearance of water (Reid & Williamson, 2010). Unlike ostracods,
673 copepods generally reproduce sexually, but often have very fast development times, as little
674 as one week to reach the adult stage, depending on prevailing environmental conditions
675 (Williams, 2006). Similarly to ostracods, however, eggs are passively dispersed via wind or
676 attached to mobile fauna (Dahms, 1995). Very little other ecological work has been directed
677 specifically at copepods, with the exception of Wasserman et al. (2016a), Dalu et al. (2017b),
678 Dalu et al. (2017c) and Wasserman et al. (2018), who all investigated aspects of the trophic

679 ecology of the paradiaptominids *L. raynerae* and *P. lamellatus* in the Eastern Cape province
680 of SA.

681 *Crustaceans: Cladocera*

682 In contrast to copepods, most of the world's over 600 species (Forró et al., 2008) of
683 Cladocera ('water fleas') occur in freshwater environments, where they often dominate the
684 lentic fauna in terms of abundance. Southern Africa harbours a mixture of Afrotropical,
685 Palaearctic, circumtropical and local endemic elements in its cladoceran fauna (Smirnov,
686 2008; Van Damme et al., 2013). The taxonomy of cladocerans in the region requires an
687 urgent overhaul, particularly the Chydoridae and the Daphniidae (Seaman et al., 1999; Van
688 Damme et al., 2013), thus an estimate of the number of species in the region is not presently
689 available. Smirnov (2008) reported 112 cladoceran species from SA, with a wide
690 representation of Palearctic species. Traditionally, the region was considered to possess low
691 levels of cladoceran endemism (Frey, 1993; Korovchinsky, 2006; Smirnov, 2008), but more
692 recently this has been attributed to the limited state of knowledge of the fauna rather than
693 genuinely low endemism (Van Damme et al., 2013). As is the pattern for Ostracoda and
694 Copepoda, the greatest cladoceran species richness and endemism in the region occurs in
695 temporary wetland environments, particularly montane rock pools in the east (e.g.
696 Drakensberg) and endemics of Western Cape lowland pools (Van Damme et al., 2013). The
697 families Chydoridae and Macrothricidae, the sub-genus *Daphnia* (*Ctenodaphnia*) and the
698 genera *Ceriodaphnia* and *Moina* are common inhabitants of temporary wetlands in the region
699 (Seaman et al., 1999). A literature search of taxonomic and ecological records of cladocerans
700 from temporary wetlands in the region was attempted, but given the urgent need of
701 taxonomic revision of species in this region (Van Damme et al., 2013), it was later
702 abandoned.

703 Survival strategies of cladocerans in temporary wetlands are broadly similar to those of the
704 other microcrustaceans, with highly desiccation-resistant eggs being produced as water levels
705 draw down (Vandekerckhove et al., 2005). Cladocerans favour parthenogenetic (asexual)
706 cloning during favourable times when the wetland is inundated and males are rare for much
707 of the hydroperiod (Williams, 2006). Males appear more frequently towards the end of the
708 hydroperiod and sexual reproduction prevails as water disappears from the wetland. Whilst
709 parthenogenetic cloning produces soft eggs for rapid development, sexually produced eggs
710 are encased in a hardened ephippium and can withstand drying, freezing and passive dispersal

to other waterbodies (Bilton et al., 2001; Vandekerckhove et al., 2005; Williams, 2006). As has been found in many aquatic environments globally, cladocerans have also been identified as important trophic components in systems of the region (Dalu et al., 2016; Wasserman et al., 2016b; Dalu et al., 2017b; Dalu et al., 2017c).

Hydrachnidia: Water mites

Water mites (suborder Hydrachnidia) are abundant and ubiquitous in temporary wetlands and over 6 000 species have been described worldwide from freshwater environments, including 160 species from southern Africa (Di Sabatino et al., 2008). Since then, more species have been added to the southern African records for Torrenticolidae (Goldschmidt & Smit, 2009), Hygrobatidae (Pešić et al., 2013), Mideopsidae (Pešić et al., 2013), Arrenuridae (Smit, 2012) and Hydryphantidae (Gerecke, 2004). Jansen van Rensburg (1976) provided a taxonomic key to identify the 22 water mite families from the region (updated by Jansen van Rensburg & Day, 2002). Throughout the key, and indeed also for more recent species-level accounts (see above), no mention is made in the habitat descriptions for each taxon as to the preference for temporary versus permanent water bodies. Hence, we are unable to provide further information from the taxonomic literature regarding those taxa specifically known to inhabit temporary wetlands in the region. The following taxa have been recorded from temporary wetlands in the south-western Cape Mediterranean-climate region of SA (records from Mlambo et al., 2011 unless otherwise specified), with the help of Dr R. Gerecke, Tübingen, Germany, and Dr E. Ueckerman, ARC, Pretoria: *Arrenurus* spp. (Arrenuridae), *Eylais* spp. (Eylaidae), *Hydrachna fissigera* Koenike, 1898, *Hydryphantes parmulatus* Koenike, 1912, *Diplodontus schaubi* (Koenike, 1893), *Mamersa testudinata* Koenike, 1898, *Limnochares crinita* Koenicke, 1898, *Piona* sp. (Pionidae), and *Neumania* spp. (Unionicolidae). In addition, hygrophilous representatives of several terrestrial mite taxa were reported by Bird et al. (2013) from the cohorts Gamasina (*Macrocheles* sp., Macrochelidae), Astigmatina (Oribatida), and Parasitengona (Trombididae).

The great diversity of water mites in temporary habitats is remarkable, given that most do not possess the typical adaptations of temporary wetland invertebrates (either diapause of eggs/adults or active dispersal as adults) (Bohonak et al., 2004). Instead, the radiation of water mites in temporary wetlands has been facilitated by parasitic larvae that, once hatched from eggs laid within a waterbody, parasitize adult insects and are thus dispersed to other waterbodies (Bilton et al., 2001; Bohonak et al., 2004). Parasitism is obligate in all species

studied so far that colonize astatic water bodies, and mite larvae die if they fail to penetrate an insect host (Bohonak et al., 2004). However, not all water mites survive the dry phase via parasitism, and certain taxa (e.g. some members of the Hydryphantidae) are capable of surviving dry phases as deutonymphs or adults that enter an inactive state in microhabitats with persistent moisture, such as moist mud or damp crevices in logs (Wiggins et al., 1980; Bohonak et al., 2004).

Insecta: Ephemeroptera (mayflies)

The order Ephemeroptera have the distinction of being the most ancient lineage of extant winged insects, dating back to the late Carboniferous or early Permian periods (Brittain & Sartori, 2003; Barber-James et al., 2008). Despite over 3 500 species worldwide and 435 species from the Afrotropical region (Sartori & Barber-James, 2018), most mayfly diversity is represented in running waters and this ancient group is relatively depauperate in lentic environments (Newell & Hossack, 2009). A baetid species, *Cloeodes hydatum* McCafferty & Lugo-Ortiz, 1995, from temporary wetlands (rock pools, ephemeral pools and springs) of the Pantanal wetland system was the first mayfly species to be experimentally shown to withstand repeated drying conditions of up to nine hours at a time (Nolte et al., 1996). On the contrary, a known inhabitant of intermittent waters, *Daleatidium* spp. (Leptophlebiidae), from New Zealand dies only after three hours of drying exposure (Robinson & Buser, 2007). However, Nolte et al. (1996) made a distinction between *C. hydatum* and the chironomid *Polypedilum vanderplanki* Hinton, 1951, calling it drying-tolerant as opposed to drying-resistant, which the latter is. Although the genus *Cloeodes* occurs in the region, a similar characteristic of drying-tolerance has not been demonstrated, and it has also not been reported from temporary wetlands. In southern Africa, only members of the genera *Cloeon* Leach, 1815 and *Procloeon* Bengtsson, 1915 (Baetidae) are known from truly lentic waterbodies such as dams and wetlands and appear to be capable of reaching high abundance in temporary waterbodies (Barber-James & Lugo-Ortiz, 2003). Similar results have been reported in Irish turloughs (Reynolds, 2000).

Although the aquatic nymphal phase of these baetid mayflies is rather delicate and desiccation-intolerant, populations persist in temporary wetlands due to the nymphs being extremely fast-growing (reaching maturity in 2-3 weeks) and the adults apparently producing desiccation-resistant eggs (Williams, 2006). Dispersal is via short-lived winged adults or via passive dispersal of eggs (Bilton et al., 2001; Williams, 2006). Barber-James & Lugo-Ortiz

(2003) list 22 Afrotropical species of *Cloeon* (nine from SA) and three species of *Procloeon* (one from SA). It is not yet known which of these species inhabit temporary wetland environments, which reflects the desperate need for more species-level studies of mayflies in Africa (Barber-James & Gattolliat, 2012). *Cloeon* is widespread in the region and has been recorded from temporary wetlands throughout the region, including the Western Cape (Mlambo et al., 2011; Bird et al., 2013) and Mpumalanga (Foster et al., 2015) provinces of SA, Namibia (Day, 1990), Zimbabwe (Chakona et al., 2008) and Botswana (Jocqué et al., 2006).

Gillies (1990) reported the unusual habitat of the baetid *Acanthiops erepens* (Gillies, 1990), which occurs in Tanzania on rock in semi-permanent vertical trickles and in the splash zone next to small cascades. The nymphs cling to moss growing on the rock face that is kept moist by spray from the adjacent waterfall, and if disturbed they plummet into the waterfall pool below.

Insecta: Odonata

The order Odonata, comprising some 5,680 species worldwide (Kalkman et al., 2008), is divided into the suborders Zygoptera (damselflies) and Epiprocta, which is further divided into the infraorders Anisoptera (dragonflies) and Aniszygoptera (Asian species only) (Rehn, 2003). As with the Ephemeroptera, the odonates are an ancient order, dating back well into the Permian (Kalkman et al., 2008). The Odonata are relatively well studied, especially in the Afrotropical region where their diversity is depauperate compared to other tropical regions (Dijkstra, 2003) and thus their taxonomy is relatively well known in comparison to other freshwater invertebrate groups (but still not without some confusion, see Dijkstra, 2003). That said, Dijkstra et al. (2015) recently described a further 60 new African species as adults, from regions north of the area under review in this paper. Therefore, if brightly coloured, charismatic creatures such as adult Odonata are still incompletely known, how much less do we yet know about the more obscure freshwater invertebrates, especially those inhabiting temporary waterbodies in Africa.

Unlike many of the African freshwater invertebrates, IUCN Red List information has been compiled for many of the southern African odonate species, with most of the threatened species being habitat specialists occurring in highland areas of the Western and Eastern Cape provinces (Samways, 2004), many frequenting temporary water bodies, but most as

opportunistic visitors rather than being dependent upon these habitats. Odonate species exploiting temporary habitats depend on dispersal for survival and tend to have large home ranges and are habitat generalists, being tolerant of ecological change (Clausnitzer et al., 2012). Thus, temporary wetland species are generally not threatened. In southern African arid regions, no desert endemics are known and most of the widespread desert-inhabiting species are powerful fliers belonging to the Anisoptera that are not restricted to deserts, being more common in savannah wetlands (Suhling et al., 2003). The 889 known odonate species of the Afrotropical region (Kalkman et al., 2008) all have aquatic larvae, which are not capable of withstanding desiccation themselves (for a rare exception from Brazil, see Van Damme & Dumont, 1999). Odonate populations are however, able to persist in temporary habitats through production of drought-resistant eggs combined with fast-growing larvae, multivoltine life cycles and strong active dispersal of adults (for more detailed reviews of odonate adaptations to temporary habitats, see Johansson & Suhling, 2004; Suhling et al., 2005; De Block et al., 2008; Hassall & Thompson, 2008). Timing of hatching may also be important. Schiel & Buchwald (2015) showed that for 15 species of odonates across three genera (*Lestes* – Lestidae, *Aeshna* – Aeshnidae and *Sympetrum* – Libellulidae), species inhabiting only temporary ponds hatched significantly earlier in the year than congeners inhabiting a combination of permanent and temporary ponds. Despite the harsh environmental conditions often presented in temporary waterbodies, a major reward for exploiting such habitats is often the reduced predation pressure on larvae (lack of fish for instance) allowing them to forage actively for food, whereas in permanent waters larvae generally need to remain concealed and rely on ambush predation (Hassall & Thompson, 2008).

Several checklists are available covering various sub-regions of southern Africa, from which habitat preferences of the various species can be discerned (if such information is available at all). Martens et al. (2003) listed 102 odonate species from Namibia. They listed the following taxa as facultative inhabitants of temporary wetlands: *Lestes pallidus* Rambur, 1842, *Ischnura senegalensis* (Rambur, 1842), *Anax ephippiger* (Burmeister, 1839), *Pantala flavescens* (Fabricius, 1798), and *Sympetrum fonscolombii* (Selys, 1840). Suhling et al. (2003), in their investigation of desert-adapted odonates, listed the following species from ephemeral pools in the Namib desert of Namibia: *Africallagma glaucum* (Burmeister, 1839), *Ischnura senegalensis* (Rambur, 1842), *Anax ephippiger*, *A. imperator* Leach, 1815, *Paragomphus genei* (Selys, 1841), *Crocothemis erythraea* (Brullé, 1832), *Orthetrum chrysostigma* (Burmeister, 1839), *Palpopleura lucia* (Drury, 1773), *P. flavescens*, *S.*

839 *fonscolombii*, *Trithemis annulata* (Palisot de Beauvois, 1807) and *T. kirbyi ardens*
840 (Gerstäcker, 1891). Mlambo et al. (2011) recorded the anisopteran genera *Anax* (Aeshnidae)
841 and *Trithemis* (Libellulidae) and the zygopteran genera *Africallagma*, *Ischnura* and
842 *Pseudagrion* (all Coenagrionidae) from temporary wetlands of the Western Cape. Jocqué et
843 al. (2006) reported the larvae of *P. flavescens* as top predators in temporary rock pools of
844 south-eastern Botswana. Simaika et al. (2016) investigated the benefits of artificial ponds as
845 biodiversity refugia for Western Cape dragonflies and listed the following taxa from a
846 temporary pond: *A. imperator*, *C. erythraea*, *I. senegalensis*, *Orthetrum chrysostigma*
847 (Burmeister, 1839), *O. trinacria* (Selys, 1841) and *Trithemis arteriosa* (Burmeister, 1839).
848 Samways (1999) noted *L. pallidus* as a frequent inhabitant of temporary wetlands in SA,
849 while Clark & Samways (1996) recorded *Palpopleura deceptor* (Calvert, 1899), *P. jucunda*
850 Rambur, 1842, *P. lucia*, *Bradinopyga cornuta* Ris, 1911, *T. kirbyi ardens* and *P. flavescens*
851 from ephemeral rock pools in KNP. Interestingly, larvae of the libellulid *Hemistigma*
852 *albipunctum* (Rambur, 1842), which inhabits marshes and swamps, can remain alive in the
853 mud as these habitats dry out, surviving with minimal water and emerging as an adult before
854 the local area completely dries out (Samways, 2008).

855 Kipping (2010) provided a checklist of the odonates of Botswana and noted that most of
856 the country's odonates are widespread African savannah species that, although having a
857 preference for more stable habitats, are able to disperse over wide distances and breed in
858 temporary waterbodies in arid areas such as the Kalahari Desert. Suhling et al. (2009)
859 reported that true obligate migrant species of the arid Kalahari region include *P. flavescens*,
860 *S. fonscolombii* and *A. ephippiger*. Kipping (2010) recorded the following species as
861 facultative inhabitants of temporary wetlands of Botswana: *Lestes dissimulans* Fraser, 1955,
862 *L. pallidus*, *L. pinheyi* Fraser, 1955, *A. glaucum*, *Azuragrion nigradorsum* (Selys, 1876),
863 *Ceriagrion suave* Ris, 1921, *I. senegalensis*, *A. ephippiger*, *A. tristis* Hagen, 1867,
864 *Brachythemis leucosticta* (Burmeister, 1839), *B. cornuta*, *Diplacodes luminans* (Karsch,
865 1893), *Orthetrum brachiale* (Palisot de Beauvois, 1817), *O. caffrum caffrum* (Burmeister,
866 1839), *O. chrysostigma chrysostigma* (Burmeister, 1839), *O. trinacria*, *P. deceptor*, *P.*
867 *flavescens*, *Rhyothemis semihyalina* (Desjardins, 1832), *S. fonscolombii*, *Tramea basilaris*
868 (Palisot de Beauvois, 1807), *T. kirby ardens* and *Urothemis assignata* (Selys, 1872). For
869 more detailed information on the distribution, habitat and Red List status of these species, the
870 reader is referred to Kipping (2010) and Samways & Simaika (2016).

In some odonate species there are temporal changes in geographic range in accordance with the wet and dry cycles of the El Niño Southern Oscillation. An extreme case was for *Aciagrion dondoense* Dijkstra, 2007, which was not known in SA prior to the year 2000 when it appeared in great numbers on the KwaZulu-Natal coast, having come down the coast with flooding farther north. A few years later it was gone from the country with the dry phase appearing (Samways, 2010). Several South African species are adapted to these cycles, with species like *Orthetrum robustum* Balinsky, 1965 retreating from temporary pools in the dry phase to permanent lakes, to return later to the temporary pools in the wet cycle (Samways, 2008). One last factor to consider is that rivers that stop flowing and have only pools ('kuile') in the dry season. Such pools in these rivers may attract species temporarily until the river flows again. There may even be a succession of species as the pools gradually dry out, with *Pseudagrion commoniae* (Förster, 1902) and *P. salisburyense* Ris, 1921 being the last to survive before the pools finally dry out (Samways, 2008).

Insecta: Hemiptera

The hemimetabolous insect order Hemiptera is usually considered to contain four suborders: Sternorrhyncha (aphids, whiteflies, and scale insects); Auchenorrhyncha (cicadas and hoppers); Coleorrhyncha (moss bugs); and Heteroptera (true bugs) (Capinera, 2008; Forero, 2008). It comprises some 82,000 described species, and possibly almost 200,000 species in total worldwide. The bugs associated with water all belong to the Heteroptera, particularly the semi-aquatic Gerromorpha and the predominantly aquatic Nepomorpha. Gerromorphs have hydrophobic tarsi and gracile bodies that allow them to stand on the water surface supported by the surface tension of the meniscus, while nepomorphs generally crawl or swim underwater and typically have reduced antennae and ocelli. The Nepomorpha have a fossil record dating back about 250 million years (Grimaldi & Engel, 2005).

Most species have winged adults that disperse well and can therefore colonise temporary wetlands quickly. The nepomorph families all breathe air through plastrons or respiratory tubes, and the gerromorph families are epipleustonic, living just above the water surface rather than in the water, so dissolved oxygen does not limit the potential of a water body for colonisation by water bugs. The species in most families are generalist predators, and can therefore inhabit a wide variety of waters with flow rates that are slow or absent. Many gerromorphs associate with floating or emergent vegetation that helps them to avoid predators and being washed away in flowing waters (Reavell, 2003). Immature nymphs have

biologies similar to their adults, but cannot fly, so they complete their development only in temporary wetlands that take more than (generally) four to ten weeks to dry up. Adults of some species of gerromorph may be winged or apterous (regulated by temperature and photoperiod); when apterous adults inhabit a temporary wetland, they indicate that at least one generation has completed its life cycle in the wetland. The high mobility and broader habitat preferences of most aquatic bugs means that populations of temporary wetland species are generally not threatened (Collinson et al., 1995).

The morphology, taxonomy and general ecology of the southern African water bug fauna was summarised by Reavell (2003), who stated that little work had been done on the fauna of the region, and that it needed urgent taxonomic revision. Reavell (2003) provided a key to the families of aquatic Hemiptera in the region, and a list of the known genera, but keys to the genera and species are still lacking. Ecological and life history studies of temporary wetland water bugs in the region include Hutchinson (1933), Hynes (1955), Weir (1966), Tawfik (1969), Boettger (1974), Mlambo et al. (2011), Ferreira et al. (2012), Bird et al. (2013) and Nhiwatiwa et al. (2017a). Hemiptera from six families constituted an average of 14% of the individual invertebrates surveyed in a large number of temporary waters in the Western Cape (Mlambo et al., 2011), and Hemiptera were co-dominant with Coleoptera (beetles) in nine temporary pans in Mpumalanga (Ferreira et al., 2012). The dominant families are usually Corixidae and Notonectidae. The water bugs tend to be associated with warmer, alkaline water in temporary pans (Ferreira et al., 2012), and larger-bodied species are more common in the absence of fishes in ephemeral pans (Nhiwatiwa et al., 2017a; Nhiwatiwa et al., 2017b). However, because of their high vagility and relatively broad habitat tolerances, water bugs may be poor indicators of anthropogenic effects on temporary wetlands in SA (cf. Bird et al., 2013), an effect that can be compounded by artefacts of the sampling method (de Klerk & Wepener, 2013). Their instantaneous diversity may also be related to the stage of the wetland's hydroperiod (Lahr et al., 1999; Dalu et al., 2017b; Dalu et al., 2017c), since older wetlands have had more time to become colonised and develop prey populations, thus obscuring the effects of human activities.

Insecta: Trichoptera

Trichoptera are holometabolous insects, mostly with an aquatic larval and a terrestrial adult phase. They have been recorded worldwide with the exception of some oceanic islands and the polar regions (de Moor & Scott, 2003). The Trichoptera World Checklist (Morse, 2011)

lists 14 548 species of 616 genera in 49 families recorded globally thus far. Of these, 1 193 species (8.2%) in 83 genera from 21 families are known to occur in southern Africa (Tobias & Tobias, 2008). Recently, some species were collected in temporary wetlands, despite the earlier assertion of Scott (1970) that Trichoptera do not inhabit temporary lentic waters of the region: *Oxyethria velocipes* (Barnard, 1934) (Jones, 2002; Day et al., 2010) and *Athripsodes* sp. (Bird, 2012), both from the Western Cape province; and *Ecnomus thomasseti* Mosely, 1932, *Oecetis* sp. and *Oxyethira* sp. from Mpumalanga province (Ferreira et al., 2012). Trichoptera found in temporary waters show several adaptations to this habitat in different life stages, but little is known about this in southern Africa as research has mainly been conducted on species from other regions, discussed below. Larvae use silk to construct transportable or fixed cases from material readily available, such as sand, small stones, sticks, leaves and algae. The ability to build cases has allowed trichopteran larvae to adapt to a wide range of environmental conditions: spring sources, mountain streams, splash zones of waterfalls, large rivers, lakes and temporary wetlands (de Moor & Scott, 2003). Many species are represented in cooler and faster-moving waters as dissolved oxygen is a limiting factor for habitat colonisation (de Moor & Scott, 2003).

Trichopteran larvae are apneustic, which means that they lack spiracles; they take up dissolved oxygen through the thin abdominal membrane by direct diffusion. Their abdominal surfaces are often expanded with tubular or branched gills that increase the surface area of the abdominal membrane. To ensure sufficient uptake of dissolved oxygen, larvae can create a current through their cases by abdominal undulation. This aids gas exchange when oxygen concentrations are low, as is the case in temporary wetlands (Williams, 1987; de Moor & Scott, 2003).

As water levels decrease during drying, faunal density increases and cases offer a protective body covering and camouflage to protect larvae from predation. A combination of case material (mineral rather than organic), shape (short, wide case with “hedgehog” structure rather than long tubular cases or flat purses) and defensive behaviour (retreating deep into the case and remaining motionless) are adaptive strategies employed by *Limnephilus externus* Hagen, 1861 against predation by diving beetle larvae (*Dytiscus* sp.) (Wissinger et al., 2006). Another species, *L. picturatus* McLachlan, 1875, evades predation of pupae by burrowing into soft substrates to pupate, whereas larvae of *Asynarchus nigriculus* (Limnephilidae) (Banks, 1908) build stronger mineral cases prior to pupation. Furthermore, both species have

967 faster developmental times, which reduce temporal overlap with predators (Wissinger et al.,
968 2003). Pupae can develop while the water column has dried up and the soil is still moist
969 (Wiggins, 1973). The case material is important with regard to pond-drying. Larvae of *L.*
970 *coenosus* Curtis, 1834 construct cases using organic material whereas *L. vittatus* (Fabricius,
971 1798) larvae use mineral material. Organic material has been shown to retain moisture better
972 than mineral material, preventing drying of larvae (Zamora-Muñoz & Svensson, 1996).

973 Cases also protect caddisfly larvae from inter- and intraspecific aggression as water levels
974 decrease (Wissinger et al., 2004b). Aggressive behaviour includes fighting (foreleg wrestling,
975 biting, case shaking) and mobbing of individuals (involving 5-10 larvae) which often results
976 in death and cannibalism of conspecifics (Lund et al., 2016). Wissinger et al. (2004b) have
977 shown that cases reduced or prevented cannibalism significantly among larvae of *A.*
978 *nigriculus*. Cases also reduced vulnerability of three *Limnephilus* species (*L. externus*, *L.*
979 *picturatus*, *L. secludens* Banks, 1914) to intraguild predation by *A. nigriculus* (Banks, 1908).
980 Aggression among *A. nigriculus* in mesocosm experiments was higher in high-density
981 treatment and even higher in the absence of dietary protein supplement (Lund et al., 2016).
982 Although aggressive behaviour has trade-offs, such as injury and subsequently becoming a
983 secondary victim, cannibalism provides an important protein- or lipid-rich dietary supplement
984 that offers advantages later in the life-cycle.

985 Dietary experiments on *L. coenotus* and *L. vittatus* supplementing detritus diet with
986 protein have shown that larger larvae survive drying better as they contain more water
987 compared to smaller ones (Zamora-Muñoz & Svensson, 1996). Furthermore, body mass of
988 adult *L. externus* significantly increased when ambient food treatment was supplemented with
989 a protein-rich supplement (Jannot et al., 2008). Larval diet can have a significant long-term
990 effect on adult fitness (Jannot, 2009) as female body size is correlated with fecundity and
991 male body size with mating success (Wissinger et al., 2004a).

992 Adults of *L. externus*, *L. picturatus* and *A. nigriculus* emerge prior to pond-drying and
993 females enter an ovarian diapause after mating (Wissinger et al., 2003). Summer diapause is
994 an important adaptation to survive dry periods (Wiggins, 1973; Wiggins et al., 1980), which
995 also protects eggs from predation, drying and freezing (Wissinger et al., 2003). Oviposition in
996 temporary pools usually occurs above or near water, not directly in water as is normal for
997 most trichopteran species (Wiggins, 1973). Eggs are deposited in a gelatinous egg-matrix,
998 which is highly resistant to extremes like desiccation and freezing (Wiggins, 1973). Eggs are

deposited under rocks and logs, where larvae hatch and develop further within a gelatinous mass until re-wetting occurs (Wissinger et al., 2003). *Lectrides varians* Mosely, 1953 show a bimodal response – some larvae become active with the first onset of re-wetting or re-immersion whereas others remain aestivating and only become active when surface water levels are more stable. Although this adaptation ensures population survival, it is at the cost of larval survivorship (Wickson et al., 2012). Larvae enter the temporary ponds with, for example, melt water, develop further and pupate prior to pond-drying (Wissinger et al., 2003).

Insecta: Diptera

Virtually all African temporary waters support the larvae of at least one species of fly, and flies boast of a large array of adaptive mechanisms for this variable environment (McLachlan & Ladle, 2001). In the region, the families Chironomidae, Ceratopogonidae and Culicidae commonly occur even in the Kalahari and Namib deserts, where rain may fall only at intervals of several years. Although dipterans are ubiquitous in these systems, very few species are able to survive desiccation (Williams, 2006). Unlike many branchiopod crustaceans, which withstand dry periods as desiccation-resistant cysts, most flies that occur in temporary waters survive the dry period as adults or, in the case of some culicids, as larvae in perennial water bodies. Hinton (1953) indicated that some African Stratiomyidae can withstand desiccation, though this is not recorded in southern Africa.

Chironomids mostly survive dry periods as adults. Frouz et al. (2003) reviewed the strategies employed by chironomids to survive the variable environment of temporary wetlands. Given that larvae of several species are found in hyper-arid conditions in the Namib Desert, where pools may not form for 5 to 10 years at a stretch, the question yet to be answered is, “How are the *adults* able to survive the dry years in these environments?” A few other species survive for some time in stout pupal cocoons during dry periods (Armitage, 1995), but presumably these periods are relatively short. When it comes to survival under harsh conditions, larvae of the ‘sleeping chironomid’, *Polypedilum vanderplanki* Hinton, 1951, hold the record. The species breeds in small unshaded rock pools in Central and West Africa. Larvae can undergo up to 17 years of desiccation and, when wetted, will ‘wake up’ and behave normally, feeding and growing until the next time they are desiccated. Hinton (1960) performed various experiments on these remarkable organisms, showing that they can withstand temperatures between -270°C and +102°C and can survive through a

1031 wetting/drying cycle at least ten times. An additional species of ‘sleeping chironomid’, *P.*
1032 *ovahimba* Cranston, 2014, has been recently described from rock pools of the Waterberg
1033 Plateau, Namibia (Cranston, 2014).

1034 Other chironomids, such as *Chironomus pulcher* Wiedemann, 1830, also breed in rain
1035 pools on rocks throughout much of Africa and have rapid life cycles. Larvae of the
1036 chironomid subfamily Podonominae (*Archaeochlus* sp. and *Afrochlus harrisoni* Freeman,
1037 1964) are found in temporary trickles over rocks and boulders in the Drakensberg, on granite
1038 inselbergs near Harare, and in Namibia (Harrison, 2002). Similarly, *Harrisonina petricola*
1039 Freeman, 1956 was originally found breeding in the thin film of water running over stones in
1040 an ephemeral stream in the Olifants River Valley. It is not known how these survive dry
1041 periods. Rock pools in Africa are dominated by dipteran larvae (up to 300 000 per pool),
1042 which are often the only macroinvertebrate inhabitants. If the pools are very short-lived (1-2
1043 days), the only inhabitants are larvae of *P. vanderplanki*. Pools lasting several weeks often
1044 support *Chironomus imicola* Kieffer, 1913, which has little tolerance of desiccation, but is
1045 able to complete its life cycle in 10-12 days (McLachlan & Cantrell, 1980).

1046 Larvae of a few ceratopogonid species are found in extremely arid regions (de Meillon &
1047 Wirth, 2002) and yet they seem not to be particularly adapted to withstanding desiccation.
1048 There are no African examples. Larvae of *Culicoides sonorensis* Wirth & Jones, 1957 from
1049 the Sonoran Desert in North America, for instance, are able to survive drying, but only for a
1050 day or two (McDermott & Mullens, 2014). Pools lasting several days often contain larvae of
1051 the ceratopogonid *Dasyhelea thompsoni* de Meillon, 1936 that burrow into mud at the bottom
1052 of the pond when water evaporates (McLachlan & Cantrell, 1980). Similarly, Dodson (1987)
1053 found that *Dasyhelea subletti* Wirth, 1987 was the most common taxon in short-lived
1054 ephemeral rock pools in the Moab desert, USA.

1055 Numerous species of culicid are associated with temporary waters in Africa (Coetzee,
1056 2002), and many of them are vectors of parasites causing disease in humans and other
1057 vertebrates. Several of the vectors prefer to breed in very small containers such as old tyres,
1058 pots, and in footprints, or in phytotelmata: pools of water held by plants, for instance in the
1059 axils of leaves. *Aedes aegypti* (Linnaeus, 1762) is the vector of yellow fever. While the
1060 disease does not occur in southern Africa, *Ae. aegypti* does. It breeds in tiny containers, and
1061 the eggs can withstand desiccation for years (Coetzee, 2002). Farnesi et al. (2015), comparing
1062 the degree of resistance to desiccation of eggs of three species of mosquito, have shown that

1063 eggs of *Ae. Aegypti* can survive in a dry form for months, those of *Anopheles aquasalis*
1064 Curry, 1932 for a day, and of *Culex quinquefasciatus* Say, 1823 for a few hours, these times
1065 relating to the thickness of the egg shell. The members of ‘floodwater Aedes’, which can be
1066 vectors of Rift Valley Fever, such as *Ae. (Neomelanicion) mcintoshi* Huang, 1985, *Ae.*
1067 *(Neo.) circumluteolus* (Theobald, 1908), *Aedes (Neo.) luridus* McIntosh, 1971, *Aedes (Neo.)*
1068 *lineatopennis* (Ludlow, 1905), *Aedes (Ochlerotatus) caballus* (Theobald, 1912) and *Aedes*
1069 *(Och.) juppi* McIntosh, 1973, aestivate as eggs for up to 20 months (Jupp et al., 1980).

1070 Several members of the genus *Anopheles* are vectors of malarial parasites. Although some
1071 anophelines prefer to oviposit in temporary waters, neither the eggs nor the larvae are
1072 resistant to desiccation. First-instar larvae of *An. gambiae* s.l. are said to be amphibious
1073 (Miller et al., 2007), in that they can survive in the film of water on the surface of mud. Both
1074 *An. gambiae* s.s. and *An. arabiensis* Patton, 1905 (two South African vectors of malaria)
1075 prefer sunny temporary pools and puddles, the major adaptation to this biotope being very
1076 rapid larval development. Mosquitoes appear to overwinter as larvae in permanent waters
1077 (Appleton et al., 1995).

1078 A few species of other dipteran families are known to occur in temporary waters in
1079 southern Africa, but very little is known about their systematics or their biology. We
1080 hypothesise that those occurring in the very arid western areas, such as the Namib Desert,
1081 have some specific adaptations that allow them to persist in areas where rain pools are the
1082 only surface freshwaters and these may fill only every few years.

1083 *Insecta: Coleoptera*

1084 Worldwide, beetles are one of the most abundant and speciose macroinvertebrate groups in
1085 inland waters, occurring across the entire hydrological spectrum, from the smallest puddles
1086 and phytotelmata, to large lakes and rivers. Water beetles are not a single taxonomic group,
1087 and instead result from around 15-20 transitions between land and water during the
1088 evolutionary history of the Coleoptera (Jäch & Balke, 2008; Mckenna et al., 2015; Short,
1089 2018). Beetles are diverse and frequent inhabitants of temporary waterbodies, where
1090 carnivorous taxa such as diving beetles (Dytiscidae) are often the top predators. Temporary
1091 pool inhabitants are typically a mix of specialist and more generalist species, some of which
1092 may only occur there temporarily as adults. In many cases, specialising in temporary pools
1093 may have arisen as a life-history strategy to evade fish predation, and in the northern

1094 hemisphere some taxa characteristic of temporary ponds (e.g. *Agabus labiatus* (Brahm,
1095 1790)) are also found in permanent, acidic, fish-free sites (Foster et al., 2016). Unlike most
1096 freshwater insect groups, the majority of water beetles are aquatic as both larvae and adults,
1097 but in many cases, the precise ecology of these life-history stages differs to some extent.
1098 Some beetles utilise temporary waters primarily for reproduction, with adults occupying
1099 different habitats. In the Mediterranean, *Acilius duvergeri* Gobert, 1874 for example,
1100 breeding occurs in temporary lagoons, where larvae feed on microcrustacea in open water (a
1101 strategy which would make them vulnerable to fish predation in permanent waters), whilst
1102 adults occupy permanent ponds during summer drought (Dettner, 1982).

1103 Temporary pond water beetles have a variety of strategies to survive during the dry phase,
1104 reflecting the diverse origins of aquatic beetles themselves, and the repeated evolution of
1105 temporary water specialists from within these aquatic lineages. Temporary water species may
1106 have shorter larval duration compared to their permanent water relatives, allowing
1107 development to be completed during relatively short hydroperiods (e.g. Peters, 1972), and a
1108 range of approaches exists to surviving dry phases, including migration to permanent waters
1109 or adult/egg diapause *in situ* (e.g. Nilsson, 1986a, b, c). The most temporary of water bodies
1110 appear to be devoid of breeding water beetles, suggesting that the extent to which larval
1111 duration can be truncated may be limited. In addition, beetle larvae do not seem to have the
1112 same resistance to desiccation as many temporary-water Crustacea (see above), but do have
1113 the advantage that adults of most species can fly between habitats (e.g. Bilton et al., 2001).

1114 In southern Africa, members of a number of beetle families can be found in temporary
1115 water bodies, although there are no formal publications dealing with their biology and
1116 ecology. As a consequence, the following information is based on Bilton (pers. obs.), unless
1117 otherwise stated. The faunas of vegetated temporary pools and rock pools appear to be quite
1118 distinct, and in addition brackish water bodies have their own specialised beetle fauna.
1119 Haliplidae are relatively rare in southern Africa, and members of *Haliphus* and *Peltodytes* are
1120 mostly found in larger, more permanent waters. The rare Cape endemic, *Algophilus*
1121 *lathridioides* Zimmermann, 1924 is poorly known ecologically, but appears to be a specialist
1122 of slightly mineralised waters. The only recent record of this beetle is from the Berg River
1123 floodplain close to Hopefield, where it occurs abundantly in slightly brackish temporary
1124 flood pools in grassland. *Algophilus* is wingless, and is likely to be dispersed during floods,
1125 having been found in floodwater along the Berg River in spring.

Whereas the Noteridae are largely permanent water inhabitants, a number of Dytiscidae are temporary pond specialists, and others are found there occasionally or temporarily as adults. *Canthyporus* species occur across a spectrum of freshwater habitats, including seepages, streams and ponds, both permanent and temporary (Biström & Nilsson, 2006; Bilton et al., 2015). This genus is particularly diverse in the Cape, where most of its species are endemic. Many of the species in this genus, including *C. aenigmaticus* Biström & Nilsson 2006 and *C. lateralis* (Boheman, 1848), are characteristic inhabitants of small mountain rock pools, where both adults and presumed larvae can be found. Some inhabited pools form parts of temporary stream/seepage systems during periods of high rainfall, whereas others constitute completely isolated basins. Cape vernal pools are also inhabited by *Canthyporus*, including the relatively generalist *C. hottentottus* (Gemminge & Harold, 1868) and *C. petulans* Guignot, 1951, as well as *C. exilis* (Boheman, 1848), a species otherwise found in temporary seeps and marshes. Besides the widespread genus *Hyphydrus*, the South African Hyphydrini includes five endemic genera (see Ribera & Balke, 2007), three of which occur in temporary ponds. *Darwinhydrus solidus* Sharp is found from the Cederberg to Port Elizabeth, in both permanent and temporary waters, but is most frequent in densely vegetated Cape vernal pools. *Primospes suturalis* Sharp, 1882 is largely restricted to the far southwest of the Western Cape, and occurs in temporary pools and rivers, and two *Hydropeplus* species are inhabitants of both stream pools and temporary lentic pools, both vegetated and rock pools. Species of the genus *Rhantus* can also be found in vegetated temporary ponds, as both adults and presumed larvae, as can the phylogenetically isolated Cape endemic *Caperhantus cicurius* (Fabricius, 1787) (Bilton, 2017). The Critically Endangered Cape endemic *Capelatus prykei* Turner & Bilton 2015 has been found in vernal pools on the Cape Flats (Bilton et al., 2015).

To date, only a single species of the largely Holarctic genus *Helophorus* has been reported from southern Africa. *Helophorus aethiops* Balfour-Browne, 1954 is found throughout much of the winter rainfall zone, and is common in a variety of temporary pools, particularly shallow sites with exposed silt or sand substrates. The species-level taxonomy of southern African *Hydrochus* remains incompletely resolved, but a number of species are found in vegetated temporary ponds, as can many members of the Hydrophilidae, including *Enochrus hartmanni* Hebauer, 1998; *Enochrus (Methydrus)* spp.; *Helochaetes* spp; *Paracymus* spp; *Berosus* spp. and *Limnoxenus sjoestedti* Knisch, 1924, the latter being restricted to the South African Cape (Short & Liebheer, 2007).

The regional hydraenid beetle fauna is one of the most diverse on earth, most species in the region being Cape endemics. Here the family also attains its maximum ecological and morphological diversity, with both aquatic and (semi)terrestrial representatives (e.g. Perkins & Balfour-Browne, 1994). Species of the genus *Prosthetops*, which includes some of the largest hydraenids on earth, are characteristic inhabitants of temporary rock pools, where larvae and adults can be abundant, grazing biofilms. Bilton (2013) noted that large larvae of *P. wolfbergensis* Bilton, 2013 became active underwater in previously dry rock pools one day after rains, suggesting some desiccation resistance mechanism in these insects. During dry periods, *Prosthetops* adults can be found in rocky streams, although this does not appear to be their breeding habitat. Temporary vegetated freshwater ponds are inhabited by a number of species of *Hydraena* (*Hydraenopsis*), *Ochthebius* and *Parhydraena*, whilst temporary saline pools are home to species of *Ochthebius*, including *O. capicola* (Péringuey, 1892), a specialist of supralittoral coastal rock pools (Sabatelli et al., 2013), a habitat independently colonised by members of this genus on a number of occasions worldwide (Sabatelli et al., 2016).

Discussion

Much of southern Africa is characterised by a semi-arid or arid climate (Davis, 2011) and small temporary wetlands, though not widely acknowledged, are one of the most abundant and ubiquitous aquatic features of such landscapes (Silberbauer & King, 1991; Brendonck & Williams, 2000; Williams, 2006). The invertebrates found in these wetlands, whilst not widely acknowledged, constitute a significant proportion of the total aquatic biodiversity of the region. Whilst some recent research has highlighted the ecological and biodiversity importance of temporary wetland invertebrates (e.g. Spencer et al., 2002; Nicolet et al., 2004; Urban, 2004; Jeffries, 2005; Williams, 2006; Vanschoenwinkel et al., 2007; Ruhí et al., 2013; Batzer & Boix, 2016; Kneitel, 2016; Dalu et al., 2017a), baseline biodiversity data on these faunas are often lacking. Here we present such a synthesis for southern Africa, in the hope that this will aid further ecological investigations of temporary wetlands in the region.

A clear finding of this work is that, with the exception of a few groups (e.g. gastropod molluscs, odonates and certain dipteran families that act as disease vectors), we are often unable to reliably name many of the invertebrate organisms inhabiting these waterbodies. This becomes most obvious for the ‘lower’ invertebrate groups covered in this review, which is perhaps not surprising given their small size and the practical difficulties which may be

associated with collection and identification. For some of these groups, notably gastrotrichs, bryozoans and tardigrades, there is very little information about the freshwater species that occur throughout the region, including temporary wetlands. Although there is better information in this regard for the Northern Hemisphere, the ‘lower’ invertebrate phyla of freshwater environments have been relatively neglected worldwide (see Balian et al., 2008). Perhaps more surprisingly, the status quo reveals that in some cases we cannot put reliable names on species for groups represented by larger and more ‘charismatic’ organisms such as the ‘clam shrimps’ (Laevicaudata, Cyclestherida and Spinicaudata, but see Rogers et al., 2012 for revision of the family Limnadiidae), ephemeropterans and many of the hemipterans from temporary freshwater wetlands of the region. Harpacticoid copepods are another example of a taxon completely unstudied in the region, even from permanent freshwater systems.

To get a better hold on patterns of richness and endemism, one first needs to know what species are present in these wetland systems. A case in point is the Cladocera, which have traditionally been considered to contain few endemics in the region (Frey, 1993; Korovchinsky, 2006; Smirnov, 2008). More recent work suggests that the apparent low endemism is most likely an artefact of the limited state of knowledge of the fauna, rather than genuinely low endemism (Van Damme et al., 2013). Thus, statements on patterns of richness and endemism are of a preliminary nature at present for many of the invertebrate groups. A few such patterns are worthy of comment here. First, only for a few of the groups is evidence presented that they reach higher richness and/or endemism in temporary wetlands than for their permanent wetland counterparts. These taxa include large branchiopods, ostracods, copepods, cladocerans, certain dipterans (notably culicids) and coleopterans (some dytiscids). Thus, for the vast majority of freshwater invertebrate taxa in the region, permanent waterbodies would appear to hold a greater richness and/or endemism than temporary ones. Despite this, temporary wetlands support unique and specialised taxa. For instance, the temporary wetland ostracod fauna of northern and central Namibia and Botswana has a generic endemism rivalled worldwide only by East African paleolakes (Martens, 2001). Another feature of the ostracod fauna is the specific diversity in the family Lymnocytheridae for rock pools of the Drakensberg (SA), which is unmatched anywhere else in the world (Martens et al., 1998). Southern Africa is considered one of the world’s hotspots of large branchiopod diversity (Tuytens et al., 2015), almost all of which is housed in temporary wetlands. When considering the region’s area, the anostracan fauna is one of the richest worldwide and also has a very high endemism (~80% at the time of the review of Hamer &

Brendonck, 1997). Other than for the anostracans and ostracods, further commentary on which of the major invertebrate groups covered in this review stand out globally for their richness/endemism is limited due to the taxonomic uncertainty associated with most of the groups in the region, as highlighted above.

Identifications to genus and species level of specimens mentioned in this review are mostly based on morphological characteristics (unless specifically mentioned otherwise) and accuracy greatly depends on reliable identification keys and taxonomic expertise, which is considered a scarce skill in southern Africa (Hamer, 2013). The few region-specific keys that exist are in need of updating (Barber-James & Gattolliat, 2012). Species identification using molecular systematics has only become prominent in the region recently (da Silva & Willows-Munro, 2016) and will greatly contribute to more accurate identification of species recorded in temporary wetlands in future.

Another general trend picked up from this review is that virtually all of the major invertebrate groups possess some sort of adaptation for survival in temporary wetlands. Although only certain taxa can truly withstand desiccation (e.g. branchiopods, ostracods and copepods), many taxa have dispersal mechanisms which allow them to leave the waterbody before it dries up (e.g. leeches, water mites and odonates), or are able to survive the drought phase as adults (e.g. dipterans). Many of the taxa inhabiting these environments are habitat generalists that may opportunistically exploit temporary waterbodies for the advantages afforded by these environments, such as reduced levels of predation and competition (although this notion is in itself controversial, see Brendonck et al., 2002). A case in point is the Odonata and Hemiptera, which are common and widespread in temporary wetlands of the region, but tend to be opportunistic colonisers of these environments, depending on dispersal for survival and having large home ranges, and thus are generally not threatened. This generalistic nature of many of the inhabitants of temporary wetlands in the region is hypothesized as one of the reasons why many of the taxa do not display high levels of endemism. There are certain exceptions, such as for rock pools of the Drakensberg in the east of the region, and the lowland pools of the Western Cape province, both of which are hotspots of ostracod and cladoceran endemism and richness (Martens et al., 1998; Martens, 2001; de Moor & Day, 2013; Van Damme et al., 2013).

In terms of the best known of the temporary wetland specialists, ‘fairy shrimp’ (anostracan) species richness is depicted as an array of hotspots throughout the region, not

being centered on any particular area and does not appear to be governed by any particular climatic factor (Hamer & Brendonck, 1997). The large branchiopod crustaceans show the greatest potential as ‘flagships’ for conservation of temporary wetlands in the region, given their relatively large size and uniqueness in terms of adaptations to life in the temporary aquatic realm. The odonates, molluscs and anostracans are the only groups covered in this review to have an appreciable amount of IUCN Red List information in the region. The freshwater molluscs appear to have few threatened species in the region (10 species, 8.6% of the fauna, Kristensen et al., 2009). Anostracan Red List information is still far from comprehensive. For instance, De Roeck et al. (2007) reported that only two of the 14 anostracan species known to occur in the Western Cape province of SA were listed by IUCN and that insufficient data were available to determine the IUCN Red Data Category of six of these species. Of the 46 anostracan species reported for southern Africa by Hamer & Brendonck (1997), two were considered by IUCN (1994) as ‘Critically Endangered’, four were ‘Endangered’, three were ‘Vulnerable’ and 24 were ‘Least Concern’, whilst data were deficient for the categorisation of 13 species. As is the trend worldwide, the anostracans of the region are threatened by agriculture and urbanization, due to both indirect (pollution and pesticides) and direct (physical) habitat alteration arising from these activities (Brendonck et al., 2008). These factors are thought to have driven the extinction of *Streptocephalus gracilis* Sars, 1898 and *Branchipodopsis karroensis* Barnard, 1929 (De Roeck et al., 2007).

We hypothesize that temporary wetland specialists such as the branchiopods (including cladocerans), ostracods and copepods, which tend to be more habitat-restricted and less capable of escaping degradation, are more likely to be affected by human impacts than for taxa that are able to easily disperse to other more favourable areas (e.g. dipterans, hemipterans, odonates and coleopterans). The sensitivity of the region’s temporary wetland invertebrates to human impacts such as pollution and habitat alteration is a topic still under investigation, with some evidence indicating a resilient fauna (Bird et al., 2013), whilst other evidence is in favour of a more sensitive fauna (Bird & Day, 2016), especially when effects are combined with climate change impacts (Dalu et al., 2017a). Whatever the case may be, what is certain is that temporary wetland habitats worldwide have been degraded and destroyed at an alarming rate (Semlitsch & Bodie, 1998), and various studies indicate that the situation is no different in southern Africa (De Roeck et al., 2007; Bird & Day, 2014; Dalu et al., 2017a).

In terms of the way forward, we cannot expect to properly understand and conserve temporary wetlands if we don't know what lives in them. As things stand, this is certainly the case in southern Africa in that we lack updated taxonomic information for many groups that inhabit these systems. This review shows that we have a base of knowledge on which to work, but the now famous stagnation of taxonomic expertise both locally and abroad (see Godfray, 2002; Wheeler et al., 2004; Agnarsson & Kuntner, 2007; Bebbier et al., 2014) is a major threat hindering the further progression of this knowledge. In addition to providing a synthesis for southern Africa, this work highlights the dearth of fundamental taxonomic information on the invertebrate fauna of temporary wetlands in the region, and constitutes a plea for more foundational taxonomic work to be initiated.

Acknowledgements

We acknowledge use of infrastructure and equipment provided by the NRF-SAIAB Research Platform and the funding channelled through the NRF-SAIAB Institutional Support system. This study was partially funded by the National Research Foundation (Grant No. 110507). Any opinions, findings, conclusions or recommendations expressed in this material are those of the authors, and the NRF does not accept any liability in this regard.

References

- Abebe, E., I. Andrassy & W. Traunspurger, 2006. Freshwater nematodes: ecology and taxonomy. CABI Publishing, Wallingford, UK.
- Abebe, E., W. Decraemer & P. De Ley, 2008. Global diversity of nematodes (Nematoda) in freshwater. *Hydrobiologia* 595(1): 67-78.
- Agnarsson, I. & M. Kuntner, 2007. Taxonomy in a changing world: seeking solutions for a science in crisis. *Systematic Biology* 56(3): 531-539.
- Anlauf, A., 1990. Cyst formation of *Tubifex tubifex* (Müller)—an adaptation to survive food deficiency and drought. *Hydrobiologia* 190(1): 79-82.
- Anusa, A., H. Ndagurwa & C. Magadza, 2012. The influence of pool size on species diversity and water chemistry in temporary rock pools on Domboshawa Mountain, northern Zimbabwe. *African Journal of Aquatic Science* 37(1): 89-99.
- Appleton, C., 2002a. Platyhelminthes. In Day, J. & I. de Moor (eds) Guides to the freshwater invertebrates of southern Africa Volume 5: Non-Arthropods. WRC Report No. TT 167/02, Water Research Commission, Pretoria, 88-110.
- Appleton, C., 2002b. Mollusca. In Day, J. & I. de Moor (eds) Guides to the freshwater invertebrates of southern Africa Volume 6: Arachnida and Mollusca. WRC Report No. TT 182/02, Water Research Commission, Pretoria, pp. 42-125.
- Appleton, C., B. Sharp & D. le Sueur, 1995. Wetlands and water-related parasitic diseases of man in Southern Africa. In Cowan, G. I. (ed) Wetlands of South Africa. Department of Environmental Affairs and Tourism, Pretoria, South Africa., 227-246.
- Appleton, C. C., B. A. Curtis, L. E. Alonso & J. Kipping, 2003. Freshwater invertebrates of the Okavango Delta, Botswana. In: Alonso, L. E. & L.-A. Nordin (eds) RAP Bulletin of Biological Assessment. vol 27, 58-69.
- Armitage, P. D., 1995. Behaviour and ecology of adults. In Armitage, P. D., P. S. Cranston & L. C. V. Pinder (eds) The Chironomidae: biology and ecology of non-biting midges. Springer Netherlands, Dordrecht, 194-224.
- Artois, T., W. Willems, E. De Roeck, M. Jocqué & L. Brendonck, 2004. Freshwater Rhabdocoela (Platyhelminthes) from ephemeral rock pools from Botswana, with the description of four new species and one new genus. *Zoological Science* 21(10): 1063-1072.
- Balian, E., H. Segers, K. Martens & C. Lévêque, 2008. The freshwater animal diversity assessment: an overview of the results. *Hydrobiologia* 595(1): 627-637.

- 1338 Barber-James, H. M. & J.-L. Gattolliat, 2012. How well are Afrotropical mayflies known?
 1339 Status of current knowledge, practical applications, and future directions. *Inland*
 1340 *Waters* 2(1): 1-9.
- 1341 Barber-James, H. M., J.-L. Gattolliat, M. Sartori & M. D. Hubbard, 2008. Global diversity of
 1342 mayflies (Ephemeroptera, Insecta) in freshwater. *Hydrobiologia* 595(1): 339-350.
- 1343 Barber-James, H. M. & C. Lugo-Ortiz, 2003. Ephemeroptera. In de Moor, I., J. Day & F. de
 1344 Moor (eds) *Guides to the Freshwater Invertebrates of southern Africa*. vol 7: Insecta I
 1345 - Ephemeroptera, Odonata and Plecoptera. WRC Report no. TT 207/03, Water
 1346 Research Commission, Pretoria, 16-142.
- 1347 Barnard, K., 1926. A study of the freshwater isopodan and amphipodan Crustacea of South
 1348 Africa. *Transactions of the Royal Society of South Africa* 14(1): 139-215.
- 1349 Batzer, D. & D. Boix, 2016. *Invertebrates in freshwater wetlands: an international*
 1350 *perspective on their ecology*. Springer, Switzerland.
- 1351 Batzer, D. P., R. B. Rader & S. A. Wissinger, 1999. *Invertebrates in freshwater wetlands of*
 1352 *North America: ecology and management*. John Wiley and Sons.
- 1353 Baxevanis, A. D., S. Maniatsi, D. Kouroupis, K. Marathiotis, I. Kappas, H. Kaiser & T. J.
 1354 Abatzopoulos, 2014. Genetic identification of South African *Artemia* species:
 1355 invasion, replacement and co-occurrence. *Journal of the Marine Biological*
 1356 *Association of the United Kingdom* 94(04): 775-785.
- 1357 Bebbier, D. P., J. R. Wood, C. Barker & R. W. Scotland, 2014. Author inflation masks global
 1358 capacity for species discovery in flowering plants. *New Phytologist* 201(2): 700-706.
- 1359 Belk, D., H. Dumont & N. Munuswamy, 1991. *Studies on large branchiopod biology and*
 1360 *aquaculture*. Kluwer Academic, Belgium.
- 1361 Bilton, D. T., 2013. *Prosthetops wolfbergensis* sp. nov.—a giant amongst the ‘minute moss
 1362 beetles’, with new data on other members of the genus (Coleoptera, Hydraenidae).
 1363 *Zootaxa* 3666(3): 345-357.
- 1364 Bilton, D. T., 2017. Water beetles from the Bokkeveld Plateau: a semi-arid hotspot of
 1365 freshwater biodiversity in the Northern Cape of South Africa. *Zootaxa* 4268(2): 191-
 1366 214.
- 1367 Bilton, D. T., J. R. Freeland & B. Okamura, 2001. Dispersal in freshwater invertebrates.
 1368 *Annual Review of Ecology and Systematics* 32(1): 159-181.
- 1369 Bilton, D. T., E. F. Toussaint, C. R. Turner & M. Balke, 2015. *Capelatus prykei* gen. et sp.
 1370 n.(Coleoptera: Dytiscidae: Copelatinae) – a phylogenetically isolated diving beetle
 1371 from the Western Cape of South Africa. *Systematic Entomology* 40(3): 520-531.

1372 Bird, M. & J. Day, 2014. Wetlands in changed landscapes: the influence of habitat
 1373 transformation on the physico-chemistry of temporary depression wetlands. PLoS
 1374 ONE 9(2): e88935.

1375 Bird, M. S., 2012. Effects of habitat transformation on temporary wetlands in the south-
 1376 western Cape, South Africa. PhD thesis, Zoology Department, University of Cape
 1377 Town.

1378 Bird, M. S. & J. A. Day, 2016. Impacts of terrestrial habitat transformation on temporary
 1379 wetland invertebrates in a sclerophyllous Sand fynbos landscape. Hydrobiologia
 1380 782(1): 169-185.

1381 Bird, M. S., M. C. Mlambo & J. A. Day, 2013. Macroinvertebrates as unreliable indicators of
 1382 human disturbance in temporary depression wetlands of the south-western Cape,
 1383 South Africa. Hydrobiologia 720: 19-37 doi:10.1007/s10750-013-1618-2.

1384 Biström, O. & A. N. Nilsson, 2006. Taxonomic revision of the Ethiopian genus *Canthyporus*
 1385 (Coleoptera Dytiscidae). Memorie della Societa Entomologica Italiana 85(1): 209-
 1386 306.

1387 Boettger, K., 1974. The biology of *Sphaerodema grassei ghesquierei*. Studies of central
 1388 African belostomatids (Heteroptera, Insecta) Archiv für Hydrobiologie 74: 100-122.

1389 Bohonak, A. J., B. P. Smith & M. Thornton, 2004. Distributional, morphological and genetic
 1390 consequences of dispersal for temporary pond water mites. Freshwater Biology 49(2):
 1391 170-180.

1392 Boxshall, G. A. & D. Defaye, 2008. Global diversity of copepods (Crustacea: Copepoda) in
 1393 freshwater. Hydrobiologia 595(1): 195-207.

1394 Brain, C., 2002. Rotifera. In Day, J. & I. de Moor (eds) Guides to the freshwater invertebrates
 1395 of southern Africa Volume 5: Non-Arthropods. WRC Report No. TT 167/02, Water
 1396 Research Commission, Pretoria, 116-135.

1397 Brendonck, L., 1995. A new branchipodid genus and species (Crustacea: Branchiopoda:
 1398 Anostraca) from South Africa. Zoological Journal of the Linnean Society 115(4): 359-
 1399 372.

1400 Brendonck, L., 1996. Diapause, quiescence, hatching requirements: what we can learn from
 1401 large freshwater branchiopods (Crustacea: Branchiopoda: Anostraca, Notostraca,
 1402 Conchostraca). Hydrobiologia 320(1-3): 85-97.

1403 Brendonck, L., 1999. Conchostraca. In Day, J., I. de Moor, B. Stewart & A. Louw (eds)
 1404 Guides to the freshwater invertebrates of southern Africa Volume 2: Crustacea I -

1405 Notostraca, Anostraca, Conchostraca and Cladocera. WRC Report No. TT 121/00,
 1406 Water Research Commission, Pretoria, 59-80.

1407 Brendonck, L. & A. Coomans, 1994. Egg morphology in African Streptocephalidae
 1408 (Crustacea: Branchiopoda: Anostraca). Part 1: South of Zambezi and Kunene rivers.
 1409 Archiv für Hydrobiologie 3(Suppl. 99 (Monographische Beiträge)): 313-334.

1410 Brendonck, L. & L. De Meester, 2003. Egg banks in freshwater zooplankton: evolutionary
 1411 and ecological archives in the sediment. Hydrobiologia 491(1-3): 65-84.

1412 Brendonck, L., L. De Meester & B. J. Riddoch, 2000a. Regional structuring of genetic
 1413 variation in short-lived rock pool populations of *Branchipodopsis wolffi* (Crustacea:
 1414 Anostraca). Oecologia 123(4): 506-515.

1415 Brendonck, L., M. Hamer, B. Riddoch & M. Seaman, 2000b. *Branchipodopsis* species—
 1416 specialists of ephemeral rock pools. African Journal of Aquatic Science 25(1): 98-
 1417 104.

1418 Brendonck, L., E. Michels, L. De Meester & B. Riddoch, 2002. Temporary pools are not
 1419 'enemy-free'. Hydrobiologia 486(1): 147-159.

1420 Brendonck, L. & B. Riddoch, 1997. Anostracans (Branchiopoda) of Botswana: morphology,
 1421 distribution, diversity, and endemism. Journal of Crustacean Biology 17(1): 111-134.

1422 Brendonck, L., B. Riddoch, V. Van de Weghe & T. Van Dooren, 1998. The maintenance of
 1423 egg banks in very short-lived pools—a case study with anostracans (Branchiopoda).
 1424 Archiv für Hydrobiologie 52(Special issues)): 141-161.

1425 Brendonck, L. & B. J. Riddoch, 1999. Wind-borne short-range egg dispersal in anostracans
 1426 (Crustacea: Branchiopoda). Biological Journal of the Linnean Society 67(1): 87-95.

1427 Brendonck, L., D. C. Rogers, J. Olesen, S. Weeks & W. R. Hoeh, 2008. Global diversity of
 1428 large branchiopods (Crustacea: Branchiopoda) in freshwater. Hydrobiologia 595(1):
 1429 167-176.

1430 Brendonck, L. & W. D. Williams, 2000. Biodiversity in wetlands of dry regions (drylands).
 1431 In Gopal, B., W. J. Junk & J. A. Davis (eds) Biodiversity in wetlands: assessment,
 1432 function and conservation Volume 1. Backhuys Publishers, Leiden, The
 1433 Netherlands, 181-194.

1434 Brittain, J. E. & M. Sartori, 2003. Ephemeroptera. In Resh, V. H. & R. T. Carde' (eds)
 1435 Encyclopedia of Insects. Academic Press, Amsterdam, 373-380.

1436 Brown, D. S., 1994. Freshwater snails of Africa and their medical importance. Taylor &
 1437 Francis Ltd, London, UK.

- 1438 Brucet, S., D. Boix, R. López-Flores, A. Badosa, R. Moreno-Amich & X. D. Quintana, 2005.
1439 Zooplankton structure and dynamics in permanent and temporary Mediterranean salt
1440 marshes: taxon-based and size-based approaches. *Archiv für Hydrobiologie* 162(4):
1441 535-555.
- 1442 Calhoun, A. J., D. M. Mushet, K. P. Bell, D. Boix, J. A. Fitzsimons & F. Isselin-Nondedeu,
1443 2017. Temporary wetlands: challenges and solutions to conserving a
1444 ‘disappearing’ ecosystem. *Biological Conservation* 211: 3-11.
- 1445 Capinera, J. L., 2008. *Encyclopedia of Entomology*. 2nd ed. Springer Science & Business
1446 Media. 4346 pp.
- 1447 Chakona, A., C. Phiri, C. H. Magadza & L. Brendonck, 2008. The influence of habitat
1448 structure and flow permanence on macroinvertebrate assemblages in temporary rivers
1449 in northwestern Zimbabwe. *Hydrobiologia* 607(1): 199-209.
- 1450 Clark, T. E. & M. J. Samways, 1996. Dragonflies (Odonata) as indicators of biotope quality
1451 in the Kruger National Park, South Africa. *Journal of Applied Ecology* 33(5): 1001-
1452 1012.
- 1453 Clausnitzer, V., K.-D. B. Dijkstra, R. Koch, J.-P. Boudot, W. R. Darwall, J. Kipping, B.
1454 Samraoui, M. J. Samways, J. P. Simaika & F. Suhling, 2012. Focus on African
1455 freshwaters: hotspots of dragonfly diversity and conservation concern. *Frontiers in*
1456 *Ecology and the Environment* 10(3): 129-134.
- 1457 Coetzee, M., 2002. Culicidae. In Day, J. & I. de Moor (eds) *Guides to the freshwater*
1458 *invertebrates of southern Africa Vol 9: Diptera*. WRC Report No. TT 201/02, Water
1459 Research Commission, Pretoria, 57-74.
- 1460 Collinson, N., J. Biggs, A. Corfield, M. Hodson, D. Walker, M. Whitfield & P. Williams,
1461 1995. Temporary and permanent ponds: an assessment of the effects of drying out on
1462 the conservation value of aquatic macroinvertebrate communities. *Biological*
1463 *Conservation* 74(2): 125-133.
- 1464 Corrêa, D. D., 1951. Freshwater Nemertines from the Amazon region and from South Africa.
1465 Universidade de Sao Paulo Boletins da Faculdade de Filosofia, Ciencias e Letras 16:
1466 257-269.
- 1467 Cranston, P. S., 2014. A new putatively cryptobiotic midge, *Polypedilum ovahimba* sp.
1468 nov.(Diptera: Chironomidae), from southern Africa. *Austral Entomology* 53(4): 373-
1469 379.
- 1470 Cumberlidge, N., 2009. The status and distribution of freshwater crabs. In Smith, K. G., M.
1471 D. Diop, M. Niane & W. R. T. Darwell (eds) *The status and distribution of freshwater*

1472 biodiversity in West Africa: IUCN and Wetlands International Report. Gland,
 1473 Switzerland and Cambridge, UK, 56-72.
 1474 Curtis, B., 1991. Freshwater macro-invertebrates of Namibia. *Madoqua* 17(2): 163-187.
 1475 Dahms, H.-U., 1995. Dormancy in the Copepoda—an overview. *Hydrobiologia* 306(3): 199-
 1476 211.
 1477 Dalu, T., R. J. Wasserman & M. T. Dalu, 2017a. Agricultural intensification and drought
 1478 frequency increases may have landscape-level consequences for ephemeral
 1479 ecosystems. *Global Change Biology* 23(3): 983-985.
 1480 Dalu, T., R. J. Wasserman, P. W. Froneman & O. L. Weyl, 2017b. Trophic isotopic carbon
 1481 variation increases with pond's hydroperiod: evidence from an Austral ephemeral
 1482 ecosystem. *Scientific Reports* 7: 7572.
 1483 Dalu, T., R. J. Wasserman, T. J. Vink & O. L. Weyl, 2017c. Sex and species specific isotopic
 1484 niche specialisation increases with trophic complexity: evidence from an ephemeral
 1485 pond ecosystem. *Scientific Reports* 7: 43229.
 1486 Dalu, T., O. L. Weyl, P. W. Froneman & R. J. Wasserman, 2016. Trophic interactions in an
 1487 austral temperate ephemeral pond inferred using stable isotope analysis.
 1488 *Hydrobiologia* 768(1): 81-94.
 1489 Daniels, S. R., E. E. Phiri & J. Bayliss, 2014. Renewed sampling of inland aquatic habitats in
 1490 southern Africa yields two novel freshwater crab species (Decapoda: Potamonautidae:
 1491 *Potamonautes*). *Zoological Journal of the Linnean Society* 171(2): 356-369.
 1492 da Silva, J.M. & S. Willows-Munro, 2016. A review of over a decade of DNA barcoding in
 1493 South Africa: A faunal perspective. *African Zoology* 51(1): 1-12.
 1494 Davies, R. W., L. Linton & F. Wrona, 1982. Passive dispersal of four species of freshwater
 1495 leeches (Hirudinoidea) by ducks. *Freshwater Invertebrate Biology* 1: 40-44.
 1496 Davis, C. L., 2011. Climate risk and vulnerability: a handbook for southern Africa. Council
 1497 for Scientific and Industrial Research, Pretoria, South Africa, 92pp.
 1498 Day, J., 1990. Environmental correlates of aquatic faunal distribution in the Namib Desert. In
 1499 Seely, M. K. (ed) *Namib ecology: 25 years of Namib research*. Transvaal Museum
 1500 Monograph No. 7. Transvaal Museum, Pretoria, 99-107.
 1501 Day, J., E. Day, V. Ross-Gillespie & A. Ketley, 2010. The assessment of temporary wetlands
 1502 during dry conditions. WRC Report TT434/09, Water Research Commission,
 1503 Pretoria.

- Day, J. & J. Day, 2002. Polychaeta. In Day, J. & I. de Moor (eds) Guides to the Freshwater Invertebrates of southern Africa. vol Volume 5: Non-Arthropods. WRC Report No. TT 167/02, Water Research Commission, Pretoria, 193-202.
- De Block, M., M. A. McPeck & R. Stoks, 2008. Stronger compensatory growth in a permanent-pond *Lestes* damselfly relative to temporary-pond *Lestes*. *Oikos* 117(2): 245-254.
- de Klerk, A. R. & V. Wepener, 2013. Macroinvertebrate assemblage changes as an indicator of water quality of perennial endorheic reed pans on the Mpumalanga highveld, South Africa. *Journal of Environmental Protection* 4: 10-21.
- de Meillon, B. & W. Wirth, 2002. Ceratopogonidae. In Day, J. & I. de Moor (eds) Guides to the freshwater invertebrates of southern Africa Vol 9: Diptera. WRC Report No. TT 201/02, Water Research Commission, Pretoria, 50-56.
- de Moor, F. & J. Day, 2013. Aquatic biodiversity in the mediterranean region of South Africa. *Hydrobiologia* 719(1): 237-268.
- de Moor, F. & K. Scott, 2003. Trichoptera. In de Moor, I., J. Day & F. C. De Moor (eds) Guides to the freshwater invertebrates of southern Africa Volume 8: Insecta II - Hemiptera, Megaloptera, Neuroptera, Trichoptera and Lepidoptera. vol 8. WRC Report No. TT 214/03, 84-181.
- de Necker, L., M. Ferreira, J. J. van Vuren & W. Malherbe, 2016. Aquatic invertebrate community structure of selected endorheic wetlands (pans) in South Africa. *Inland Waters* 6(3): 303-313.
- De Roeck, E., A. Waterkeyn & L. Brendonck, 2010. Life-history traits of *Streptocephalus purcelli* Sars, 1898 (Branchiopoda, Anostraca) from temporary waters with different phenology. *Water SA* 36(3): 323-328.
- De Roeck, E. R., T. Artois & L. Brendonck, 2005. Consumptive and non-consumptive effects of turbellarian (*Mesostoma* sp.) predation on anostracans. *Hydrobiologia* 542(1): 103-111.
- De Roeck, E. R., B. J. Vanschoenwinkel, J. A. Day, Y. Xu, L. Raitt & L. Brendonck, 2007. Conservation status of large branchiopods in the Western Cape, South Africa. *Wetlands* 27(1): 162-173.
- Dettner, K., 1982. Description of the larvae of *Acilius duvergeri* (Coleoptera, Dytiscidae), with keys to larvae of European species of genus *Acilius* and of the European genera of subfamily Dytiscinae. *Aquatic Insects* 4(2): 81-88.

- 1537 Di Sabatino, A., H. Smit, R. Gerecke, T. Goldschmidt, N. Matsumoto & B. Cicolani, 2008.
 1538 Global diversity of water mites (Acari, Hydrachnidia; Arachnida) in freshwater.
 1539 Hydrobiologia 595: 303-315.
- 1540 Dijkstra, K.-D. B., 2003. A review of the taxonomy of African Odonata: finding ways to
 1541 better identification and biogeographic insight. Cimbebasia 18: 191-206.
- 1542 Dijkstra, K.-D. B., J. Kipping & N. Mézière, 2015. Sixty new dragonfly and damselfly
 1543 species from Africa (Odonata). Odonatologica 44(4): 447-678.
- 1544 Dodson, S. I., 1987. Animal assemblages in temporary desert rock pools: aspects of the
 1545 ecology of *Dasyhelea sublettei* (Diptera: Ceratopogonidae). Journal of the North
 1546 American Benthological Society 6(1): 65-71.
- 1547 Dumont, H. & S. Negrea, 2002. Introduction to the class Branchiopoda. In Dumont, H. J. (ed)
 1548 Guides to the Identification of the Microinvertebrates of the Continental Waters of the
 1549 World 19. Backhuys Publishers, Leiden.
- 1550 Dumont, H. J., 1983. Biogeography of rotifers. Hydrobiologia 104: 19-30.
- 1551 Escalera-Vázquez, L. H. & L. Zambrano, 2010. The effect of seasonal variation in abiotic
 1552 factors on fish community structure in temporary and permanent pools in a tropical
 1553 wetland. Freshwater Biology 55(12): 2557-2569.
- 1554 Farnesi, L. C., R. F. S. Menna-Barreto, A. J. Martins, D. Valle & G. L. Rezende, 2015.
 1555 Physical features and chitin content of eggs from the mosquito vectors *Aedes aegypti*,
 1556 *Anopheles aquasalis* and *Culex quinquefasciatus*: Connection with distinct levels of
 1557 resistance to desiccation. Journal of Insect Physiology 83: 43-52.
- 1558 Ferreira, M., V. Wepener & J. J. Van Vuren, 2011. The occurrence of large branchiopod
 1559 crustaceans in perennial pans: a research note. African Zoology 46(1): 176-178.
- 1560 Ferreira, M., V. Wepener & J. J. van Vuren, 2012. Aquatic invertebrate communities of
 1561 perennial pans in Mpumalanga, South Africa: a diversity and functional approach.
 1562 African Invertebrates 53(2): 751-768.
- 1563 Forero, D., 2008. The systematics of the Hemiptera. Revista Colombiana de Entomologia
 1564 34(1): 1-21.
- 1565 Forró, L., N. Korovchinsky, A. Kotov & A. Petrusek, 2008. Global diversity of cladocerans
 1566 (Cladocera; Crustacea) in freshwater. Hydrobiologia 595(1): 177-184.
- 1567 Foster, G. N., D. T. Bilton & B. Nelson, 2016. Atlas of the predaceous water beetles
 1568 (Hydradephaga) of Britain and Ireland. Field Studies Council, UK.

1569 Foster, L., W. Malherbe, M. Ferreira & J. J. van Vuren, 2015. Macroinvertebrate variation in
 1570 endorheic depression wetlands in North West and Mpumalanga provinces, South
 1571 Africa. *African Journal of Aquatic Science* 40(3): 287-297.
 1572 Frey, D., 1993. The penetration of cladocerans into saline waters. *Hydrobiologia* 267: 233-
 1573 248.
 1574 Frost, T. M., 1991. Porifera. In Thorpe, J. H. & A. P. Covich (eds) *Ecology and Classification*
 1575 *of North American Freshwater Invertebrates*. Academic Press, New York, 95-124.
 1576 Frouz, J., J. Matěna & A. Ali, 2003. Survival strategies of chironomids (Diptera:
 1577 Chironomidae) living in temporary habitats: a review. *European Journal of*
 1578 *Entomology* 100(4): 459-466.
 1579 Fryer, G., 1988. Studies on the functional morphology and biology of the Notostraca
 1580 (Crustacea: Branchiopoda). *Philosophical Transactions of the Royal Society of*
 1581 *London B: Biological Sciences* 321(1203): 27-124.
 1582 Garey, J. R., S. J. McInnes & P. B. Nichols, 2008. Global diversity of tardigrades
 1583 (Tardigrada) in freshwater. *Hydrobiologia* 595(1): 101-106.
 1584 Gerecke, R., 2004. Taxonomy and phylogeny in African water mites of the genus
 1585 *Diplodontus* Dugés, 1834 (Acari, Hydrachnidia, Hydryphantidae). *Annales de*
 1586 *Limnologie-International Journal of Limnology* 40(1): 71-85.
 1587 Gillies, M., 1990. A revision of the African species of *Centroptilum* Eaton (Baetidae,
 1588 Ephemeroptera). *Aquatic Insects* 12(2): 97-128.
 1589 Godfray, H. C. J., 2002. Challenges for taxonomy. *Nature* 417(6884): 17-19.
 1590 Goldschmidt, T. & H. Smit, 2009. Studies on torrenticolid water mites mainly from South
 1591 Africa—*Torrenticola* Piersig, 1896 and *Monatractides* K. Viets, 1926 (Acari:
 1592 Hydrachnidia: Torrenticolidae). *International Journal of Acarology* 35(3): 179-234.
 1593 Goudie, A. & G. Wells, 1995. The nature, distribution and formation of pans in arid zones.
 1594 *Earth-Science Reviews* 38(1): 1-69.
 1595 Gouws, G., B. A. Stewart & C. A. Matthee, 2005. Lack of taxonomic differentiation in an
 1596 apparently widespread freshwater isopod morphotype (Phreatoicidea:
 1597 Mesamphisopidae: Mesamphisopus) from South Africa. *Molecular Phylogenetics and*
 1598 *Evolution* 37(1): 289-305.
 1599 Grimaldi, D. & M. S. Engel, 2005. *Evolution of the Insects*. Cambridge University Press.
 1600 Gueriau, P., N. Rabet, G. Clément, L. Lagebro, J. Vannier, D. E. Briggs, S. Charbonnier, S.
 1601 Olive & O. Béthoux, 2016. A 365-million-year-old freshwater community reveals

1602 morphological and ecological stasis in branchiopod crustaceans. *Current Biology*
 1603 26(3): 383-390.

1604 Hamer, M., 1999. Anostraca. In Day, J., I. de Moor, B. Stewart & A. Louw (eds) *Guides to*
 1605 *the freshwater invertebrates of southern Africa Volume 2: Crustacea I - Notostraca,*
 1606 *Anostraca, Conchostraca and Cladocera.* WRC Report No. TT 121/00, Water
 1607 Research Commission, Pretoria, 14-58.

1608 Hamer, M., 2013. A national strategy for zoological taxonomy (2013-2020). South African
 1609 National Biodiversity Institute, Pretoria, South Africa: 1-53.

1610 Hamer, M. & C. Appleton, 1991. Physical and chemical characteristics and phyllopod fauna
 1611 of temporary pools in north-eastern Natal, Republic of South Africa. *Hydrobiologia*
 1612 212: 95-104.

1613 Hamer, M. & L. Brendonck, 1997. Distribution, diversity and conservation of Anostraca
 1614 (Crustacea: Branchiopoda) in southern Africa. *Hydrobiologia* 359: 1-12.

1615 Hamer, M., L. Brendonck, A. Coomans & C. Appleton, 1994. A review of African
 1616 Streptocephalidae (Crustacea: Branchiopoda: Anostraca). Part 1: South of Zambezi
 1617 and Kunene rivers. *Archiv für Hydrobiologie* 3(Suppl. 99 (Monographische
 1618 Beiträge)): 235-277.

1619 Hamer, M. & K. Martens, 1998. The large Branchiopoda (Crustacea) from temporary habitats
 1620 of the Drakensberg region, South Africa. *Hydrobiologia* 384(1-3): 151-165.

1621 Hamer, M. & N. Rayner, 1995. A note on the taxonomy and distribution of *Triops* Schrank
 1622 (Crustacea: Branchiopoda: Notostraca) in southern Africa. *Annals of the Natal*
 1623 *Museum* 36: 9-19.

1624 Hamer, M. & N. Rayner, 1996. A note on the unusual crustacean community of a temporary
 1625 pool in the Northern Cape. *Southern African Journal of Aquatic Science* 22(1-2): 100-
 1626 104.

1627 Hamer, M. L., 1989. Studies on the phyllopod fauna of ephemeral pools in north-eastern
 1628 Natal. MSc thesis, University of Natal, Pietermaritzburg, South Africa. 192 pp.

1629 Harris, P. M., B. R. Roosa & L. Norment, 2002. Underground dispersal by amphipods
 1630 (*Crangonyx pseudogracilis*) between temporary ponds. *Journal of Freshwater*
 1631 *Ecology* 17(4): 589-594.

1632 Harrison, A. D., 2002. Chironomidae. In Day, J. & I. de Moor (eds) *Guides to the freshwater*
 1633 *invertebrates of southern Africa Volume 9: Diptera.* WRC Report No. TT 201/02,
 1634 Water Research Commission, Pretoria.

- 1635 Hassall, C. & D. J. Thompson, 2008. The effects of environmental warming on Odonata: a
1636 review. *International Journal of Odonatology* 11(2): 131-153.
- 1637 Heeg, J., 2002a. Porifera. In Day, J. & I. de Moor (eds) *Guides to the Freshwater*
1638 *Invertebrates of southern Africa*. vol 5: Non-Arthropods, WRC Report No. TT
1639 167/02, Water Research Commission, Pretoria, 59-73.
- 1640 Heeg, J., 2002b. Gastrotricha. In Day, J. & I. de Moor (eds) *Guides to the Freshwater*
1641 *Invertebrates of southern Africa*. vol 5: Non-Arthropods. WRC Report No. TT
1642 167/02, Water Research Commission, Pretoria, 166-172.
- 1643 Heeg, J., 2002c. Bryozoa. In Day, J. & I. de Moor (eds) *Guides to the Freshwater*
1644 *Invertebrates of southern Africa*. vol 5: Non-Arthropods. WRC Report No. TT
1645 167/02, Water Research Commission, Pretoria, 173-188.
- 1646 Heyns, J., 2002. Checklist of free living nematodes recorded from freshwater habitats in
1647 Southern Africa. *Water SA* 28(4): 449-456.
- 1648 Hinton, H. E., 1953. Some adaptations of insects to environments that are alternately dry and
1649 flooded, with some notes on the habits of the Stratiomyidae. *Transactions of the*
1650 *Society for British Entomology* 11: 209-227.
- 1651 Hinton, H. E., 1960. A fly larva that tolerates dehydration and temperatures of -270° to +
1652 102° C. *Nature* 188: 336-337.
- 1653 Holsinger, J. R. & G. W. Dickson, 1977. Burrowing as a means of survival in the troglobitic
1654 amphipod crustacean *Crangonyx antennatus* Packard (Crangonyctidae).
1655 *Hydrobiologia* 54(3): 195-199.
- 1656 Horne, D. & K. Martens, 1998. An assessment of the importance of resting eggs for the
1657 evolutionary success of Mesozoic non-marine cypridoidean Ostracoda (Crustacea).
1658 *Archiv für Hydrobiologie* 52: 549-561.
- 1659 Houben, A. M., N. Van Steenkiste & T. Artois, 2014. Revision of *Phaenocora* Ehrenberg,
1660 1836 (Rhabditophora, Typhloplanidae, Phaenocorinae) with the description of two
1661 new species. *Zootaxa* 3889(3): 301-354.
- 1662 Hutchinson, G. E., 1933. The zoo-geography of the African aquatic Hemiptera in relation to
1663 past climatic change. *Internationale Revue der gesamten Hydrobiologie und*
1664 *Hydrographie* 28: 436-488.
- 1665 Hynes, H., 1955. Biological notes on some East African aquatic Heteroptera. *Physiological*
1666 *Entomology* 30(4-6): 43-54.
- 1667 Jäch, M. A. & M. Balke, 2008. Global diversity of water beetles (Coleoptera) in freshwater.
1668 *Hydrobiologia* 595(1): 419-442.

1669 Jannot, J. E., 2009. Life history plasticity and fitness in a caddisfly in response to proximate
1670 cues of pond-drying. *Oecologia* 161(2): 267-277.

1671 Jannot, J. E., S. A. Wissinger & J. R. Lucas, 2008. Diet and a developmental time constraint
1672 alter life-history trade-offs in a caddis fly (Trichoptera: Limnephilidae). *Biological*
1673 *Journal of the Linnean Society* 95(3): 495-504.

1674 Jansen van Rensburg, C., 1976. An identification key to the water mite families of the
1675 Ethiopian region. *Journal of the Limnological Society of Southern Africa* 2(1): 11-20.

1676 Jansen van Rensburg, C. & J. Day, 2002. Water mites In Day, J. & I. de Moor (eds) *Guides to*
1677 *the Freshwater Invertebrates of southern Africa. vol 6: Arachnida and Mollusca.*
1678 *WRC Report no. TT 182/02, Water Research Commission, Pretoria, 23-41.*

1679 Jeffries, M., 2005. Small ponds and big landscapes: the challenge of invertebrate spatial and
1680 temporal dynamics for European pond conservation. *Aquatic Conservation: Marine*
1681 *and Freshwater Ecosystems* 15(6): 541-547.

1682 Jocqué, M., L. Brendonck, B. J. Riddoch & K. Martens, 2010. On *Amphibolocypis arida* sp.
1683 nov.(Crustacea, Ostracoda), from rock pools in Botswana (southern Africa). *Zootaxa*
1684 2408: 47-58.

1685 Jocqué, M., K. Martens, B. Riddoch & L. Brendonck, 2006. Faunistics of ephemeral rock
1686 pools in southeastern Botswana. *Archiv für Hydrobiologie* 165(3): 415-431.

1687 Johansson, F. & F. Suhling, 2004. Behaviour and growth of dragonfly larvae along a
1688 permanent to temporary water habitat gradient. *Ecological Entomology* 29(2): 196-
1689 202.

1690 Jones, M., 2002. Developing a classification system for Western Cape wetlands. MSc thesis,
1691 Zoology Department, University of Cape Town.

1692 Jupp, P., B. McIntosh & E. Nevill, 1980. A survey of the mosquito and Culicoides faunas at
1693 two localities in the Karoo region of South Africa with some observations of
1694 bionomics. *The Onderstepoort journal of veterinary research* 47(1): 1-6.

1695 Kaiser, H., A. Gordon & T. Paulet, 2006. Review of the African distribution of the brine
1696 shrimp genus *Artemia*. *Water SA* 32(4): 597-603.

1697 Kalkman, V. J., V. Clausnitzer, K.-D. B. Dijkstra, A. G. Orr, D. R. Paulson & J. van Tol,
1698 2008. Global diversity of dragonflies (Odonata) in freshwater. *Hydrobiologia* 595(1):
1699 351-363.

1700 Kipping, J., 2010. The dragonflies and damselflies of Botswana: an annotated checklist with
1701 notes on distribution, phenology, habitats and Red List status of the species (Insecta:
1702 Odonata). *Mauritiana (Altenberg)* 21: 126-204.

1703 Kneitel, J. M., 2016. Climate-driven habitat size determines the latitudinal diversity gradient
1704 in temporary ponds. *Ecology* 97(4): 961-968.

1705 Korn, M. & A. K. Hundsdoerfer, 2006. Evidence for cryptic species in the tadpole shrimp
1706 *Triops granarius* (Lucas, 1864)(Crustacea: Notostraca). *Zootaxa* 1257: 57-68.

1707 Korovchinsky, N., 2006. The Cladocera (Crustacea: Branchiopoda) as a relict group.
1708 *Zoological Journal of the Linnean Society* 147(1): 109-124.

1709 Kristensen, T. K., C. C. Appleton, B. Curtis & A.-S. Stensgaard, 2009. The status and
1710 distribution of freshwater molluscs. In Darwall, W., D. Tweddle, K. Smith & P.
1711 Skelton (eds) *The status and distribution of freshwater biodiversity in southern Africa*.
1712 IUCN, Gland, Switzerland, 38-47.

1713 Lahr, J., 1997. Ecotoxicology of organisms adapted to life in temporary freshwater ponds in
1714 arid and semi-arid regions. *Archives of Environmental Contamination and Toxicology*
1715 32(1): 50-57.

1716 Lahr, J., K. B. Ndour, A. Badji & P. S. Diouf, 1999. Phenology of invertebrates living in a
1717 sahelian temporary pond. *Hydrobiologia* 405: 189-205.

1718 Lund, J. O., S. A. Wissinger & B. L. Peckarsky, 2016. Caddisfly behavioral responses to
1719 drying cues in temporary ponds: implications for effects of climate change.
1720 *Freshwater Science* 35(2): 619-630.

1721 Manconi, R. & R. Pronzato, 2008. Global diversity of sponges (Porifera: Spongillina) in
1722 freshwater. *Hydrobiologia* 595(1): 27-33.

1723 Martens, A., R. Jödicke & F. Suhling, 2003. An annotated checklist of the Odonata of
1724 Namibia. *Cimbebasia* 18: 139-160.

1725 Martens, K., 2001. Ostracoda. In Day, J., I. de Moor, B. Stewart & L. AE (eds) *Guides to the*
1726 *freshwater invertebrates of southern Africa Volume 3: Crustacea II - Ostracoda,*
1727 *Copepoda and Branchiura*. WRC Report no. TT 148/01, Water Research Commission,
1728 Pretoria, 78-123.

1729 Martens, K., 2003. On a remarkable South African giant ostracod (Crustacea, Ostracoda,
1730 Cyprididae) from temporary pools, with additional appendages. *Hydrobiologia* 500:
1731 115-130.

1732 Martens, K., 2007. On a new species and genus in the Cypridini (Crustacea, Ostracoda,
1733 Cyprididae) from South Africa, with a phylogenetic analysis of the tribe and a
1734 discussion on the genus concept in this group. *Journal of Natural History* 41(5-8):
1735 381-399.

- 1736 Martens, K., B. Davies, A. Baxter & M. Meadows, 1996. A contribution to the taxonomy and
1737 ecology of the Ostracoda (Crustacea) from Verlorenvlei (Western Cape, South
1738 Africa). *African Zoology* 31(1): 23-36.
- 1739 Martens, K., M. Hamer & M. Coke, 1998. A preliminary account of the diversity of non-
1740 marine Ostracoda (Crustacea) in KwaZulu-Natal, South Africa. *Lammergeyer* 45: 17-
1741 31.
- 1742 Martin, P., E. Martinez-Ansemil, A. Pinder, T. Timm & M. J. Wetzel, 2008. Global diversity
1743 of oligochaetous clitellates ("Oligochaeta"; Clitellata) in freshwater. *Hydrobiologia*
1744 595(1): 117-127.
- 1745 Mathers, T. C., R. L. Hammond, R. A. Jenner, B. Hänfling & A. Gomez, 2013. Multiple
1746 global radiations in tadpole shrimps challenge the concept of 'living fossils'. *PeerJ* 1:
1747 e62.
- 1748 Matzke-Karasch, R. & K. Martens, 2007. On *Afrocypris barnardi* GO Sars, 1924 (Ostracoda),
1749 a second giant ostracode with additional appendages. *Crustaceana* 80(5): 603-623.
- 1750 McDermott, E. G. & B. A. Mullens, 2014. Desiccation tolerance in the eggs of the primary
1751 North American bluetongue virus vector, *Culicoides sonorensis* (Diptera:
1752 Ceratopogonidae), and implications for vector persistence. *Journal of Medical*
1753 *Entomology* 51(6): 1151-1158.
- 1754 McKee, P. & G. Mackie, 1980. Desiccation resistance in *Sphaerium occidentale* and
1755 *Musculium securis* (Bivalvia: Sphaeriidae) from a temporary pond. *Canadian Journal*
1756 *of Zoology* 58(9): 1693-1696.
- 1757 McKenna, D. D., A. L. Wild, K. Kanda, C. L. Bellamy, R. G. Beutel, M. S. Caterino, C. W.
1758 Farnum, D. C. Hawks, M. A. Ivie & M. L. Jameson, 2015. The beetle tree of life
1759 reveals that Coleoptera survived end-Permian mass extinction to diversify during the
1760 Cretaceous terrestrial revolution. *Systematic Entomology* 40(4): 835-880.
- 1761 McLachlan, A. & M. Cantrell, 1980. Survival strategies in tropical rain pools. *Oecologia*
1762 47(3): 344-351.
- 1763 McLachlan, A. & R. Ladle, 2001. Life in the puddle: behavioural and life-cycle adaptations
1764 in the Diptera of tropical rain pools. *Biological Reviews* 76(3): 377-388.
- 1765 Meintjes, S., 1996. Seasonal changes in the invertebrate community of small shallow
1766 ephemeral pans at Bain's Vlei, South Africa. *Hydrobiologia* 317(1): 51-64.
- 1767 Meyer, H. A. & J. G. Hinton, 2009. The Tardigrada of southern Africa, with the description
1768 of *Minibiotus harrylewisi*, a new species from KwaZulu-Natal, South Africa
1769 (Eutardigrada: Macrobiotidae). *African Invertebrates* 50(2): 255-268.

1770 Miller, J. R., J. Huang, J. Vulule & E. D. Walker, 2007. Life on the edge: African malaria
1771 mosquito (*Anopheles gambiae* s.l.) larvae are amphibious. *Naturwissenschaften* 94(3):
1772 195-199.

1773 Mlambo, M. C., M. S. Bird, C. C. Reed & J. A. Day, 2011. Diversity patterns of temporary
1774 wetland macroinvertebrate assemblages in the south-western Cape, South Africa.
1775 *African Journal of Aquatic Science* 36(3): 299-308.

1776 Møbjerg, N., K. A. Halberg, A. Jørgensen, D. Persson, M. Bjørn, H. Ramløv & R. M.
1777 Kristensen, 2011. Survival in extreme environments—on the current knowledge of
1778 adaptations in tardigrades. *Acta Physiologica* 202(3): 409-420.

1779 Montalto, L. & M. Marchese, 2005. Cyst formation in Tubificidae (Naidinae) and
1780 Opistocystidae (Annelida, Oligochaeta) as an adaptive strategy for drought tolerance
1781 in fluvial wetlands of the Paraná River, Argentina. *Wetlands* 25(2): 488-494.

1782 Morse, J. C., 2011. The Trichoptera world checklist. *Zoosymposia* 5(1): 372-380.

1783 Newell, R. L. & B. R. Hossack, 2009. Large, wetland-associated mayflies (Ephemeroptera)
1784 of Glacier National Park, Montana. *Western North American Naturalist* 69(3): 335-
1785 342.

1786 Nhiwatiwa, T., L. Brendonck & T. Dalu, 2017a. Understanding factors structuring
1787 zooplankton and macroinvertebrate assemblages in ephemeral pans. *Limnologia-
1788 Ecology and Management of Inland Waters* 64: 11-19.

1789 Nhiwatiwa, T., L. Brendonck, A. Waterkeyn & B. Vanschoenwinkel, 2011. The importance
1790 of landscape and habitat properties in explaining instantaneous and long-term
1791 distributions of large branchiopods in subtropical temporary pans. *Freshwater Biology*
1792 56(10): 1992-2008.

1793 Nhiwatiwa, T. & T. Dalu, 2017. Seasonal variation in pans in relation to limno-chemistry,
1794 size, hydroperiod, and river connectivity in a semi-arid subtropical region. *Physics
1795 and Chemistry of the Earth, Parts A/B/C* 97: 37-45.

1796 Nhiwatiwa, T., T. Dalu & L. Brendonck, 2017b. *Streptocephalus sangoensis* n.
1797 sp.(Anostraca, Streptocephalidae), a new large branchiopod species for southern
1798 Africa. *Crustaceana* 90(6): 673-683.

1799 Nhiwatiwa, T., A. Waterkeyn, B. Riddoch & L. Brendonck, 2014. A hotspot of large
1800 branchiopod diversity in south-eastern Zimbabwe. *African Journal of Aquatic Science*
1801 39(1): 57-65.

- 1802 Nicolet, P., J. Biggs, G. Fox, M. J. Hodson, C. Reynolds, M. Whitfield & P. Williams, 2004.
1803 The wetland plant and macroinvertebrate assemblages of temporary ponds in England
1804 and Wales. *Biological Conservation* 120: 261-278.
- 1805 Nilsson, A. N., 1986a. Life cycles and habitats of the northern European Agabini
1806 (Coleoptera, Dytiscidae). *Entomologica Basiliensia* 11: 391-417.
- 1807 Nilsson, A. N., 1986b. Larval morphology and phenology of four Fennoscandian species of
1808 *Hydroporus* Clairville (Coleoptera: Dytiscidae), with a preliminary key to the known
1809 larvae. *Aquatic Insects* 8(3): 141-153.
- 1810 Nilsson, A. N., 1986c. Community structure in the Dytiscidae (Coleoptera) of a northern
1811 Swedish seasonal pond. *Annales Zoologici Fennici* 23(1): 39-47.
- 1812 Nolte, U., R. S. Tietbühl & W. P. McCafferty, 1996. A mayfly from tropical Brazil capable
1813 of tolerating short-term dehydration. *Journal of the North American Benthological*
1814 *Society* 15(1): 87-94.
- 1815 Noreña, C., F. Brusa & A. Faubel, 2003. Census of “Microturbellarians” (free-living
1816 Platyhelminthes) of the zoogeographical regions originating from Gondwana. *Zootaxa*
1817 146(1): 34.
- 1818 Oosthuizen, J. & M. Siddall, 2002. Hirudinea. In Day, J. & I. de Moor (eds) *Guides to the*
1819 *Freshwater Invertebrates of southern Africa*. vol 5: Non-Arthropods. WRC Report
1820 No. TT 167/02, Water Research Commission, Pretoria, 237-263.
- 1821 Paraense, W. L., 1996. Neotropical planorbid snails with apertural lamellae: I. *Biomphalaria*
1822 *helophila* (Orbigny, 1835). *Memorias do Instituto Oswaldo Cruz* 91(2): 177-186.
- 1823 Peer, N., R. Perissinotto, G. Gouws & N. A. Miranda, 2015. Description of a new species of
1824 *Potamonautes* MacLeay, 1838, from the iSimangaliso Wetland Park, South Africa.
1825 *Zookeys* 503: 23-43.
- 1826 Perkins, P. D. & J. Balfour-Browne, 1994. A contribution to the taxonomy of aquatic and
1827 humicolous beetles of the family Hydraenidae in southern Africa. *Fieldiana Zoology*
1828 77: 1-159.
- 1829 Pešić, V., D. Cook, R. Gerecke & H. Smit, 2013. The water mite family Mideopsidae (Acari:
1830 Hydrachnidia): a contribution to the diversity in the Afrotropical region and
1831 taxonomic changes above species level. *Zootaxa* 3720(1): 1-75.
- 1832 Peters, J., 1972. The ecology of Tarn Dub. *Vasculum* 57: 42-50.
- 1833 Pinceel, T., B. Vanschoenwinkel, J. Uten & L. Brendonck, 2013. Mechanistic and
1834 evolutionary aspects of light-induced dormancy termination in a temporary pond
1835 crustacean. *Freshwater Science* 32(2): 517-524.

- 1836 Rayner, N., 1999. Notostraca. In Day, J., I. de Moor, B. Stewart & A. Louw (eds) Guides to
1837 the freshwater invertebrates of southern Africa Volume 2: Crustacea I - Notostraca,
1838 Anostraca, Conchostraca and Cladocera. WRC Report no. TT 121/00, Water Research
1839 Commission, Pretoria, 7-13.
- 1840 Rayner, N., 2000. Distribution and biogeography of the Paradiaptominae (Copepoda:
1841 Calanoida: Diaptomidae). African Journal of Aquatic Sciences 25(1): 93-97.
- 1842 Rayner, N., 2001. Copepoda. In Day, J., I. de Moor, B. A. Stewart & A. Louw (eds) Guides
1843 to the freshwater invertebrates of southern Africa Volume 3: Crustacea II - Ostracoda,
1844 Copepoda and Branchiura. WRC Report No. TT 148/01, Water Research
1845 Commission, Pretoria, 78-123.
- 1846 Rayner, N., 2002. Tardigrada Guides to the Freshwater Invertebrates of southern Africa. vol
1847 5: Non-Arthropods. WRC Report No. TT 167/02, Water Research Commission,
1848 Pretoria, 189-192.
- 1849 Rayner, N. & C. Appleton, 2002. Nematomorpha. In Day, J. & I. de Moor (eds) Guides to the
1850 Freshwater Invertebrates of southern Africa. vol 5: Non-Arthropods. WRC Report
1851 No. TT 167/02, Water Research Commission, Pretoria, 162-165.
- 1852 Rayner, N., C. Appleton & N. Millard, 2002. Cnidaria. In Day, J. & I. de Moor (eds) Guides
1853 to the Freshwater Invertebrates of southern Africa. vol 5: Non-Arthropods. WRC
1854 Report No. TT 167/02, Water Research Commission, Pretoria, 74-87.
- 1855 Rayner, N. & J. Heeg, 1994. Distribution patterns of the Diaptomidae (Calanoida: Copepoda)
1856 in southern Africa. Hydrobiologia 272: 47-75.
- 1857 Rayner, N. A., 1998. *Paradiaptomus peninsularis*, *P. hameri* and *P. warreni*, three new
1858 species of *Paradiaptomus* (Calanoida: Diaptomidae) from South Africa.
1859 Hydrobiologia 391(1-3): 87-98.
- 1860 Reavell, P., 2003. Hemiptera. In de Moor, I., J. Day & F. de Moor (eds) Guides to the
1861 freshwater invertebrates of southern Africa Volume 8: Insecta II Hemiptera,
1862 Megaloptera, Neuroptera, Trichoptera and Lepidoptera. WRC Report No. TT 214/03,
1863 Water Research Commission, Pretoria, 16-71.
- 1864 Rehn, A. C., 2003. Phylogenetic analysis of higher-level relationships of Odonata. Systematic
1865 Entomology 28(2): 181-240.
- 1866 Reid, J. & C. Williamson, 2010. Copepoda. In Thorp, J. & A. Covich (eds) Ecology and
1867 classification of North American freshwater invertebrates. Academic Press, 915-954.

- 1868 Reynolds, J., 2000. Invertebrate communities of turloughs (temporary lakes) in southeast
1869 Galway, Ireland. Internationale Vereinigung für theoretische und angewandte
1870 Limnologie: Verhandlungen 27(3): 1679-1684.
- 1871 Riato, L., C. Van Ginkel & J. C. Taylor, 2014. Zooplankton and diatoms of temporary and
1872 permanent freshwater pans in the Mpumalanga Highveld region, South Africa.
1873 African Zoology 49(1): 113-127.
- 1874 Ribera, I. & M. Balke, 2007. Recognition of a species-poor, geographically restricted but
1875 morphologically diverse Cape lineage of diving beetles (Coleoptera: Dytiscidae:
1876 Hyphydrini). Journal of Biogeography 34(7): 1220-1232.
- 1877 Ricci, C. & M. Balsamo, 2000. The biology and ecology of lotic rotifers and gastrotrichs.
1878 Freshwater Biology 44(1): 15-28.
- 1879 Robinson, C. & T. Buser, 2007. Density-dependent life history differences in a stream mayfly
1880 (*Deleatidium*) inhabiting permanent and intermittent stream reaches. New Zealand
1881 Journal of Marine and Freshwater Research 41(3): 265-271.
- 1882 Rogers, D., 2009. Branchiopoda (Anostraca, Notostraca, Laevicaudata, Spinicaudata,
1883 Cyclestherida). Encyclopedia of inland waters 2: 242-249.
- 1884 Rogers, D. C., 2014. Larger hatching fractions in avian dispersed anostracan eggs
1885 (Branchiopoda). Journal of Crustacean Biology 34(2): 135-143.
- 1886 Rogers, D. C., 2015. A conceptual model for anostracan biogeography. Journal of Crustacean
1887 Biology 35(5): 686-699.
- 1888 Rogers, D. C., N. Rabet & S. C. Weeks, 2012. Revision of the extant genera of Limnadiidae
1889 (Branchiopoda: Spinicaudata). Journal of Crustacean Biology 32(5): 827-842.
- 1890 Rogers, D. C., N. Rabet & S. C. Weeks, 2016. *Gondwanalimnadia* (Branchiopoda:
1891 Spinicaudata), replacement name for *Afrolimnadia* (Limnadiidae), junior homonym of
1892 *Afrolimnadia* (Lioestheriidae). Journal of Crustacean Biology 36(1): 105-105.
- 1893 Rubaba, O., M. Chimbari & S. Mukaratirwa, 2016. The role of snail aestivation in
1894 transmission of schistosomiasis in changing climatic conditions. African Journal of
1895 Aquatic Science 41(2): 143-150.
- 1896 Ruhí, A., D. Boix, S. Gascón, J. Sala & D. P. Batzer, 2013. Functional and phylogenetic
1897 relatedness in temporary wetland invertebrates: current macroecological patterns and
1898 implications for future climatic change scenarios. PLoS ONE 8(11): e81739.
- 1899 Sabatelli, S., P. Audisio, G. Antonini, E. Solano, A. Martinoli & M. Trizzino, 2016.
1900 Molecular ecology and phylogenetics of the water beetle genus *Ochthebius* revealed

1901 multiple independent shifts to marine rockpools lifestyle. *Zoologica Scripta* 45(2):
 1902 175-186.

1903 Sabatelli, S., P. Audisio, M. Trizzino & A. Di Giulio, 2013. Description of the larva of
 1904 *Ochthebius capicola* (Coleoptera: Hydraenidae) from marine rockpools of South
 1905 Africa. *Zootaxa* 3683(3): 280-288.

1906 Samways, M., 1999. Diversity and conservation status of South African dragonflies
 1907 (Odonata). *Odonatologica* 28(1): 13-62.

1908 Samways, M. J., 2004. Critical species of Odonata in southern Africa. *International Journal*
 1909 *of Odonatology* 7(2): 255-262.

1910 Samways, M. J., 2008. Dragonflies and damselflies of South Africa. Pensoft Publishers.

1911 Samways, M. J., 2010. Extreme weather and climate change impacts on South African
 1912 dragonflies. In Ott, J. (ed) *Monitoring Climate Change with Dragonflies*. Pensoft,
 1913 Sophia, Bulgaria, pp. 73-84.

1914 Samways, M. J. & J. P. Simaika, 2016. Manual of freshwater assessment for South Africa:
 1915 Dragonfly Biotic Index, South African National Biodiversity Institute, Pretoria.

1916 Sartori, M. & H. Barber-James, 2018. World checklist on Ephemeroptera. In: *The Freshwater*
 1917 *Animal Diversity Assessment (FADA) Belgium*
 1918 <http://fadabiodiversitybe/group/show/35> Updated February 2018.

1919 Schiel, F.-J. & R. Buchwald, 2015. Hatching phenology of Odonata species inhabiting
 1920 temporary and permanent water bodies (Odonata: Lestidae, Aeshnidae, Libellulidae).
 1921 *International Journal of Odonatology* 18(2): 105-123.

1922 Schmidt-Rhaesa, A. & R. Perissinotto, 2016. *Chordodes ferox*, a new record of horsehair
 1923 worms (Nematomorpha, Gordiida) from South Africa. *Zookeys* 566: 1-11.

1924 Scott, K. M. F., 1970. Some notes on the trichoptera of standing waters in Africa, mainly
 1925 south of the Zambezi. *Hydrobiologia* 35(2): 177-195.

1926 Seaman, M., D. Kok, B. Von Schlichting & A. Kruger, 1991. Natural growth and
 1927 reproduction in *Triops granarius* (Lucas)(Crustacea: Notostraca). *Hydrobiologia* 212:
 1928 87-94.

1929 Seaman, M., D. Kok & M. Watson, 1999. Cladocera. In Day, J., I. de Moor, B. Stewart & A.
 1930 Louw (eds) *Guides to the freshwater invertebrates of southern Africa Volume 2:*
 1931 *Crustacea I - Notostraca, Anostraca, Conchostraca and Cladocera*. WRC Report No.
 1932 TT 121/00, Water Research Commission, Pretoria, 81-110.

1933 Semlitsch, R. D. & J. R. Bodie, 1998. Are small, isolated wetlands expendable? *Conservation*
 1934 *Biology* 12(5): 1129-1133.

- 1935 Short, A. E. & J. K. Liebherr, 2007. Systematics and biology of the endemic water scavenger
1936 beetles of Hawaii (Coleoptera: Hydrophilidae, Hydrophilini). Systematic Entomology
1937 32(4): 601-624.
- 1938 Short, A. E. Z., 2018. Systematics of aquatic beetles (Coleoptera): current state and future
1939 directions. Systematic Entomology 43: 1-18.
- 1940 Silberbauer, M. J. & J. M. King, 1991. The distribution of wetlands in the south-western
1941 Cape Province, South Africa. South African Journal of Aquatic Science 17(1-2): 65-
1942 81.
- 1943 Simaika, J. P., M. J. Samways & P. P. Frenzel, 2016. Artificial ponds increase local
1944 dragonfly diversity in a global biodiversity hotspot. Biodiversity and Conservation
1945 25(10): 1921-1935.
- 1946 Simovich, M. A. & S. A. Hathaway, 1997. Diversified bet-hedging as a reproductive strategy
1947 of some ephemeral pool anostracans (Branchiopoda). Journal of Crustacean Biology
1948 17(1): 38-44.
- 1949 Smirnov, N., 2008. Check-list of the South-African Cladocera (Crustacea: Branchiopoda).
1950 Zootaxa 1788: 47-56.
- 1951 Smit, H., 2012. New records of the water mite family Arrenuridae from the Afrotropical
1952 region, with the description of 11 new species and two new subspecies (Acari:
1953 Hydrachnidia). Zootaxa 3187: 1-31.
- 1954 Spencer, M., S. S. Schwartz & L. Blaustein, 2002. Are there fine-scale spatial patterns in
1955 community similarity among temporary freshwater pools? Global Ecology and
1956 Biogeography 11(1): 71-78.
- 1957 Strachan, S. R., E. T. Chester & B. J. Robson, 2015. Freshwater invertebrate life history
1958 strategies for surviving desiccation. Springer Science Reviews 3(1): 57-75.
- 1959 Suárez-Morales, E. & N. A. Rayner, 2004. New records for the South African diaptomid
1960 fauna with a complementary description of *Paradiaptomus lamellatus* Sars
1961 (Copepoda, Diaptomidae). Journal of Natural History 38(22): 2803-2819.
- 1962 Suárez-Morales, E., R. Wasserman & T. Dalu, 2015. A new species of *Lovenula* Schmeil
1963 (Copepoda, Calanoida, Diaptomidae) from the Eastern Cape Province of South
1964 Africa. Crustaceana 88(3): 324-342.
- 1965 Suhling, F., R. Jödicke & W. Schneider, 2003. Odonata of African arid regions—are there
1966 desert species. Cimbebasia 18: 207-224.

- 1967 Suhling, F., G. Sahlén, J. Kasperski & D. Gaedecke, 2005. Behavioural and life history traits
1968 in temporary and perennial waters: comparisons among three pairs of sibling
1969 dragonfly species. *Oikos* 108(3): 609-617.
- 1970 Suhling, F., M. Samways, J. Simaika & J. Kipping, 2009. The status and distribution of
1971 dragonflies (Odonata). In Darwall, W., K. Smith, D. Tweddle & P. Skelton (eds) *The*
1972 *status and distribution of freshwater biodiversity in southern Africa*. IUCN,
1973 Gland/Cambridge, pp. 48-65.
- 1974 Tawfik, M., 1969. Life-history of the giant water-bug, *Lethocerus niloticus* Stael (Hemiptera:
1975 Belostomatidae). *Bulletin de la Société entomologique d’Egypte* 53: 299-310.
- 1976 Tobias, D. & W. Tobias, 2008. *Trichoptera africana*. Available from:
1977 <http://trichoptera.insects-online.de/Trichoptera%20africana/index.htm>.
- 1978 Todaro, M., R. Perissinotto & S. Bownes, 2013. Neogosseidae (Gastrotricha, Chaetonotida)
1979 from the iSimangaliso Wetland Park, KwaZulu-Natal, South Africa. *Zookeys* 315: 77-
1980 94.
- 1981 Tuytens, K., B. Vanschoenwinkel, B. Clegg, T. Nhiwatiwa & L. Brendonck, 2015. Exploring
1982 links between geology, hydroperiod, and diversity and distribution patterns of
1983 anostracans and notostracans (Branchiopoda) in a tropical savannah habitat in SE
1984 Zimbabwe. *Journal of Crustacean Biology* 35(3): 309-318.
- 1985 Tuytens, K., B. Vanschoenwinkel, A. Waterkeyn & L. Brendonck, 2014. Predictions of
1986 climate change infer increased environmental harshness and altered connectivity in a
1987 cluster of temporary pools. *Freshwater Biology* 59(5): 955-968.
- 1988 Urban, M. C., 2004. Disturbance heterogeneity determines freshwater metacommunity
1989 structure. *Ecology* 85(11): 2971-2978.
- 1990 Van Damme, K., E. I. Bekker & A. A. Kotov, 2013. Endemism in the Cladocera (Crustacea:
1991 Branchiopoda) of Southern Africa. *Journal of Limnology* 72(3): 440-463.
- 1992 Van Damme, K. & H. J. Dumont, 1999. A drought-resistant larva of *Pantala flavescens*
1993 (Fabricius, 1798)(Odonata: Libellulidae) in the Lençóis Maranhenses, NE-Brazil.
1994 *International Journal of Odonatology* 2(1): 69-76.
- 1995 van Hoven, W. & J. Day, 2002. Oligochaeta. In Day, J. & I. de Moor (eds) *Guides to the*
1996 *Freshwater Invertebrates of southern Africa*. vol 5: Non-Arthropods. WRC Report
1997 No. TT 167/02, Water Research Commission, Pretoria, 203-236.
- 1998 Van Steenkiste, N., P. Davison & T. Artois, 2010. *Bryoplana xerophila* n.g.n. sp., a new
1999 limnoterrestrial microturbellarian (Platyhelminthes, Typhloplanidae, Protoplanellinae)
2000 from epilithic mosses, with notes on its ecology. *Zoological Science* 27(3): 285-291.

2001 Vandekerckhove, J., G. Louette, L. Brendonck & L. De Meester, 2005. Development of
 2002 cladoceran egg banks in new and isolated pools. *Archiv für Hydrobiologie* 162(3):
 2003 339-347.

2004 Vanschoenwinkel, B., F. Buschke & L. Brendonck, 2013. Disturbance regime alters the
 2005 impact of dispersal on alpha and beta diversity in a natural metacommunity. *Ecology*
 2006 94(11): 2547-2557.

2007 Vanschoenwinkel, B., C. De Vries, M. Seaman & L. Brendonck, 2007. The role of
 2008 metacommunity processes in shaping invertebrate rock pool communities along a
 2009 dispersal gradient. *Oikos* 116(8): 1255-1266.

2010 Vanschoenwinkel, B., S. Gielen, M. Seaman & L. Brendonck, 2009. Wind mediated dispersal
 2011 of freshwater invertebrates in a rock pool metacommunity: differences in dispersal
 2012 capacities and modes. *Hydrobiologia* 635(1): 363-372.

2013 Vanschoenwinkel, B., S. Gielen, H. Vandewaerde, M. Seaman & L. Brendonck, 2008.
 2014 Relative importance of different dispersal vectors for small aquatic invertebrates in a
 2015 rock pool metacommunity. *Ecography* 31(5): 567-577.

2016 Vanschoenwinkel, B., T. Pinceel, M. P. Vanhove, C. Denis, M. Jocque, B. V. Timms & L.
 2017 Brendonck, 2012. Toward a global phylogeny of the "living fossil" crustacean order
 2018 of the Notostraca. *PLoS ONE* 7(4): e34998.

2019 Vanschoenwinkel, B., M. Seaman & L. Brendonck, 2010a. Hatching phenology, life history
 2020 and egg bank size of fairy shrimp *Branchipodopsis* spp.(Branchiopoda, Crustacea) in
 2021 relation to the ephemerality of their rock pool habitat. *Aquatic Ecology* 44(4): 771-
 2022 780.

2023 Vanschoenwinkel, B., A. Waterkeyn, M. Jocqué, L. Boven, M. Seaman & L. Brendonck,
 2024 2010b. Species sorting in space and time-the impact of disturbance regime on
 2025 community assembly in a temporary pool metacommunity. *Journal of the North*
 2026 *American Benthological Society* 29(4): 1267-1278.

2027 Vanschoenwinkel, B., A. Waterkeyn, T. Nihwatiwa, T. Pinceel, E. Spooren, A. Geerts, B.
 2028 Clegg & L. Brendonck, 2011. Passive external transport of freshwater invertebrates
 2029 by elephant and other mud-wallowing mammals in an African savannah habitat.
 2030 *Freshwater Biology* 56(8): 1606-1619.

2031 Vera, C., P. Bremond, R. Labbo, F. Mouchet, E. Sellin, D. Boulanger, J. Pointier, B. Delay &
 2032 B. Sellin, 1995. Seasonal fluctuations in population densities of *Bulinus senegalensis*
 2033 and *B. truncatus* (Planorbidae) in temporary pools in a focus of *Schistosoma*

2034 *haematobium* in Niger: implications for control. Journal of Molluscan Studies 61(1):
2035 79-88.

2036 Wasserman, R. J., M. E. Alexander, D. Barrios-O'Neill, O. L. Weyl & T. Dalu, 2016a. Using
2037 functional responses to assess predator hatching phenology implications for
2038 pioneering prey in arid temporary pools. Journal of Plankton Research 38(1): 154-
2039 158.

2040 Wasserman, R. J., M. E. Alexander, O. L. Weyl, D. Barrios-O'Neill, P. W. Froneman & T.
2041 Dalu, 2016b. Emergent effects of structural complexity and temperature on predator–
2042 prey interactions. Ecosphere 7(2): e01239.

2043 Wasserman, R. J., M. Weston, O. L. Weyl, P. W. Froneman, R. J. Welch, T. J. Vink & T.
2044 Dalu, 2018. Sacrificial males: the potential role of copulation and predation in
2045 contributing to copepod sex-skewed ratios. Oikos doi: 10.1111/oik.04832.

2046 Weir, J. S., 1966. Ecology and zoogeography of aquatic Hemiptera from temporary pools in
2047 Central Africa. Hydrobiologia 28(1): 123-128.

2048 Wharton, D. A., 2004. Survival strategies. In Gaugler, R. & A. L. Bilgrami (eds) Nematode
2049 Behaviour. CABI Publishing, Wallingford, UK, 371-399.

2050 Wheeler, Q. D., P. H. Raven & E. O. Wilson, 2004. Taxonomy: Impediment or Expedient?
2051 Science 303(5656): 285.

2052 Wickson, S., E. Chester & B. Robson, 2012. Aestivation provides flexible mechanisms for
2053 survival of stream drying in a larval trichopteran (Leptoceridae). Marine and
2054 Freshwater Research 63(9): 821-826.

2055 Wiggins, G. B., 1973. Contribution to the biology of caddisflies (Trichoptera) in temporary
2056 pools. Royal Ontario Museum Life Sciences Contributions 88: 1-28.

2057 Wiggins, G. B., R. J. Mackay & I. M. Smith, 1980. Evolutionary and ecological strategies of
2058 animals in annual temporary pools. Archiv für Hydrobiologie 58 (Suppl): 97-206.

2059 Williams, D. D., 1987. The Biology of Temporary Waters. Timber Press, Portland, Oregon.

2060 Williams, D. D., 2006. The biology of temporary waters. Oxford University Press, Oxford.

2061 Wissinger, S., W. Brown & J. Jannot, 2003. Caddisfly life histories along permanence
2062 gradients in high-altitude wetlands in Colorado (USA). Freshwater Biology 48(2):
2063 255-270.

2064 Wissinger, S., J. Steinmetz, J. S. Alexander & W. Brown, 2004a. Larval cannibalism, time
2065 constraints, and adult fitness in caddisflies that inhabit temporary wetlands. Oecologia
2066 138(1): 39-47.

2067 Wissinger, S. A., C. Eldermire & J. C. Whissel, 2004b. The role of larval cases in reducing
2068 aggression and cannibalism among caddisflies in temporary wetlands. *Wetlands*
2069 24(4): 777-783.

2070 Wissinger, S. A., J. C. Whissel, C. Eldermire & W. S. Brown, 2006. Predator defense along a
2071 permanence gradient: roles of case structure, behavior, and developmental phenology
2072 in caddisflies. *Oecologia* 147(4): 667-678.

2073 Womersley, C. & C. Ching, 1989. Natural dehydration regimes as a prerequisite for the
2074 successful induction of anhydrobiosis in the nematode *Rotylenchulus reniformis*.
2075 *Journal of Experimental Biology* 143(1): 359-372.

2076 Young, J. O., 1976. The freshwater Turbellaria of the African continent. *Zoologischer*
2077 *Anzeiger* 197: 419-432.

2078 Yozzo, D. & R. Diaz, 1999. Tidal freshwater wetlands: invertebrate diversity, ecology, and
2079 functional significance. In Batzer, D. P., R. B. Rader & S. A. Wissinger (eds)
2080 *Invertebrates in freshwater wetlands of North America: ecology and management*.
2081 Wiley, Hoboken, New Jersey, 889-918.

2082 Zamora-Muñoz, C. & B. Svensson, 1996. Survival of caddis larvae in relation to their case
2083 material in a group of temporary and permanent pools. *Freshwater Biology* 36(1): 23-
2084 31.

2085 Zawierucha, K., Ł. Michalczyk & Ł. Kaczmarek, 2012. The first record of Tardigrada from
2086 Zambia, with a description of *Doryphoribius niedbalai* n. sp. (Eutardigrada:
2087 Isohypsibiidae, the evelinae group). *African Zoology* 47(2): 275-284.

2088