Faculty of Health: Medicine, Dentistry and Human Sciences

School of Health Professions

2019-06-01

The oral nitrate-reducing capacity correlates with peak power output and peak oxygen uptake in healthy humans.

Thomas, B

http://hdl.handle.net/10026.1/13578

10.1016/j.niox.2019.03.001 Nitric Oxide: Biology and Chemistry Elsevier

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Manuscript Details

Manuscript number	NOX_2018_270					
Title	Nitrate-reducing capacity of oral bacteria correlates with aerobic exercise performance in healthy humans					
Article type	Research Paper					

Abstract

Interest in inorganic nitrate and nitrite has grown substantially over the past decade as research has revealed the role of these anions in enhancing nitric oxide (NO) availability through an oral pathway. Nitrite synthesis in the mouth has been shown to be a key source of the circulatory levels of this anion. This is interesting since greater plasma nitrite concentration has been associated with better exercise capacity in humans, but this question has not been investigated in relation to salivary nitrite concentration. Additionally, no previous study has investigated the nitratereducing activity of oral bacteria in regards to exercise performance in humans. Thus, the main goal of this study was to investigate whether salivary nitrite and nitrate concentration and the nitrate-reducing capacity of oral bacteria were associated with aerobic exercise capacity in healthy humans. Fifty individuals (22 females and 28 males; 38.8 ± 14.3 years/old; BMI= 22.8 ± 3.9) performed a graded exercise test on a cycle ergometer to assess their maximum aerobic capacity (VOpeak). Unstimulated salivary samples were taken before and 20 min after exercise to measure nitrate/nitrite, pH and lactate. The nitrate-reducing capacity of oral bacteria was also assessed in 25 subjects before and after exercise. Nitrate-reducing capacity of oral bacteria was positively associated with muscle power (rs= 0.64; P=0.001) and the VO2peak (rs= 0.54; P=0.005). Similar correlations were found when these variables were analysed after exercise. In addition, a significant decrease in salivary pH (pre:7.28 ± 0.361; post-exercise: 7.16 ± 0.33; P = 0.003) accompanied by an increase of salivary lactate (pre: 0.17 ± 0.14 mmol/L; post-exercise: 0.48 ± 0.38 ; P < 0.001) was found after exercise. However, these changes did not have any impact on salivary nitrate/nitrite concentration and the nitrate-reducing activity of oral bacteria after exercise. In conclusion, this is the first evidence showing a link between the nitrate-reducing activity of oral bacteria and aerobic fitness levels in healthy humans.

Keywords nitrate, nitrite, exercise, oral bacteria					
Manuscript category	Nitrite/Nitrate				
Corresponding Author	Raul Bescos Garcia				
Corresponding Author's Institution	University of Plymouth				
Order of Authors	Bernie Thomas, Stephen Smallwood, Craig Cutler, Raul Bescos Garcia				
Suggested reviewers	Chris Easton, Simone Porcelli, Eddie Weitzberg				

Submission Files Included in this PDF

File Name [File Type]

Coverletter.docx [Cover Letter]

Manuscript_RB_submitted_1.docx [Manuscript File]

Figure 1_Salivary_lactate_pH.JPG [Figure]

Figure 2_Salivary_NO3_NO2_ONRC.JPG [Figure]

Figure 3_Correlations Performance.JPG [Figure]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Research Data Related to this Submission

There are no linked research data sets for this submission. The following reason is given: Data will be made available on request



13th November 2018

To the Editor—NITRIC OXIDE JOURNAL

Dear Prof. Hogg:

Please find uploaded our manuscript entitled '*Nitrate-reducing capacity of oral bacteria correlates with aerobic exercise performance in healthy humans*'. This is the largest study (*n* = 50) that has been performed to date looking at salivary nitrate and nitrite concentration, and the nitrate-reducing activity of oral bacteria in regards to exercise capacity in humans. Interestingly, we found an strong correlation between the nitrate-reducing activity of oral bacteria and the main two parameters of aerobic exercise performance (power and maximal oxygen uptake). This finding is very novel, and it may explain, at least partially, the lack of effect of nitrate supplements in well-trained individuals. On the other hand, we did not find an association between salivary nitrate and nitrite and exercise performance in this study. Differences between gender were not observed either. The manuscript is original, is not under concurrent consideration elsewhere, and was prepared in accordance with the requirements and style of the NITRIC OXIDE. All authors have participated in this study and they have read the manuscript being agreed with the submitted work.

Yours sincerely,

Dr. Raul Bescos Garcia Faculty of Health & Human Sciences University of Plymouth Plymouth United Kingdom

Nitrate-reducing capacity of oral bacteria correlates with aerobic exercise performance in healthy humans

Thomas B¹, Smallwood S¹, Cutler C¹, Bescos R^{1,2*}

¹ School of Health Professions, Faculty of Health & Human Sciences, University of Plymouth (UK) ² Institute of Health & Community, University of Plymouth (UK)

Keywords: nitrate, nitrite, exercise, oral bacteria

*Corresponding author Raul Bescos Garcia Institute of Health & Community University of Plymouth Plymouth United Kingdom Raul.Bescos@plymouth.ac.uk

1 Abstract

2 Interest in inorganic nitrate and nitrite has grown substantially over the past decade as research has 3 revealed the role of these anions in enhancing nitric oxide (NO) availability through an oral pathway. 4 Nitrite synthesis in the mouth has been shown to be a key source of the circulatory levels of this anion. 5 This is interesting since greater plasma nitrite concentration has been associated with better exercise 6 capacity in humans, but this question has not been investigated in relation to salivary nitrite 7 concentration. Additionally, no previous study has investigated the nitrate-reducing activity of oral 8 bacteria in regards to exercise performance in humans. Thus, the main goal of this study was to 9 investigate whether salivary nitrite and nitrate concentration and the nitrate-reducing capacity of oral bacteria were associated with aerobic exercise capacity in healthy humans. 10

Fifty individuals (22 females and 28 males; 38.8 ± 14.3 years/old; BMI= 22.8 ± 3.9) performed a graded exercise test on a cycle ergometer to assess their maximum aerobic capacity (VO_{peak}). Unstimulated salivary samples were taken before and 20 min after exercise to measure nitrate/nitrite, pH and lactate. The nitrate-reducing capacity of oral bacteria was also assessed in 25 subjects before and after exercise.

Nitrate-reducing capacity of oral bacteria was positively associated with muscle power (r_s = 0.64; P=0.001) and the VO_{2peak} (r_s = 0.54; P=0.005). Similar correlations were found when these variables were analysed after exercise. In addition, a significant decrease in salivary pH (pre:7.28 ± 0.361; postexercise: 7.16 ± 0.33; P = 0.003) accompanied by an increase of salivary lactate (pre: 0.17 ± 0.14 mmol/L; post-exercise: 0.48 ± 0.38; P < 0.001) was found after exercise. However, these changes did not have any impact on salivary nitrate/nitrite concentration and the nitrate-reducing activity of oral bacteria after exercise.

In conclusion, this is the first evidence showing a link between the nitrate-reducing activity of oral
bacteria and aerobic fitness levels in healthy humans.

25

26 Introduction

Nitrate and nitrite have emerged over the last decade as key regulators of nitric oxide (NO) metabolism in humans (1). Nitrate is the main circulatory anion in healthy humans, and it has been estimated that about 25% of circulatory nitrate is absorbed in the salivary glands (2). There, nitrate is excreted with saliva into the mouth where different species of oral bacteria can reduce it to nitrite. Once nitrite is swallowed, a small part is rapidly absorbed into the circulation across the upper gastrointestinal tract (2). Circulating nitrite can be reduced to NO through different pathways (3).

33

34 The main sources of nitrate in the body are diet (mainly vegetables) and endogenous synthesis of nitric 35 oxide through the enzymatic L-Arginine/Nitric Oxide Synthase (NOS) pathway (2). Substantial research 36 over the last decade has focused on dietary nitrate from two different perspectives: 1) as a potential 37 ergogenic aid in sport, and 2) as a potential blood pressure-lowering dietary compound in health and 38 disease (4, 5). Taking the first approach, recent data suggest that supplementation of dietary inorganic 39 nitrate may be useful to enhance exercise capacity especially in low or moderate trained individuals, 40 but this is not evident in well-trained athletes (6). This lack of effect has been attributed to several factors such as physiological and metabolic adaptations of chronic exercise training (6). Years of 41 42 training may lead to reconditioning responses on cardiovascular function, skeletal muscle 43 vascularization, and mitochondrial efficiency, resulting in greater metabolic and mechanical efficiency 44 (7). These exercise-induced adaptations can also enhance NO synthesis through the L-Arginine/NOS 45 pathway during exercise increasing nitrate and nitrite bioavailability, and lowering any potential 46 ergogenic benefit from supplementation (5). Two previous studies by Rassaf et al (8) and Totzeck et al 47 (9) found a positive and significant correlation between circulatory nitrite and exercise capacity in 48 healthy humans under fasting conditions. However, this question has not been investigated using 49 salivary nitrite and nitrate. This is interesting since recent evidence (10), but not all (11), suggests that 50 synthesis of oral nitrite is a key mechanism in sustaining the circulatory levels of this anion, and to support NO availability in the body (10). With salivary-nitrite test-strip manufacturers claiming that 51 52 greater concentration of salivary nitrite enhances exercise performance, there is a need for research 53 to investigate this question more in depth.

54

55 Another key question is the differences in NO metabolism between men and women. A recent study 56 by Kapil et al (12) has shown that the nitrate-reducing activity of oral bacteria differs between males 57 and females, but this has not been previously investigated in regards to gender and exercise capacity 58 in humans. Thus, the main aims of this study were to investigate whether salivary nitrate and nitrite

3

59 concentrations were associated with maximum aerobic capacity in males and females. In a second 60 part of this study, we investigated whether the nitrate-reducing ability of oral bacteria was related to 61 parameters of aerobic exercise performance in a similar cohort of subjects. First, we hypothesised that 62 higher exercise performance would be associated with greater salivary nitrate/nitrite levels in both 63 males and females. Then, we hypothesised that greater salivary nitrite could be related to higher 64 activity of oral bacteria and, consequently, better exercise capacity.

65

66 Materials and methods

67 Participants

68 Fifty-three participants were recruited for this study, but two participants could not complete the 69 protocol due to physical problems, and another participant did not follow the full protocol. Thus, fifty 70 participants were included in this study (Table 1). They were recruited from local cycling, triathlon and 71 University sports clubs. A medical questionnaire was completed prior to the study, and individuals 72 with any pathology that could affect the oral cavity, such as gingivitis or periodontitis, or which could 73 affect vascular function such as diabetes, hypertension, smoking or psychological conditions were 74 excluded. Furthermore, individuals under medication or treatment potentially affecting oral bacteria 75 (e.g antibacterial mouthwash) were excluded. All participants gave their written informed consent to 76 participate in the study. The procedures carried out were approved by the human ethics committee 77 of the Faculty of Health and Human Sciences (17/18-415) (University of Plymouth, UK) in accordance 78 with the Declaration of Helsinki.

79

80 Protocol

81 Participants were required to report to the laboratory on one occasion in a fully rested state having 82 not participated in strenuous exercise in the preceding 24 hours. They were encouraged to refrain 83 from eating at least 3 hours before the exercise test. Food rich in nitrate and caffeine were restricted 84 24 and 12 hours before the trial, respectively. On their arrival at the laboratory, body fat percentage 85 and body mass were measured using a body composition analyser (Tanita, TBF-300MA, Tokyo, Japan). 86 Height was measured using a stadiometer (Seca, Hamburg, Germany). Then, salivary nitrite was 87 measured with a test strip (Nitric Oxide test strips, Berkeley, Chicago, US) before a whole non-88 stimulated saliva sample (~3 mL) was collected to analyse nitrate, nitrite, pH and lactate. Following 89 this, the nitrate-reducing capacity of oral bacteria was measured in 25 participants (Table 1). They 90 rinsed their mouth with a 10 mL solution of distilled water containing 80 µmol of sodium nitrate 91 (Sodium Nitrate, Fisher Chemical, UK) for 5 minutes. The solution was collected in sterile tubes. All

saliva samples were centrifuged at 4,000 rpm for 10 min at 4°C. The supernatant was obtained, filtered

- 93 and stored at -80°C.
- 94

95 Following sampling, subjects performed a graded exercise test until exhaustion on a bicycle ergometer 96 (Lode Corival, Groningen, Netherlands) to assess their maximal aerobic capacity (VO_{peak}) and peak 97 power (W_{peak}). The test commenced with a five-minute warm-up with resistance set at 70W for 98 females and 100W for males. Pedalling cadence was kept within the range of 80-100 rpm and on 99 completion of the warm-up exercise intensity increased 20W every minute for females and 25W for 100 males. The exercise test finished when the subject reached physical exhaustion or was unable to 101 maintain a minimum pedalling cadence of 70 rpm. Oxygen uptake (VO₂) and carbon dioxide 102 production (VCO₂) were measured at 10-second intervals by a computerized gas analyser (Cortex 103 Metalyzer 3B, Leipzig, Germany), which was calibrated in accordance with the manufacturer's 104 instructions prior to each test. Heart rate was measured via a wireless transmitter (Polar, Kempele, 105 Finland). VO_2 and power output (W) averaged over the last 30-second period of the test, was defined 106 as VO_{2peak} and W_{peak} respectively.

107

At the cessation of the exercise test the subject was seated and allowed to rest for 20 min. During this period water was permitted however food and other drinks were restricted. Then, salivary nitrite was again measured using a test strip (Nitric Oxide test strips, Berkeley, Chicago, US) before taking another salivary sample (~ 3 mL). Finally, the oral nitrate-reducing capacity was measured using the same procedure as before exercise.

113

114 Analyses

115 Salivary nitrate and nitrite

Samples were thawed and 100 μL of supernatant was collected and diluted 10 times with a solution containing 10% methanol, 0.15 M NaCl/NH₄Cl, and 0.5 g/L 4Na-EDTA. Then, 10 μL was directly injected in a HPLC system (ENO-30, Eicom, Japan) to measure nitrate and nitrite. The same procedure was followed to measure salivary nitrite from the rinsing solutions used to assess the nitrate-reducing activity of oral bacteria.

- 121 Change in colour of salivary strips (Nitric Oxide test strips, Berkeley, Chicago, US) was categorised on
 122 5 different levels according to the manufacturer's color chart.
- 123

124 Salivary pH, lactate and glucose

125 One mL of saliva was used to measure the pH (25°C) using an electronic pH meter (PH-208, Lutron,

- 126 Taiwan) calibrated in accordance with the manufacturer's instructions. Salivary lactate and glucose
- 127 were analysed by a biochemistry analyser (YSI 2300 Stat Plus, YSI Life Sciences, US).
- 128

129 Statistical analyses

130 Data are presented as means \pm SD. All data were analyzed to determine the normal distribution using 131 the Shapiro Wilk's test. Comparison between groups (males / females) were performed using the 132 Mann-Whitney U-test as data were not normally distributed. Comparison between pre and post 133 exercise measurements in saliva parameters were carried out using a Wilcoxon Signed Ranks Test for 134 all the variables except for salivary pH as this was normally distributed, and a paired t-test was used in this case. The association between salivary markers, power and VO_{2peak} was analysed using two-135 136 tailed Spearman's rank correlation analyses (95% confidence). Additionally, Spearman's rank 137 correlation analyses (95% confidence) were used to assess the level of agreement between the 138 salivary strips and nitrite laboratory measurements.

139

140 Results

141 Baseline characteristics of study subjects

Table 1 lists the baseline characteristics of the subjects. Significant differences between males and females were found in age (P = 0.048), anthropometrical (body mass P < 0.001; BMI P = 0.036; body fat P < 0.001) and physiological parameters (VO_{2peak} P = 0.002; W_{peak} P = 0.008) as expected, but no differences were observed (P > 0.05) in any salivary parameter between males and females (P < 0.001) (Table 1).

147 148

-Table 1-

149 Changes in salivary parameters between pre and post-exercise measurements

A significant increase (P < 0.001) in salivary lactate was observed in females (Pre: 0.16 ± 0.15; Post: 0.45 ± 0.37; P < 0.001) and males (Pre: 0.17 ± 0.15; Post: 0.53 ± 0.40; P < 0.001) after exercise (Figure 1). This response was associated with a significant reduction of salivary pH in males (Pre: 7.26 ± 0.33; Post: 7.11±0.25; P < 0.001). Although a similar trend was observed in females, the reduction in salivary pH was not statistically significant (Pre: 7.29 ± 0.39; Post: 7.18 ± 0.37; P = 0.64). However, when all

155	participants were analysed together salivary pH was significantly lower after exercise compared to the
156	pre-exercise values (Pre: 7.28 \pm 0.36; Post: 7.16 \pm 0.33; P = 0.003) (Figure 1).
157	
158	In contrast, the concentration of salivary nitrate, nitrite and the nitrate-reducing capacity of oral
159	bacteria did not differ (P > 0.05) between pre and post-exercise measurements in either group (Figure
160	2).
161	-Figure 1-
162	-Figure 2-
163	
164	
165	Association between salivary strips and HPLC measurements on nitrite concentration
166	A moderate ($r_s = 0.43$) and significant (P = 0.002) association was found between salivary strips
167	(Berkeley test $^{\circ}$) and salivary nitrite laboratory measurements (Table 2). Although this association was
168	still significant between measurements after exercise, it was weakened compared to the pre-exercise
169	values (Table 2).
170	-Table 2-
171	
172	Associations between salivary markers and exercise capacity
173	No associations were found between salivary nitrate and exercise capacity in males or females (Table
174	3). A moderate and significant association (r_s = 0.44, P= 0.04) between salivary nitrite and power was
175	only found in females. However, this was not associated with a correlation between salivary nitrite
176	and VO_{2peak} ($r_s = 0.31$, $P= 0.16$). Taking all participants together, power was significantly associated
177	with salivary lactate ($r_s = 0.33$, $P= 0.02$).
178	-Table 3-
179	
180	The nitrate-reducing capacity of oral bacteria was significantly associated with power (r_s = 0.64, P=
181	0.01) and VO_{2peak} ($r_s = 0.54$, $P= 0.04$) in males before exercise (Table 3). While the association between
182	the nitrate-reducing capacity and power did not change after exercise (r_s = 0.64, P = 0.01), a lower and
183	non-significant correlation was observed with the VO_{2peak} ($r_s = 0.47$, $P= 0.08$). In females, the oral
184	nitrate-reducing capacity was not associated with power or VO_{2peak} before and after exercise (Table
185	3). On the other hand, a positive and significant association was found between the nitrate-reducing
186	capacity of oral bacteria and power and VO_{2peak} before and after exercise when all the participants
187	(males and females) were analysed together (Figure 3).

188

-Figure 3-

189 Discussion

190 The main finding of this study was the positive and significant association between the nitrate-191 reducing capacity of oral bacteria and aerobic exercise performance in terms of power (W) and 192 respiratory capacity (VO_{2peak}) when all subjects were analysed together. To the best of our knowledge, 193 this is the first evidence linking exercise capacity with the activity of oral bacteria in healthy humans. 194 These results suggest that exercise training may modulate the activity and/or diversity of the oral 195 microbiome. From this viewpoint, recent studies on the gut microbiome have shown a positive 196 association between exercise levels and the diversity of bacteria (13-16), however no previous study 197 has investigated a possible link between oral bacteria and exercise capacity. Thus, this new evidence 198 may explain the lack of ergogenic effect of dietary nitrate supplements in well-trained individuals (6, 199 17). Higher nitrate-reducing capacity of oral bacteria in trained individuals may be an adaptive 200 mechanism that helps sustain nitrite availability through endogenous sources under different 201 physiological conditions.

202

203 On the other hand, this study did not show an association between concentration of salivary nitrite and nitrate, and aerobic exercise capacity. A recent study has found that salivary nitrite production is 204 205 well associated with higher abundance of oral nitrate-reducing capacity (18). According to this 206 evidence and current results, it may suggest that highly trained individuals have similar levels of 207 nitrate-reducing oral bacteria, but the activity of these species may differ between low and well-208 trained individuals. In order to answer this, there is a need to examine the oral biome of individuals 209 with different fitness levels. On the other hand, our results in saliva were contrary to two previous 210 studies showing a positive correlation between plasma nitrite concentration and aerobic exercise 211 capacity in a similar cohort of individuals (8, 9). We based our hypothesis on these previous findings 212 as well as on recent evidence showing that salivary nitrite correlated positively with the intensity and 213 load of training (19). However, previous studies had some limitations requiring further examination. 214 For instance, in the first study by Rassaf et al (8) the correlation coefficient was only 0.37 which 215 suggests a large inter-individual variability in the cohort. In a second study by the same group, a 216 stronger association was found (r= 0.65) between plasma nitrite and heart rate at lactate anaerobic 217 threshold (9). From a physiological perspective this association is weak since heart rate at lactate 218 anaerobic threshold cannot be considered a valid marker of aerobic exercise capacity in humans (20). 219 Despite this, we found a positive and significant association between salivary nitrite and power in 220 females after exercise, this association was moderate (r_s = 0.44), and we did not see the same effect 8

between these two variables before exercise. Additionally, we did not observe a similar correlation between salivary nitrite and the VO_{2peak}, which is another key marker of aerobic capacity analysed in this study. Thus, the observed correlation between salivary nitrite and power in females after exercise, should be treated with caution, as it may be a casual association.

225

226 As expected, this study found significant differences between males and females on anthropometrical 227 and physiological variables, however no differences were found regarding salivary markers at 228 baseline. These results differ from a recent study indicating that females had higher salivary nitrite 229 concentrations than males which was linked to higher nitrate-reducing capacity, but not to a different 230 composition of the oral microbiome (12). Regarding this discrepancy, a limitation of our study was 231 that only 10 females took part in the study assessment of the nitrate-reducing capacity of oral bacteria 232 while Kapil et al (12) investigated 25 females. In addition, the combination of this small sample size, 233 and the lack of highly trained females (e.g VO_{2peak} > 60 mL/kg/min) may also explain the lack of 234 association between the oral nitrate-reducing capacity and exercise performance in our group of 235 females. This is supported by the fact that when collating all the subjects (males and females) together in order to increase the range of fitness levels we found that oral nitrate-reducing capacity was 236 237 strongly correlated with W_{peak} as well as VO_{2peak} before and after exercise.

238

239 A significant increase of salivary lactate was observed in both groups after exercise. In males, this was 240 accompanied by a significant decrease of salivary pH after exercise, but despite a similar pattern being 241 observed in females, differences were not statistically different in this group. Perhaps, a reason that 242 can explain, at least partially, these differences between males and females is the exercise-induced 243 stress levels. Limitations in female leg muscle strength as opposed to cardiovascular capacity could 244 limit their performance on the cycle ergometer test and explain the differences in the biological and 245 physiological markers observed in this study. However, further research would be needed to elucidate 246 this question. For instance, a previous study did not find an increase in salivary pH in a combined group 247 of males and females (21). However, these authors did not take into account gender differences, and 248 they measured salivary pH much earlier after exercise (5 min vs 20 min), which may reflect the 249 different kinetics of this metabolite.

250

Another interesting finding of this study was the lack of changes in salivary nitrite levels as well as the oral nitrate-reducing capacity after exercise. It has been suggested that the oral nitrate/nitrite pathway is enhanced under acidic conditions (2), but in spite of the low salivary pH and higher lactate concentrations observed after exercise, this did not induce any change on salivary nitrate and nitrite.
It is possible that these variations on the oral pH balance were too small to stimulate the nitrate/nitrite
reduction response compared to the changes that have been observed in the stomach under much
more acidic conditions (22).

258

259 We also compared the level of agreement between salivary strips and a standard laboratory assay in 260 order to assess salivary nitrite concentration. A stronger correlation was found between pre-exercise 261 measurements compared to post-exercise measurements. However, the level of agreement of both 262 correlations (pre and post-exercise) was much lower than values (r= 0.76 and r= 0.59) reported by 263 other two previous studies (23, 24). Perhaps, this discrepancy may be explained by differences in the 264 population and methodology of studies. For instance, Modi et al (23) recruited twenty overweight and 265 obese individuals with an average BMI of 34, and showing low levels of salivary nitrite compared to 266 our participants. Another study by McDonagh et al (24) found a significant correlation between 267 laboratory measurements and the salivary strip indicators following an increase of salivary nitrite due 268 to the ingestion of beetroot at different doses in 10 subjects. This could provide high readings that are 269 much easier to compare with reference values from the salivary strips than low readings. However, 270 from a statistical viewpoint the current study was much more powerful than previous studies by Modi 271 et al (23) and McDonagh et al (24) as measurements were taken from 50 individuals compared to 20 272 and 10, respectively.

273

274 This study has some limitations that are important to highlight. For instance, a direct screening to 275 assess oral health was not performed. While participants completed a medical questionnaire to assess 276 their medical and oral health before the start of the study, this did not exclude participants carrying 277 some degree of gum disease. It has been estimated that prevalence of gum disease in adults living in 278 the UK is over 45% (25). This is relevant since periodontal disease can seriously disrupt the equilibrium 279 of the oral ecosystem (26), and can also increase the levels of salivary nitrate and nitrite due to 280 immune-mediated inflammatory response (27). Thus, future studies should examine the oral health 281 of participants more in depth. Additionally, we did not analyse the salivary flow rate and salivary 282 buffering capacity. It may be interesting in future studies to assess whether a higher salivary flow rate 283 after exercise is a key mechanism for increasing the availability of circulatory nitrite. Furthermore, we 284 did not assess the diversity of the oral microbiome due to funding constraints and because this 285 question was not originally planned. Last but not least, it would be interesting to assess other non-

- 286 invasive cardiovascular parameters such as blood pressure and flow-mediated arterial dilation. The
- 287 questions arising from the study are shaping our current research.

288 Conclusion

- 289 This is the first study to reveal that the nitrate-reducing activity of oral bacteria is related to the aerobic
- exercise capacity in healthy humans. Further research is needed to investigate more in depth the
- 291 effect of exercise on the activity and diversity of the oral microbiome as well as inherited factors that
- can also play a key role on the oral ecosystem.
- 293

294 Acknowledgements

- 295 We would like to acknowledge Liz Preston, Vicki Cammack and Natalie Sweet for their technical
- support during this study. In addition, we would like to thank to all the participants their contribution
- to this study. A research grant from the Institute of Health & Community (University of Plymouth, UK)
- supported this study.

299 References

- Lundberg JO, Carlström M, Weitzberg E. Metabolic Effects of Dietary Nitrate in Health and
 Disease. Cell Metabolism. 2018;28(1):9-22.
- Lundberg JO, Weitzberg E, Gladwin MT. The nitrate-nitrite-nitric oxide pathway in physiology
 and therapeutics. Nat Rev Drug Discov. 2008;7(2):156-67.
- 304 3. Omar SA, Webb AJ, Lundberg JO, Weitzberg E. Therapeutic effects of inorganic nitrate and 305 nitrite in cardiovascular and metabolic diseases. Journal of Internal Medicine. 2015:n/a-n/a.
- Ashworth A, Bescos R. Dietary nitrate and blood pressure: evolution of a new nutrient?
 Nutrition Research Reviews. 2017:1-12.
- 3085.Bescos R, Sureda A, Tur JA, Pons A. The effect of nitric-oxide-related supplements on human309performance. Sports Med. 2012;42(2):99-117.
- Jonvik KL, Nyakayiru J, van Loon LJC, Verdijk LB. Can elite athletes benefit from dietary nitrate
 supplementation?2015 2015-05-21 20:11:45.
- Jensen L, Bangsbo J, Hellsten Y. Effect of high intensity training on capillarization and presence
 of angiogenic factors in human skeletal muscle. J Physiol. 2004;557(Pt 2):571-82.
- Rassaf T, Lauer T, Heiss C, Balzer J, Mangold S, Leyendecker T, et al. Nitric oxide synthasederived plasma nitrite predicts exercise capacity. Br J Sports Med. 2007;41(10):669-73; discussion 73.
- Totzeck M, Hendgen-Cotta UB, Rammos C, Frommke LM, Knackstedt C, Predel HG, et al. Higher
 endogenous nitrite levels are associated with superior exercise capacity in highly trained
 athletes. Nitric Oxide. 2012;27(2):75-81.
- Kapil V, Haydar SMA, Pearl V, Lundberg JO, Weitzberg E, Ahluwalia A. Physiological role for nitrate-reducing oral bacteria in blood pressure control. Free Radical Biology and Medicine.
 2013;55:93-100.
- 11. Clodfelter WH, Basu S, Bolden C, Dos Santos PC, King SB, Kim-Shapiro DB. The relationship
 between plasma and salivary NOx. Nitric Oxide. 2015;47:85-90.
- Kapil V, Rathod KS, Khambata RS, Bahra M, Velmurugan S, Purba A, et al. Sex differences in the
 nitrate-nitrite-NO• pathway: Role of oral nitrate-reducing bacteria. Free Radical Biology and
 Medicine. 2018;126:113-21.
- Allen JM, Mailing LJ, Niemiro GM, Moore R, Cook MD, White BA, et al. Exercise Alters Gut
 Microbiota Composition and Function in Lean and Obese Humans. Med Sci Sports Exerc. 2017.
- 14. Clarke SF, Murphy EF, O'Sullivan O, Lucey AJ, Humphreys M, Hogan A, et al. Exercise and
 associated dietary extremes impact on gut microbial diversity. Gut. 2014;63(12):1913-20.
- Mach N, Fuster-Botella D. Endurance exercise and gut microbiota: A review. Journal of Sport
 and Health Science. 2017;6(2):179-97.
- McFadzean. Exercise can help modulate human gut microbiota.
 <u>https://scholar.colorado.edu/cgi/viewcontent.cgi?article=1154&context=honr_theses</u>:
 University of Colorado Boulder; 2014.
- Porcelli S, Ramaglia M, Bellistri G, Pavei G, Pugliese L, Montorsi M, et al. Aerobic fitness affects
 the exercise performance responses to nitrate supplementation. Med Sci Sports Exerc.
 2015;47(8):1643-51.
- 18. Chen H, Peng S, Dai L, Zou Q, Yi B, Yang X, et al. Oral microbial community assembly under the
 influence of periodontitis. PLoS One. 2017;12(8):e0182259.
- 19. Díaz Gómez MM, Bocanegra Jaramillo OL, Teixeira RR, Espindola FS. Salivary surrogates of
 plasma nitrite and catecholamines during a 21-week training season in swimmers. PloS one.
- 344 2013;8(5):e64043-e.

- Kuipers H, Keizer HA, de Vries T, van Rijthoven P, Wijts M. Comparison of heart rate as a non invasive determinant of anaerobic threshold with the lactate threshold when cycling. European
 Journal of Applied Physiology and Occupational Physiology. 1988;58(3):303-6.
- Frese C, Frese F, Kuhlmann S, Saure D, Reljic D, Staehle HJ, et al. Effect of endurance training on
 dental erosion, caries, and saliva. Scandinavian Journal of Medicine & Science in Sports.
 2015;25(3):e319-e26.
- Benjamin N, O'Driscoll F, Dougall H, Duncan C, Smith L, Golden M, et al. Stomach NO synthesis.
 Nature. 1994;368(6471):502.
- Modi A, Morou-Bermudez E, Vergara J, Patel RP, Nichols A, Joshipura K. Validation of two point of-care tests against standard lab measures of NO in saliva and in serum. Nitric Oxide.
 2017;64:16-21.
- McDonagh ST, Wylie LJ, Webster JM, Vanhatalo A, Jones AM. Influence of dietary nitrate food
 forms on nitrate metabolism and blood pressure in healthy normotensive adults. Nitric Oxide.
 2018;72:66-74.
- 359 25. Steele J, O'Sullivan A. Adult Dental Health Survey 2009—Summary report. London: National
 360 Health Service Health and Social Care Information Centre. 2011.
- 361 26. Kilian M, Chapple I, Hannig M, Marsh P, Meuric V, Pedersen A, et al. The oral microbiome-an
 362 update for oral healthcare professionals. British Dental Journal. 2016;221(10):657-66.
- 363 27. Sánchez GA, Miozza VA, Delgado A, Busch L. Total salivary nitrates and nitrites in oral health and
 364 periodontal disease. Nitric Oxide. 2014;36(0):31-5.

365	Figures & tables
366	
367 368	Figure 1 : Salivary concentration of lactate and in males ($n = 28$), females ($n = 22$) and in all ($n = 50$) the participants together before and 20 min after exercise (mean ± SD).
369 370	
371 372	Figure 2 : Salivary concentration of nitrate in males (<i>n</i> = 28) and females (<i>n</i> = 22) as well as nitrate reducing capacity (<i>n</i> males = 15; <i>n</i> females = 10) before and after exercise (mean ± SD).
373	
374	
375 376	Figure 3 : Spearman's rank correlation between the oral nitrate-reducing capacity of bacteria and maximum power (A) and aerobic capacity (B) in males and females (<i>n</i> = 25).
377	

378	Table 1: Main characteristics of the participants (mean ± SD).
-----	--

		Males			Females		All
	Without	With	Overall	Without	With	Overall	
	ONRC	ONRC		ONRC	ONRC		
Participants (n)	13	15	28	12	10	22	50
Age (y/o)	43.9 ± 14.2	40.6 ± 15.4	38.9 ± 14.8	38.8 ± 13.2	29.3 ± 10.7	38.1 ± 13.9*	38.8 ± 14.3
Body mass (kg)	76.0 ± 15.1	76.6 ± 9.3	72.1 ± 13.4	62.0 ± 9.5	59.6 ± 7.8	66.8 ± 13.6*	69.7 ± 13.2
BMI	23.3 ± 2.7	23.7 ± 3.5	23.0 ± 3.0	22.3 ± 2.7	21.6 ± 2.4	22.5 ± 2.6	22.8 ± 2.9
Body fat (%)	16.1 ± 4.8	16.4 ± 5.4	17.9 ± 5.8	25.7 ± 6.0	22.5 ± 5.5	21.5 ± 6.3*	19.8 ± 6.7
VO _{2peak}	50.7 ± 10.7	52.7 ± 12.7	49.3 ± 11.5	42.2 ± 9.2	42.2 ± 8.1	42.7 ± 9.0*	47.6 ± 11.4
(mL/kg/min)							
W _{peak} (W/kg)	4.5 ± 1.0	4.4 ± 1.1	4.3 ± 1.1	3.8 ± 0.9	3.6 ± 0.8	$4.0 \pm 0.9^{*}$	4.2 ± 1.0
Salivary NO ³	189 ± 255	225 ± 234	225 ± 278	360 ± 355	154 ± 106	223 ± 281	225 ± 264
(μM/L)							
Salivary NO ²	142 ± 110	74 ± 63	100 ± 88	154 ± 106	78 ± 69	128 ± 101	113 ± 95
(μM/L)							
ONRC (nitrite	-	275 ± 138		-	231 ± 143		257 ± 138
μM)							
Salivary glucose	0.04 ± 0.03	0.04 ± 0.03	0.03 ± 0.03	0.03 ± 0.02	0.02 ± 0.01	0.03 ± 0.02	0.03 ± 0.03
(μM/L)							
Salivary lactate	0.21 ± 0.20	0.18 ± 0.13	0.17 ± 0.15	0.13 ± 0.09	0.11 ± 0.09	0.16 ± 0.15	0.17 ± 0.14
(μM/L)							
Salivary pH	7.41 ± 0.37	7.25 ± 0.31	7.26 ± 0.33	7.34 ± 0.46	7.07 ± 0.23	7.29 ± 0.39	7.28 ± 0.36

 VO_{2peak} : maximum oxygen consumption; W_{peak} : peak power in watts; NO³: nitrate; NO²: nitrite; ONRC: Oral-nitrate reducing capacity; * Statistical difference (P < 0.05) 380

Table 2: Correlations between salivary nitrite estimated with salivary strips and laboratory measurements using high-performance liquid chromatography (HPLC) system (mean ± SD).

	Salivary Nitrite HPLC Pre-exercise	Salivary Nitrite HPLC Post-exercise		
n = 50 (28 M; 22 F)				
Salivary Nitrite (Berkeley Strips)	0.43 (0.002)*	0.32 (0.02) *		

* Statistical difference (P < 0.05)

Table 3: Correlations between salivary markers (nitrate [NO³]; nitrite [NO²]; lactate; pH; and the oral nitrate-reducing capacity [ONRC]) and aerobic exercise performance parameters in males and females (mean ± SD).

	Males $(n = 28)$					Females				AII (n = 50)			
	Pre-Exercise Post-Exercise			Pre-Exercise Post-Exercise			Pre-Exercise Post-Exercise			xercise			
	Power (W/kg) r _s (p)	VO _{2peak} (mL/Kg/min) r _s (p)	Power (W/kg) r _s (p)	VO _{2peak} (mL/Kg/min) r _s (p)	Power (W/kg) r _s (p)	VO _{2peak} (mL/Kg/min) r _s (p)	Power (W/kg) r _s (p)	VO _{2peak} (mL/Kg/min) r _s (p)	Power (W/kg) r _s (p)	VO _{2peak} (mL/Kg/min) r _s (p)	Power (W/kg) r _s (p)	VO _{2peak} (mL/Kg/min) r _s (p)	
Salivary NO ³	0.16 (0.41)	0.15 (0.45)	0.07 (0.78)	0.05 (0.81)	0.38 (0.08)	0.19 (0.40)	0.37 (0.09)	0.19 (0.41)	0.18 (0.21)	0.11 (0.46)	0.09 (0.56)	0.01 (0.95)	
Salivary NO ²	0.22 (0.27)	0.21 (0.28)	0.09 (0.65)	0.12 (0.55)	0.26 (0.24)	0.16 (0.49)	0.44 (0.04)*	0.31 (0.16)	0.20 (0.17)	0.16 (0.28)	0.19 (0.18)	0.18 (0.22)	
Salivary lactate	0.19 (0.34)	0.14 (0.48)	0.02 (0.93)	-0.06 (0.75)	0.40 (0.07)	0.21 (0.35)	0.41 (0.06)	0.33 (0.14)	0.33 (0.02)*	0.26 (0.07)	0.19 (0.20)	0.14 (0.34)	
Salivary pH	0.03 (0.86)	-0.90 (0.66)	0.21 (0.29)	0.15 (0.46)	0.12 (0.61)	0.19 (0.41)	0.11 (0.64)	0.14 (0.52)	0.04 (0.81)	-0.01 (0.99)	0.04 (0.80)	-0.01 (0.99)	
	(<i>n</i> = 15)				(<i>n</i> = 10)			(<i>n</i> = 25)					
ONRC	0.64 (0.01)*	0.54 (0.04)*	0.64 (0.01)*	0.47 (0.08)	0.50 (0.14)	0.43 (0.22)	0.41 (0.24)	0.51 (0.13)	0.64 (0.001)*	0.54 (0.005)*	0.63 (0.001)*	0.54 (0.003)*	

* Statistical difference (P < 0.05)









Oral Nitrate-Reducing Capacity (µM)



Oral Nitrate-Reducing Capacity (µM)