2011-04

Inhalational or total intravenous anaesthesia: is total intravenous anaesthesia useful and are there economic benefits?

Sneyd, John

<http://hdl.handle.net/10026.1/1466>

10.1097/aco.0b013e328343f3ac

Current Opinion in Anaesthesiology

Ovid Technologies (Wolters Kluwer Health)

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Inhalational or total intravenous anaesthesia: is total intravenous anaesthesia useful and are there economic benefits?

J. Robert Sneyd^a and Katherine A. Holmes^b

^aPeninsula Medical School, University of Plymouth and

^bPlymouth Hospitals NHS Trust, Plymouth, UK

Correspondence to J.R. Sneyd, Peninsula College of Medicine and Dentistry, The John Bull Building, Research Way, Tamar Science Park, Plymouth, PL6 8BU, UK

Tel: +44 1752 437355; fax: +44 1752 517842;

e-mail: Robert.sneyd@pms.ac.uk

Current Opinion in Anaesthesiology 2011, 24:000–000

Purpose of review

The comparison of inhalational and intravenous anaesthesia has been the subject of many controlled trials and meta-analyses. These reported diverse endpoints typically including measures of the speed and quality of induction of anaesthesia, haemodynamic changes, operating conditions, various measures of awakening, postoperative nausea and vomiting and discharge from recovery and hospital as well as recovery of psychomotor function. In a more patient-focused Health Service, measures with greater credibility are overall patient satisfaction, time to return to work and long-term morbidity and mortality. In practice, studies using easier to measure proxy endpoints dominate – even though the limitations of such research are well known.

Recent findings

Recent study endpoints are more ambitious and include impact on survival from cancer and the possibility of differential neurotoxic impact on the developing brain and implications for neuro-behavioural performance.

Summary

Economic analysis of anaesthesia is complex and most published studies are naive, focusing on drug acquisition costs and facility timings. Real health economics are much more difficult. Preferred outcome measures would be whole institution costs or the ability to reliably add an extra case to an operating list, close an operating room and reduce the number of operating sessions offered or permanently decrease staffing. Alongside this, however, potential long-term patient outcomes should be considered.

Keywords

cost-benefit, economics, inhalational, intravenous, propofol, total intravenous anaesthesia

Curr Opin Anaesthesiol 24:000–000

© 2011 Wolters Kluwer Health | Lippincott Williams & Wilkins
0952-7907

Introduction

In daily practice, clinicians typically choose drugs with which they are familiar, tailoring anaesthetic techniques to perceived patient benefit and personal preference rather than economic benefit. Are these aims conflicting? Are they even different? Making best use of finite resources in the face of increasing demand is a priority given the recession and public sector funding cuts in the UK and changes in the structure of healthcare provision in the USA.

The review explores the potential economic impact of choice of general anaesthetic, comparing intravenous and inhalational agents along the patient pathway from operating room to discharge and beyond. We have focused on the contemporary general anaesthetics propofol, sevoflurane, isoflurane and desflurane.

Aspects of economics

Consideration of economics and anaesthesia is not new. Rowe [1], and Kettler and Crozier [2,3] have explored the

complexity and scope of the subject, and identified the need to analyse costs with a global perspective to properly assess cost-effectiveness. What does this imply? Different types of economic analysis [1] applicable to health-care are summarized in Table 1 [1]. The choice of analysis depends on how benefits or outputs are to be treated. A cost-identification analysis is an element of all medical economic studies but can be seen as synonymous with cost-minimization studies [4].

Costs can be described in a number of different ways (Table 2) [1,5,6]. This can include total costs (fixed and variable) and average costs (total cost divided by total number of cases). Total costs can also be direct costs and indirect costs. Health economists would refer to the latter as those from loss of productivity [5].

Cost considerations

In the anaesthetic literature the approach to economic comparison of total intravenous anaesthesia (TIVA) and inhalational anaesthesia is often by their direct cost

2 Ethics, economics and outcome

[7–10]. There is disagreement regarding the regimen with least drug cost, which may reflect differences in caseload, the use of low-flow anaesthesia [10,11], variable acquisition costs in different parts of the world [2] or the expiry of patents. On average, the use of TIVA gave higher drug costs [7,9,12–15], particularly if wastage [16] or cost of consumables such as syringes and pumps were included. This excess cost declined with longer operations [17]. Many studies looked beyond direct costs, assessing cost-effectiveness rather than minimization, recognizing that a cheaper anaesthetic that results in postoperative pain or emesis may incur additional costs and delay patient discharge [18]. Opportunities for savings in the operating room include time, cost of other consumables and operating conditions, which in turn may affect blood loss, surgical outcome and length of stay. Cole *et al.* [19] report equal haemodynamic stability with both inhalational and intravenous anaesthesia for craniotomy but suggest propofol may reduce intracranial pressure and increase cerebral perfusion pressure. There was also no haemodynamic difference found in septorhinoplasty patients, using remifentanyl in both groups [20].

In cardiac surgery volatile anaesthetics may be cardioprotective [21]. However, Flier *et al.* [22] found no benefit on cardiac morbidity and mortality or troponin concentrations after surgery. In general surgery, a faster recovery of bowel function and earlier discharge was reported when propofol was compared to desflurane for laparoscopic cholecystectomy [23].

Areas of economic impact in the recovery area or postanesthesia care unit (PACU) include length of stay, costs, drug use and the ability to ‘step down’ to a less intensive, and therefore less costly clinical area.

The most common finding is that TIVA reduces the incidence of postoperative nausea and vomiting, in the early recovery period [23–27], and the decreased requirement for antiemetics may off-set a higher initial drug cost. Further, patient dissatisfaction is often attributed to nausea and vomiting – complications that both patients and their anaesthetists wish to avoid [28,29] although Fisher warns against the use of surrogate outcomes in this regard [30].

Key points

- There is increased scrutiny of all operating theatre costs and drug budgets are easily targeted.
- Total intravenous anaesthesia appears to provide decreased emesis and good recovery but often at higher direct costs.
- Anaesthetic agents have been implicated as affecting both long-term neurological development and cancer recurrence and survival and further investigative trials are underway.
- True health economics is complex and should include all costs and benefits relating to the health-care institution, the patient and society.

Some studies report a faster or better quality recovery with TIVA [8]; however, this seems to have little impact on subsequent patient progress [31]. Other investigators described faster recovery when inhalational agents were used [16] but without difference in recovery of cognitive function [25,32]. Some recent studies suggest increased postoperative pain following isoflurane and sevoflurane anaesthesia when compared to propofol [33,34] although the suggestion that propofol has an analgesic effect has been disputed [35] and the possibility of an antianalgesia effect of residual volatile anaesthetic proposed instead.

Although anaesthetic drugs cost can vary between TIVA and inhaled anaesthesia, any saving from cheaper drugs is minimal compared to operating room or hospital episode costs [36,37]. Despite this, although anaesthetic drug costs are comparatively small they are, from a managerial viewpoint, very visible [5,38].

Organizational issues

Any potential reduction in cost from faster and better recovery from anaesthesia may be negated by a delay in starting theatre cases, prolonged turnover time [39] and downstream factors such as PACU staffing patterns. Staffing is typically the greatest cost for a recovery area and if a shorter recovery room stay does not result in the reduction of staff hours or numbers then there is no real economic saving [40,41]. Restructuring of PACU patient inflow may have more impact; if patients are suitable to either bypass PACU or be ‘fast-tracked’ to a nominated area for faster discharge [42].

Table 1 Types of economic analysis applicable to anaesthesia

	Input	Output	Output examples
Cost minimization (cost identification)	Direct costs	Not applicable	Outcomes assumed equal
Cost-benefit	All costs	Economic benefits (benefits as monetary units)	Money saved, production gains or return to work
Cost-effectiveness	All costs	Natural units (measured outcomes used directly)	Numbers free from nausea, successfully treated cases
Cost-utility	All costs	Utility units (outcomes converted to common unit)	Quality adjusted life years

Adapted from [1].

Table 2 Methods of describing costs in economic analysis

Direct costs	Material and disposables used/wasted
Indirect costs	Resulting from event, for example loss of production
Fixed costs	Remain the same, for example building rent
Variable costs	Increases with case numbers, for example drugs
Marginal costs	Cost of one more/ fewer case
Intangible costs	Nonphysical, for example goodwill

Adapted from [1] (originally adapted from [5,6]).

Beyond hospital discharge

Current economic pressures make explicit the requirement that healthcare should be evidence-based and delivered efficiently. Anaesthetists need to understand the economics of anaesthesia care [43] as 'medical decision-making cannot be divorced from cost' [44]. This is a complex undertaking.

Potential economic benefits may be found along the entire peri-operative track and beyond it. Improvements in patient wellbeing after discharge may impact on return to work, dependency and social functioning; however, whilst these likely have economic impact to the patient, improving them is unlikely to have financial benefits to the healthcare provider. But benefits and costs can be looked at from the different perspectives of the patient, provider or society [4]. Commonly medical studies consider patient benefits alone. So, what of the patient once they are discharged? Few data describe anaesthetic drug effects on patients' experience after discharge. Endpoints of interest would be the need to re-access healthcare including visits to primary care or emergency department attendance, perhaps re-admission.

Carvalho *et al.* [45] explored functional recovery during the week following discharge after laparoscopic sterilization. Functional recovery was slow in both inhalational and TIVA groups and not affected by anaesthetic technique [45]. Sung *et al.* [46] reported that patients anaesthetized with propofol group required less nursing care and returned to work earlier those receiving isoflurane during breast biopsy.

Neurological effects

True health economics considers the impact on society as a whole and this takes us from ill-health to wellbeing. Once our patients stop being patients, how long is it before they return to their normal life? Do they return to work or become a financial burden on the state? What is their quality of life? Steinmetz *et al.* [47] reported that elderly patients with cognitive dysfunction 1 week after noncardiac surgery had increased dependency on social transfer payments and increased likelihood of leaving the workforce prematurely. Those in whom the cognitive

dysfunction persisted to 3 months experienced increased mortality [47]. What we do not know is whether the type of general anaesthetic used affected the risk of this postoperative cognitive dysfunction; other risk factors may be more important [48]. Recent investigations in rats linked cognitive impairment to hippocampus pro-inflammatory cytokine release precipitated by surgery rather than anaesthesia [49] with greater effects in older animals. Inhalational general anaesthesia can lead to amyloid deposition as seen in Alzheimer's disease [50[•]], but in animal studies isoflurane impacts negatively on neonatal rats rather than adult ones, by reducing neuro-genesis and causing memory deficit [51,52].

The impact of anaesthesia on long-term outcomes is an important contemporary research area and an enlarging and concerning body of evidence. Anaesthesia, as whole, is relatively well tolerated [53] and our drugs have a short duration of effect with a generally reliable swift recovery. Our impact on our patients is brief and beneficial. How uncomfortable is it then, to consider the latest articles looking at neurotoxicity from anaesthesia in children or the differential risk of cancer metastasis dependent on anaesthetic technique? Are we actually doing harm?

Emergence agitation in children after sevoflurane is well known, appears short-lived [54] and was minimally reduced by switching to propofol [55]. However, children undergoing urological surgery before the age of 2 years showed a trend towards an increased incidence of behavioural disturbance [56] and children undergoing multiple general anaesthetics before the age of 4 were more likely to develop learning disabilities later [57]. It is not clear if some anaesthetics are better tolerated than others [58[•]] and separating the long-term effects of the condition necessitating surgery and the effects of the procedure itself are problematic. In any case there are times when the requirement for a general anaesthetic cannot be avoided, hence further studies are needed [59] and many are already underway [60^{••}].

Cancer outcomes

Of serious concern are the possible interactions of anaesthetic technique and cancer recurrence [61]. Surgery for tumour removal is associated with the release of tumour cells and the balance between the ability of these to seed and the body's immune defence determines the development of clinical metastases [62[•]]. The peri-operative factors that can contribute to this process are surgery which causes release of tumour cells and growth factors, reduction of antiangiogenic factors and depression of the cellular immune system, volatile anaesthesia-related impairment of natural killer (NK) cells (amongst others) and opioid-induced inhibition of both cellular and humoral immunity [63^{••},64[•]].

4 Ethics, economics and outcome

Biki *et al.* [65] reported that patients undergoing radical prostatectomy showed less biochemical cancer recurrence when a general anaesthetic was combined with using an epidural in place of opioids. Similarly, the use of paravertebral block to provide analgesia for breast cancer surgery reduced the risk of tumour recurrence [66]. Gottschalk *et al.* [67**] found epidural use gave no decrease in cancer recurrence after colorectal surgery but there was a potential benefit in older patients, who may have had a different tumour type.

Regional anaesthesia might improve cancer outcomes by reduction of surgical stress, decreasing need for opioids (morphine promotes angiogenesis) and by allowing the administration of lower amounts of volatile anaesthetic. In contrast to volatile agents, propofol attenuated the surgical stress response, not suppressing NK cell activity or promoting lung tumour metastases and *in vitro* inhibited breast tumour cell proliferation [68,69].

The clinical trials so far have been retrospective and descriptive but randomized trials are in progress [70].

Conclusion

Whilst in the past the use of total intravenous anaesthesia has appeared to incur higher drug and equipment acquisition cost, it has favourable effects such as reduced postoperative nausea and vomiting and less immunosuppression in the perioperative period. Nevertheless with only limited economic analysis of true outcome measures and at present only retrospective evidence of potentially improved cancer survival, as yet we do not have the evidence to provide a definitive answer to a complex question.

There is continuing development of novel anaesthetic compounds [71] and these need to be evaluated for cost-effectiveness and any effect on long-term outcomes. Providing robust evidence will be the challenge. In the meantime it is likely that a careful technique, tailored to the individual patient's needs, operation and outcome is unlikely to have a drastic impact on institutional spending but we should still do our part, however small. If new evidence shows that our choice of anaesthetic has real impact on postoperative function or patient survival then we will need to balance small-scale economics with the bigger picture and first, do no harm.

Acknowledgements

No funding or grants were received for this article.

J.R.S. acted as a paid adviser to AstraZeneca, the inventors of propofol during 2010.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

Additional references related to this topic can also be found in the Current World Literature section in this issue (pp. 000–000).

- 1 Rowe WL. Economics and anaesthesia. *Anaesthesia* 1998; 53:782–788.
- 2 Kettler D, Crozier T. Cost-effectiveness of anaesthesia. *Curr Opin Anaesthesiol* 2001; 14:569–572.
- 3 Crozier TA, Kettler D. Cost effectiveness of general anaesthesia: inhalation vs. i.v. *Br J Anaesth* 1999; 83:547–549.
- 4 Sperry RJ. Principles of economic analysis. *Anesthesiology* 1997; 86:1197–1205.
- 5 Watcha MF, White PF. Economics of anesthetic practice. *Anesthesiology* 1997; 86:1170–1196.
- 6 Davidson S, Stickney CP, Weil RL. Accounting: the language of the business. In 7th Sun Lakes, Arizona: Thomas Horton and Daughters; 1987.
- 7 Stevanovic PD, Petrova G, Miljkovic B, *et al.* Low fresh gas flow balanced anesthesia versus target controlled intravenous infusion anesthesia in laparoscopic cholecystectomy: a cost-minimization analysis. *Clin Therapeut* 2008; 30:1714–1725.
- 8 Chan SM, Horng HC, Huang ST, *et al.* Drug cost analysis of three anesthetic regimens in prolonged lumbar spinal surgery. *J Med Sci* 2009; 29:075–080.
- 9 Horng HC, Kuo CP, Ho CC, *et al.* Cost analysis of three anesthetic regimens under auditory evoked potentials monitoring in gynecologic laparoscopic surgery. *Acta Anaesthesiol Taiwan* 2007; 45:205–210.
- 10 Boldt J, Muller M, Uphus D, Hempelmann G. Cost aspects in anesthesia. Propofol versus isoflurane anesthesia [German] Kostenaspekte in der Anästhesie. Propofol- versus Isoflurananästhesie. *Der Anaesthesist* 1996; 45:731–736.
- 11 Bangaari A, Bangia R, Puri GD. Prevalence of low flow anesthesia and its cost comparison with other anesthesia techniques in an Indian tertiary care center. *Acta Anaesthesiol Taiwan* 2005; 43:141–145.
- 12 Lomartire N, Caracciolo F. Cost-effectiveness of balanced and total intravenous anaesthesia with or without Navigator device: preliminary results. *Eur J Anaesthesiol* 2009; 26:41–42.
- 13 Siampalioti A, Aretha A, Monantera G, *et al.* Does sevoflurane offer advantages over intravenous anaesthetic agents in morbid obese patients undergoing major surgery? *Eur J Anaesthesiol* 2009; 26.
- 14 Smith I, Terhoeve PA, Hennart D, *et al.* A multicentre comparison of the costs of anaesthesia with sevoflurane or propofol. *Br J Anaesth* 1999; 83:564–570.
- 15 Smith I, Thwaites AJ. Target-controlled propofol vs. sevoflurane: a double-blind, randomised comparison in day-case anaesthesia. *Anaesthesia* 1999; 54:745–752.
- 16 Dolk A, Cannerfelt R, Anderson RE, Jakobsson J. Inhalation anaesthesia is cost-effective for ambulatory surgery: a clinical comparison with propofol during elective knee arthroscopy. *Eur J Anaesthesiol* 2002; 19:88–92.
- 17 Biro P, Suter G, Alon E. Intravenous anaesthesia with propofol versus thiopentone/enflurane: an analysis of consumption and costs [German]. *Anaesthesist* 1995; 44:163–170.
- 18 Nathan N, Rezzoug A, Dolan P, *et al.* Propofol versus isoflurane: a comparison of costs [French]. *Annales Francaises d'Anesthesie et de Reanimation* 1993; 12:571–574.
- 19 Cole CD, Gottfried ON, Gupta DK, Couldwell WT. Total intravenous anesthesia: advantages for intracranial surgery. *Neurosurgery* 2007; 61:369–377; discussion 377–378.
- 20 Gokce BM, Ozkose Z, Tuncer B, *et al.* Hemodynamic effects, recovery profiles, and costs of remifentanyl-based anesthesia with propofol or desflurane for septorhinoplasty. *Saudi Med J* 2007; 28:358–363.
- 21 Landoni G, Fochi O, Tritapepe L, *et al.* Cardiac protection by volatile anesthetics. A review. *Minerva Anesthesiol* 2009; 75:269–273.
- 22 Flier S, Post J, Concepcion AN, *et al.* Influence of propofol-opioid vs. isoflurane-opioid anaesthesia on postoperative troponin release in patients undergoing coronary artery bypass grafting. *Br J Anaesth* 2010; 105:122–130.
- 23 Akkurt BCO, Temiz M, Inanoglu K, *et al.* Comparison of recovery characteristics, postoperative nausea and vomiting, and gastrointestinal motility with total intravenous anesthesia with propofol versus inhalation anesthesia with desflurane for laparoscopic cholecystectomy: a randomized controlled study. *Curr Therapeut Res Clin Experiment* 2009; 70:94–103.

- 24 Vari A, Gazzanelli S, Cavallaro G, *et al.* Postoperative nausea and vomiting (PONV) after thyroid surgery: a prospective, randomized study comparing totally intravenous versus inhalational anesthetics. *Am Surg* 2010; 76:325–328.
 - 25 Rohm KD, Piper SN, Suttner S, *et al.* Early recovery, cognitive function and costs of a desflurane inhalational vs. a total intravenous anaesthesia regimen in long-term surgery. *Acta Anaesthesiol Scand* 2006; 50:14–18.
 - 26 Gupta A, Stierer T, Zuckerman R, *et al.* Comparison of recovery profile after ambulatory anesthesia with propofol, isoflurane, sevoflurane and desflurane: a systematic review. *Anesth Analg* 2004; 98:632–641.
 - 27 Sneyd JR, Carr A, Byrom WD, Bilski AJ. A meta-analysis of nausea and vomiting following maintenance of anaesthesia with propofol or inhalational agents. *Eur J Anaesthesiol* 1998; 15:433–445.
 - 28 Macario A, Weinger M, Carney S, Kim A. Which clinical anesthesia outcomes are important to avoid? The perspective of patients. *Anesth Analg* 1999; 89:652–658.
 - 29 Macario A, Weinger M, Truong P, Lee M. Which clinical anesthesia outcomes are both common and important to avoid? The perspective of a panel of expert anesthesiologists. *Anesth Analg* 1999; 88:1085–1091.
 - 30 Fisher D. Surrogate outcomes: they don't get it. *Anesth Analg* 2009; 109:994; author reply 994–995.
 - 31 Mukherjee K, Seavell C, Rawlings E, Weiss A. A comparison of total intravenous with balanced anaesthesia for middle ear surgery: effects on postoperative nausea and vomiting, pain, and conditions of surgery. *Anaesthesia* 2003; 58:176–180.
 - 32 Lauter E, Abbinante C, Del Gaudio A, *et al.* Emergence times are similar with sevoflurane and total intravenous anesthesia: results of a multicenter RCT of patients scheduled for elective supratentorial craniotomy. *J Neurosurg Anesthesiol* 2010; 22:110–118.
 - 33 Cheng SS, Yeh J, Flood P. Anesthesia matters: patients anesthetized with propofol have less postoperative pain than those anesthetized with isoflurane. *Anesth Analg* 2008; 106:264–269; table of contents.
 - 34 Tan TMF, Bhinder RM, Carey MMDF, Briggs LMDF. Day-surgery patients anesthetized with propofol have less postoperative pain than those anesthetized with sevoflurane. *Anesth Analg* 2010; 111:83–85.
- The finding from this study that postoperative pain management can be affected simply by the choice of general anaesthetic agent is particularly relevant to day case anaesthesia and may have potential to affect case throughput.
- 35 White PF. CON: anesthesia versus analgesia: assessing the analgesic effects of anesthetic drugs. *Anesth Analg* 2010; 111:88–89.
 - 36 Macario A, Vitez TS, Dunn B, McDonald T. Where are the costs in perioperative care? Analysis of hospital costs and charges for inpatient surgical care. *Anesthesiology* 1995; 83:1138–1144.
 - 37 Demeere JL, Merckx C, Demeere N. Cost minimisation and cost effectiveness in anaesthesia for total hip replacement surgery, in Belgium? A study comparing three general anaesthesia techniques. *Acta Anaesthesiol Belg* 2006; 57:145–151.
 - 38 Macario A, Glenn D, Dexter F. What can the postanesthesia care unit manager do to decrease costs in the postanesthesia care unit? *J PeriAnesth Nurs* 1999; 14:284–293.
 - 39 White PF, Watcha MF. Pharmacoeconomics in anaesthesia: what are the issues? *Eur J Anaesthesiol Suppl* 2001; 18:10–15.
 - 40 Smith I. Cost considerations in the use of anaesthetic drugs. *Pharmacoeconomics* 2001; 19:469–481.
 - 41 Smith I. Total intravenous anaesthesia: is it worth the cost? *CNS Drugs* 2003; 17:609–619.
 - 42 White PF, Rawal S, Nguyen J, Watkins A. PACU fast-tracking: an alternative to 'bypassing' the PACU for facilitating the recovery process after ambulatory surgery. *J PeriAnesth Nurs* 2003; 18:247–253.
 - 43 Fisher DM. The 'Big Little Problem' of postoperative nausea and vomiting: do we know the answer yet? *Anesthesiology* 1997; 87:1271–1273.
 - 44 Eisenberg JM. Clinical economics. A guide to the economic analysis of clinical practices. *J Am Med Assoc* 1989; 262:2879–2886.
 - 45 Carvalho B, Benton JI, Vickery PJ, *et al.* Longterm functional recovery following day-case laparoscopic sterilisation: inhalational versus TIVA maintenance. *Ambulat Surg* 2002; 10:45–51.
 - 46 Sung YF, Reiss N, Tillette T. The differential cost of anesthesia and recovery with propofol-nitrous oxide anesthesia versus thiopental sodium-isoflurane-nitrous oxide anesthesia. *J Clin Anesth* 1991; 3:391–394.
 - 47 Steinmetz J, Christensen KB, Lund T, *et al.* Long-term consequences of postoperative cognitive dysfunction. *Anesthesiology* 2009; 110:548–555; 10.1097/ALN.0b013e318195b569.
 - 48 Maze M, Cibelli M, Grocott HP. Taking the lead in research into postoperative cognitive dysfunction. *Anesthesiology* 2008; 108:1–2.
 - 49 Cao XZ, Ma H, Wang JK, *et al.* Postoperative cognitive deficits and neuroinflammation in the hippocampus triggered by surgical trauma are exacerbated in aged rats. *Prog Neuropsychopharmacol Biol Psychiatry* 2010; 34:1426–1432.
 - 50 Fodale V, Santamaria LB, Schifilliti D, Mandal PK. Anaesthetics and postoperative cognitive dysfunction: a pathological mechanism mimicking Alzheimer's disease. *Anaesthesia* 2010; 65:388–395.
- This study suggests a potential mechanism of postoperative cognitive dysfunction related to the beta amyloid plaque oligomerization seen in Alzheimer's disease, which may lead to future clinical management options.
- 51 Zhu C, Gao J, Karlsson N, *et al.* Isoflurane anesthesia induced persistent, progressive memory impairment, caused a loss of neural stem cells, and reduced neurogenesis in young, but not adult, rodents. *J Cereb Blood Flow Metab* 2010; 30:1017–1030.
 - 52 Stratmann G, Sall JW, Bell JS, *et al.* Isoflurane does not affect brain cell death, hippocampal neurogenesis, or long-term neurocognitive outcome in aged rats. *Anesthesiology* 2010; 112:305–315.
 - 53 Gaba DM. Anaesthesiology as a model for patient safety in healthcare. *Br Med J* 2000; 320:785–788.
 - 54 Huddy N. Emergence agitation in children. *Br J Anaesth* 2010; 105:95; author reply 95–96.
 - 55 Pieters BJ, Penn E, Nicklaus P, *et al.* Emergence delirium and postoperative pain in children undergoing adenotonsillectomy: a comparison of propofol vs. sevoflurane anesthesia. *Paediatr Anaesth* 2010; 20:944–950.
 - 56 Kalkman CJ, Peelen L, Moons KG, *et al.* Behavior and development in children and age at the time of first anesthetic exposure. *Anesthesiology* 2009; 110:805–812.
 - 57 Wilder RT, Flick RP, Sprung J, *et al.* Early exposure to anesthesia and learning disabilities in a population-based birth cohort. *Anesthesiology* 2009; 110:796–804.
 - 58 Schifilliti D, Grasso G, Conti A, Fodale V. Anaesthetic-related neuroprotection: intravenous or inhalational agents? *CNS Drugs* 2010; 24:893–907.
- This study covers a variety of neurological situations, considering risk benefits in recent evidence and useful for decision-making in clinical practice.
- 59 Sun LS, Li G, DiMaggio C, *et al.* Anesthesia and neurodevelopment in children: time for an answer? *Anesthesiology* 2008; 109:757–761.
 - 60 Durieux M, Davis PJ. The safety of key inhaled and intravenous drugs in pediatrics (SAFEKIDS): an update. *Anesth Analg* 2010; 110:1265–1267.
- This editorial explains the SAFEKIDS Initiative, considers the current evidence with regard to neurotoxicity of general anaesthesia and observes that at present there is no evidence to postpone necessary surgery in children – an important point for clinicians and parents.
- 61 Arain MR, Buggy DJ. Anaesthesia for cancer patients. *Curr Opin Anaesthesiol* 2007; 20:247–253.
 - 62 Sessler DI. Long-term consequences of anesthetic management. *Anesthesiology* 2009; 111:1–4.
- This editorial is a concise summary of anaesthesia effects on surgical site infection and cancer recurrence but also usefully directs towards other relevant editorials on other types of outcomes.
- 63 Snyder GL, Greenberg S. Effect of anaesthetic technique and other perioperative factors on cancer recurrence. *Br J Anaesth* 2010; 105:106–115.
- This is a review article containing a comprehensive review of the pathophysiological basis of cancer recurrence. It looks at the place of surgery, perioperative medications and interventions within this valuable for reflection on current practice.
- 64 Bovill JG. Surgery for cancer: does anesthesia matter? *Anesth Analg* 2010; 110:1524–1526.
- Covering the evidence so far with regard to anaesthesia and cancer surgery, this editorial describes in particular the potential contribution of nonsteroidal anti-inflammatory drugs.
- 65 Biki B, Mascha E, Moriarty DC, *et al.* Anesthetic technique for radical prostatectomy surgery affects cancer recurrence: a retrospective analysis. *Anesthesiology* 2008; 109:180–187.
 - 66 Exadaktylos AK, Buggy DJ, Moriarty DC, *et al.* Can anesthetic technique for primary breast cancer surgery affect recurrence or metastasis? *Anesthesiology* 2006; 105:660–664.
 - 67 Gottschalk A, Ford JG, Regelin CC, *et al.* Association between epidural analgesia and cancer recurrence after colorectal cancer surgery. *Anesthesiology* 2010; 113:27–34.
- Whilst still retrospective in design, this study found a differential effect of epidurals on the recurrence of colorectal cancer, suggesting a more complex clinical picture of benefit than previous studies.

6 Ethics, economics and outcome

- 68** Melamed R, Bar-Yosef S, Shakhar G, *et al.* Suppression of natural killer cell activity and promotion of tumor metastasis by ketamine, thiopental, and halothane, but not by propofol: mediating mechanisms and prophylactic measures. *Anesth Analg* 2003; 97:1331–1339.
- 69** Deegan CA, Murray D, Doran P, *et al.* Effect of anaesthetic technique on oestrogen receptor-negative breast cancer cell function in vitro. *Br J Anaesth* 2009; 103:685–690.
- 70** Sessler DI, Ben-Eliyahu S, Mascha EJ, *et al.* Can regional analgesia reduce the risk of recurrence after breast cancer? Methodology of a multicenter randomized trial. *Contemp Clin Trials* 2008; 29:517–526.
- 71** Sneyd JR, Rigby-Jones AE. New drugs and technologies, intravenous anaesthesia is on the move (again). *Br J Anaesth* 2010; 105:246–254.

ACO

Manuscript No. 200719

Current Opinion in Anaesthesiology
Sort- Curr Opin
Anaesthesiol
Typeset by Thomson Digital
for Lippincott Williams & Wilkins

Dear Author,

During the preparation of your manuscript for typesetting, some queries have arisen. These are listed below. Please check your typeset proof carefully and mark any corrections in the margin as neatly as possible or compile them as a separate list. This form should then be returned with your marked proof/list of corrections to the Production Editor.

QUERIES: to be answered by AUTHOR/EDITOR

QUERY NO.	QUERY DETAILS	RESPONSE
NO QUERY		