2013-08-08

Brief guided imagery and body scanning interventions reduce food cravings

Hamilton, J

<http://hdl.handle.net/10026.1/1604>

10.1016/j.appet.2013.08.005

Appetite

Elsevier BV

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

This is the Authors' copy of the paper to be published as:
Hamilton, J., Fawson, S., May, J., Andrade, J., & Kavanagh, D.J. (in press) Brief guided
imagery and body scanning interventions reduce food cravings. *Appetite*.

Brief guided imagery and body scanning
interventions reduce food cravings

Jonathan Hamilton¹, Sophie Fawson¹,
Jon May¹, Jackie Andrade¹ &
David Kavanagh²

¹ School of Psychology, Cognition Institute, Plymouth University, UK

² Institute of Health & Biomedical Innovation and School of Psychology & Counselling,
Queensland University of Technology, Australia

Corresponding author: Jon May, School of Psychology, Plymouth University Drake Circus
Plymouth PL4 8AA. jon.may@plymouth.ac.uk, +44 1752 584839.

Keywords: mindfulness, , cognition, food, intrusive thoughts, acceptance, eating

Conflict of Interest: None

Running head: Brief guided imagery interventions

Word count: 3004; 1 Table; 2 Figures

1 Abstract

2 Elaborated Intrusion (EI) Theory proposes that cravings occur when involuntary thoughts
3 about food are elaborated; a key part of elaboration is affectively-charged imagery. Craving
4 can be weakened by working memory tasks that block imagery. EI Theory predicts that
5 cravings should also be reduced by preventing involuntary thoughts being elaborated in the
6 first place. Research has found that imagery techniques such as body scanning and guided
7 imagery can reduce the occurrence of food thoughts. This study tested the prediction that
8 these techniques also reduce craving. We asked participants to abstain from food
9 overnight, and then to carry out 10 minutes of body scanning, guided imagery, or a control
10 mind wandering task. They rated their craving at ten points during the task on a single item
11 measure, and before and after the task using the Craving Experience Questionnaire. While
12 craving rose during the task for the mind wandering group, neither the guided imagery nor
13 body scanning group showed an increase. These effects were not detected by the CEQ,
14 suggesting that they are only present during the competing task. As they require no devices
15 or materials and are unobtrusive, brief guided imagery strategies might form useful
16 components of weight loss programmes that attempt to address cravings.

17 [203 words]

18

20 Desires for food are common, everyday experiences, and are highly adaptive in
21 signalling the need to eat sufficient food to fuel activity and maintain healthy physiological
22 systems. However, when desires become intense cravings and cannot immediately be
23 satisfied, they are aversive, and when they trigger significant distress or excessive eating or
24 impair our ability to perform other important cognitive tasks, they may become highly
25 dysfunctional.

26 The Elaborated Intrusion Theory of Desire (EI Theory; Kavanagh, Andrade & May,
27 2005) explicates processes underpinning the onset, exacerbation and termination of desires
28 for appetitive targets including food. Initially, desire-related cognitions typically appear
29 spontaneous, because they are triggered by associated environmental or internal cues
30 rather than being deliberately elicited (Kavanagh et al., 2005). If other later cognitions
31 capture attention, intrusive desire-related thoughts may not be processed further
32 (Kavanagh, May & Andrade, 2009). However, if the pleasure or relief from these cognitions
33 gives them greater salience than other concurrent experiences, they are consciously
34 elaborated, further increasing their affective power. Central to the theory is that sensory
35 imagery is especially likely, and because it simulates the actual experience, it conveys a
36 stronger affective pull than other cognition and is therefore more likely and more vivid
37 when craving is intense (Kavanagh, Statham, Feeney, et al., 2012; May, Andrade, Kavanagh
38 & Hetherington, 2012; Andrade, May & Kavanagh, 2012; Statham, Connor, Kavanagh, et al.,
39 2011).

40 According to EI Theory, the elaboration of craving imagery places demands on
41 limited-capacity working memory systems, and in particular, on the visuospatial sketchpad,
42 which is required for construction of visual images (Baddeley & Andrade, 2000). Concurrent

tasks that require attention to other visuospatial information reduce the vividness of imagery and blunt the intensity of associated craving, especially if they impose a working memory load. So, Andrade, Pears, May and Kavanagh (2012) found that constructing shapes from clay that was hidden behind a screen reduced craving for chocolate: This task requires repeatedly comparing visual imagery about a desired product with imagery constructed from tactile information, as well as co-ordination of hand movements based on this imagery. Instructions to create competing, emotionally neutral imagery also reduce cravings, including ones for coffee (Kemps & Tiggemann, 2009), cigarettes (May, Andrade, Panabokke, & Kavanagh, 2010; Versland & Rosenberg, 2007), chocolate (Kemps & Tiggemann, 2007) and other food (Harvey, Kemps, & Tiggemann, 2005; Kemps & Tiggemann, 2007). In contrast, since food-related craving rarely requires auditory imagery, tasks that load on the phonological loop have only weak effects at best (Kemps & Tiggemann, 2007; May, Andrade, Panabokke & Kavanagh, 2010), unless they impose heavy general cognitive (or 'central executive') load (Andrade et al., 2012).

Another approach to the management of cravings uses mindfulness (Alberts, Mulkens, Sweets & Thewissen, 2010; Alberts, Thewissen & Raes, 2012), which involves enhancement of non-judgemental awareness of the present moment or experience (Brown & Ryan, 2003). The absence of a necessity either to control or elaborate thoughts, and an attention to ever-changing experiences, join to liberate the person from entanglement in rumination (Kabat-Zinn, 2003).

Consistent with this idea, in Berry, May, Andrade & Kavanagh (2010) people who were naturally mindful had less distress in response to intrusive desire-related thoughts and had lower craving than less mindful people. Similarly, Alberts et al. (2012) found that people who undertook 8 weeks of mindfulness-based cognitive-behaviour therapy had

greater decreases in food cravings and in their eating in response to sensory cues or negative emotional states than did others on a wait list. Jenkins and Tapper (2013) found that people who were trying to reduce their chocolate consumption ate less chocolate when following a mindfulness-based cognitive defusion strategy (seeing their thoughts as merely thoughts) than a control (muscle relaxation) group, or a group who accepted chocolate thoughts (by 'urge surfing', using the knowledge that urges are transient, to help with resisting desires to indulge). Papies, Barsalou and Custers (2012) also found that brief training in 'mindful attention', viewing thoughts as transient mental events, reduced approach responses to appetitive foods.

Body scanning is a technique commonly used in mindfulness training, where the focus of attention is consciously directed around the body, while concurrently maintaining an awareness of breathing (Cropley, Ussher & Charitou, 2007). As in other mindfulness exercises, the person is encouraged to attend to changing experiences without questioning, suppressing or being judgemental of ones that come to mind (Kristeller & Hallett, 1999). If other thoughts and emotions arise, they are noted briefly before returning attention to the body.

While mindfulness exercises including body scanning are derived from ancient Buddhist traditions, their expected effects on craving are highly consistent with EI Theory. For example, the attentional demands of body scanning would reduce cognitive capacity for the elaboration of craving imagery, while a focus on new, emerging experiences would introduce craving-irrelevant content. Accepting desire-related thoughts rather than trying to suppress them would avoid the risk that monitoring the suppression would ironically provide a trigger for their occurrence (Wegner & Erber, 1992), while adopting the stance of

a curious bystander would blunt the affective intensity of the thoughts (and their ability to maintain a cyclical elaboration of craving cognitions).

There are indications that body scanning may indeed help people to address food cravings. A small randomised controlled trial by Alberts et al. (2010) with attendees at a weight reduction program reported lower food cravings after a 7-week mindfulness-based intervention where body scanning was an important component. In a more controlled setting, May, Andrade, Batey, Berry & Kavanagh (2010, Experiment 2) examined the effect of 10 mins of Body Scanning instructions on thoughts and craving for snack foods by students who were attempting to reduce consumption of snack foods and had not eaten for 2 hours. Effects were compared with those from Guided Imagery and Control (mind wandering) instructions. During the experimental period, Body Scanning and Guided Imagery reduced concurrent thoughts about snack foods, relative to 10-min Baseline and Post-task periods, but the control instructions did not. However, there were no Condition by Time effects on single-item ratings of craving for snack foods that were taken at the end of each period. In contrast, May, Andrade, Willoughby and Brown (2012) found that both craving and thought frequency about smoking were reduced in the experimental period by body scanning, relative to mind wandering. That study applied a 2-session within-subjects design with smokers who were asked to abstain from smoking for 2 hours, and measured craving by Factor 1 of the Questionnaire on Smoking Urges (Tiffany & Drobes, 1991), but otherwise had a similar procedure to May, Andrade, Batey et al. (2010).

The current study attempted to address two potential reasons for the lack of effects on craving for snack foods in May, Andrade, Batey et al. (2010). Firstly, it increased the period of food deprivation from 2 hours to at least 9 hours (in recognition of the need to induce a stronger sense of deprivation, such as we saw after abstention from smoking).

Secondly, it sampled craving during the baseline, experimental and post-task periods rather than relying on craving assessments after completion of the instructional task (when competing tasks were no longer present). The study retained the three-group Between-subjects design and instructional procedures used by May, Andrade, Batey et al. (2010), allowing it to examine whether body scanning had similar effects to those from a task that should interfere with imagery-based aspects of craving (Guided Imagery), and whether Body Scanning and Guided Imagery produced differential effects from a Mind Wandering control.

Method

Participants

Participants were recruited from Plymouth University's Psychology student pool and received participation credit for their participation, which they could use in their own research. The experiment was conducted in accordance with the ethical guidelines of the British Psychological Society and had approval from the University's Faculty of Science and Technology Ethics Committee. Ninety-eight participants (75 female, 23 male, M age = 20, range = 18-45) took part.

Assessment Instruments

A single-item craving intensity measure (0, no craving, to 10, intense craving) was used for repeated measurements during the experimental phase of the study. Before and after the imagery task, the *Craving Experience Questionnaire* (CEQ; May, Andrade, Kavanagh et al., submitted) was used to measure the strength of food craving (CEQ-S), and the frequency of craving thoughts (CEQ-F). The CEQ-S and CEQ-F each have 10 items, over three subscales measuring craving intensity, use of imagery and intrusiveness. Using data from twelve studies on chocolate, other food, alcohol and cigarettes, May et al. (submitted)

found that the internal structure of the CEQ was robust across substances and timeframes over which desires were assessed, and that internal consistencies of the total CEQ-S and CEQ-F were high. CEQ-S items focused on current craving. At Baseline, participants completed the CEQ-F on the frequency of cravings since they last ate: at the end of the session, they rated the frequency of cravings during the session.

The *Eating Attitudes Test* Factor 1 (EAT26; Garner, Olmsted, Bohr & Garfinkel, 1982) was used to identify whether any participants screened positive for an eating disorder, while a *Brief Mindfulness Measure* (BMM; Berry et al. 2010) and an *Emotional and Behavioural Reactions to Intrusions Questionnaire* (EBRIQ; Berry, et al. 2010) checked whether levels of trait mindfulness and usual reactions to intrusive thoughts were equivalent across conditions.

Procedure

Sessions were held between 9am and noon. Participants were asked to abstain from eating since midnight the night before, so that they were food-deprived for at least 9 hours. They were asked to bring a breakfast item to the session. After providing informed consent, they placed their breakfast item alongside a selection of other breakfast bars on the table, in order to augment their craving at Baseline. If they did not bring an item, they were offered a bar to take at the end of the study. They then completed the CEQ, EAT26, BMM and EBRIQ, and were asked a series of questions about when they last ate, their usual breakfast habits and favourite foods, in order to reinvoke craving. The breakfast bars were then covered, to reduce distractions during the session's experimental phase.

Random allocation to conditions controlled for Baseline CEQ-S scores by using separate random sequences for those above and below the running average. All conditions involved a 10-min audio recording being played through headphones, while a craving scale

was displayed on a computer screen. An instructional statement was heard every 20 seconds: This instructional frequency kept them on task, while allowing them time to undertake the task between statements. A bell sounded ten times within each recording (at intervals of between 38 to 80 seconds, using the same pseudo-random sequence across conditions), at which points participants reported their craving on the 0-10 scale.

The audiotapes were the same as those used by May, Andrade, Batey et al. (2010). In *Mind Wandering*, participants were instructed to think about ‘anything or nothing at all’ and let their ‘mind wander wherever it will go’. In *Body Scan*, they focused on specific parts of the body, starting with their toes and moving to the top of their head, relaxing, while noticing and accepting thoughts. Instructions included ‘notice any sensations here right now’ and ‘focus on breathing’. *Guided imagery* involved imagining a forest walk, using multiple senses (e.g. ‘brightly coloured birds call from the wood’; ‘feel the path beneath your feet as you travel through the wood’). The random allocation gave 33 participants in Mind Wandering, 34 in Body Scan and 31 in Guided Imagery.

At the end of the recording, participants again completed the CEQ-S and CEQ-F, were debriefed and then were allowed to see and take their breakfast item.

Results

Four participants did not comply with the task requirements (two Mind Wandering and two Guided Imagery), giving minimum or maximum ratings to all items on one or more scales, and their data were removed from the analyses. One way ANOVA on the remaining 94 participants showed that the three groups (see Table 1) did not differ on Mindfulness (BMM: $F(2,91) = 1.70$, $p = .19$, $\eta^2 = .04$), Reactivity (EBRIQ), eating attitudes (EAT) or baseline craving (CEQ) totals (all $F_s < 1$).

184

Table 1 About Here

185

186

187

188

189

190

191

192

193

A repeated measures ANOVA on the ten single-item craving ratings obtained during the imagery showed no main effect of time ($F(9,819) = 1.05, \eta^2 = .011, p = .40$) or of condition ($F(1, 91) = 2.11, \eta^2 = .044, p = .13$), but a significant interaction ($F(18, 819) = 1.90, \eta^2 = .04, p = .013$), as shown in Figure 1. Follow-up ANOVAs on each group showed no effects of time for the bodyscan or guided imagery groups ($F_s < 1$), but a significant effect of time for the control group ($F(9,270) = 3.31, \eta^2 = .10, p = .001$). Contrasts showed significant Linear ($F(1,30) = 4.59, \eta^2 = .13, p = .04$) and Quadratic ($F(1,30) = 4.51, \eta^2 = .13, p = .04$) effects, indicating a rise in craving over the first half of the mind wandering task, which then levelled off.

194

Figure 1 About Here

195

196

197

198

199

200

Repeated measures ANOVA was also conducted on the before and after task totals from the CEQ. There were no statistically significant effects for the scale totals or any of the subscales (Figure 2), although for the CEQ-S Imagery subscale the effect of time approached significance ($F(1,91) = 3.43, p = .067, \eta^2 = .04$), as did the interaction of time x condition ($F(2,91) = 2.39, p = .098, \eta^2 = .05$), as the mind wandering group's scores rose while the other two groups' fell.

201

Figure 2 About Here

202

Discussion

203

204

The single item craving measures obtained during the experimental task rose for participants who allowed their mind to wander, but remained constant for those in the

guided imagery and body scanning conditions. We have therefore extended the findings of May, Andrade, Batey et al. (2010) to show that these attentional control tasks can prevent the occurrence of cravings for food as well as reducing the frequency of intrusive thoughts, as it did for cravings and thoughts about smoking in May et al. (2012). As in May, Andrade, Batey et al. (2010), the before and after measures of craving did not show an effect, suggesting that craving is only suppressed while the imagery interventions are being conducted.

In practical terms, this research suggests that the body scanning and guided imagery tasks could be helpful for people who are trying to resist the cravings that occur during abstinence or reduction attempts, and which lead to relapse (Sitton, 1991; Massey & Hill, 2012). While body scanning is a core mindfulness technique because of its links to Buddhist meditation practice and the idea that thoughts are, like physical sensations, transitory, the guided imagery script used here had identical effects. In fact, many mindfulness tapes available commercially also make use of a 'forest walk' script, directing the listener's imagery to different sensory modalities. Guided imagery shares many facets with mindfulness exercises, such as relaxation and directed mental activity, without explicitly including key aspects such as thought acceptance and non-judgementalism. In our work we use it as a form of comparison task to assess the contribution that body scanning has beyond relaxation and the control of mental imagery, and in this study, as in May, Andrade, Batey et al. (2010) and May et al. (2012), find that the two are identical in effectiveness.

This indicates that suppressing food related imagery during a potential craving episode is the most important factor in preventing a rise in craving intensity, in line with the predictions of Elaborated Intrusion Theory. An alternative explanation, of course, is that the mind wandering condition, intended as a control condition, is exacerbating craving rather

than just allowing it to occur. Allowing the mind to wander allows any intrusive thought about food to be elaborated, but it does not mean that it was elaborated using visual imagery: in this experimental situation, having missed breakfast, a food related thought might just be the most interesting thing to think about. Although a body of other research has also shown that visual imagery is a central component of craving for a range of substances (e.g., Salkovskis & Reynolds, 1994; May, Panabokke, Andrade & Kavanagh, 2004; Harvey et al., 2005; Statham et al., 2011), and that visual imagery tasks can also weaken craving and prevent craving developing (May, Andrade, Panabokke & Kavanagh, 2010; Versland & Rosenberg 2007), further research could compare unrestricted mind wandering with a mind wandering condition in which visual imagery is restricted, perhaps through a visual monitoring task, or through exposure to a dynamic visual noise display (Quinn & McConnell, 1996), which has been shown to interfere with craving (May et al, 2010; Kemps, Tiggemann, Woods & Soekov, 2004).

In our research we have used recorded prompts to instruct our participants in the body scanning and guided imagery, but people can learn to do both tasks without such prompts. As an entirely cognitive strategy, requiring no devices or materials, taking a few seconds to imagine oneself in a sensorially rich environment is an easy and practical self-help technique that could form part of dietary restraint and weight-loss programmes. It is unobtrusive and can be employed repeatedly, whenever needed, until the temptation has passed. Knäuper, Pillay, Lacaille, McCollam & Kelso (2011) asked people to deal with cravings over a four-day period by vividly imagining engaging in a favourite activity and found that it reduced craving intensity, whereas three control conditions had no effect.

It is likely that the positive affective nature of the scenes imagined in our work and in that of Knäuper et al. (2011) would motivate people to continue to use the strategy, but it

must also be recognised that the effects upon craving might be due more to the mood and relaxation effects than to the demands of imagery and attentional control, and further work should contrast these conditions with positive mood enhancement that does not include these cognitive aspects. Evaluation could also usefully focus upon the need for novelty in the imagined scenes, to assess the contribution of higher-order cognitive processes in the generation of imagery, as opposed to the retrieval of familiar scenes from memory.

References

- Alberts, H.J.E.M., Mulkens, S., Smeets, M. & Thewissen, R. (2010) Coping with food cravings. Investigating the potential of a mindfulness-based intervention. *Appetite*, 55, 160-163. doi:10.1016/j.appet.2010.05.044
- Alberts, H.J.E.M., Thewissen, R. & Raes, L. (2012) Dealing with problematic eating behaviour. The effects of a mindfulness-based intervention on eating behaviour, food cravings, dichotomous thinking and body image concern. *Appetite*, 58, 847-851. doi: 10.1016/j.appet.2012.01.009
- Andrade, J., May, J., & Kavanagh, D.J. (2012) Sensory imagery in craving: From cognitive psychology to new treatments for addiction. *Journal of Experimental Psychopathology*, 3(2) 127-145. doi: 10.5127/jep.024611
- Andrade, J., Pears, S., May, J., & Kavanagh, D.J. (2012) Use of a clay modeling task to reduce chocolate craving. *Appetite*, 58, 955-963. doi: 10.1016/j.appet.2012.02.044

272 Baddeley, A.D. & Andrade, J. (2000) Working memory and the vividness of imagery. *Journal*
 273 *of Experimental Psychology: General*, 129(1), 126-145. Doi:10.1037/0096-
 274 3445.129.1.126

275 Berry, L.-M., May, J. Andrade, J. & Kavanagh, D. (2010) Emotional and behavioural reaction
 276 to intrusive thoughts. *Assessment*, 17(1), 126-137. doi:10.1177/1073191109344694

277 Brown, K.W. & Ryan, R.M. (2003). The Benefits of Being Present: Mindfulness and Its Role in
 278 Psychological Well-Being. *Journal of Personality and Social Psychology*, 84(4), 822–848.
 279 doi:10.1037/0022-3514.84.4.822

280 Cropley, M., Ussher, M., & Charitou, E. (2007). Acute effects of a guided relaxation routine
 281 (body scan) on tobacco withdrawal symptoms and cravings in abstinent smokers.
 282 *Addiction*, 102, 989–993. doi:10.1111/j.1360-0443.2007.01832.x

283 Garner, D. M, Olmsted, M. P., Bohr, Y., & Garfinkel, P. E. (1982) The Eating Attitudes Test:
 284 psychometric features and clinical correlates. *Psychological Medicine* 12, 871-8.

285 Harvey, K., Kemps, E. & Tiggemann, M. (2005) The nature of imagery processes underlying
 286 food cravings. *British Journal of Health Psychology*, 10, 49-56. doi:
 287 10.1348/135910704X14249

288 Jenkins, K.T. & Tapper, K. (2013) Resisting chocolate temptation using a brief mindfulness
 289 strategy. *British Journal of Health Psychology*, 18. retrieved 17/5/2013.

290 Kabat-Zinn J. (2003). Mindfulness-based interventions in context: Past, present, and future.
 291 *Clinical Psychology: Science and Practice*, 10, 144-156. doi: 10.1093/clipsy/bpg016

292 Kavanagh, D.J., Andrade, J., & May, J. (2005) Imaginary relish and exquisite torture: The
 293 elaborated intrusion theory of desire. *Psychological Review*, 112, 446-467.
 294 doi:10.1037/0033-295X.112.2.446

295 Kavanagh, D.J., May, J., & Andrade, J. (2009). Tests of the Elaborated Intrusion Theory of
 296 craving and desire: Features of alcohol craving during treatment for an alcohol
 297 disorder. *British Journal of Clinical Psychology*, 48, 241–254.
 298 doi:10.1348/014466508X387071

299 Kavanagh, D.J., Statham, D.J., Feeney, G.F.X., Young, R. McD., May, J., Andrade, J. & Connor,
 300 J. (2012) Measurement of alcohol craving. *Addictive Behaviors*, 38, 1572-1584 . Doi:
 301 10.1016/j.addbeh.2012.08.004

302 Kemps, E. & Tiggemann, M., (2007). Modality-specific imagery reduces cravings for food: An
 303 application of the Elaborated Intrusion theory of desire to food craving. *Journal of*
 304 *Experimental Psychology: Applied*, 13(2), 95-104. doi:10.1037/1076-898X.13.2.95

305 Kemps, E. & Tiggemann, M., (2009) Competing Visual and Olfactory Imagery Tasks Suppress
 306 Craving for Coffee. *Experimental and Clinical Psychopharmacology*, 17(1), 43–50. doi:
 307 10.1037/a0014962

308 Kemps, E., Tiggemann M., Woods, D., & Soekov, B. (2004). Reduction of food cravings
 309 through concurrent visuo-spatial processing. *International Journal of Eating Disorders*,
 310 36, 31-40. doi: 10.1002/eat.20005

311 Knäuper, B., Pillay, R., Lacaille, J., McCollam, A. & Kelso, E. (2011) Replacing craving imagery
312 with alternative pleasant imagery reduces craving intensity. *Appetite*, 57(1) 173-178.
313 doi: 10.1016/j.appet.2011.04.021

314 Kristeller, J. L., & Hallett, C. B. (1999). An exploratory study of a meditation-based
315 intervention for binge eating disorder. *Journal of Health Psychology*, 4, 357–363. doi:
316 10.1177/135910539900400305

317 Massey, A., & Hill, A.J. (2012) Dieting and food craving. A descriptive, quasi-prospective
318 study. *Appetite*, 58(3), 781-785. doi:10.1016/j.appet.2012.01.020

319 May, J., Andrade, J., Batey, H., Berry, L.-M. & Kavanagh, D. (2010) Less food for thought:
320 Impact of Attentional Instructions on Intrusive Thoughts about Snack Foods. *Appetite*,
321 55(2) 279-287. doi:10.1016/j.appet.2010.06.014

322 May, J., Andrade, J., Kavanagh, D.J., Feeney, G.F.X., Gullo, M., Statham, D.J., ... Connor, J.P.
323 (submitted). The Craving Experience Questionnaire: A brief, theory-based measure of
324 consummatory desire and craving.

325 May, J., Andrade, J., Kavanagh, D.J. & Hetherington, M. (2012) Elaborated Intrusion theory:
326 A cognitive-emotional theory of food craving. *Current Obesity Reports*, 1(2), 114-121.
327 Doi: 10.1007/s13679-012-0010-22

328 May, J., Andrade, J., Panabokke, N., & Kavanagh, D. (2010). Visual imagery tasks suppress
329 craving for cigarettes. *Behaviour Research and Therapy*, 48, 476-485.
330 doi:10.1016/j.brat.2010.02.001

331 May, J., Andrade, J., Willoughby, K. & Brown, C. (2012) An attentional control task reduces
 332 intrusive thoughts about smoking. *Nicotine & Tobacco Research*, 14(4), 472-478. doi:
 333 10.1093/ntr/ntr238

334 May, J., Panabokke, N., Andrade, J., & Kavanagh, D. (2004). Images of desire: cognitive
 335 models of craving. *Memory*, 12, 447–461. doi: 10.1080/09658210444000061

336 Papies, E.K., Barsalou, L.W. & Custers, R. (2012). Mindful Attention Prevents Mindless
 337 Impulses. *Social Psychological and Personality Science*, 3(3), 291-299. doi:
 338 10.1177/1948550611419031

339 Quinn, J. G. & McConnell, J. (1996a). Irrelevant pictures in visual working memory. *Quarterly*
 340 *Journal of Experimental Psychology*, 49A, 200-215. doi: 10.1080/713755613

341 Salkovskis, P. M., & Reynolds, M. (1994). Thought suppression and smoking cessation.
 342 *Behaviour Research and Therapy*, 32(2): 193-291. doi: 10.1016/0005-7967(94)90112-0

343 Sitton, S. C. (1991). Role of craving for carbohydrates upon completion of a protein-sparing
 344 fast. *Psychological Reports*, 69, 683–686. doi: 10.2466/pr0.1991.69.2.683

345 Statham, D. J., Connor, J. P., Kavanagh, D. J., Feeney, G. F. X., Young, R. McD., May, J., &
 346 Andrade, J. (2011). Measuring Alcohol Craving: Development of the Alcohol Craving
 347 Questionnaire. *Addiction* 106(7), 1230-1238. doi: 10.1111/j.1360-0443.2011.03442.x

348 Tiffany, S. T., & Drobes, D. J. (1991). The development and initial validation of a
 349 questionnaire on smoking urges. *British Journal of Addiction*, 86, 1467–1476.
 350 doi:10.1111/j.1360-0443. 1991.tb01732.x

351 Versland, M. & Rosenberg, H. (2007) Effects of brief imagery interventions on craving in
352 college student smokers. *Addiction Research and Theory*, 15(2), 177-187. doi:
353 10.1080/16066350701200582

354 Wegner, D. & Erber, R. (1994) The hyperaccessibility of suppressed thoughts. *Journal of*
355 *Personality and Social Psychology*, 63(6), 903-912. doi: 10.1037/0022-3514.63.6.903.

Table 1: Means (standard deviations) for the Eating Attitudes Test (EAT), Brief Measure of Mindfulness (BMM) ,Emotional & Behavioural Reaction to Intrusions Questionnaire (EBRIQ), and baseline Craving Experience Questionnaire Strength (CEQ-S) and Frequency (CEQ-F).

Imagery Condition	N	EAT	BMM	EBRIQ	CEQ-S	CEQ-F
Control	31	7.8 (7.1)	3.1 (0.5)	2.9 (0.7)	4.4 (1.9)	4.2 (1.6)
Body Scan	34	8.5 (5.6)	2.9 (0.5)	2.7 (0.6)	4.8 (2.2)	4.2 (2.1)
Guided Imagery	29	9.0 (7.0)	3.1 (0.5)	2.7 (0.6)	4.8 (1.9)	4.3 (1.9)

Figure 1. During the imagery tasks, craving did not change for the Body Scanning (dotted line) and Guided Imagery (dashed line) groups, but rose for the Mind Wandering control group (Solid line).

Figure 2. Although CEQ-Strength Imagery and Intensity scores declined for the Body Scan (dotted line) and Guided Imagery (dashed line) groups, compared to the Mind Wandering control group (Solid line), no interactions were statistically significant. CEQ-Frequency scores were unchanged for all three groups.