2020-12-15

Dynamic partitioning of tropical Indian Ocean surface waters using ocean colour data management and modelling applications

Schwarz, Jill

<http://hdl.handle.net/10026.1/16413>

10.1016/j.jenvman.2020.111308

Journal of Environmental Management

Elsevier BV

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Dynamic partitioning of tropical Indian Ocean surface waters using ocean colour data - management and modelling applications

Jill N. Schwarz

*School of Biological & Marine Sciences, University of Plymouth, Drake Circus,
Plymouth, PL4 8AA, UK*

Abstract

Over the past few decades, partitioning of the surface ocean into ecologically-meaningful spatial domains has been approached using a range of data types, with the aim of improving our understanding of open ocean processes, supporting marine management decisions and constraining coupled ocean-biogeochemical models. The simplest partitioning method, which could provide low-latency information for managers at low cost, remains a purely optical classification based on ocean colour remote sensing. The question is whether such a simple approach has value. Here, the efficacy of optical classifications in constraining physical variables that modulate the epipelagic environment is tested for the tropical Indian Ocean, with a focus on the Chagos marine protected area (MPA). Using remote sensing data, it was found that optical classes corresponded to distinctive ranges of wind speed, wind stress curl, sea surface temperature, sea surface slope, sea surface height anomaly and geostrophic currents (Kruskal-Wallis and post-hoc Tukey honestly significantly different

Email address: `jill.schwarz@plymouth.ac.uk` (Jill N. Schwarz)

tests, $\alpha = 0.01$). Between-class differences were significant for a set of sub-domains that resolved zonal and meridional gradients across the MPA and Seychelles-Chagos Thermocline Ridge, whereas between-domain differences were only significant for the north-south gradient (PERMANOVA, $\alpha = 0.01$). A preliminary test of between-class differences in surface CO₂ concentrations from the Orbiting Carbon Observatory-2 demonstrated a small decrease in mean pCO₂ with increasing chl, from 418 to 398 ppm. Simple optical class maps therefore provide an overview of growth conditions, the spatial distribution of resources – from which habitat fragmentation metrics can be calculated, and carbon sequestration potential. Within the 18 year study period, biotic variables were found to have decreased at up to 0.025%a⁻¹ for all optical classes, which is slower than reported elsewhere (Mann-Kendall-Sen regression, $\alpha = 0.01$). Within the MPA, positive Indian Ocean Dipole conditions and negative Southern Oscillation Indices were weakly associated with decreasing chl, fluorescence line height (FLH), eddy kinetic energy, easterly wind stress and wind stress curl, and with increasing FLH/chl, sea surface temperature, SSH gradients and northerly wind stress, consistent with reduced surface mixing and increased stratification. The optical partitioning scheme described here can be applied in Google Earth Engine to support management decisions at daily or monthly scales, and potential applications are discussed.

Keywords: Remote sensing, biogeography, habitat fragmentation, Orbiting Carbon Observatory-2 (OCO-2), Marine Protected Area, epipelagic

1. Introduction

Physical drivers such as light, temperature and dynamical mixing shape the epipelagic ecosystem, and the biota of deeper water layers is determined, at least in part, by the productivity of upper layers (Jerlov, 1976; Longhurst, 2007; Spalding et al., 2012; Kavanaugh et al., 2016; Proud et al., 2017). Dividing the oceans into geographical areas with common physical conditions has been approached using a range of methods and suites of data (reviewed by Krug et al., 2017; Kavanaugh et al., 2016), and referred to variously as ecological geography, partitioning, biogeography, biohydrography, biogeographical provinces and seascapes. Most schemes include nested spatial scales with slightly different nomenclature for different elements of hierarchical structure. Partitioning the oceans is similar to habitat mapping and species distribution modelling in the sense that a geographical representation of resources is produced. However, it does not relate to specific organisms, and no model of the relationships between predictor and response variables is produced (c.f. Blanco et al., 2015; Coelho et al., 2013; Scales et al., 2014; Zydulis et al., 2011). Partitioning schemes have found application to two key challenges. Firstly, they provide static and dynamical geographical boundaries to guide management planning (over the long term) and intervention (in the short term). Management tasks include monitoring ecosystem health, assessing risk and implementing control measures such as fishery closure, and information is required at multiple spatial scales and depth ranges to support these actions (Rice et al., 2011; Spalding et al., 2012; Caldow et al., 2015; Roberson et al., 2017). Short-term responses can only be supported when the frequency at which new data is available exceeds the rate of critical fluctuations occur-

ring within the ecosystem. The second use is to provide spatial context for the evaluation of climate model reliability (Vichi et al., 2011; Kavanaugh et al., 2014; Fay & McKinley, 2014). The focus here is on the exchange of climate-relevant gases across the air-sea interface and there is no requirement for low-latency information. Partitions serve as a proxy for ecosystem function with the inference that they constrain rates of CO₂ diffusion, biotic carbon uptake and the efficiency with which carbon is removed from the surface layer. Earth system models (ESM), in which the representation of biogeochemical cycles remains quite simplistic (Hense et al., 2017; Jung et al., 2019; Sreeush et al., 2018) and which are sensitive to feedback between biotic and abiotic components (Lim & Kug, 2017; Park & Kug, 2014; Romanou et al., 2014) can then be evaluated in the context of static or dynamic seascapes (Landschützer et al., 2019; Lovenduski et al., 2019).

Satellite remote sensing data, which currently provides the best compromise between area coverage and temporal resolution for surface ocean studies, has increasingly been used for partitioning, as the disparate worlds of marine management and oceanography converge on how this rich data source can best be used (Kachelriess et al., 2014; Maxwell et al., 2015; Miloslavich et al., 2018). Figure 1 illustrates some of the pelagic abiotic and biotic factors that can be derived from remote sensing data, and characteristics relevant for modelling and management decisions.

[FIGURE 1 HERE: SINGLE-COLUMN; COLOUR ONLY ONLINE]

Of the variables amenable to remote sensing, only ocean colour is directly affected by the pelagic ecosystem at short time-scales. Partitioning schemes mostly use an ocean-colour-derived variable, chlorophyll-a concen-

51 tration (hereafter chl), as a measure of biomass and combine it with other
52 key physical drivers such as temperature. However, being a function of the
53 concentrations of dissolved and particulate substances in the surface ocean,
54 ocean colour itself represents a response to the physical drivers. Using ocean
55 colour to derive chl incurs spatially-varying errors (Jackson et al., 2017) and
56 ignores other coloured variables of interest that may, or may not, covary with
57 chl (O'Reilly et al., 1998; Werdell et al., 2018).

58 Classifications based on water colour alone have shown that, whilst chl is
59 generally constrained by these classifications, information about other water
60 mass tracers is also present (Alvain et al., 2005, 2008; Vantrepotte et al., 2012;
61 Trochta et al., 2015; Krug et al., 2018; Monolisha et al., 2018; Dierssen, 2010).
62 Taken together with the knowledge that physical dynamics at all scales com-
63 bine to control the growth environment, this suggests an opportunity to use
64 pure optical classes as the smallest scale in a seascape heirarchy, with the
65 advantage that it is a low-latency product which could feed into decision-
66 making flows on a daily basis (e.g. using GoogleEarthEngine; Gorelick et al.,
67 2017) where coverage allows. This possibility is explored here, with a focus
68 on the Chagos marine protected area (MPA). If optical classes are found to
69 constrain abiotic drivers as well as biotic response, then a further question
70 arises of whether they can also be used to estimate carbon flows (extending
71 Kavanaugh et al., 2014) without the need to identify individual elements,
72 such as phytoplankton function type, as an intermediate step.

73 *1.1. The study area*

74 The Chagos marine protected area occupies 640,000 km² in the tropical
75 Indian Ocean, with the Chagos Archipelago system of islands and atolls at it

76 centre (Figure 2; UNEP-WCMC, 2016). The coastline of the Indian Ocean
77 is shared amongst over twenty countries, representing a substantial fraction
78 of the human population with variable socio-economic status and strong re-
79 liance on coastal and open-ocean fisheries (Hermes et al., 2019). The MPA
80 is of particular value because of its coral health, resilience and diversity,
81 extensive seagrass beds, potential support for the wider Indian Ocean fish-
82 eries and related benefits (Koldewey et al., 2010; Ateweberhan et al., 2018;
83 Gravestock & Sheppard, 2015; Esteban et al., 2018). In common with shallow
84 tropical corals around the world, reefs in the Chagos MPA are vulnerable to
85 temperature increases associated with climate change as well as to increases
86 in extreme high energy dynamics. Their relative resilience compared with
87 other reef systems is associated with protection from human disturbance as
88 well as to geographical location. Although extensive coral bleaching has oc-
89 curred (Sheppard et al., 2008), interactions between dynamical processes at
90 a range of scales and topographic diversity may alleviate temperature stress
91 (Sheppard, 2012; Hosegood et al., 2019). Understanding whether this nat-
92 ural protection will continue under ongoing climate change is important in
93 terms of economic as well as natural resource value.

94 The Chagos Archipelago lies at the edge of the South Indian tropical
95 gyre within the influence of the Indian Ocean monsoon. In austral summer,
96 northerly winds drive an anticyclonic cell and the gyre contracts, so that
97 the archipelago is at the northern edge. For the rest of the year, the MPA
98 lies at or near the southern edge of the gyre. The location and large-scale
99 circulation features are summarised in Figure 2. In the west, divergent winds
100 and negative wind stress curl lift the thermocline along 5-12°S creating the

101 Seychelles-Chagos Thermocline Ridge (SCTR), with a thermocline depth
102 around of 50 m and no surface signature, which extends around the MPA
103 throughout the year (Hermes & Reason, 2008, 2009; Aguiar-Gonzalez et al.,
104 2016; Xie et al., 2002).

105 [FIGURE 2 HERE: SINGLE or 1.5 COLUMN; COLOUR ONLY ON-
106 LINE]

107 The South Equatorial Current (SEC), flowing to the south of the MPA,
108 denotes the boundary between relatively nutrient replete but O₂-poor surface
109 waters to the north and southern sub-tropical gyre waters to the south which
110 are nutrient-depleted throughout the water column but represent a CO₂ sink
111 (Garcia et al., 2018; Landschützer et al., 2016). *In situ* biogeochemical data
112 are sparse across the tropical Indian Ocean and considerable deviations from
113 the mean conditions in the World Ocean Atlas have been reported (Subha
114 Anand et al., 2017; Chinni et al., 2019). Whilst the tropics are generally
115 considered to be oligotrophic, year-round elevated phytoplankton biomass
116 is observed close to the archipelago as well as over the Mascarene Plateau
117 to the west and broadly over the SCTR (Wilson & Qiu, 2008; Levy et al.,
118 2007). *In situ* measurements of net primary production in this region range
119 from close to zero up to 20 mgCm⁻²d⁻¹ and can be exceeded by bacterial
120 production (Subha Anand et al., 2017; Fernandes et al., 2008; Veldhuis et al.,
121 1997). A few high temporal resolution datasets from moored fluorometers
122 have shown high frequency, high magnitude fluctuations in phytoplankton
123 biomass (Hosegood et al., 2019; Strutton et al., 2015). Phytoplankton as-
124 semblages have been found to be dominated by *Prochlorococcus* and *Syneco-*
125 *coccus* as expected in the oligotrophic gyres, but substantial fractions of di-

atoms, dinoflagellates and prymnesiophytes have also been reported in the TIO (Thorrington-Smith, 1971; Veldhuis et al., 1997; Soares et al., 2015). To my knowledge, there are no long-term biogeochemical monitoring efforts in the pelagic SCTR, despite detailed repeat monitoring in the shallow reef waters of the MPA (e.g. Sheppard, 2012). Many studies have used coupled ocean-biogeochemical models, together with available *in situ* or remote sensing data, to elucidate biophysical coupling in the tropical Indian Ocean (TIO) (e.g. Wiggert et al., 2006; Jin et al., 2012; Liu et al., 2013; Resplandy et al., 2009; George et al., 2018). Of particular interest here are the results of George et al. (2018), Dilmahamod et al. (2016) and Wiggert et al. (2006), who explore meridional and zonal gradients in the SCTR.

The epipelagic growth environment is directly modulated by entrainment and advection of nutrients and plankton, fluctuations in mixed layer temperature and depth, the relative euphotic to mixed layer depths, turbulence and varying illumination conditions. Conversely, feedback effects have been demonstrated between chl and shortwave heating, SST, surface convergence and basin-scale dynamical features (Back & Bretherton, 2009; Park & Kug, 2014). At the seasonal scale, the eastward extent of the SCTR and westward extent of Indonesian Throughflow (ITF) respond to monsoon wind weakening and reversal (Aguiar-Gonzalez et al., 2016). Two of the eight Madden-Julian Oscillation (MJO) phases are centred in the TIO (Hendon & Salby, 1994), with a westward-propagating Rossby wave (Seiki et al., 2013) impacting SST, evaporation, precipitation, cloud cover, rainfall, salinity gradients (Guan et al., 2014; Jin et al., 2013; McPhaden & Foltz, 2013) and wind-driven entrainment of nutrients into the mixed layer (Jin et al., 2012b). At the in-

151 terannual scale, El Nino and the Southern Oscillation (ENSO) and Indian
152 Ocean Dipole (IOD) events have been reported to affect surface conditions
153 and surface chlorophyll concentrations. Nino conditions affect mixed layer
154 temperatures through precipitation, downwelling (anticyclonic) winds and
155 westward propagating Rossby waves (Santoso et al., 2010; Dilmahamod et
156 al., 2016; Ma et al., 2014; Racault et al., 2017). Positive IOD phases coincide
157 with cooler surface temperatures in the eastern TIO and warmer conditions
158 to the west (SST anomalies of 0.1 to 0.3 °C or more; Currie et al., 2013;
159 Vialard et al., 2009). The impact of ENSO and IOD events is amplified when
160 they coincide, and both are expected to increase in frequency (IPCC, 2013;
161 Sheppard et al., 2008; Currie et al., 2013; Cai et al., 2014).

162 Despite the scarcity of data in the epipelagic in and around the Cha-
163 gos MPA, the importance of conditions in these waters has recently been
164 highlighted by electronic tagging of seabirds and high trophic level pelagic
165 feeders, which has been used to document foraging at considerable distances
166 (Pecoraro et al., 2017; Danckwerts et al., 2014; Le Corre et al., 2012), with
167 measurable positive impacts on reef health through nutrient redistribution
168 (Graham et al., 2018). The near-shore pelagic biome is also a critical food re-
169 source for corals (Houlbrque & Ferrier-Pags, 2009) and the strength of this
170 relationship has now been demonstrated using remotely sensed ocean colour
171 data (Fox et al., 2019). Elevated chl related to the wind-driven circulation
172 around the Chagos MPA is therefore a potential resource both for the reef
173 and for pelagic organisms.

174 The underlying hypothesis for this study is that water colour represents
175 the evolving trophic status of the upper ocean and is characteristic of the suite

176 of physical drivers of ecosystem function as illustrated in Figure 1 (e.g. Jerlov,
177 1976), making it a candidate as a low-latency, fine-scale level in a seascape
178 heirarchy. This paper addresses how methodological choices affect optical
179 classifications, whether pure optical classes provide useful information about
180 the physical environment and potential applications of optical class maps.

181 2. Methods

182 2.1. Remote sensing data

183 This study spans August 2002 to October 2018. Remote sensing re-
184 flectance (Rrs), chlorophyll-a concentration (chl), normalised fluorescence
185 line height (nFLH) and sea surface temperature (SST) data from the NASA
186 Moderate Resolution Imaging Spectrometer aboard the Aqua satellite (MODIS-
187 Aqua), at Level 2 and Level 3, were acquired from the Ocean Biology Pro-
188 cessing Group (oceancolor.gsfc.nasa.gov). Rrs is the ocean colour product
189 with the least degree of processing and therefore the lowest uncertainty, with
190 errors on the order of 0.001 sr^{-1} but varying with waveband and water type
191 (Franz et al., 2007; IOCCG, 2019). Reflectances from the seven 1 km reso-
192 lution, 10 nm wavebands in the visible domain were augmented by band 1,
193 with 250 m resolution (50 nm waveband) and bands 3 and 4, with 500 m res-
194 olution (20 nm wavebands), and these three bands were spatially averaged to
195 match the 1 km wavebands. Globally, chl is the best-validated ocean colour
196 product, with mean errors of ca. $\pm 33 \text{ mgm}^{-3}$ (Hu, et al., 2012; O'Reilly et
197 al., 1998). Little product validation data is available for nFLH in the tropi-
198 cal Indian Ocean, but the MODIS-Aqua and MODIS-Terra products perform
199 well against *in situ* data in the Southern and Atlantic Oceans (Erickson et

200 al., 2019; Hoge et al., 2003).

201 Daily sea surface height (SSH), height anomalies (SSHA) and geostrophic
202 current velocities (denoted eastwards u and northwards v), from merged al-
203 timeter datasets, were acquired at ca. 30 km resolution from the Coper-
204 nicus Marine Environment Monitoring Service (CMEMS). Reported errors
205 on these products range from < 1 cm to 30 cm, with higher uncertainties
206 under more dynamic conditions (CMEMS, 2020). Eddy kinetic energy was
207 calculated as $EKE = 1/2(u^2 + v^2)$. Sea surface slope was calculated pixel-
208 wise as $\nabla SSH = \partial SSH / \partial x + \partial SSH / \partial y$, with no smoothing. Each product
209 was subsampled to 4 km resolution before applying the optical class masks
210 (section 2.7).

211 Daily surface wind fields from the SeaWinds and ASCAT scatterometer
212 sensors were acquired at 25 km resolution from the Jet Propulsion Labo-
213 ratory Physical Oceanography Distributed Archive Center (SeaPAC, 2006;
214 EUMETSAT/OSI SAF, 2018). Errors in these products are of order 0.1
215 ms^{-1} (Verhoef et al., 2017). Wind stress curl was calculated pixelwise as
216 $\nabla \times \tau = -\partial \tau_N / \partial x - \partial \tau_E / \partial y$, where τ_N and τ_E are the north-south and east-
217 west components of the wind stress and ∂x and ∂y are the pixel dimensions.
218 Monthly averages were calculated before subsampling to 4 km resolution and
219 applying the optical class masks (section 2.7)

220 Profiles of the partial pressure of CO_2 derived from Orbiting Carbon
221 Observatory 2 (OCO-2) data using the full physics model version 7.3 (O'Dell
222 et al., 2018) were acquired at native resolution (ca. 3 km^2) from the Goddard
223 Earth Sciences Data and Information Services Center (OCO-2 Science Team,
224 2016, GES DISC). Only estimates of pCO_2 from the lowest model altitude

225 were used and the ramifications of this are discussed in section 3.4. No spatial
226 or temporal compositing was applied.

227 Bathymetry data were acquired at 1 arc-second resolution from the Na-
228 tional Centers for Environmental Information (ETOPO1, 2019) and regrid-
229 ded using nearest-neighbour gridding to match the Level 3 Rrs data.

230 Data products, sources and abbreviations are summarised in table 1.

231 [TABLE 1 HERE]

232 2.2. *Dynamic partitioning based on ocean colour*

233 Fuzzy classification was applied to the Indian Ocean domain surrounding
234 the BIOT MPA (after Moore et al., 2001; Jackson et al., 2017). This method
235 was chosen for its potential to allow a single pixel to have multiple class
236 memberships, which is likely in natural phytoplankton populations, particu-
237 larly at the relatively coarse spatial scales of remote sensing data (1 to >30
238 km) in waters where mesoscale and submesoscale processes may be at play.
239 The study bounds were 40° to 100° E, -20° to 15° N, spanning the central
240 tropical Indian Ocean with the BIOT MPA roughly central to the domain
241 (Figure 3; UNEP-WCMC, 2016). In the absence of *in situ* data with which
242 to verify class memberships or interpret class composition, only the dominant
243 class assigned to each pixel at any given time was retained (multiple class
244 memberships were removed, to be considered in future work when validation
245 data are available). Biovolumes calculated from microscopy analysis on five
246 stations within the MPA were used as a preliminary test of whether differ-
247 ent classes represented different phytoplankton biomass (Kruskal-Wallis test;
248 Schwarz, 2020).

Variable	Abbreviation	Source
Remote sensing reflectance, Level 2	L2 Rrs	Ocean Biology Processing Group Level 2 data ocean-color.gsfc.nasa.gov
Remote sensing reflectance, Level 3	L3 Rrs	
Surface chlorophyll-a concentration	chl	
Surface normalised fluorescence line height	nflh	
Ratio of fluorescence line height to chlorophyll-a	flh:chl	MODIS-Aqua Level 3 monthly data, Giovanni data portal giovanni.gsfc.nasa.gov
Sea surface temperature	SST	
Metric for within-class spectral variability (chlorophyll-like pigments): ratio of residual reflectances (Rrs - class mean) at 555 nm to 443 nm	R555:443	
Metric for within-class spectral variability (accessory pigments): ratio of residual reflectances (Rrs - class mean) at 555 nm to 488 nm	R555:488	
Sea surface height (absolute dynamic topography)	SSH	AVISO Level 4 reprocessed gridded sea surface heights and derived variables (product suite SEALEVEL GLO_PHY_L4
Sea surface slope	∇ SSH	REP-OBSERVATIONS_008_047),
Sea surface height anomaly	SLA	marine.copernicus.eu, $0.25^\circ \times 0.25^\circ$ regridded to 4 x 4 km
Eastward component of the geostrophic current	u	
Northward component of the geostrophic current	v	
Eddy kinetic energy	EKE	
Eastward component of the surface wind field	τ_E	Quikscat and ASCAT Level 3 gridded wind fields, podaac.jpl.nasa.gov, 25 x 25 km regridded to 4 x 4 km
Northward component of the surface wind field	τ_N	
Wind stress curl	$\nabla \times \tau$	
Water depth	z	ETOPO 1 arc-minute bathymetry regridded to 4 x 4 km (Amante & Eakins, 2009)
Indian Ocean Dipole index	IOD	Dipole mode index (Saji & Yamagata, 2003) esrl.noaa.gov/psd/gcos_wgsp/Timeseries/DMI
Madden Julian Oscillation index	MJO	Kilidas et al. (2014) esrl.noaa.gov/pas/mjo/mjoindex
Southern Oscillation Index	SOI	Ropelewski & Jones (1987) esrl.noaa.gov/psd/data/20thC_Rean/time-series/monthly/SOI
Surface partial pressure of CO ₂	pCO ₂	Level 2 OCO-2 physical model surface pCO ₂ , release 9, GES DISC (Boesch et al., 2019)

Table 1: Optical, biological and physical remote sensing products used to explore optical classes

249 [FIGURE 3 HERE: SINGLE-COLUMN or 1.5 COLUMN; PLEASE QUOTE
250 FOR COLOUR PRINTING COSTS]

251 *2.3. Impact of spatial and temporal compositing on optical classification*

252 As noted by Jackson et al. (2017, hereafter JSM), averaging of the re-
253 motely sensed reflectance spectra from instantaneous, Level 2 (ca. 1 x 1
254 km resolution) to Level 3 (ca. 4 x 4 km resolution, weekly or monthly)
255 data incurs a risk of smoothing out phytoplankton dynamics associated with
256 growth/decay/advection events that are short-lived, and it increases the dif-
257 ference between any available *in situ* reflectance data used in characterising or
258 evaluating optical classes from the already spatially-averaged remote sensing
259 pixel values. Daily data may also be most appropriate for some management
260 applications. However, cloud-cover and the reduced overpass rates at low
261 latitudes make the use of daily data for capturing spatial patterns challeng-
262 ing in this region. The impact of spectral aliasing (from averaging) on the
263 classification was tested by creating classes firstly from all the Level 2 (L2)
264 data for 2003, with no averaging or regridding. The resulting classes were
265 compared against the same suite of classification procedures applied to the
266 Level 3 (L3) monthly, 4 km gridded data for 2003 and against the hybrid
267 classification scheme produced by JSM and Moore et al. (2001) who used *in*
268 *situ* data.

269 Both the L2 and L3 data were classified in three forms: Remote sensing
270 reflectance (Rrs), Rrs with the mean 2003 Rrs subtracted ($Rrs - \overline{Rrs_{2003}}$)
271 and Rrs normalized to 488 nm (Rrs/Rrs_{488}). Training data were selected
272 randomly in space and time from the study domain. For comparability with
273 previous studies, three sizes of training dataset were used. The smallest

dataset contained 42,000 pixels, corresponding approximately to the volume of data used by JSM for two of their study sites - they used two central Indian Ocean sites referenced to Longhurst (2007) provinces. Additional datasets of double and fifty times the original size were added to test for sensitivity of the classification scheme to dataset size. Each dataset was classified using the Matlab fcm function (Bezdek, 1981) with between 2 and 9 classes and the weighting exponent m was varied between 1.05 and 2.0. Class separability and compactness were assessed using the partition coefficient (F) and compactness and separation index (S) as in Moore et al. (2001); Windham (1982); Xie & Beni (1991). In contrast to previous studies, all ten available MODIS visible wavebands were used in the fuzzy classification procedure.

2.4. Interpretation of within-class spectral variability

The use of ten wavebands for an optical classification allows limited exploration of within-class spectral variability, which may be related to pigmentation and size differences caused by change in phytoplankton community composition or physiology, differences in backscatter related to the viral, bacterial and phytoplankton communities, variability in the relative quantities of coloured, dissolved organic matter or inorganic particulate matter, variability in the depth distributions of coloured materials and noise in the satellite signal (Kirk, 1994; Brown et al., 2008; Defoin-Platel & Chami, 2007; Alvain et al., 2005; Lain & Bernard, 2018; Brewin et al., 2011b). Having excluded water depths shallower than 200 m and in the absence of *in situ* validation data, the main focus here is on testing whether R_{rs} spectra varied uniformly with optical class. Residual reflectance ratios $R_{rs}(555)/R_{rs}(443)$

299 and $Rrs(555)/Rrs(488)$ were calculated after subtracting the dominant class
300 mean reflectance spectrum (Table 4) at each pixel, as indicators of spectral
301 variability. In a full application of the fuzzy classification scheme (Moore et
302 al., 2001), this step could be pre-empted by allowing multiple class member-
303 ships at each pixel.

304

305 2.5. Comparison against other classifications

306 Class spectra produced in this study and by JSM were mapped to other
307 interpretations for comparison. Firstly, the standard NASA OC3M-CI algo-
308 rithms were applied to each central class spectrum to produce chl concentra-
309 tion (O'Reilly et al., 1998; Hu, et al., 2012). This was passed to abundance-
310 based algorithms for phytoplankton size class published by Brewin et al.
311 (2010, Atlantic Ocean), Brewin et al. (2011a, global *in situ* data), Brewin et
312 al. (2012, eastern Indian Ocean) and Devred et al. (2011) (North Atlantic
313 and global *in situ* data; IOCCG, 2014). Chl was also used to select the
314 closest stratified water trophic class from Uitz et al. (2006). For comparison
315 against Alvain et al. (2005, 2008), every 10th Level 2 MODIS Rrs file from
316 2003 was used to generate local PHYSAT-equivalent mean-chl spectra (39 chl
317 divisions from 0.01 to 4.00 mgm^{-3} in intervals of $\ln(0.15)$; between 67016 and
318 5488110 pixels per chl interval with a total of 64733448 pixels). These were
319 subtracted from each mean optical class spectrum and the residual compared
320 against the criteria provided by Alvain et al. (2008).

Name	Weighting coefficient m	Partition coefficient F	Separation index S	Source	
N5	1.05	10	0.986	0.211	This study
N8	1.05	10	0.982	0.342	This study
JSM	Unknown	6	Unknown	Unknown	Jackson et al. (2017)

Table 2: Classification schemes used for further analysis. See supplementary online material, Table S1, for full classification evaluation metrics.

2.6. Classification of the 17 year Level 3 dataset

Following evaluation of the L2 and L3 classifications, a suite of three classification schemes was chosen and applied to 17 full years of MODIS-Aqua Level 3 Rrs version 2018.0 (Table 2). Water depths shallower than 200 m were excluded from analysis. This provided the framework for testing whether other remotely sensed parameters were well constrained by the optical classes.

[TABLE 2 HERE]

2.7. Relationship between colour class and biophysical parameters

Summary statistics of remotely sensed biological and physical variables were produced to characterize each class in each of seven geographical domains (Figure 3): SCTR-W (10° to 5° S, 50° to 62° E), SCTR-E (10° to 5° S, 63° to 75° E), SCTR-C (10° to 5° S, 76° to 88° E), Wiggert-N (7° to 2° S, 60° to 85° E), Wiggert-S (12° to 7° S, 60° to 85° E), the MPA, z200 (the whole study domain at depths greater than 200 m) and z1000 (the whole study domain at depths greater than 1000 m).

The potential of purely optical classifications as indicators of the pelagic growth environment was explored using within- and between-class statistics. Non-parametric analysis of variance was used to test whether the optical classifications reflected differences in each of the the biological and physical

340 variables (Kruskal-Wallis, applied to one variable at a time, assuming that
 341 underlying distributions of each variable within each class subset of each geo-
 342 graphical domain are the same, followed by post-hoc Tukey’s honestly signifi-
 343 cantly difference, HSD, tests between pairs of classes; Ruxton & Beauchamp,
 344 2008) for each of the geographical domains. The distinctiveness of smaller
 345 domains for which contrasting processes across zonal or meridional gradients
 346 have been discussed in the literature (SCTR-W,E,C; Wiggert-N, S Wiggert
 347 et al., 2006; George et al., 2018; Dilmahamod et al., 2016) was tested using
 348 permuted multivariate analysis of variance (PERMANOVA; Anderson, 2001,
 349 2017) on standardised variables (z-scores), with all fifteen biological and phys-
 350 ical variables. Anomaly time-series were used in addition to simple z-scores
 351 where clear seasonal cycles were present. Because of the large dataset sizes,
 352 100 subsets of 1000 pixels were selected randomly through time within each
 353 geographical domain for bootstrapped testing; pixels with missing data were
 354 excluded at each iteration. Domain was taken as the first, fixed, factor and
 355 class as the second, nested factor. Euclidean distance, correlation distance
 356 and squared correlation distance gave similar PERMANOVA hypothesis test
 357 results, whilst χ and χ^2 results varied; euclidean distance and 1000 permuta-
 358 tions were used for all reported results (McCune & Grace, 2002; Anderson,
 359 2017; Pillar, 2013). The fathom Matlab toolbox was used for PERMANOVA
 360 tests (Jones, 2012). Mann-Kendall-Sen correlation coefficients (Sen, 1968;
 361 Hamed & Rao, 1998) were used to identify which biological and physical
 362 variables covaried within each class. Correlations were tested firstly using all
 363 data and secondly using bootstrapping to insure against spatial and temporal
 364 autocorrelation effects (100 random subsamples of 1000 pixels).

365 A full comparison of optical classes with OCO-2 pCO₂ data was beyond
366 the scope of this study - OCO-2 was launched in July, 2014, and the data
367 therefore do not span the study period considered here. OCO-2 data are also
368 lower in spatial and temporal coverage, although the recent launch of OCO-3
369 will mitigate this. As a proof of concept, a single month of OCO-2 surface
370 pCO₂ data were matched to MODIS Level 2 optical classes for January 2015
371 (same-day match-ups only), and between-class differences evaluated using
372 Kruskal-Wallis and post-hoc HSD tests, for the z1000 study domain.

373 Assuming that optical classifications do partition pixels dynamically in
374 space and time according to the physical and biological variables that can
375 be derived from remote sensing data, two additional tests were applied to
376 establish whether conditions within pixels assigned to each class changed
377 during the study period, and whether they are correlated with basin- and
378 global-scale circulation patterns. Mann-Kendall-Sen trend tests were ap-
379 plied in regressions of the class-averaged time-series of each remote sensing
380 variable against climate indices that characterize the Indian Ocean Dipole
381 (Dipole Mode Index, referred to hereafter as IOD; Saji & Yamagata, 2003),
382 Madden-Julian Oscillation (MJO; Kilidas et al., 2014) and El Nino-Southern
383 Oscillation, Southern Oscillation index (SOI; Ropelewski & Jones, 1987).
384 The SOI was chosen because it represents variability in the Walker circula-
385 tion, rather than directly in SST or combinations of variables. Habitat frag-
386 mentation metrics were used to characterise the distribution of lower trophic
387 level resources in the MPA, Wiggert-S and Wiggert-N domains to investigate
388 whether changes related to climate patterns can be detected in the pelagic
389 growth environment using available remote sensing data between 2002 and

2018. Patches of the same optical class were created using each month's class map as a binary image, grouping adjacent pixels of like-class (Matlab `bwconncomp`) and finding the perimeter, patch centre and number of pixels contained within each class patch (Matlab `regionprops`). Average patch area, distance between patches and patch density were calculated (Wang et al., 2014) for each month for temporal regression and regression against climate indices (Mann-Kendall-Sen test). Finally, correlations between the total area occupied by each optical class within each subdomain and climate indices were tested (Mann-Kendall-Sen trend test).

To test whether optical classifications could form a useful basis for fisheries management and enforcement, fishing catch and effort data from the Indian Ocean Tuna Commission were acquired at monthly temporal resolution, in 1° and 5° grids (IOTC, 2020, 2014). The most common optical class was assigned to each of these coarse fisheries grid cells, matched by year and month, and between-class differences in catch, effort and catch per unit effort (CPUE) were evaluated (Kruskal-Wallis and post-hoc HSD). This comparison was applied across the z1000 domain and within the IOTC data gridcells that contain the MPA as well as $15^\circ \times 15^\circ$ and $25^\circ \times 25^\circ$ domains centred on the MPA and the full z1000 area. Long-line fishing effort was reported in hooks, whereas surface fishery effort was reported in hours; the two datasets were analysed separately.

All data analysis and visualisation was carried out using Matlab 2018a running under MacOS10.12.6.

3. Results & Discussion

3.1. Comparison of classification results with Level 2 and Level 3 input data

Classification results were similar for L2 and L3 data for the two smaller datasets, but classes were less compact when the largest dataset was used. Partition coefficients (F) remained above 0.9 for all three Rrs treatments with weighting coefficients 1.05, 1.1 and 1.2, but separation coefficient S increased from 0.15 to 0.3 as the number of classes, N, was increased from two to nine, and the best separability was obtained for 2 classes with all data treatments. Beyond N=8 classes, processing time increased significantly and F and S values were unstable between repeat runs, therefore no classification with 9 or more classes was pursued. Classification performance metrics (F and S) for all L2 and L3 classifications are provided in full in the Supplementary Online Material, Table S1, sheet ‘L2 L3 Classification F S’. Rrs treatment made little difference to F and S for Rrs and $\overline{Rrs(2003)}$ (<1% variability in performance metrics for $m < 1.3$), but separation index increased by a factor of 20 for Rrs/Rrs(488).

Figure 4 shows the JSM classes and the L2 and L3 class spectra produced using Rrs with $m = 1.05$ and two to eight classes, and Table 3 shows the proportions of pixels mapped to each class, with N=5 and N=8 class schemes mapped to JSM classes using Euclidean distance between Rrs values at the six common wavebands. Fewer of the JSM classes were reproduced in L3 classifications than in L2 classifications, with the extreme blue-water spectra (highest and lowest Rrs(412) values) lost in an N=8-class scheme (Table 3). Class spectra fell within or at the edges of the JSM classes, with most deviation in spectral shape (relative to the closest JSM class) in the blue:green

438 wavebands. Only one of the three JSM coastal water classes was found in
439 the N=8 classifications produced here, and none with N=5.

440 [FIGURE 4 HERE, SINGLE-COLUMN; COLOUR ONLINE ONLY]

441 [TABLE 3 HERE]

442 As a compromise between F and S metrics and representation of the JSM
443 classes, and to enable direct comparison of results against similar studies,
444 the 5- and 8-class schemes (henceforth N5 and N8) produced using $m=1.05$,
445 42,000 training pixels and absolute Rrs spectra were used for further analy-
446 sis, together with the 6-waveband JSM classification. These Level 2 and 3
447 classifications for 2003 are summarized in Table 3 and the class spectra are
448 given in Table 4.

449 [TABLE 4 HERE]

450 Mapping of the N5 and N8 classifications from Level 2 and Level 3 data
451 to abundance-based and reflectance-based PFT algorithms is shown in Fig-
452 ure 5. PFT algorithms consistently interpreted the lower-OC3-CI chl classes
453 as being dominated by picoplankton, with the contribution of nano- and
454 microplankton fractions increasing with increasing chl. Eutrophic-type dis-
455 tributions, dominated by microplankton, were only produced in the N8 clas-
456 sification, but this was the only difference between 5- and 8-class schemes.
457 Since the proportion of pixels assigned to the 8th class was 0.3% (Table 3),
458 the abundance-based PFT algorithms generally classify these optical classes
459 as dominated by small cells. In contrast, switching from Level 2 to Level 3
460 data produced more PHYSAT-type spectra that fell into the pseudo-diatom
461 class (similar spectral shape to PHYSAT-diatom, but higher Rrs values).
462 The N5 and N8 schemes correspond to JSM classes between 1 and 9, and

Class	Level 2 % pixels assigned to each class			Level 3 % pixels assigned to each class		
	N5	N8	JSM	N5	N8	JSM
1		5.7	1.9			0.5
2	10.7		6.7	11.9	8.1	5.6
3		14.8	14.6		14.0	13.2
4	24.7	19.7	18.0	20.9	16.3	17.2
5	28.0	20.9	16.6	24.7	17.7	17.2
6		18.5	13.4		14.8	12.9
7	23.9		9.7	19.7	10.9	8.3
8		14.0	7.7			6.0
9	12.8		4.6	12.0	6.0	3.5
10		6.3	3.6			2.3
11			2.6			0.8
12			0.3			0.3
13		0.3	0.2		0.3	0.2
14			0.1			0.1

Table 3: Total number of Level 2 and Level 3 dominant classes mapped to each fuzzy cluster in the 10-waveband classifications (5 classes and 8 classes, N5, N8) and in the Jackson-Moore 6-waveband classification. The 10-waveband classes are mapped to the JSM classification by Euclidean distance (Figure 4).

Classification / class no.	Rrs(412) x1000	Rrs(443) x1000	Rrs(469) x1000	Rrs(488) x1000	Rrs(531) x1000	Rrs(547) x1000	Rrs(555) x1000	Rrs(645) x1000	Rrs(667) x1000	Rrs(678) x1000
N5: 1	15.032	11.065	9.366	7.298	2.621	1.929	1.535	0.009	0.204	0.256
N5:2	12.186	9.035	7.988	6.455	2.523	1.857	1.489	0.076	0.197	0.248
N5:3	9.620	7.332	6.782	5.672	2.447	1.814	1.458	0.059	0.191	0.247
N5:4	6.974	5.602	5.482	4.787	2.388	1.807	1.468	0.058	0.196	0.262
N5:5	3.961	3.476	3.748	3.480	2.215	1.768	1.474	0.085	0.224	0.324
N8:1	15.530	11.400	9.556	7.381	2.558	1.861	1.467	0.077	0.183	0.236
N8:2	13.131	9.683	8.426	6.718	2.519	1.842	1.469	0.070	0.189	0.241
N8:3	11.193	8.355	7.507	6.138	2.462	1.808	1.445	0.057	0.184	0.236
N8:4	9.375	7.166	6.656	5.583	2.418	1.787	1.433	0.048	0.183	0.241
N8:5	7.500	5.948	5.743	4.965	2.375	1.778	1.435	0.045	0.186	0.250
N8:6	5.514	4.593	4.680	4.196	2.313	1.779	1.457	0.054	0.195	0.271
N8:7	3.184	2.913	3.268	3.103	2.146	1.746	1.467	0.096	0.236	0.348
N8:8	7.662	9.253	11.578	12.541	14.101	14.027	13.220	4.781	3.947	3.880

Table 4: Class Rrs spectra for the N5 and N8 classifications produced in this study using Level 3 data.

with a finer gradation of Rrs spectra corresponding to 0.03 to 0.5 mgm^{-3} , the JSM classes include more scope for identifying mixed size-class waters with the PFT algorithms included here. The PHYSAT classification was designed to identify cases in which a single PFT dominates water colour and, correspondingly, not all classes could be mapped to a PHYSAT class (Figure 5). Of those that did, *Synechococcus*-like cyanobacteria was the most common designation (3 classes at Level 2, N5; 4 classes at Level 2, N8; 1 class at Level 3, N5 and JSM and 2 classes at Level 3, N8).

[FIGURE 5 HERE: SINGLE-COLUMN; ONLINE COLOUR ONLY]

Class maps for every second month of 2003 (Figure 6) show more fragmented spatial distribution of classes in the L2 data, corresponding to the higher spatial resolution of the data and higher spectral separability. Broader spatial patterns (01000 km) are consistent between all classifications and include a limited seasonal north to south shift in class, distinction of coastal, Somali Current and Arabian Sea waters from the central domain and coher-

ent and elongated patches of higher class (4-5 in the N5 classification) waters in the central Indian Ocean, often enveloping the Chagos MPA. These zonal bands are more evident in Figure 7, which summarises class diversity at each pixel, as corresponding to the seasonally-varying SCTR, SEC and SECC domains. Figure 8 shows a subset of the remote sensing climatologies, for comparison.

[FIGURE 6, 7, 8 CLOSE TO HERE: SINGLE-COLUMN or 1.5 COLUMN; PLEASE QUOTE FOR COLOUR PRINTING]

Whether or not the smaller features correspond to ecologically meaningful variations in the microbial biome or carbon cycling can only be answered definitively with *in situ* data. The microscopy stations lay within classes 3 and 4, and a significant difference in phytoplankton biovolume was confirmed (class 3 mean biovolume = $8.1 \times 10^5 \pm 2.6 \times 10^5 \mu m^3 l^{-1}$, N=3; class 4 mean biovolume = $2.3 \times 10^6 \pm 8.4 \times 10^5 \mu m^3 l^{-1}$, N=9; Kruskal Wallis $p = 0.0126$). There was also an order of magnitude difference in the ratio between phytoplankton and zooplankton biovolume (Schwarz, 2020). In the absence of a larger *in situ* dataset with which to evaluate the full classification, comparison between these classifications and previous studies is helpful. In spatial and temporal variability, these optical classifications are most similar to previous studies that use Rrs or radiance, as expected, and to some of the mesopelagic biogeographies. The degree of patchiness is consistent with examples given by JSM, and the classes assigned to the SCTR in their example of July 2004 are in direct agreement with the N5 classification produced here (JSM classes 7/8, based on 6 wavebands, correspond to N5 class 4, based on 10 wavebands; Table 3). A similar degree of patchiness is reported by Oliver & Irwin (2008)

503 using nLw412, nLw551 and SST, and their approach, allowing the number
 504 of classes to emerge from the data, assigned up to ten classes over the TIO,
 505 which supports the richer N8 or JSM (N=14) classifications tested here in
 506 terms of spectral separability. George et al. (2013) reported multiple patches
 507 of elevated chl extending some 200 km along 67° E in the SCTR, as well as
 508 suppression of surface chl by eddies further south. In contrast, classifications
 509 that used chl as the only ocean colour variable, together with other physical
 510 drivers, have been less spatially diverse (e.g. Longhurst, 2007; Spalding et
 511 al., 2012; Reygondeau et al., 2013; Fay & McKinley, 2014; Sayre et al., 2017,
 512 surface zone) and do not distinguish the SEC or SCTR domains clearly.

513 Mesopelagic classifications using a range of approaches including derived
 514 ocean colour variables, acoustic data, World Ocean Atlas data and species
 515 abundance mostly do distinguish the SCTR and SEC zones (Proud et al.,
 516 2017; Sutton et al., 2017; Sutton & Beckley, 2017; Sayre et al., 2017, 200 to
 517 800 m zones) although in some cases the distinction between coastal influ-
 518 ences and SEC/SCTR features is unclear (Costello et al., 2017; Reygondeau
 519 et al., 2018). Differences in the spatial richness of mixed-input epipelagic
 520 classifications relate partly to the scales and methods used, but may also
 521 reflect subtle changes in the growth environment that are related to phyto-
 522 plankton community composition that are not detected in the chl algorithms,
 523 or to chl variability being outweighed in a classification by the contribution
 524 of SST, producing spatially coarser structures because of dynamics that have
 525 no surface signature in SST (e.g. Santoso et al., 2010; Drushka et al., 2012;
 526 Strutton et al., 2015), or both. Previous analysis of phytoplankton bloom
 527 dynamics in the TIO, based on satellite-derived chl and biogeochemical mod-

528 elling, suggested a summer bloom spanning the full breadth of the basin at
 529 SEC latitudes, but no winter bloom (Levy et al., 2007), and the modelled
 530 emergent biogeography of Follows & Dutkiewicz (2011) predicted a band
 531 of *Prochlorococcus* analogs and low species richness in this region which is
 532 consistent with the PFT algorithm interpretation of classes 1 to 3 with N5.
 533 However, Wiggert et al. (2006) predict larger phytoplankton cells between
 534 January and May in the deep chlorophyll maximum in the SCTR and Jeffries
 535 et al. (2015) found eukaryotes contributed >10% to relative cell abundance
 536 at a deep water site within the Chagos MPA. Similarly, Thorrrington-Smith
 537 (1971) found diatom and dinoflagellate communities in water samples from
 538 100 m depth across the western TIO - a signal that is consistent with the
 539 higher-chl PFT interpretations of classes 5 (N5) and 6-8 (N8) which could be
 540 expected to be detected in satellite data in zones of strong vertical mixing,
 541 such as the tropical gyre boundaries.

542 The loss, at L3, of classes representing high and very low chl values,
 543 may be important for monitoring carbon export and ecosystem resources and
 544 Duarte et al. (2013) suggested that a chlorophyll concentration of 0.44 mgm^{-3}
 545 represents a transition between heterotrophic and autotrophic communities.
 546 Application of the Duarte et al. (2013) threshold to L3 chl values in this area
 547 is consistent with Fernandes et al. (2008), who reported net heterotrophic
 548 production between 1° N and 5° S at 83° E , but the appearance of higher
 549 ranges in L2 data, the prevalence of higher classes for most of the year and
 550 the paucity of *in situ* data for the pelagic MPA domain area renders this
 551 use of the classification results uncertain. Level 2 data classifications are
 552 therefore potentially valuable for modelling and monitoring tasks (Tweddle

et al., 2018), despite the low daily spatial coverage. For the purposes of exploring spatiotemporal variability over the MPA and wider TIO domain, Level 3 classifications are explored further.

3.2. Are biological and physical variables distinct for each optical class ?

The class-specific ranges of remote sensing variables are summarised for the MPA and z1000 domains in Figure 9 and Appendix 1. The N5 optical classification constrained all variables most effectively. For the MPA, only sea surface height (classes 4 and 5), eddy kinetic energy (classes 3 and 5) and the R443:R555 reflectance ratio (classes 1 and 5) were ambiguous (Kruskal-Wallis with post-hoc HSD tests, $p < 0.01$, $N > 10000$). Increasing class number in the MPA was associated with increasing chl, FLH, ∇ SSH, u , τ_N and depth, and with decreasing FLH:chl ratio, SST, SSH, v and τ_E . Westerly and northerly currents, and westerly wind stress, associated with SWM wind reversals, were associated with class 1 only (lowest chl).

[FIGURE 9 HERE; SINGLE-COLUMN; COLOUR ONLINE ONLY]

Greater between-class ambiguity in optical and physical variables was found with N8 and JSM, both of which included sparse classes ($N < 10000$). For N8, class 8, representing OC3M-CI chl = 2.39 mgm^{-3} , was always sparse and class 7 (chl = 0.46 mgm^{-3}) was sparse in the SCTR-E and -C domains. Similarly, JSM classes 10 to 14, representing chl $> 0.62 \text{ mgm}^{-3}$, were always sparse or empty, and class 9 (chl = 0.47 mgm^{-3}) was sparse in all the sub-domains except z1000 and SCTR-W. Between-class variability was not significant in the MPA for the majority of physical variables in N8 and JSM (SSH, EKE, u , v , τ , $\nabla \times \tau$) or for FLH, but chl and the residual reflectance ratios were significantly different in all 8 classes. This could be interpreted as

578 a smaller number of physically-distinct conditions hosting a larger number
 579 of optically distinct conditions, consistent with growth, decay and succes-
 580 sion occurring within each physical 'province' over the averaging period of 1
 581 month. In JSM, only 8 classes were well-populated (No. pixels > 10000) and
 582 all variables except $\nabla \times \tau$ were ambiguous for two or more classes. In the
 583 wider z1000 domain, between-class variability was significant for most vari-
 584 ables in each of the classifications (exceptions were N5: SST for classes 2 and
 585 3; N8: ∇ SSH for classes 4 and 7, u for classes 6 and 7; JSM: SST for classes
 586 1 and 8, v for classes 7 and 8 and τ_E for classes 1 and 2), reflecting the much
 587 greater size of this dataset. In this wider domain, the tendency of windstress
 588 variables with increasing class number was reversed so that increasing class
 589 and chl were associated with increasing τ_E and decreasing τ_N .

590 All between-class test results are given in Supplementary Online Mate-
 591 rial Table S1, sheet 'Variable Ranges by Class'. The optical classifications do
 592 correspond to distinct ranges of biotic and abiotic variables, suggesting their
 593 potential value in providing a useful diagnostic for management and mod-
 594 elling applications. Five optical classes produces least ambiguity in physical
 595 variables, although significant residual ocean colour differences are detected
 596 in up to 7 classes. However, the biophysical relationships vary within the
 597 wider domain, as may be expected from the known oceanographic processes
 598 in the region, suggesting that the use of optical classes may be most appro-
 599 priate within a heirarchical scheme (c.f. Kavanaugh et al., 2014; Oliver et al.,
 600 2004; Oliver & Irwin, 2008). Exploration of between-class variability within
 601 different sub-domains is addressed in the next section.

602 To test whether seasonal variability in winds and associated mixing and

603 entrainment (Halkides & Lee, 2011; Wiggert et al., 2006) produces between-
 604 class ambiguity, Kruskal-Wallis tests were applied to the MPA data for each
 605 month (Online Supplementary Figures S2, S3; Tables S2, S3). Seasonal wind
 606 stress reversals were detected with the annual average conditions (negative
 607 τ_E decreasing with increasing class number) found during austral winter and
 608 the opposite trend (positive τ_E increasing with increasing class number) dur-
 609 ing austral summer (Figure S2). The winter months were significantly differ-
 610 ent at $\alpha=0.01$ for classes 3, 4 and 5 (higher chl), whereas austral summer
 611 months were distinct in classes 2 and 4, and τ_N trends generally mirrored τ_E
 612 trends. Less pronounced seasonal reversals were found for u, v and ∇SSH .
 613 One heirarchical partitioning approach could therefore be to use the optical
 614 classifications with a monthly or seasonal interpretation to constrain vari-
 615 ability in the epipelagic growth environment, but a more objective approach
 616 using the physical variables at higher levels of the heirarchy avoids the need
 617 to assume a regular seasonal cycle. In either case, the correspondence be-
 618 tween optical class and biotic environmental conditions needs to be explored
 619 using *in situ* biogeochemical data if the optical classification is to be used to
 620 deduce ecological function.

621 3.3. Do optical classifications capture zonal or meridional differences around 622 the Chagos MPA?

623 Between-class variability in remotely sensed variables for each of the sub-
 624 domains is shown in Figure 10. Most remotely sensed variables tended to
 625 increase or decrease monotonically with increasing optical class number, and
 626 spatially segmenting the dataset had little effect on these tendencies. Of the
 627 three SCTR domains, the westernmost area had the lowest SST values in

all classes, as well as lower SST and higher τ_N in the highest-chl class 5. The central TIO domain, SCTR-C, which is the easternmost of the three, had the highest SST and EKE values and much lower nFLH:chl values in all classes. Stronger contrasts were evident between the Wiggert-N and -S domains. nFLH, v , τ_N and nFLH:chl were lower in the northern than the southern domain in some or all classes, whilst SST, ∇SSH , EKE, u and τ_E were higher. Whereas the depth-resolved modelling studies of George et al. (2018) and Dilmahamod et al. (2016) suggested east-west gradients in biophysical mechanisms operating across the SCTR, sub-domains SCTR-E, -W and -C could not be distinguished in the surface remote sensing variables studied here (PERMANOVA, $p > 0.1$, $N > 691 \times 100$; Table ??), although the optical classes were significantly different in all domains and classifications ($p < 0.005$, $N > 692 \times 100$). In contrast, differences were detected between classes and domains for the north-south division discussed by Wiggert et al. (2006) ($p < 0.037$, $N > 692$; Table 5). Excluding the shelf slope depths between 200 and 1000 m had no impact on class ambiguities in any variable (Figure 10).

[FIGURE 10 HERE; SINGLE-COLUMN; ONLINE COLOUR ONLY]

[TABLE 5 HERE]

Correlations between physical and biological variables within each class and domain are shown in Figure 8. Chl was negatively correlated with SST in all but the z200/z1000 domains. In the MPA, increasing SST was associated with lower EKE and τ_N and with higher v , τ_E and $\nabla \times \tau$ in the lowest-chl class (class 1) only, with the reverse relationships found in classes 2 to 5. The lower chl class 1 is therefore associated with conditions in which τ_E and $\nabla \times \tau$

Table 5: Results of PERMANOVA tests for differences between regions. a) Fixed factor 1 is location in the western or eastern SCTR or adjacent central Indian Ocean domain; nested factor 2 is optical class. All 15 remotely sensed variables were included in the test as z-scores. Average values are from 100 random draws of 1000 data points from each domain, time-matched. Pixels with any missing data were excluded, yielding residual degrees of freedom between 600 and 800 (denoted 'Resid'). p-values were calculated using 999 permutations. b) Fixed factor 1 is location in the northern or southern central Indian Ocean sectors; other details as for a).

a) Seychelles-Chagos Thermocline Ridge W / E / C (George et al., 2018)					
	Factor	df	$\overline{F}$	$\overline{p}$	p range
N5	F1: Domain	2	1.096	0.406	0.104 to 0.847
	F2: Class	10.9 (10-12)	15.066	0.0012	0.001 to 0.006
	Resid:	[691-766]			
N8	F1: Domain	2	1.256	0.271	0.048 to 0.700
	F2: Class	15.8 (14-18)	13.785	0.001	0.001 to 0.003
	Resid:	[703-765]			
JSM	F1: Domain	2	1.515	0.142	0.015 to 0.424
	F2: Class	18.3 (16-22)	12.825	0.001	0.001 to 0.001
	Resid:	[688-771]			
b) Central tropical Indian Ocean N / S (Wiggert et al., 2006)					
	Factor	df	$\overline{F}$	$\overline{p}$	p range
N5	F1: Domain	1	3.420	0.0093	0.001 to 0.037
	F2: Class	7.4 (6-8)	20.871	0.0011	0.001 to 0.005
	Resid:	[692-755]			
N8	F1: Domain	1	4.030	0.003	0.001 to 0.014
	F2: Class	10.2 (9-11)	18.127	0.001	0.001 to 0.001
	Resid:	[708-775]			
JSM	F1: Domain	1	4.415	0.0023	0.001 to 0.013
	F2: Class	12.1 (10-15)	16.044	0.001	0.001 to 0.001
	Resid:	[692-755]			

653 do not correspond to surface cooling and the northward geostrophic current
 654 component is positive. Chl and nFLH were positively correlated everywhere
 655 except in class 1 (lowest chl) in the MPA, SCTR-C and Wiggert-N,S areas.
 656 Higher nFLH:chl ratios were associated with lower τ_E , $\nabla \times \tau$, u, EKE, ∇ SSH,
 657 SSH and SST in the MPA class 5, and to a lesser extent for classes 3 and 4,
 658 suggesting the importance of the tropical gyre strength for surface chl. These
 659 relationships are similar, but less pronounced, in the SCTR-W, Wiggert-N
 660 and Wiggert-S domains, and in z1000 classes 3 and 4. Assuming relative
 661 homogeneity of the phytoplankton community within a given optical class,
 662 nFLH:chl can be interpreted as a proxy for relatively high iron limitation
 663 (as opposed to other limiting factors, Behrenfeld et al., 2009). Although
 664 Chinni et al. (2019) and Wiggert et al. (2006) suggest Fe-limitation in some
 665 seasons within and north of the SCTR, George et al. (2013) reported that
 666 the deep chlorophyll maximum (DCM) followed the nitricline and did not
 667 measure iron concentrations, so the interpretation of nFLH:chl requires more
 668 *in situ* data in this region. SLA, which is related to westward-propagating,
 669 downwelling Rossby waves in the SCTR (George et al., 2018), decreased
 670 with increasing optical class and was negatively correlated with chl in at
 671 least 4 classes in all domains, including z1000 (Figure 11). The use of class-
 672 specific correlations across different sub-domains captured other contrasts in
 673 physical relationships, such as a switch from positive to negative coupling
 674 between u and EKE in MPA class 1, SCTR-W classes 1,2,3 and 5; SCTR-C
 675 classes 1 and 2; Wiggert-N class 1, whilst the relationship was negative for
 676 all classes in Wiggert-S, suggesting dominance of the westward SEC in the
 677 southern domain and more varied interactions in the northern SCTR (e.g.

mesoscale and fine-scale processes, George et al., 2013; Hosegood et al., 2019). For all domains, the greatest between-class differences were found between classes [1,2] and classes [3,4,5], suggesting that the different physical processes driving the growth environment can be distinguished by the chl 0.08 mgm³ isoline (Figure 5). Biophysical coupling in the Chagos MPA was most similar to that in the Wiggert-S domain (Figure 10; Supplementary Material Table 1, sheet ‘Variable Ranges by Class’).

[FIGURE 11 HERE; SINGLE-COLUMN; COLOUR ONLINE ONLY]

3.4. *Modelling applications of optical partitioning*

Assessment of the global carbon cycle and sequestration of anthropogenic CO₂ emissions underpins the Paris Agreement (UN, 2015). The oceanic biogeochemical models used for global carbon cycle assessment remain fairly simplistic and uncertainties are high (Lim & Kug, 2017; Le Quéré et al., 2013, 2018; Gruber et al., 2019); increasing the complexity of ecosystem dynamics in models without rendering them unstable is challenging (Anderson, 2005) and different approaches are still being developed (e.g. Hense et al., 2017; Wanninkhof et al., 2013). Optical classes offer an empirical constraint on ecosystem models and provide a dynamic framework for aggregating model outputs and assessing model skill, for example in the prediction of CO₂ uptake or sequestration rates. The preliminary comparison of pCO₂ between optical classes supports both of these applications (Figure 12). Between-class differences in surface partial pressure of CO₂ were significant ($p < 0.01$, $N > 13,000$) for all N5 classes. pCO₂ distributions were mostly bi-modal, reflecting a background latitudinal gradient in class (increasing class number, reflecting increasing chl, to the north) with patches of higher classes to

703 the south. The average $p\text{CO}_2$ value decreased slightly with increasing class
704 number (416 ± 49 , 418 ± 48 , 404 ± 44 , 406 ± 48 , 398 ± 48 ppm), in contrast with
705 Nagelkerken et al. (2015) who reported no simple relationship between pri-
706 mary production and CO_2 uptake in this area. Between-class differences in
707 this study were within the version 8 OCO-2 model error for X_{CO_2} , but uncer-
708 tainties in the profile retrieval are not specified (O'Dell et al., 2018). Because
709 of these uncertainties, the scope of this comparison and because too few data
710 were available to test between-area differences, a more complete comparison
711 is reserved for a future study.

712 [FIGURE 12 HERE; TWO-COLUMN; COLOUR ONLINE ONLY]

713 In a fully-realised fuzzy optical classification, multiple colour class mem-
714 berships enable mixed phytoplankton communities (or mixed water types)
715 to be represented and this has been shown to be useful in enhancing the in-
716 terpretation of ocean colour data as well as constraining chl algorithm errors
717 (e.g. Moore et al., 2001, 2009). Better remote sensing information about the
718 ocean surface microbial community feeds into the Conservation on Biodi-
719 versity as well as monitoring and understanding the ecosystem services they
720 provide (CBD, 2010; Tweddle et al., 2018; Roberts et al., 2017). In this study,
721 the Rrs class residuals did not show promise for identifying details of phyto-
722 plankton community composition, but ensemble class biophysical agreement
723 suggests that ecosystem function may be constrained by optical class.

724 3.5. *Management and conservation applications of optical partitioning*

725 Information is lost when Rrs spectra are partitioned into discrete classes,
726 rather than applying an algorithm to produce a continuous biological variable
727 such as chl, but a distinct advantage of this is the possibility of using habitat

728 fragmentation metrics on the class maps. Fragmentation metrics applied to
 729 the N5 class maps are shown in Figure 13 for the MPA and the Wiggert-
 730 N and -S domains. The lower-chl classes 1 and 2 behaved differently in
 731 the three domains: Class 1 patches were of order 5000 km² and generally
 732 largest to the south, where they were separated by ca. 200 km. In the
 733 MPA, between-patch distances were shorter and stable through time, whereas
 734 they fluctuated between 20 and 100 km to the south. The 2011 La Nina
 735 period coincided with particularly high separation distances in Wiggert-N
 736 (separation ca. 500 km) and with high patch sizes in Wiggert-S (areas up to
 737 ca. 15 000 km²; Figure 13). In contrast, class 2 patches were larger overall
 738 (O20 000 km²) with higher values to the north, and a maximum coinciding
 739 with the 2015 El Nino (mean patch size ca. 50 000 km²). Class 2 patch
 740 separation distances fluctuated in Wiggert-N as for class 1 but with a lower
 741 range (20 to 70 km). The 2015 El Nino coincided with the highest class 2
 742 patch sizes in Wiggert-N (up to ca. 50 000 km²). Classes 1 and 2 represent
 743 clear, warm water, which is a foraging habitat used by seabirds preying on
 744 flying fish and squid, often in association with subsurface predators (e.g.
 745 Weimerskirch et al., 2005; Catry et al., 2009b; Le Corre et al., 2012). The
 746 absence of seasonal cycles in the fragmentation metrics for these classes is
 747 marked: Prey occurrence, driven by cetaceans and tuna, is stochastic, but
 748 patches of similar foraging conditions are predictable at the monthly scales
 749 used here, with patch separations that are within the known range of some
 750 seabirds (Weimerskirch et al., 2007; Nel et al., 2001; Pinaud & Weimerskirch,
 751 2007).

752 [FIGURE 13 HERE: SIGNLE-COLUMN; BLACK AND WHITE]

Patch sizes were smaller in the higher-chl classes 3 and 4 (mostly within 5000 km²), with separation distances between 20 and 200 km and marked seasonality. For classes 1 to 4 the MPA clearly straddles the Wiggert-N and Wiggert-S conditions, potentially providing stability of resource within foraging range in the event of extreme (Nino/Nina/IOD) conditions. The highest-chl class 5 was not always present (Figure 13e, j, o) and represents the smallest but most intense resource patches (10 to 300 km² in size), separated by 20 to 100 km in the MPA, 40 to 300 km in Wiggert-S and 20 to 1000 km in Wiggert-N. Scott et al. (2010) and Trevail et al. (2019) highlight the importance of fine spatial and temporal scales in prey resource; Level 2 (ca. daily, 1 km) or higher spatial resolution data are therefore also of interest.

Class 5 patches were largest within the MPA domain up to 2014, after which increasingly large class 5 patches appear in Wiggert-N (Figure 13j). However, the time-series is too short to confirm whether this is a robust trend. Significant temporal trends in fragmentation metrics were only found for fragmentation distance in the MPA, where the average distance between class 2 patches increased over the study period, whilst the distance between class 3 patches decreased ($p < 0.05$, $N = 204$). There is evidence of a shift in the spatial distribution of resources over the study period, but without loss of any of the colour classes, suggesting that the range of niches that seabirds exploit has been maintained across the Wiggert-N and -S domains (Waugh & Weimerskirch, 2003; Catry et al., 2009a; Le Corre et al., 2012). In this study, water depths shallower than 200 m were excluded to avoid land adjacency and bottom reflectance effects, so the MPA domain metrics do not include the near-shore and lagoon waters which may augment class 5.

Although mesopelagic biomes have been shown to reflect the spatial distribution of primary production (Proud et al., 2017), the only direct link between optical classes and large, commercially-fished species that could be expected is through water clarity for foraging. To test whether optical classes could provide useful fisheries management information, class maps were compared against fisheries records. The Chagos MPA has been a no-take zone since it was established in 2010, and IOTC fishing records amalgamated over the study period (2002 to 2018) are correspondingly lower in waters immediately adjacent to the MPA and increase further away (Table 6). However, 284 surface fishery records and 43 longline records were reported after 2010 in the IOTC gridcells that contain the MPA (-2.3 to -10.8°S, 67.9 to 79.4°E). Figure 14 shows the distribution of average effort, catch and CPUE at increasing distances from the MPA for the surface and longline fisheries. In the wider domain (z1000), surface fishery catch and CPUE mostly increased with increasing N5 class, as expected (e.g. Solanki et al., 2015, 2017; Mohamed et al., 2018), with significant between-class differences between low- and high-chl waters (Kruskal-Wallis, $p < 0.01$), and effort focussed in class 5 waters. Longline fishery effort and catch were highest in class 3, and although a trend for increasing CPUE with increasing class was apparent, it was not statistically significant. Few significant between-class differences were found in the MPA, but CPUE was highest in class 4, whereas effort was decreased from class 2 to 4 (Figure 14d, f). These patterns reflect reported catches, but may be used to infer behaviours of illegal fisheries and so target monitoring resources. This management application of the optical classification is easy to apply using L2 data, but is limited by cloud cover and ca. 1 day data

Table 6: Number of fishing records submitted to the IOTC within the 1° and 5° IOTC gridcells that contain the MPA, then incrementally as boxes of 3 x 3 and 5 x 5 of the 5° IOTC gridcells are added around the MPA area, and finally for entire z1000 domain. LL=long-line fisheries; Surf=surface fisheries; + denotes an increment from the MPA number and Total is the number of records reported within the full z1000 domain.

N5 class	MPA		3x3 x5°		5x5 x5°		z1000		Total	
	LL	Surf	LL	Surf	LL	Surf	LL	Surf	LL	Surf
1	0	73	+15	+131	+84	+254	+792	+948	891	1406
2	14	588	+527	+1375	+650	+2821	+1691	+12078	3152	16862
3	23	668	+648	+1381	+1681	+4674	+3264	+20870	5616	27593
4	6	82	+63	+178	+701	+1665	+2614	+20746	3384	22671
5	0	0	0	0	+25	+100	+1266	+11457	1291	11557

latency.

[TABLE 6 ROUGHLY HERE]

Knowledge of fishing behaviours is also relevant to MPA design and, potentially, to the use of dynamic MPA designations. Dynamic protection boundaries serve as a compromise between static MPA boundaries, which protect relatively small areas of rare or valuable habitat (e.g. Oppel et al., 2018; Heerah et al., 2019; Handley et al., 2020; Williamson et al., 2019), and the very large foraging domains that are intractable to static protection but where seabirds, cetaceans and large predatory fishes such as tuna are known to collocate in the tropical Indian Ocean (Anderson, 2014; Letessier et al., 2017, 2019; Pinheiro et al., 2019; Hobday et al., 2010).

[FIGURE 14 ROUGHLY HERE: 1-COLUMN or 1.5-COLUMN; COLOUR ONLINE ONLY]

3.6. Are there class-specific temporal trends in biological/physical variables?

All conservation and management efforts must consider temporal variability and the possible impacts of climate change (IPCC, 2013). The 17-year

819 study period used here is sufficient to characterise temporal variability but
820 not to detect long-term change in this region (Henson et al., 2016). However,
821 where biophysical conditions are coupled to large-scale climate indices for
822 which robust, long-term time-series are available, the patterns observed can
823 be extrapolated backwards and, with modelling, forward in time.

824 Time-series of selected variables are shown in Figure 15 for the MPA and
825 Figure S4 for the Wiggert-S domain. A weak tendency for increased chl
826 values in the austral winter was observed, but with no clear seasonal cycle
827 either for the MPA as a whole or within a given class. This is consistent with
828 removal of seasonality by class-switching (c.f. Figure 6). The class 4 and
829 5 chl values overlapped (Figure 15a), with class 4 representing an elevated
830 background level of chl compared with classes 1 to 3, superimposed with
831 stochastic, higher chl events in class 5, which often coincided with increases
832 in nFLH. The strongest chl peak, in 2011, coincided with higher ∇ SSH and a
833 protracted period of positive SOI index (Nina conditions). Higher nFLH:Chl
834 ratios and SSH values were evident for class 1 in both the MPA and Wiggert-S
835 domains, as were lower SST values for class 5. Class 5 chl peaks in the MPA
836 were not synchronised (or time-lagged, judging by visual inspection) with
837 those in the Wiggert-S domain, suggesting small-scale, rather than basin-
838 wide processes are being captured, despite the use of composited Level 3
839 data. Significant trends are not shown on Figures 15 and S4, for clarity, but
840 are summarised in Figure 16a. Chl, nFLH and nFLH:Chl decreased over the
841 study period in most classes and most sub-domains (when appraised using
842 both absolute values and with anomaly time-series). For chl, the rate of
843 change was between -7.5×10^{-6} and $-1.2 \times 10^{-4} \text{ mgm}^{-3}\text{a}^{-1}$ (up to $0.025\%\text{a}^{-1}$

844 in the MPA and $0.055\%a^{-1}$ in z1000, compared with decreases of $0.7\%a^{-1}$
845 reported by Gregg et al., 2017, for the Indian Ocean).

846 Of the abiotic variables, significant trends in all five N5 classes were only
847 found for the MPA, z1000 and Wiggert-N domains, with weaker positive
848 trends in some areas (domain-integrated) and classes for SST, ∇ SSH, u and
849 v, and a positive trend in $\nabla \times \tau$ in Wiggert-S only. Full temporal trend
850 results, including 95% confidence intervals on the rates of change, are given
851 in Table S1, sheet ‘Trends Correlations’.

852 [FIGURE 15 ROUGHLY HERE: SINGLE-COLUMN; PLEASE QUOTE
853 FOR COLOUR PRINTING]

854 [FIGURE 16 ROUGHLY HERE: 1.5 OR SINGLE-COLUMN; PLEASE
855 QUOTE FOR COLOUR PRINTING]

856 No significant correlations were found between the MJO and remotely
857 sensed variables, consistent with the brief residence of MJO events over the
858 TIO as well as the short time-scales of response of chl to MJO events which
859 precludes detection of MJO effects in this Level 3 data analysis (order of
860 days to weeks; Vialard et al., 2009; Jin et al., 2012; Wheeler & Hendon,
861 2004). Surface cooling and enhanced surface primary production have been
862 documented in response to the MJO (Vialard et al., 2009; Resplandy et al.,
863 2009, Supplementary material Figure S5), making the use of higher temporal
864 resolution data desirable where coverage allows. The monthly-averaged MJO
865 index was not correlated with the IOD or SOI indices over the study period.
866 However, a weak, negative correlation was found between the IOD and SOI
867 ($SOI = -0.39 IOD + 0.27$; $n=202$, $p = 0.084$), in contrast to the decoupling of
868 these cycles found using EOF analysis by Saji et al. (1999). Fragmentation

869 metrics for N5 class 3 were related to the SOI (negative relationship for patch
 870 density; positive relationship for patch area) and IOD (positive relationship
 871 for patch density only). No relationships between fragmentation metrics
 872 and any climate index were found for the Wiggert-S domain, whereas for
 873 Wiggert-N, patch density increased with increasing IOD for classes 1 and
 874 3, and decreased with increasing IOD for classes 4 and 5. Similarly, patch
 875 area increased with increasing IOD for classes 1 and 2 and decreased for
 876 classes 3 and 4. Fewer significant relationships were found with the SOI, but
 877 they mirrored the IOD relationships, consistent with a negative relationship
 878 between the two climate indices. These results suggest that within the MPA,
 879 if the frequency of Nino events increases as predicted, incurring more negative
 880 SOI conditions, the higher chl N5 classes 3 and 4 will yield to larger and more
 881 closely spaced patches of lower N5 class 1. If positive IOD events increase
 882 in frequency, fewer, smaller N5 class 4 and 5 patches separated by greater
 883 distances are predicted. The observed North-South diversity in epipelagic
 884 conditions and temporal trends may contribute to resilience of reefs and
 885 mobile species in the MPA. However, only surface effects are characterised
 886 and, in this study, processes lasting days to weeks, such as MJO events and
 887 cyclones that have an impact on vertical mixing (Jin et al., 2012; Webster et
 888 al., 2005), may be averaged out.

889 Positive IOD events are associated with elevated SST in the western TIO
 890 as the easterly wind driving the SEC converges further west (Saji et al.,
 891 1999). Domain-specific correlations between IOD and SST (Table 16) were
 892 in agreement, with positive correlations in all but the SCTR-C (eastern-most)
 893 domain. Positive relationships were also found for SSH, τ_N and nFLH:chl,

whilst negative relationships were found with Chl, FLH, R555:488, EKE and τ_E . Mostly positive, class-specific relationships only were found for ∇ SST, suggesting stronger gradients at the sub-domain spatial scale which can not be explained by the regressions undertaken here as the spatially coarse altimetry products used to calculate ∇ SSH and EKE, which might indicate small-scale processes, tended to decrease with increasing IOD. The SST product (4 x 4 km resolution) is more sensitive to mesoscale and perhaps submesoscale processes.

Negative impacts of Nino and (positive) IOD conditions on biotic remotely sensed variables is consistent with surface warming and deepening of the mixed layer in the western TIO, corresponding to down-mixing of phytoplankton within a strong, westward SEC current extending several hundreds of metres below the surface (Vialard et al., 2009). The response of elevated nFLH:chl ratios (Figure 15) under such a deeply-mixed layer could indicate nutrient stress (e.g. Fe, NO₃) or possibly a thin, surface freshening related to precipitation that is isolating a surface, light-stressed population (Behrenfeld et al., 2009; Chinni et al., 2019; George et al., 2013). The satellite data used for this study can not distinguish between these possibilities and are further limited by their short time-span (Dilmahamod et al., 2016; Landschützer et al., 2019), coarse spatial and temporal resolution (Hosegood et al., 2019; Vialard et al., 2009) and lack of information about depth variability. *In situ* data are needed to interpret many of the possible biophysical interactions in the MPA domain in terms of management application. For example, deepening of the mixed layer in the SCTR has been found to be associated with deepening of the DCM with an increase in chl owing to nutrient entrainment

919 (George et al., 2013); or with no net impact on water-column productivity
920 owing to redistribution of light (Resplandy et al., 2009) or with a decrease in
921 chl owing to reduced nutrient entrainment (Ma et al., 2014). Mesopelagic bio-
922 geographies suggest a strong positive relationship between satellite-derived
923 primary production and zooplankton scattering layers (Proud et al., 2017),
924 which is consistent with two of those scenarios. Depending on the response
925 of grazers, a deepened DCM may have a protective effect on pelagic fish
926 that are forced to forage further from the surface (Vialard et al., 2009) but
927 a negative effect for surface-foraging seabirds, so that any future increase in
928 the frequency of these events may have unexpected ramifications at higher
929 trophic levels and for the Chagos MPA reefs (Graham et al., 2018; Fox et al.,
930 2019).

931 4. Conclusions

932 Pure, optical classifications of unnormalised satellite remotely sensed re-
933 flectance data have been shown here to constrain physical variables that
934 shape the epipelagic growth environment, making them a potentially useful
935 source of management-relevant information at low- to medium latency. Con-
936 straint of most remotely sensed variables was best when used within spatial
937 sub-domains such as the MPA area, suggesting their use within a seascape
938 heirarchy. Potential applications discussed here include monitoring ecosys-
939 tem services, including CO₂ uptake, and resource distribution, but in all cases
940 *in situ* validation data are needed to elucidate optical biome composition and
941 function.

942 Optical classification provides a snapshot at monthly (or better) time-

scales of spatial variability of epipelagic resources that are amenable to habitat fragmentation analysis, which suggested a change in the spacing and size of richer surface food resources in response to Nino and IOD events in this study. However, ecological interpretation of the trends and interactions between remotely sensed variables requires knowledge of higher trophic level responses. Inclusion of fragmentation metrics in species distribution models might help to address this, where target species observations are sparse. Although ocean colour remote sensing is among the most finely resolved satellite products in space and time, it is limited by cloud cover and does not capture the full range of dynamical interactions that are relevant to habitat structure and use. Increased spatiotemporal coverage is needed and may be provided by combining information from all available sensors (though this is problematic) and by increased *in situ* monitoring using moorings and perhaps unmanned devices. The improved spectral resolution of NASAs forthcoming PACES mission may provide better discrimination of microbial community composition.

The suggested applications of optical classifications are globally applicable, but the need for more *in situ* data is not restricted to the tropical Indian Ocean. Data requirements include repeat vertical profiles (reflectance, temperature, salinity, vertical mixing, nutrient concentrations and carbon cycle parameters) and spatial fields of surface bio-optical and oceanographic conditions that resolve sub-pixel variability, so that appropriate spatial scales can be identified for a given question. This scope of fieldwork has been attempted in a few international, inter-disciplinary projects and it is to be hoped that more will take shape under the biogeosciences programme (?).

968 **Acknowledgements**

969 I would like to thank Gillian Glegg, Ken Kingston and two anonymous
970 reviewers for helpful comments on the manuscript.

971 **References**

- 972 Aguiar-Gonzalez, B., L. Ponsoni, H. Ridderinkhof, H. M. van Aken, W.P.M.
973 de Ruijter, L.R.M. Maas, 2016. Seasonal variation of the South Indian trop-
974 ical gyre. *Deep-Sea Research I*, 110, 123-140, doi:10.1016/j.dsr.2016.02.004.
- 975 Alvain, S., C. Moulin, Y. Dandanneau, F.M. Bréon, 2005. Remote sensing
976 of phytoplankton groups in case 1 waters from global SeaWiFS imagery.
977 *Deep-Sea Research Part I* 52, 1989-2004, doi:10.1016/j.dsr.2005.06.015.
- 978 Alvain, S., C. Moulin, Y. Dandanneau, H. Loisel, 2008. Seasonal distri-
979 bution and succession of dominant phytoplankton groups in the global
980 ocean: A satellite view. *Global Biogeochemical Cycles* 22, GB3001,
981 doi:10.1029/2007GB003154.
- 982 Amante, C. and B.W. Eakins, 2009. ETOPO1 1 arc-minute global relief
983 model: Procedures, data sources and analysis. NOAA Technical Memo-
984 randum NESDIS NGDC-24. National Geophysical Data Centre, NOAA,
985 doi:10.7289/V5C8276M.
- 986 Anderson, M.J., 2001. A new method for non-parametric multivariate
987 analysis of variance. *Austral Ecology* 26(1), 32-46, doi:10.1111/j.1442-
988 9993.2001.01070.pp.x.

- 989 Anderson, T.R., 2005. Plankton functional type modelling: running be-
990 fore we can walk? *Journal of Plankton Research* 27(11), 1073-1081,
991 doi:10.1093/plankt/fbi076.
- 992 Anderson, R. C., 2014. Cetaceans and Tuna Fisheries in the Western and
993 Central Indian Ocean. IPNLF Technical Report 2, International Pole and
994 Line Foundation, London, 133 pp.
- 995 Anderson, M.J., 2017. Permutational Multivariate Analysis of Variance
996 (PERMANOVA). *Wiley StatsRef: Statistics Reference Online*, 2014-17,
997 doi:10.1002/9781118445112.stats07841.
- 998 Ateweberhan, M., T.R. McClanahan, J. Maina, C. Sheppard, 2018. Thermal
999 energy and stress properties as the main drivers of regional distribution
1000 of coral species richness in the Indian Ocean. *Journal of Biogeography* 45,
1001 1355-1366, doi:10.1111/jbi.13224.
- 1002 Back, L.E., C.S. Bretherton, 2009. On the relationship between SST gra-
1003 dients, boundary layer winds and convergence over the tropical oceans.
1004 *Journal of Climate* 22, 4182-4196, doi:10.1175/2009JCLI2392.1.
- 1005 Behrenfeld, M.J., T.K. Westberry, E.S. Boss, R.T. O'Malley, D.A. Siegel et
1006 al., 2009. Satellite-detected fluorescence reveals global physiology of ocean
1007 phytoplankton. *Biogeosciences* 6, 779-794, doi:10.5194/bg-6-779-2009.
- 1008 Bezdek, J.C., 1981. *Pattern Recognition with Fuzzy Objective Function Al-*
1009 *gorithms*. Plenum Press, NY, doi:10.1007/978-1-4757-0450-1.
- 1010 Blanco, G.S., J.P. Pisoni, F. Quintana, 2015. Characterization of the
1011 seascape used by juvenile and wintering adult Southern Giant Petrels from

1012 Patagonia Argentina. *Estuarine, Coastal and Shelf Science* 153, 135–144,
1013 doi:10.1016/j.ecss.2014.12.007.

1014 Boesch, H., L. Brown, R. Castano, M. Christi, B. Connor et al., 2019. Orbit-
1015 ing Carbon Observatory (OCO)-2 Level 2 Full Physics Algorithm Theoret-
1016 ical Basis Document, version 2.0, Rev. 3. National Aeronautics and Space
1017 Administration, Jet Propulsion Laboratory, OCO D-55207, 92pp.

1018 Brewin, R.J.W., S. Sathyendranath, T. Hirata, S.J. Lavender, R. Barciela,
1019 N.J. Hardman-Mountford, 2010. A three-component model of phytoplank-
1020 ton size class for the Atlantic Ocean. *Ecological Modelling* 221, 1472-1483,
1021 doi:10.1016/j.ecolmodel.2010.02.014.

1022 Brewin, R.J.W., E. Devred, S. Sathyendranath, S.J. Lavender,
1023 N.J. Hardman-Mountford, 2011. Model of phytoplankton absorp-
1024 tion based on three size classes. *Applied Optics* 50(2), 4535-4549,
1025 doi:10.1364/AO.50.004535.

1026 Brewin, R.J., N.J. Hardman-Mountford, S.J. Lavender, D.E. Raitsos, T. Hi-
1027 rata, J. Uitz, E. Devred, A. Bricaud, A. Ciotti, B. Gentili, 2011. An inter-
1028 comparison of bio-optical techniques for detecting dominant phytoplankton
1029 size-class from satellite remote sensing. *Remote Sensing of Environment*
1030 115(2), 325-339, doi:10.1016/j.rse.2010.09.004.

1031 Brewin, R.J.W., T. Hirata, N.J. Hardman-Mountford, S. Lavender, S.
1032 Sathyendranath, R. Barlow, 2012. The influence of the Indian Ocean
1033 Dipole on interannual variations in phytoplankton size structure as re-

1034 vealed by Earth Observation. Deep-Sea Research Part II 77-80, 117-127,
1035 doi:10.1016/j.dsr2.2012.04.009.

1036 Brown, C., Y. Huot, J. Werdell, B. Gentili, H. Claustre, 2008. The origin
1037 and global distribution of second order variability in satellite ocean color
1038 and its potential applications to algorithm development. Remote Sensing
1039 of Environment 112(12), 4186-4203, doi:10.1016/j.rse.2008.06.008.

1040 Cai, W., S. Agus, G. Wang, E. Weller, L. Wu et al., 2014. Increased frequency
1041 of extreme Indian Ocean Dipole events due to greenhouse warming. Nature
1042 510, 254-258, doi:10.1038/nature13327.

1043 Caldow, C., M.E. Monaco, S.J. Pittman, M.S. Kendall, T.L. Goedeke, C.
1044 Menza, B.P. Kinlan, B.M. Costa, 2015. Biogeographic assessments: A
1045 framework for information synthesis in marine spatial planning. Marine
1046 Policy 51, 423-432, doi:10.1016/j.marpol.2014.07.023.

1047 Catry, T., J.A. Ramos, S. Jaquemet, L. Faulquier, M. Berlincourt, A. Hausel-
1048 mann, P. Pinet, M. Le Corre, 2009. Comparative foraging ecology of a trop-
1049 ical seabird community of the Seychelles, western Indian Ocean. Marine
1050 Ecology Progress Series 374, 259 – 272, doi:10.3354/meps07713.

1051 Catry, T., J.A. Ramos, M. Le Corre, R.A. Phillips, 2009. Movements, at-sea
1052 distribution and behaviour of a tropical pelagic seabird: the wedge-tailed
1053 shearwater in the western Indian Ocean. Marine Ecology Progress Series
1054 391, 231 – 242, doi:10.3354/meps07717.

1055 Chinni, V., S. Kumar Singh, R. Bhushan, V.V.S.S. Sarma, 2019. Spa-
1056 tial variability in dissolved iron concentrations in the marginal and

1057 open waters of the Indian Ocean. *Marine Chemistry* 208, 11-28,
1058 doi:10.1016/j.marchem.2018.11.007.

1059 Convention on Biological Diversity 10, Decision X/2, 2010. Strategic plan for
1060 biodiversity 2011-2020 (Target 11, Strategic goal C).

1061 Copernicus Marine Environment Monitoring Service (CMEMS),
1062 2020. Quality information document for sea level TAC DU-
1063 ACS products. CMEMS-SL-QUID-008-032-062, 8th April, 2020,
1064 [https://resources.marine.copernicus.eu/documents/QUID/CMEMS-](https://resources.marine.copernicus.eu/documents/QUID/CMEMS-SL-QUID-008-032-062.pdf)
1065 [SL-QUID-008-032-062.pdf](https://resources.marine.copernicus.eu/documents/QUID/CMEMS-SL-QUID-008-032-062.pdf).

1066 Coelho, R., P. Infante, M.N. Santos, 2013. Application of generalized linear
1067 models and generalised estimation equations to model at-haulback mor-
1068 tality of blue sharks captured in a pelagic longline fisher in the Atlantic
1069 Ocean. *Fisheries Research* 145, 66–75, doi:10.1016/j.fishres.2013.02.010.

1070 Costello, M.J., P. Tsai, P. Shan Wong, A. Kwok Lun Cheung, Z. Basher,
1071 C. Chaudhary, 2017. Marine biogeographic realms and species endemism.
1072 *Nature Communications* 8:1057, doi:10.1038/s41467-017-01121-2.

1073 Currie, J.C., M. Lengaigne, J. Vialard, D.M. Kaplan, O. Aumont et al., 2013.
1074 Indian Ocean Dipole and El Nino/Southern Oscillation impacts on regional
1075 chlorophyll anomalies in the Indian Ocean. *Biogeosciences* 10, 6677-6698,
1076 doi:10.5194/bg-10-6677-2013.

1077 Danckwerts, D.K., C.D. McQuaid, A. Jaeger, G.K. McGregor, R. Dwight, M.
1078 Le Corre, S. Jaquemet, 2014. Biomass consumption by breeding seabirds in

1079 the western Indian Ocean: Indirect interactions with fisheries and implica-
1080 tions for management. *ICES Journal of Marine Science* 71(9), 2589-2598,
1081 doi:10.1093/icesjms/fsu093.

1082 Defoin-Platel, M., M. Chami, 2007. How ambiguous is the inverse problem
1083 of ocean color in coastal waters? *Journal of Geophysical Research: Oceans*
1084 112(C3), C03004, doi:10.1029/2006JC003847.

1085 Devred, E., S. Sathyendranath, V. Stuart, T. Platt, 2011. A three com-
1086 ponent classification of phytoplankton absorption spectra: Applications
1087 to ocean colour data. *Remote Sensing of Environment* 115(9), 2255-2266,
1088 doi:10.1016/j.rse.2011.04.025.

1089 Dierssen, H.M., 2010. Perspectives on empirical approaches for ocean
1090 color remote sensing of chlorophyll in a changing climate. *Proceed-*
1091 *ings of the National Academy of Sciences* 107(40), 17073-17078,
1092 doi:10.1073/pnas.0913800107.

1093 Dilmahamod, A.F., J.C. Hermes, C.J.C. Reason, 2016. Chlorophyll-a vari-
1094 ability in the Seychelles-Chagos Thermocline Ridge: Analysis of a
1095 coupled biophysical model. *Journal of Marine Systems* 154, 220-232,
1096 doi:10.1016/j.jmarsys.2015.10.011.

1097 Drushka, K., J. Sprintall, S.T. Gille, S. Wijffels, 2012. In situ observations of
1098 Madden-Julian Oscillation mixed layer dynamics in the Indian and western
1099 Pacific Oceans. *Journal of Climate* 25, 2306-2328, doi:10.1175/JCLI-D-11-
1100 00203.1.

- 1101 Duarte, C.M., A. Regaudie-de-Gioux, J.M. Arrieta, A. Delgado-Huertas, S.
 1102 Agustí, 2013. The oligotrophic ocean is hetertrophic. *Annual Review of*
 1103 *Marine Science* 5, 551-569, doi:10.1146/annurev-marine-121211-172337.
- 1104 Erickson, Z. K., Frankenberg, C., Thompson, D. R., Thompson, A. F., Gier-
 1105 ach, M., 2019. Remote sensing of chlorophyll fluorescence in the ocean using
 1106 imaging spectrometry: Toward a vertical profile of fluorescence. *Geophys-*
 1107 *ical Research Letters* 46, 1571–1579, doi:10.1029/2018GL081273.
- 1108 Esteban, N., R.K.F. Unsworth, J.B.Q. Gurlay, G.C. Hays, 2018. The discov-
 1109 ery of deep-water seagrass meadows in a pristine Indian Ocean wilderness
 1110 revealed by tracking green turtles. *Marine Pollution Bulletin* 134, 99-105,
 1111 doi:10.1016/j.marpolbul.2018.03.018.
- 1112 ETOPO1 Global Relief Model, www.ngdc.noaa.gov/mgg/global
 1113 doi:10.7289/V5C8276M.
- 1114 EUMETSAT/OSI SAF. 2018. MetOp-A ASCAT Level 2 25-
 1115 km Ocean Surface Wind Vector Climate Data Record. Ver.
 1116 1.0. PO.DAAC, CA, USA. Dataset accessed 2019-03 at
 1117 https://doi.org/10.15770/EUM_SAF_OSI_0006.
- 1118 Fay, A.R., G.A. McKinley, 2014. Global open-ocean biomes: mean and tem-
 1119 poral variability. *Earth System Science Data* 6, 273-284, doi:10.5194/essd-
 1120 6-273-2014.
- 1121 Fernandes, V., V. Rodrigues, N. Ramaiah, J.T. Paul, 2008. Relevance of
 1122 bacterioplankton abundance and production in the oligotrophic equatorial

- 1123 Indian Ocean. Aquatic Ecology 42(4), 511-519, doi:10.1007/s10452-007-
1124 9142-y.
- 1125 Follows, M.J., S. Dutkiewicz, 2011. Modeling diverse communities of
1126 marine microbes. Annual Review of Marine Science 3, 427-451,
1127 doi:10.1146/annurev-marine-120709-142848.
- 1128 Fox, M.D., G.J. Williams, M.D. Johnson, V.Z. Radice, B.J. Zgliczyn-
1129 ski et al., 2019. Gradients in primary production predict trophic strate-
1130 gies of mixotrophic corals across spatial scales. Current Biology (2018),
1131 doi:10.1016/j.cub.2018.08.057.
- 1132 Franz, B.A., S.W. Bailey, P.J. Werdell, C.R. McClain, 2007. Sensor-
1133 independent approach to the vicarious calibration of satellite
1134 ocean color radiometry. Applied Optics 46(22), 5068 - 5082,
1135 doi:10.1364/AO.46.005068.
- 1136 Garcia, H.E., T. Boyer, O.K. Baranova, R.A. Locarnini, A.V. Mis-
1137 honov et al., 2018. World Ocean Atlas 2018 (pre-release). Produce
1138 Documentation. A. Mishonov, Technical Editor. Available online at
1139 <https://www.nodc.noaa.gov/OC5/woa18>.
- 1140 George, J.V., M. Nuncio, R. Chacko, N. Anilkumar, S.B. Noronha et al.,
1141 2013. Role of physical processes in chlorophyll distribution in the west-
1142 ern tropical Indian Ocean. Journal of Marine Systems 113-114, 1-12,
1143 doi:10.1016/j.jmarsys.2012.12.001.
- 1144 George, J.V., M. Nuncio, N. Anilkumar, R. Chacko, D. Rajashekhar, 2018.

1145 Seasonal surface chlorophyll-a variability in the Seychelles-Chagos Ther-
1146 moclone Ridge. *Current Science* 114(4), 868-878.

1147 Gorelick, N., M. Hancher, M. Dixon, S. Ilyushchenko, D. Thau, R.
1148 Moore, 2017. Google Earth Engine: Planetary-scale geospatial anal-
1149 ysis for everyone. *Remote Sensing of Environment* 202, 18 - 27,
1150 doi:10.1016/j.rse.2017.06.031.

1151 Graham, N.A.J., S.K. Wilson, P.Carr, A.S. Hoey, S. Jennings, M.A. Mac-
1152 Neil, 2018. Seabirds enhance coral reef productivity and functioning in
1153 the absence of invasive rats. *Nature* 559, 250-253, doi:10.1038/s41586-018-
1154 0202-03.

1155 Gravestock, P., C. Sheppard, 2015. Valuing the ecosystem services of the
1156 Chagos: a review of challenges and estimates. *Marine Ecology Progress*
1157 *Series* 530, 255-270, doi:10.3354/meps11235.

1158 Gregg, W.W., C.S. Rousseaux, B.A. Franz, 2017. Global trends in ocean
1159 phytoplankton: a new assessment using revised ocean colour data. *Remote*
1160 *Sensing Letters* 8(12), 1102-1111, doi:10.1080/2150704X.2017.1354263.

1161 Gruber, N., D. Clement, B.R. Carter, R.A. Feely, S. van Heuven et al.,
1162 2019. The oceanic sink for anthropogenic CO₂ from 1994 to 2007. *Science*
1163 363:6432, 1193-1199, doi:10.1126/science.aau5153.

1164 Guan, B., T. Lee, D.J. Halkides, D.E. Waliser, 2014. Aquarius surface salinity
1165 and the Madden-Julian Oscillation: The role of salinity in surface layer
1166 density and potential energy. *Geophysical Research Letters* 41, 2858-2869,
1167 doi:10.1002/2014GL059704.

- 1168 Halkides, D.J., T. Lee, 2011. Mechanisms controlling seasonal mixed
1169 layer temperature and salinity in the Southwestern Tropical In-
1170 dian Ocean. *Dynamics of Atmospheres and Oceans* 51, 77-93,
1171 doi:10.1016/j.dynatmoce.2011.03.002.
- 1172 Hamed, K.H., A.R. Rao, 1998. A modified Mann-Kendall trend test for au-
1173 tocorrelated data. *Journal of Hydrology* 204, 182-196.
- 1174 Handley, J.M., E.J. Pearmain, S. Oppel, A.P.B. Carneiro, C. Hazin et al.,
1175 2020. Evaluating the effectiveness of a large multi-use MPA in protecting
1176 key biodiversity areas for marine predators. *Diversity and Distributions*
1177 26, 715 – 729, doi:10.1111/ddi.13041.
- 1178 Heerah, K., M.P. Dias, K. Delord, S. Oppel, C. Barbraud, H. Weimerskirch,
1179 C.A. Bost, 2019. Important areas and conservation sites for a community
1180 of globally threatened marine predators of the Southern Indian Ocean.
1181 *Biological Conservation* 234, 192 – 201, doi:10.1016/j.biocon.2019.03.037.
- 1182 Hendon, H.H., M.L. Salby, 1994. The life cycle of the Madden-Julian Oscil-
1183 lation. *Journal of the Atmospheric Sciences* 51(15), 2225-2237.
- 1184 Hense, I., I. Stemmler, S. Sonntag, 2017. Ideas and perspectives: climate-
1185 relevant marine biologically driven mechanisms in Earth system models.
1186 *Biogeosciences* 14, 403-413, doi:10.5194/bg-14-403-2017.
- 1187 Henson, S.A., C. Beaulieu, R. Lampitt, 2016. Observing climate change
1188 trends in ocean biogeochemistry: when and where. *Global Change Biology*
1189 22, 1561-1571, doi:10.1111/gbc.13152.

- 1190 Hermes, J.C., C.J.C. Reason, 2008. Annual cycle of the South Indian Ocean
1191 (Seychelles-Chagos) thermocline ridge in a regional ocean model. *Journal*
1192 *of Geophysical Research* 113, C04035, doi:10.1029/2007JC004363.
- 1193 Hermes, J.C., C.J.C. Reason, 2009. The sensitivity of the Seychelles-Chagos
1194 thermocline ridge to large-scale wind anomalies. *ICES Journal of Marine*
1195 *Science* 66, 1455-1466, doi:10.1093/icesjms/fsp074.
- 1196 Hermes, J.C., Y. Masumoto, L.M. Beal, M.K. Roxy, J. Vialard et al., 2019. A
1197 sustained ocean observing system in the Indian Ocean for climate related
1198 scientific knowledge and societal needs. *Frontiers in Marine Science* 6:355,
1199 doi:10.3389/fmars.2019.00355.
- 1200 Hobday, A., J. Hartog, T. Timmiss, and J. Fielding 2010. Dynamic spatial
1201 zoning to manage southern bluefin tuna (*Thunnus maccoyii*) capture in a
1202 multispecies longline fishery. *Fisheries Oceanography* 19:243–253.
- 1203 Hoge, F.E., P.E. Lyon, R.N. Swift, J.K. Yungel, M.R. Abbott et al., 2003.
1204 Validation of Terra-MODIS phytoplankton chlorophyll fluorescence line
1205 height. I. Initial airborne lidar results. *Applied Optics* 42(15), 2767-2771,
1206 doi:10.1364/AO.42.002767.
- 1207 Hosegood, P.J., W.A.M. Nimmo-Smith, R. Proud, K. Adams, A.S. Brier-
1208 ley et al., 2019. Internal lee waves and baroclinic bores over a trop-
1209 ical seamount shark 'hot-spot'. *Progress in Oceanography* 172, 34-50,
1210 doi:10.1016/j.pocean.2019.01.010.
- 1211 Houlbrque, F., C. Ferrier-Pags, 2009. Heterotrophy in tropical scleractinian
1212 corals. *Biological Reviews* 84, 1-17, doi:10.1111/j.1469-185X.2008.00058.x.

1213 Hu, C., Z. Lee, B. Franz, 2012. Chlorophyll a algorithms for oligotrophic
 1214 oceans: A novel approach based on three-band reflectance difference. Jour-
 1215 nal of Geophysical Research 117(C1), doi:10.1029/2011jc007395.

1216 Indian Ocean Tuna Commission, catch and effort data by month, species and
 1217 gear, by vessel flag reporting country. Data accessed on 2020-06028 from
 1218 <http://iotc.org/datasets>.

1219 Indian Ocean Tuna Commision Secretariat, 2014.Guidelines for the reporting
 1220 of fisheries statistics to the IOTC. IOTC Secretariat, Mahe, Seychelles,
 1221 January 2014, 70pp.

1222 IPCC, 2013. Climate change 2013: The physical science basis. Contribution
 1223 of working group I to the Fifth Assessment Report of the Intergovernmen-
 1224 tal Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M.
 1225 Tignor, S.K. Allen, J. Boschun, A. Nauels, Y. Xia, V. Bex, P.M. Midgley
 1226 (eds)]. Cambridge University Press, Cambridge, U.K. and New York, NY,
 1227 USA, 1535pp.

1228 International Ocean Color Coordinating Group (IOCCG), 2014. Phytoplank-
 1229 ton Functional Types from Space. Sathyendranath, S. (ed.), Reports of
 1230 the International Ocean-Color Coordinating Group No. 15, IOCCG, Dart-
 1231 mouth, Canada.

1232 International Ocean Color Coordinating Group (IOCCG), 2019. Uncertain-
 1233 ties in Ocean Colour Remote Sensing. Mélin F. (ed.), IOCCG Report Se-
 1234 ries, No. 18, International Ocean Colour Coordinating Group, Dartmouth,
 1235 Canada. <http://dx.doi.org/10.25607/OBP696>

- 1236 Jackson, T., S. Sathyendranath, F. Melin, 2017. An improved optical clas-
1237 sification scheme for the Ocean Colour Essential Climate Variable and its
1238 applications. *Remote Sensing of Environment* 203, 152-161.
- 1239 Jeffries, T.C., M. Ostrowski, R.B. Williams, C. Xie, R.M. Jensen et al., 2015.
1240 Spatially extensive microbial biogeography of the Indian Ocean provides
1241 insights into the unique community structure of a pristine coral atoll. *Sci-
1242 entific Reports* 5:15383, doi:10.1038/srep15383.
- 1243 Jerlov, N.G., 1976. *Marine Optics*. Elsevier Oceanography Series, 14. 2nd
1244 revised and enlarged edition of *Optical Oceanography* (Elsevier Oceanog-
1245 raphy Series, 5), Elsevier Scientific Publishing Company, Amsterdam, Ox-
1246 ford, New York.
- 1247 Jin, D., R. Murtugudde, D.E. Waliser, 2012. Tropical Indo-Pacific Ocean
1248 chlorophyll response to MJO forcing. *Journal of Geophysical Research:*
1249 *Oceans* 117, C11008, doi:10.1029/2012JC008015.
- 1250 Jin, D., D.E. Waliser, C. Jones, R. Murtugudde, 2012b. Modulation of trop-
1251 ical ocean surface chlorophyll by the Madden-Julian Oscillation. *Climate
1252 Dynamics* 41:1321, doi:10.1007/s00382-012-1321-4.
- 1253 Jin, D., R.G. Murtugudde, D.E. Waliser, 2013. Intraseasonal atmospheric
1254 forcing effects on the mean state of ocean surface chlorophyll. *Journal of
1255 Geophysical Research: Oceans* 118, 184-196, doi:10.1029/2012JC008256.
- 1256 Jones, D. L., 2012. *The Fathom Toolbox for Matlab: multivariate eco-
1257 logical and oceanographic data analysis*. College of Marine Science,

1258 University of South Florida, St. Petersburg, Florida, USA. Available from:
1259 https://github.com/stripathy/neuroelectro_matlab_analysis/tree/master/Fathom.

1260 Jung, H.-C., B.-K. Moon, J. Wie, H.-S. Park, J. Lee, Y.-H. Byun, 2019. A
1261 single-column ocean biogeochemistry model (GOTM-TOPAZ) version 1.0.
1262 Geoscientific Model Development 12, 699-722, doi:10.5194/gmd-12/699-
1263 2019.

1264 Kachelriess, D., M. Wegmann, M. Gollock, N. Pettorelli, 2014. The applica-
1265 tion of remote sensing for marine protected area management. Ecological
1266 Indicators 36, 169-177, doi:10.1016/j.ecolind.2013.07.003.

1267 Kavanaugh, M.T., B. Hales, M. Saraceno, Y.H. Spitz, A.E. White, R.M. Lete-
1268 lier, 2014. Hierarchical and dynamic seascapes: A quantitative framework
1269 for scaling pelagic biogeochemistry and ecology. Progress in Oceanography
1270 120, 291-304, doi:10.1016/j.pocean.2013.10.013.

1271 Kavanaugh, M.T., M.J. Oliver, F.P. Chavez, R.M. Letelier, F.E. Muller-
1272 Karger, S.C. Doney, 2016. Seascapes as a new vernacular for pelagic ocean
1273 monitoring, management and conservation. ICES Journal of Marine Sci-
1274 ence 73(7), 1839-1850, doi:10.1093/icesjms/fsw086.

1275 Kilidas, G.N., J. Dias, K.H. Straub, M.C. Wheeler, S.N. Tulich, K. Kikuchi,
1276 M. Weickmann and M.J. Ventrice, 2014. A comparison of OLR and circu-
1277 lation based indices for tracking the MJO. Monthly Weather Review 142,
1278 May 2014, 1697-1715.

1279 Kirk, J.T.O., 1994. Light and photosynthesis in aquatic ecosystems, 2nd
1280 Edition. Cambridge University Press, UK, ISBN: 0 521 45966 4, 509pp.

- 1281 Koldewey, H.J., D. Curnick, S. Harding, L.R. Harrison, M. Gol-
 1282 lock, 2010. Potential benefits to fisheries and biodiversity of
 1283 the Chagos Archipelago/British Indian Ocean Territory as a no-
 1284 take marine reserve. *Marine Pollution Bulletin* 60, 1906-1915,
 1285 doi:10.1016/j.marpolbul.2010.10.002.
- 1286 Krug, L.A., T. Platt, S. Sathyendranath, A.B. Barbosa, 2017. Ocean sur-
 1287 face partitioning strategies using ocean colour remote sensing: A review.
 1288 *Progress in Oceanography* 155, 41-53, doi:10.1016/j.pocean.2017.05.013.
- 1289 Krug, L.A., T. Platt, A.B. Barbosa, 2018. Delineation of ocean surface
 1290 provinces over a complex marine domain (off SW Iberia): An objective
 1291 abiotic-based approach. *Regional Studies in Marine Science* 18, 80-96,
 1292 doi:10.1016/j.rsma.2018.01.003.
- 1293 Lain, L.R., S. Bernard, 2018. The fundamental contribution of phytoplank-
 1294 ton spectral scattering to ocean colour: Implications for satellite detec-
 1295 tion of phytoplankton community structure. *Applied Sciences* 8, 2681,
 1296 doi:10.3390/app8122681.
- 1297 Landschützer, P., N. Gruber, D.C.E. Bakker, 2016. Decadal variations and
 1298 trends of the global ocean carbon sink. *Global Biogeochemical Cycles* 30,
 1299 1396-1417, doi:10.1002/2015GB005359.
- 1300 Landschützer, P., T. Ilyina, N.S. Lovenduski, 2019. Detecting regional modes
 1301 of variability in observation-based surface pCO₂. *Geophysical Research*
 1302 *Letters* 46, 2670-2679, doi:10.1029/2018GL081756.

1303 Le Corre, M., A. Jaeger, P. Pinet, M.A. Kappes, H. Weimerskirch, T. Catry
1304 et al., 2012. Tracking seabirds to identify potential Marine Protected Areas
1305 in the tropical western Indian Ocean. *Biological Conservation* 156, 83-93,
1306 doi:10.1016/j.biocon.2011.11.015.

1307 Le Quere, C., R.J. Andres, T. Boden, T. Conway, R.A. Houghton et al.,
1308 2013. The global carbon budget 1959-2011. *Earth System Science Data* 5,
1309 165-185, doi:10.5194/essd-5-165-2013.

1310 Le Quere, C., R.M. Andrew, P. Friedlingstein, S. Sitch, J. Hauck et al., 2018.
1311 Global Carbon Budget 2018. *Earth System Science Data* 10, 2141-2194,
1312 doi:10.5194/essd-10-2141-2018.

1313 Letessier, T.B., P.J. Bouchet, J.J. Meeuwig, 2017. Sampling mobile oceanic
1314 fishes and sharks: implications for fisheries and conservation planning.
1315 *Biological Reviews* 92, 627-646, doi:10.1111/brv.12246.

1316 Letessier, T.B., D. Mouillot, P.J. Bouchet, L. Vigliola, M.C. Fernandes
1317 et al., 2019. Remote reefs and seamounts are the last refuges for ma-
1318 rine predators across the Indo-Pacific. *PLoS Biology* 17(8):e3000366,
1319 doi:10.1371/journal.pbio.3000366.

1320 Levy, M., D. Shankar, J.-M. Andre, S.S.C. Shenoi, F. Durand, C. Boyer de
1321 Montegut, 2007. Basin-wide seasonal evolution of the Indian Ocean's phy-
1322 toplankton blooms. *Journal of Geophysical Research: Oceans* 112(C12),
1323 C12014, doi:10.1029/2007JC004090.

1324 Lim, H.G., J.S. Kug, 2017. Impact of chlorophyll bias on the tropi-

1325 cal Pacific mean climate in an earth system model. *Climate Dynamics*
1326 doi:10.1007/s00382-017-4036-8.

1327 Longhurst, A.R., 2007. *Ecological Geography of the Sea*, 2nd Edition. Aca-
1328 demic Press, Burlington, MA. ISBN 0-12-455521-7.

1329 Lovenduski, N.S., S.G. Yeager, K. Lindsay, M.C. Long, 2019. Predicting
1330 near-term variability in ocean carbon uptake. *Earth System Dynamics* 10,
1331 45-57, doi:10.5194/esd-10-45-2019.

1332 Liu, L., L. Feng, W. Yu, H. Wang, Y. Liu, S. Sun, 2013. The distribution and
1333 variability of simulated chlorophyll concentration over the tropical Indian
1334 Ocean from five CMIP5 models. *Journal of Ocean University of China*
1335 12(2), 253-259, doi:10.1007/s11802-013-2168-y.

1336 Ma, J., Y. Du, H. Zhan, H. Liu, J. Wang, 2014. Influence of oceanic Rossby
1337 waves on phytoplankton production in the southern tropical Indian Ocean.
1338 *Journal of Marine Systems* 134, 12-19, doi:10.1016/j.marsys.2014.02.003.

1339 Maldonado, M.T., A. Marchetti, M. Saito, A. Tagliabue, 2018. Biogeoscapes:
1340 Ocean Metabolism and Nutrient Cycles on a Changing Planet. Min-
1341 utes of a working group at Woods Hole, November, 2018. Accessed on-
1342 line 2020-06 at [https://drive.google.com/file/d/1EjIE4Fz2edd_hmusaaio-](https://drive.google.com/file/d/1EjIE4Fz2edd_hmusaaio-JHQQaDazGYn/view)
1343 [JHQQaDazGYn/view](https://drive.google.com/file/d/1EjIE4Fz2edd_hmusaaio-JHQQaDazGYn/view).

1344 Maxwell, S.M., E.L. Hazen, R.L. Lewison, D.C. Dunn, H. Bailey et
1345 al., 2015. Dynamic ocean management: Defining and conceptual-
1346 izing real-time management of the ocean. *Marine Policy* 58, 42-50,
1347 doi:10.1016/j.marpol.2015.03.014.

- 1348 McCune, B., J.B. Grace, 2002. Analysis of Ecological Communities. MjM
1349 Software, Gleneden Beach, Oregon, USA. ISBN: 0-9721290-0-6.
- 1350 McPhaden, M.J., G.R. Foltz, 2013. Intraseasonal variations in the surface
1351 layer heat balance of the central equatorial Indian Ocean: The importance
1352 of zonal advection and vertical mixing. *Geophysical Research Letters* 40,
1353 2737-2741, doi:10.1002/grl.50536.
- 1354 Miloslavich, P., N.J. Bax, S.E. Simmons, E. Klein, W. Appeltans et al.,
1355 2018. Essential ocean variables for global sustained observations of biodi-
1356 versity and ecosystem changes. *Global Change Biology* 24(6), 2416-2433,
1357 doi:10.1111/gcb.14108.
- 1358 Mohamed, K.S., K.K. Sajikumar, N. Ragesh, T.V. Jayasankar, K.P.
1359 Said Koya, G. Sasikumar, 2018. Relating abundance of purpleback
1360 flying squid *Sthenoteuthis oualaniensis* (Cephalopoda: Ommastrephi-
1361 dae) to environmental parameters using GIS and GAM in south-
1362 eastern Arabian Sea. *Journal of Natural History* 52(29-30), 1869 – 1882,
1363 doi:10.1080/00222933.2018.1497721.
- 1364 Monolisha, S., T. Platt, S. Sathyendranath, J. Jayasankar, G.
1365 George, T. Jackson, 2018. Optical classification of the coastal wa-
1366 ters of the northern Indian Ocean. *Frontiers in Marine Science* 5:87,
1367 doi:10.3389/fmars.2018.00087.
- 1368 Moore, T.S., J.W. Campbell and H. Feng, 2001. A fuzzy logic classification
1369 scheme for selecting and blending satellite ocean color algorithms. *IEEE*
1370 *Transactions in Geoscience and Remote Sensing* 39(8), 1764-1776.

1371 Moore, T.S., J.W. Campbell, M.D. Dowell, 2009. A class-based approach
1372 to characterizing and mapping the uncertainty of the MODIS ocean
1373 chlorophyll product. *Remote Sensing of Environment* 113(11), 2424-2430,
1374 doi:10.1016/j.rse.2009.07.016.

1375 Nagelkerken, I., S.D. Connell, 2015. Global alteration of ocean ecosys-
1376 tem functioning due to increasing human CO₂ emissions. *Proceed-*
1377 *ings of the National Academy of Sciences* 112(43), 13727 – 13777,
1378 doi:10.1073/pnas.1510561112.

1379 Nel, D.C., J.R.E. Lutjeharms, E.A. Pakhomov, I.J. Ansorge, P.G.
1380 Ryan, N.T.W. Klages, 2001. Exploitation of mesoscale oceanographic
1381 features by grey-headed albatross *Thalassarche chrysostoma* in the
1382 southern Indian Ocean. *Marine Ecology Progress Series* 217, 15-26,
1383 doi:10.3354/meps217015.

1384 OCO-2 Science Team/M. Gunson, A. Eldering, 2016. ACOS
1385 GOSAT/TANSO-FTS Level 2 Full Physics Standard Product V7.3,
1386 Greenbelt, MD, USA, Goddard Earth Sciences Data and Infor-
1387 mation Services Center (GES DISC). Dataset accessed 2019-04 at
1388 https://disc.gsfc.nasa.gov/datacollection/ACOS_L2S_7.3.html.

1389 O'Dell, C.W., A. Eldering, P.O. Wennberg, D. Crisp, M.R. Gunson et
1390 al, 2018. Improved retrievals of carbon dioxide from Orbiting Carbon
1391 Observatory-2 with the version 8 ACOS algorithm. *Atmospheric Measure-*
1392 *ment Techniques* 11, 6539-6576, doi:10.5194/amt-11-6539-2018.

1393 Oliver, M.J., S. Glenn, J.T. Kohut, A.J. Irwin, O.M. Schofield, M.A. Moline,

- 1394 W.P. Bissett, 2004. Bioinformatic approaches for objective detection of
1395 water masses on continental shelves. *Journal of Geophysical Research* 109,
1396 C07S04, doi:10.1029/2003.JC002072.
- 1397 Oliver, J.J., A.J. Irwin, 2008. Objective global ocean biogeographic provinces.
1398 *Geophysical Research Letters* 35, L15601, doi:10.1029/2008GL034238.
- 1399 Oppel, S., M. Bolton, A.P.B. Carneiro, M.P. Dias, J.A. Green et al., 2018.
1400 Spatial scales of marine conservation management for breeding seabirds.
1401 *Marine Policy* 98, 37 – 46, doi:10.1016/j.marpol.2018.08.024.
- 1402 O'Reilly, J.E., S. Maritorena, B.G. Mitchell, D.A. Siegel, K.L. Carder et al.,
1403 1998. Ocean color chlorophyll algorithms for SeaWiFS. *Journal of Geophys-*
1404 *ical Research: Oceans* 103(C11), 24937-24953, doi:10.1029/98JC02160.
- 1405 Park, J.-Y., J.-S. Kug, 2014. Marine biological feedback associated with In-
1406 dian Ocean Dipole in a coupled ocean/biogeochemical model. *Climate Dy-*
1407 *namics* 42(1-2), 329-343, doi:10.1007/s00382-012-1640-5.
- 1408 Pecoraro, C., I. Zudaire, N. Bodin, H. Murua, P. Taconet et al., 2017. Putting
1409 all the pieces together: integrating current knowledge of the biology, ecol-
1410 ogy, fisheries status, stock structure and management of yellowfin tuna
1411 (*Thunnus albacares*). *Reviews in Fish Biology and Fisheries* 27(4), 811-
1412 841, doi:10.1007/s11160-016-9460-z.
- 1413 Pinaud, D., H. Weimerskirch, 2007. At-sea distribution and scale-dependent
1414 foraging behaviour of petrels and albatrosses: a comparative study. *Journal*
1415 *of Animal Ecology* 76, 9-19, doi:10.1111/j.1365-2656.2006.01186.x.

1416 Pinheiro, H.T., J.B. Teixeira, R.B. Francini-Filho, A. Soares-Gomes, C.E.L.
 1417 Ferreira, L.A. Rocha, 2019. Hope and doubt for the world's marine
 1418 ecosystems. *Perspectives in Ecology and Conservation* 17, 19 – 25,
 1419 doi:10.1016/j.pecon.2018.11.001.

1420 Pillar, V.D., 2013. How accurate and powerful are randomization tests in
 1421 multivariate analysis of variance? *Community Ecology* 14(2), 153-163,
 1422 doi:10.1556/ComEc.14.2013.2.5.

1423 Proud, R., M.J. Cox, A.S. Brierley, 2017. Biogeography of the
 1424 global ocean's mesopelagic zone. *Current Biology* 27(1),113-119,
 1425 doi:10.1016/j.cub.2016.11.003.

1426 Racault, M.F., S. Sathyendranath, R.J. Brewin, D.E. Raitsos, T. Jackson,
 1427 T. Platt, 2017. Impact of El Nino variability on oceanic phytoplankton.
 1428 *Frontiers in Marine Science* 4:133, doi:10.3389/fmars.2017.00133.

1429 Resplandy, L., J. Vialard, M. Levy, O. Aumont, Y. Dandonneau, 2009. Sea-
 1430 sonal and intraseasonal biogeochemical variability in the thermocline ridge
 1431 of the southern tropical Indian Ocean. *Journal of Geophysical Research*
 1432 114, C07024, doi:10.1029/2008JC005246.

1433 Reygondeau, G., A. Longhurst, E. Martinez, G. Beaugrand, D. Antoine,
 1434 O. Maury, 2013. Dynamic biogeographical provinces in the global ocean.
 1435 *Global Biogeochemical Cycles* 27(1), 1046-1058, doi:10.1002/gbc.20089.

1436 Reygondeau, G., L. Guidi, G. Beaugrand, S.A. Henson, P. Koubbi et al.,
 1437 2018. Global biogeochemical provinces of the mesopelagic zone. *Journal of*
 1438 *Biogeography* 45, 500-514, doi:10.1111/jbi.13149.

- 1439 Rice, J., K.M. Gjerde, J. Ardron, S. Arico, I. Cresswell et al., 2011. Pol-
1440 icy relevance of biogeographic classification for conservation and manage-
1441 ment of marine biodiversity beyond national jurisdiction, and the GOODS
1442 biogeographic classification. *Ocean & Coastal Management* 54, 110-122,
1443 doi:10.1016/j.ocecoaman.2010.10.010.
- 1444 Roberson, L.A., E. Lagabriele, A.T. Lombard, K. Sink, T. Livingstone et al.,
1445 2017. Pelagic bioregionalisation using open-access data for better planning
1446 of marine protected area networks. *Ocean & Coastal Management* 148,
1447 214-230, doi:10.1016/j.ocecoaman.2017.08.017.
- 1448 Roberts, C.M., B.C. O’Leary, D.J. McCauley, P.M. Cury, C.M. Duarte et
1449 al., 2017. Marine reserves can mitigate and promote adaptation to climate
1450 change. *Proceedings of the National Academy of Sciences* 114(24), 6167-
1451 6175, doi:10.1073/pnas.1701262114.
- 1452 Romanou, A., J. Romanski, W.W. Gregg, 2014. Natural ocean carbon cy-
1453 cle sensitivity to parameterizations of the recycling in a climate model.
1454 *Biogeosciences* 11, 1137-1154, doi:10.5194/bg-11-1137-2014.
- 1455 Ropelewski, C.F. and P.D. Jones, 1987. An extension of the Tahiti-Darwin
1456 Southern Oscillation Index. *Monthly Weather Review* 115, 2161-2165,
1457 doi:10.1175/1520-0493(1987)115;2161:AEOTTS;2.0.CO;2.
- 1458 Ruxton, G.D., G. Beauchamp, 2008. Some suggestions about appropri-
1459 ate use of the Kruskal-Wallis test. *Animal Behaviour* 76, 1083-1087,
1460 doi:10.1016/j.anbehav.2008.04.01 1.

1461 Saji, N.H., B.N. Goswami, P.N. Vinayachandran, T. Yamagata, 1999. A
1462 dipole mode in the tropical Indian Ocean. *Nature* 401, 360-363, 23 Sept.
1463 1999.

1464 Saji, N.H. and T. Yagamaga, 2003. Possible impacts of Indian Ocean Dipole
1465 mode events on global climate. *Climate Research* 25(2), 151-169.

1466 Santoso, A., A. Sen Gupta, M.H. England, 2010. Genesis of Indian Ocean
1467 mixed layer temperature anomalies: A heat budget analysis. *Journal of*
1468 *Climate* 23(20), 5375-5403, doi:10.1175/2010JCLI2072.1.

1469 Sayre, R.G., D.J. Wright, S.P. Breyer, K.A. Butler, K. Van Graafeiland et al.,
1470 2017. A three-dimensional mapping of the ocean based on environmental
1471 data. *Oceanography* 30(1), 90-103, doi:10.5670/oceanog.2017.116.

1472 Scales, K.L., P.I. Miller, C.B. Embling, S.N. Ingram, E. Pirotta, S.C. Votier,
1473 2014. Mesoscale fronts as foraging habitats: composite front mapping re-
1474 veals oceanographic drivers of habitat use for a pelagic seabird. *Journal of*
1475 *the Royal Society Interface* 11, doi:10.1098/rsif.2014.0679.

1476 Schwarz, J.N., 2020. Phytoplankton biovolumes from light microscopy; Cha-
1477 gos Archipelago, Indian Ocean, May 2018. Data submission 2020-06-
1478 30T16:25:05Z, pangaea.de.

1479 Scott, B.E., J. Sharples, O.N. Ross, J. Wang, G.J. Pierce, C.J. Camphuy-
1480 sen, 2010. Foraging plasticity in a pelagic seabird species along a ma-
1481 rine productivity gradient. *Marine Ecology Progress Series* 408, 207 – 226,
1482 doi:10.3354/meps08552.

1483 SeaPAC. 2006. SeaWinds on QuikSCAT Level 3 Daily Gridded Ocean Wind
1484 Vectors (JPL Version 2). Ver. 2. PO.DAAC, CA, USA. Dataset accessed
1485 2019-03 at <https://doi.org/10.5067/QSXXX-L3002>.

1486 Seiki, A., M. Katsumata, T. Horii, T. Hasegawa, K.J. Richards et al., 2013.
1487 Abrupt cooling associated with the oceanic Rossby wave and lateral advec-
1488 tion during CINDY2011. *Journal of Geophysical Research: Oceans* 118,
1489 5523-5535, doi:10.1002/jgrc.20381.

1490 Sen, P.K., 1968. Estimates of the regression coefficient based on Kendall's
1491 tau. *Journal of the Americal Statistical Association* 63(324), 1379-1389.

1492 Sheppard, C.R.C., A. Harris, A.L.S. Sheppard, 2008. Archipelago-wide
1493 coral recovery patterns since 1998 in the Chagos Archipelago, cen-
1494 tral Indian Ocean. *Marine Ecology Progress Series* 362, 109-117,
1495 doi:10.3354/meps07436.

1496 Sheppard, C.R.C., M. Ateweberhan, B.W. Bowen, P. Carr, C.A. Chen et al.,
1497 2012. Reefs and islands of the Chagos Archipelago, Indian Ocean: why it
1498 is the world's largest no-take marine protected area. *Aquatic Conservation*
1499 22(2), 232-261, doi:10.1002/aqc.1248.

1500 Soares, M.A., P.V. Bhaskar, R.K. Naik, D. Dessai, J. George et al., 2015. Lat-
1501 itudinal $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ variations in particulate organic matter (POM)
1502 in surface waters from the Indian ocean sector of Southern Ocean and
1503 the Tropical Indian Ocean in 2012. *Deep-Sea Research* 118, 186-196,
1504 doi:10.1016/j.dsr2.2015.06.009.

- 1505 Solanki, H.U., D. Bhatpuria, P. Chauhan, 2015. Signature analysis of satel-
1506 lite derived SSHa, SST and chlorophyll concentration and their linkage
1507 with marine fishery resources. *Journal of Marine Systems* 150, 12 – 21,
1508 doi:10.1016/j.jmarsys.2015.05.004.
- 1509 Solanki, H.U., D. Bhatpuria, P. Chauhan, 2017. Applications of generalized
1510 additive model (GAM) to satellite-derived variables and fishery data for
1511 prediction of fishery resources distributions in the Arabian Sea. *Geocarto*
1512 *International* 32(1), 30 – 43, doi:10.1080/10106049.2015.1120357.
- 1513 Spalding, M.D., V.N. Agostini, J. Rice, S.M. Grant, 2012. Pelagic
1514 provinces of the world: A biogeographic classification of the world's
1515 surface pelagic waters. *Ocean & Coastal Management* 60, 19-30,
1516 doi:10.1016/j.ocecoaman.2011.12.016.
- 1517 Sreeush, M.G., V. Valsala, S. Pentakota, K. Venkata Siva Rama Prasad,
1518 R. Murtugudde, 2018. Biological production in the Indian Ocean up-
1519 welling zones – Part 1: refined estimation via the use of a variable com-
1520 pensation depth in ocean carbon models. *Biogeosciences* 15, 1895-1918,
1521 doi:10.5194/bg-15-1895-2018.
- 1522 Strutton, P.G., V.J. Coles, R.R. Hood, R.J. Matear, M.J. McPhaden, H.E.
1523 Phillips, 2015. Biogeochemical variability in the central equatorial In-
1524 dian Ocean during the monsoon transition. *Biogeosciences* 12, 2367-2382,
1525 doi:10.5194/bg-12-2367-2015.
- 1526 Sutton, T.T., M.R. Clark, D.C. Dunn, P.N. Halpin, A.D. Rogers et al., 2017.

- 1527 A global biogeographic classification of the mesopelagic zone. *Deep-Sea*
1528 *Research Part I* 126, 85-102, doi:10.1016/j.dsr.2017.05.006.
- 1529 Sutton, A.L., L.E. Beckley, 2017. Species richness, taxonomic distinctness
1530 and environmental influences on Euphausiid zoogeography in the Indian
1531 Ocean. *Diversity* 9, 23, doi:10.3390/d9020023.
- 1532 Subha Anand, S., R. Rengarajan, V.V.S.S. Sarma, A.K. Sudheer, R.
1533 Bhushan, S.K. Singh, 2017. Spatial variability of upper ocean POC export
1534 in the Bay of Bengal and the Indian Ocean determined using particle-
1535 reactive ^{234}Th . *Journal of Geophysical Research – Oceans* 122, 3753-3770,
1536 doi:10.1002/2016JC012639.
- 1537 Talley, L.D., G.L. Pickard, W.J. Emery, J.H. Swift, 2011. Indian Ocean.
1538 Chapter 11 in *Descriptive Physical Oceanography* (6th Ed.), Academic
1539 Press, doi:10.1016/B978-0-7506-4552-2.10033-2.
- 1540 Thorrrington-Smith, M., 1971. West Indian Ocean phytoplankton: a numeri-
1541 cal investigation of phytohydrographic regions and their characteristic phy-
1542 toplankton associations. *Marine Biology* 9, 115-137.
- 1543 Trevail, A.M., J.A. Green, J. Sharples, J.A. Poloton, J.P. Arnould, S.C.
1544 Patrick, 2019. Environmental heterogeneity amplifies behavioural response
1545 to a temporal cycle. *Oikos* 128, 517 – 528, doi:10.1111/oik.05579.
- 1546 Trochta, J.T., C.B. Mouw, T.S. Moore, 2015. Remote sensing of phys-
1547 ical cycles in Lake Superior using a spatio-temporal analysis of op-
1548 tical water typologies. *Remote Sensing of Environment* 171, 149-161,
1549 doi:10.1016/j.rse.2015.10.008.

- 1550 Tweddle, J.F., M. Gubbins, B.E. Scott, 2018. Should phytoplankton be
1551 a key consideration for marine management? *Marine Policy* 97, 1-9,
1552 doi:10.1016/j.marpol.2018.08.026.
- 1553 Uitz, J., H. Claustre, A. Morel, S.B. Hooker, 2006. Vertical distribu-
1554 tion of phytoplankton communities in open ocean: An assessment based
1555 on surface chlorophyll. *Journal of Geophysical Research* 111, C08005,
1556 doi:10.1029/2005JC003207.
- 1557 United Nations Framework Convention on Climate Change, 2015. Adoption
1558 of the Paris Agreement. Proposal by the President. UNFCCC. Con-
1559 ference of the Parties (COP), United Nations, Geneva. Available at:
1560 [https://unfccc.int/process-and-meetings/the-paris-agreement/the-paris-](https://unfccc.int/process-and-meetings/the-paris-agreement/the-paris-agreement)
1561 [agreement](https://unfccc.int/process-and-meetings/the-paris-agreement/the-paris-agreement).
- 1562 UNEP-WCMC, 2016. World database on protected areas user
1563 manual 1.4. UNEP-WCMC: Cambridge, UK. Available at:
1564 http://wcmc.io/WDPA_Manual.
- 1565 Vantrepotte, V., H. Loisel, D. Dessailly, X. Mériaux, 2012. Optical classifi-
1566 cation of contrasted coastal waters. *Remote Sensing of Environment* 123,
1567 306-323, doi:10.1016/j.rse.2012.03.004.
- 1568 Veldhuis, M.J.W., G.W. Kraay, J.D.L. van Bleijswijk, M.A. Baars, 1997.
1569 Seasonal and spatial variability in phytoplankton biomass, productivity
1570 and growth in the northwest Indian Ocean: the southwest and northeast
1571 monsoon, 1992-1993. *Deep-Sea Research I* 44(3), 425-449.

- 1572 Verhoef, A., J. Vogelzang, J. Verspeek, A. Stoffelen, 2018. Long-
1573 term scatterometer wind climate data records. *IEEE Journal of*
1574 *Selected Topics in Applied Earth Observation* 10(5), 2186-2194,
1575 doi:10.1109/JSTARS.2016.2615873.
- 1576 Vialard, J., J.P. Duvel, M.J. McPhaden, P. Bouruet-Aubertot, B. Ward et al.,
1577 2009. CIRENE: Air-sea interactions in the Seychelles-Chagos thermocline
1578 ridge region. *Bulletin of the American Meteorological Society* 90(1), 45-61,
1579 doi:10.1175/2008BAMS2499.1
- 1580 Vichi, M., J. Icarus Allen, S. Masina, N.J. Hardman-Mountford, 2011. The
1581 emergence of ocean biogeochemical provinces: A quantitative assessment
1582 and a diagnostic for model evaluation. *Global Biogeochemical Cycles* 25,
1583 GB2005, doi:10.1029/2010GB003867.
- 1584 Wang, X., F. Guillaume Blanchet, N. Koper, 2014. Measuring habitat frag-
1585 mentation: An evaluation of landscape pattern metrics. *Methods in Ecol-*
1586 *ogy & Evolution* 5, 634-646, doi:10.1111/2041-210X.12198.
- 1587 Wanningkhof, R., G.-H. Park, T. Takahashi, C. Sweeney, R. Feely et al., 2013.
1588 Global ocean carbon uptake: magnitude, variability and trends. *Biogeo-*
1589 *sciences* 10, 1983-2000, doi:10.5194/bg-10-1983-2013.
- 1590 Waugh, S.M., H. Weimerskirch, 2003. Environmental heterogeneity and the
1591 evolution of foraging behaviour in long ranging greater albatrosses. *Oikos*
1592 103, 374-384.
- 1593 Webster, P.J., G.J. Holland, J.A. Curry, H.-R. Chang, 2005. Changes in

1594 tropical cyclone number, duration and intensity in a warming environment.
 1595 Science 309(5742), 1844-1846, doi:10.1126/science.1116448.

1596 Weimerskirch, H., M. Le Corre, Y. Ropert-Coudert, A. Kato, F. Marsac,
 1597 2005. The three-dimensional flight of red-footed boobies: adaptations to
 1598 foraging in a tropical environment? Proceedings of the Royal Society B
 1599 272, 53 – 61, doi:10.1098/rspb.2004.2918.

1600 Weimerskirch, H., 2007. Are seabirds foraging for unpredictable resources?
 1601 Deep-Sea Research II 54, 211 – 223, doi:10.1016/j.dsr2.2006.11.013.

1602 Werdell, P.J., L.I.W. McKinna, E. Boss, S.G. Ackleson, S.E. Craig et al.,
 1603 2018. An overview of approaches and challenges for retrieving marine
 1604 inherent optical properties from ocean color remote sensing. Progress in
 1605 Oceanography 160, 186-212, doi:10.1016/j.pocean.2018.01.001.

1606 Wheeler, M.C., H.H. Hendon, 2004. An all-season real-time multivari-
 1607 ate MJO index: Development of an index for monitoring and pre-
 1608 diction. Monthly Weather Review 132, 1917-132, doi:10.1175/1520-
 1609 0493(2004)132;1917:AARMMI;2.0.CO;2.

1610 Williamson, M.J., E.J. Tebbs, T.P. Dawson, D.M.P. Jacoby 2019. Satellite
 1611 remote sensing in shark and ray ecology, conservation and management.
 1612 Frontiers in Marine Science 6:135, doi:10.3389/fmars.2019.00135.

1613 Wiggert, J.D., R.G. Murtugudde, J.R. Christian, 2006. Annual ecosys-
 1614 tem variability in the tropical Indian Ocean: Results of a coupled bio-
 1615 physical ocean general circulation model. Deep-Sea Research II 53, 644-
 1616 676, doi:10.1016/j.dsr2.2006.01.027.

- 1617 Wilson, C., X. Qiu, 2008. Global distribution of summer chlorophyll blooms
1618 in the oligotrophic gyres. *Progress in Oceanography* 78(2), 107-134,
1619 doi:10.1016/j.pocean.2008.05.002.
- 1620 Windham, M.P., 1982. *IEEE Transactions on Pattern Analysis and Machine*
1621 *Intelligence PAMI-4*(4), 357-363.
- 1622 Xie, X.L. and G. Beni, 1991. A validity measure for fuzzy clustering. *IEEE*
1623 *Transactions on Pattern Analysis and Machine Intelligence* 13(8), 841-847.
- 1624 Xie, S.-P., H. Annamalai, F.A. Schott, J.P. McCreary, 2002. Structure and
1625 mechanisms of south Indian Ocean climate variability. *Journal of Climate*
1626 15, 864-878.
- 1627 Zydelis, R., R.L. Lewison, S.A. Schaffer, J.E. Moore, A.M. Boustany, J.J.
1628 Roberts, M. Sims, D.C. Duncan, B.D. Best, Y. Tremblay et al., 2011.
1629 Dynamic habitat models: using telemetry data to project fisheries bycatch.
1630 *Proceedings of the Royal Society B – Biological Sciences* 278(1722), 3191
1631 – 3200, doi:10.1098/rspb.2011.0330.

5. Figure captions

Figure 1: Schematic representation of key biophysical linkages (not exhaustive). The flow of information begins with sunlight to the left. Physical variables that can be detected using remote sensing, followed by the oceanographic variables derived from them, are shown between the sun and the ocean surface processes. Oceanographic variables of interest that can be derived from remote sensing data are outlined in blue and abbreviations are explained in Table 1. The other variables shown are of interest to conservation, management or climate change applications but are not amenable to remote sensing.

Figure 2: Location of the study domain, adapted from Talley et al. (2011); Aguiar-Gonzalez et al. (2016). SCTR = Seychelles Chagos Thermocline Ridge; MPA = Chagos Marine Protected Area; SECC = South Equatorial Countercurrent

Figure 3: Bathymetry of the study domain. Shaded regions denote the sub-areas related to other published studies. Black line: coast; grey line: 200 m contour; black dashed line: 1000 m contour; thick black line: MPA boundary.

Figure 4: Central spectra for the 10-band classifications from Level 2 (upper) and Level 3 (lower) datasets with two to nine classes, compared with the 6-band JSM classification (shaded).

Figure 5: Mapping of the N5, N8 and JSM classes onto published PFT algorithms. SLC=Synechococcus-like cyanobacteria; xDiat was assigned where

1655 the residual PHYSAT-like spectrum resembled the diatom criteria in Al-
1656 vain et al. (2008) but with higher values; SynPro=spectrum matches SLC or
1657 Prochlorococcus except for one waveband, which fell in the other small-cell
1658 category. Algorithm acronyms are explained in section 3.1.

1659 Figure 6: Comparison of class maps produced with the Level 2 and Level
1660 3 Rrs data, classifications N5, N8 and JSM, in 2003. The Chagos MPA
1661 outline, 0 m and 1000 m isobaths are shown in black and the 2000 m isobath
1662 in grey.

1663 Figure 7: Number of classes held at each pixel over the 17 year study
1664 period, by month, for Level 3 data, N=5. The Chagos MPA outline is shown
1665 in black.

1666 Figure 8: Monthly 17-year climatologies of key remote sensing variables.
1667 The Chagos MPA outline and 0 m contour are shown in black, and the 3000
1668 m isobath in grey.

1669 Figure 9: Variability of remotely sensed parameters in the MPA and z1000
1670 domains: Top row = N5; Middle row = N8; Lower row = JSM; black=MPA;
1671 cyan=whole domain with depth > 1000 m (z1000).

1672 Figure 10: Variability of remotely sensed parameters for each sub-domain
1673 using the N5 classification.

1674 Figure 11: Averaged Mann-Kendall-Sen correlation coefficients between
1675 variables within each class for each domain: a) MPA, b) SCTR-W, c) SCTR-

1676 C, d) Wiggert-N, e) Wiggert-S, f) z1000. Only results with a p-value < 0.01 ,
1677 confirmed using bootstrapping to remove autocorrelation and subsampling
1678 effects, are shown.

1679 Figure 12: OCO-2 lowest altitude pCO_2 distributions within the z1000
1680 domain for the N5 classes applied to MODIS Level 2 data, January 2015.

1681 Figure 13: a) to e) Density of patches of each N5 class (a=class 1, e=class
1682 5); f) to j) Average patch area (f=class 1, j=class 5); k) to o) Average
1683 distance between patches (k=class 1, o=class 5) for the MPA, Wiggert-N
1684 and Wiggert-S domains. Note different y-axis limits are used to show detail.
1685

1686 Figure 14: IOTC fishing catch and catch per unit effort for the Indian
1687 Ocean surface (top row; a, b,c) and longline (lower row; d, e, f) fisheries.

1688 Figure 15: Time-series for each N5 class within the MPA for a) Chl, b)
1689 FLH, c) FLH:Chl, d) SST, e) SSH and f) ∇ SSH.

1690 Figure 16: a) Summary of the temporal trends found within each do-
1691 main ($p < 0.05$), including ranges of the Sen regression coefficient and rates
1692 of change. Trends for specific classes are denoted by class number; trends
1693 for the entire domain are indicated by block colour (red=positive trend;
1694 blue=negative trend). b) Summary of significant ($p < 0.05$) correlations be-
1695 tween remotely sensed variables and the Indian Dipole Mode Index, including
1696 ranges of the Sen regression coefficient. Correlations for specific classes are
1697 denoted by class number; correlations for the entire domain are indicated

1698 by block colour (red=positive trend; blue=negative trend). Full correlation
1699 results, including 95% confidence intervals on the regression slopes, are given
1700 in Table S1.

1701 Appendix A. Appendix 1

Table A.7: Characterisation of variables within each class for the N5, N8 and JSM classification schemes in the whole study site where depth >200 m (total no. pixels 177,145,574)

Variable	Mean $\pm 1\sigma$													
	N5					N8								
Class	1	2	3	4	5	1	2	3	4	5	6	7	8	
N (%)	13	25	30	22	10	10	15	21	21	17	11	5	<1	
chl	0.053	0.086	0.122	0.181	0.456	0.049	0.074	0.097	0.126	0.166	0.242	0.612	1.914	
	± 0.011	± 0.013	± 0.022	± 0.065	± 0.994	± 0.009	± 0.010	± 0.012	± 0.021	± 0.044	± 0.127	± 1.336	± 1.508	
flh	0.040	0.045	0.050	0.064	0.107	0.040	0.044	0.046	0.051	0.061	0.076	0.128	0.248	
	± 0.021	± 0.022	± 0.027	± 0.040	± 0.077	± 0.021	± 0.021	± 0.022	± 0.028	± 0.037	± 0.050	± 0.087	± 0.163	
flh/chl	0.778	0.528	0.409	0.346	0.277	0.822	0.595	0.473	0.399	0.360	0.306	0.264	0.142	
SST	0.433	0.258	0.208	0.197	0.142	0.457	0.294	0.228	0.205	0.203	0.170	0.127	0.094	
	27.62	28.58	28.65	28.02	27.30	27.55	28.22	28.77	28.61	28.10	27.76	27.01	28.67	
SLA	± 1.67	± 1.99	± 1.73	± 1.56	± 1.3	± 1.57	± 2.03	± 1.88	± 1.71	± 1.62	± 1.35	± 1.36	± 1.43	
	0.083	0.060	0.048	0.038	0.026	0.087	0.067	0.055	0.047	0.039	0.035	0.023	0.074	
∇ SSH	0.080	0.076	0.072	0.075	0.090	0.078	0.080	0.073	0.073	0.074	0.079	0.094	0.085	
	-23.1	-5.8	-0.3	-1.7	-1.3	-23.8	-13.3	-1.0	-0.4	-3.1	2.8	-1.3	-22.4	
u	± 62.0	± 52.9	± 49.1	± 55.7	± 72.5	± 62.9	± 56.5	± 49.9	± 49.0	± 54.8	± 59.2	± 80.7	± 67.5	
	-0.059	0.002	0.018	-0.010	0.020	-0.058	-0.031	0.023	0.016	-0.012	0.003	0.029	-0.000	
v	0.146	0.188	0.196	0.211	0.244	0.141	0.177	0.193	0.196	0.207	0.223	0.257	0.171	
	0.012	-0.006	-0.003	0.012	0.021	0.013	-0.000	-0.008	-0.002	0.010	0.016	0.027	-0.028	
EKE	± 0.089	± 0.111	± 0.120	± 0.147	± 0.222	± 0.087	± 0.104	± 0.115	± 0.121	± 0.138	± 0.178	± 0.245	± 0.201	
	0.017	0.024	0.027	0.034	0.056	0.016	0.022	0.026	0.027	0.032	0.042	0.066	0.037	
τ_E	0.024	0.036	0.041	0.057	0.107	0.022	0.032	0.038	0.041	0.052	0.075	0.124	0.059	
	-0.064	-0.033	-0.023	-0.021	-0.004	-0.066	-0.044	-0.026	-0.023	-0.024	-0.013	0.002	-0.012	
τ_N	0.044	0.057	0.057	0.055	0.054	0.041	0.055	0.056	0.057	0.057	0.048	0.058	0.053	
	0.031	0.024	0.023	0.022	0.008	0.031	0.028	0.023	0.024	0.025	0.011	0.013	0.003	
$\nabla \times \tau$	0.027	0.034	0.038	0.048	0.064	0.025	0.033	0.035	0.039	0.046	0.054	0.069	0.046	
	-1.43	-3.66	-3.80	-2.20	-2.14	-0.86	-3.38	-3.86	-3.76	-2.57	-1.77	-2.12	-0.76	
Depth	0.006	0.022	0.101	-0.90	0.070	0.136	0.156	0.179	0.207	0.272	0.350	0.485	-0.893	
	4350	4275	3988	3599	3309	4320	4365	4184	3953	3666	3453	3192	677	
R555:443	± 1005	± 980	± 1047	± 1222	± 1347	± 1002	± 992	± 978	± 1057	± 1197	± 1276	± 1378	± 674	
R555:488	0.006	0.022	0.101	-0.90	0.070	0.136	0.156	0.179	0.207	0.272	0.350	0.485	-0.893	
	± 8.05	± 9.75	± 23.38	± 32.55	± 30.69	± 0.028	± 0.058	± 0.082	± 0.093	± 0.103	± 0.126	± 0.152	± 0.044	
hlhne	-0.352	0.356	0.437	0.262	2.896	0.201	0.218	0.236	0.258	0.309	0.367	0.451	-0.003	
	± 88.63	± 15.80	± 50.55	± 22.06	± 482.27	± 0.095	± 0.054	± 0.077	± 0.089	± 0.098	± 0.115	± 0.127	± 16.181	

Table A.8: Characterisation of variables within each class for the N5, N8 and JSM classification schemes within the MPA (total no. pixels 5,343,309).

Variable	N5					Mean $\pm 1\sigma$					N8				
Class	1	2	3	4	5	1	2	3	4	5	6	7	8		
N (%)	8	37	39	16	1	4	20	32	26	15	3	i1	i1		
chl	0.063	0.088	0.123	0.180	0.253	0.058	0.077	0.099	0.128	0.171	0.227	0.244	0.404		
	$\pm .008$	$\pm .012$	$\pm .021$	$\pm .044$	$\pm .087$	$\pm .007$	$\pm .009$	$\pm .012$	$\pm .019$	$\pm .036$	$\pm .073$	$\pm .099$	$\pm .108$		
flh	0.049	0.049	0.060	0.094	0.117	0.050	0.048	0.050	0.062	0.089	0.114	0.081	0.092		
	$\pm .022$	$\pm .021$	$\pm .027$	$\pm .037$	$\pm .053$	$\pm .024$	$\pm .020$	$\pm .021$	$\pm .028$	$\pm .035$	$\pm .044$	$\pm .049$	$\pm .002$		
flh/chl	0.791	0.560	0.482	0.521	0.460	0.868	0.632	0.510	0.479	0.521	0.503	0.357	0.209		
	$\pm .376$	$\pm .240$	$\pm .197$	$\pm .190$	$\pm .198$	$\pm .427$	$\pm .269$	$\pm .214$	$\pm .195$	$\pm .191$	$\pm .187$	$\pm .205$	$\pm .070$		
SST	29.59	29.41	28.71	27.62	27.43	29.55	29.56	29.22	28.60	27.70	27.35	28.05	28.41		
	$\pm .77$	$\pm .89$	$\pm .99$	± 1.06	± 1.04	$\pm .72$	$\pm .87$	$\pm .89$	$\pm .99$	± 1.08	$\pm .94$	± 1.20	$\pm .41$		
SLA	0.147	0.072	0.027	0.008	0.015	0.163	0.100	0.051	0.023	0.009	0.008	0.025	0.110		
	$\pm .084$	$\pm .074$	$\pm .066$	$\pm .063$	$\pm .042$	$\pm .084$	$\pm .078$	$\pm .070$	$\pm .065$	$\pm .063$	$\pm .057$	$\pm .039$	$\pm .049$		
∇ SSH	24.3	0.83	8.52	2.66	8.35	-29.4	-8.19	6.74	8.55	2.55	5.38	14.04	9.05		
	± 39.4	± 34.3	± 33.1	± 34.6	± 32.6	± 38.7	± 37.1	± 32.2	± 33.5	± 34.9	± 33.2	± 29.3	± 15.5		
u	-0.118	0.018	0.071	0.040	0.048	-0.138	-0.038	0.058	0.071	0.041	0.046	0.048	0.110		
	$\pm .162$	$\pm .190$	$\pm .193$	$\pm .175$	$\pm .175$	$\pm .150$	$\pm .184$	$\pm .192$	$\pm .192$	$\pm .177$	$\pm .169$	$\pm .178$	$\pm .160$		
v	0.002	-0.014	-0.020	-0.010	-0.038	0.003	-0.007	-0.019	-0.020	-0.009	-0.019	-0.074	0.038		
	$\pm .089$	$\pm .090$	$\pm .088$	$\pm .093$	$\pm .111$	$\pm .086$	$\pm .090$	$\pm .090$	$\pm .088$	$\pm .092$	$\pm .102$	$\pm .110$	$\pm .051$		
EKE	0.024	0.023	0.026	0.021	0.024	0.025	0.022	0.025	0.025	0.021	0.021	0.026	0.020		
	$\pm .026$	$\pm .028$	$\pm .033$	$\pm .029$	$\pm .029$	$\pm .026$	$\pm .026$	$\pm .031$	$\pm .033$	$\pm .030$	$\pm .030$	$\pm .027$	$\pm .036$		
τ_E	0.003	-0.007	-0.027	-0.056	-0.047	0.002	-0.001	-0.014	-0.030	-0.055	-0.057	-0.016	-0.069		
	$\pm .034$	$\pm .040$	$\pm .047$	$\pm .050$	$\pm .059$	$\pm .034$	$\pm .037$	$\pm .042$	$\pm .048$	$\pm .049$	$\pm .051$	$\pm .068$	$\pm .0008$		
τ_N	0.000	0.011	0.027	0.047	0.040	-0.001	0.005	0.016	0.029	0.046	0.047	0.020	0.057		
	$\pm .017$	$\pm .025$	$\pm .032$	$\pm .034$	$\pm .039$	$\pm .017$	$\pm .022$	$\pm .027$	$\pm .032$	$\pm .034$	$\pm .034$	$\pm .046$	$\pm .003$		
$\nabla \times \tau$	-7.48	-7.04	-7.92	-7.85	-7.16	-7.86	-6.70	-7.52	-7.99	-7.77	-8.12	-3.01	0.00		
	± 17.7	± 16.4	± 16.5	± 17.3	± 26.6	± 18.5	± 16.5	± 16.4	± 16.4	± 16.6	± 22.0	± 25.4	± 8.2		
Depth	3891	3730	3598	3396	3011	3938	3785	3688	3574	3441	3129	3113	368		
	± 1143	± 1083	± 1173	± 1368	± 1433	± 1149	± 1102	± 1087	± 1193	± 1343	± 1446	± 1471	± 140		
R555:443	-0.113	0.048	0.216	0.250	0.139	0.117	0.147	0.173	0.203	0.260	0.300	0.620	6.934		
	± 8.70	± 9.88	± 21.98	± 28.20	± 22.04	$\pm .028$	$\pm .048$	$\pm .080$	$\pm .099$	$\pm .110$	$\pm .084$	$\pm .123$	± 17.700		
R555:488	-5.161	0.477	0.825	0.179	15.117	0.179	0.207	0.229	0.252	0.295	0.325	0.559	-12.526		
	± 136.2	± 16.07	± 51.55	± 21.12	± 617.4	$\pm .538$	$\pm .046$	$\pm .076$	$\pm .094$	$\pm .105$	$\pm .075$	$\pm .084$	± 40.68

Figure 1: Schematic representation of key biophysical linkages (not exhaustive). The flow of information begins with sunlight to the left. Physical variables that can be detected using remote sensing, followed by the oceanographic variables derived from them, are shown between the sun and the ocean surface processes. Oceanographic variables of interest that can be derived from remote sensing data are outlined in blue and abbreviations are explained in Table 1. The other variables shown are of interest to conservation, management or climate change applications but are not amenable to remote sensing.

Figure 2: Location of the study domain, adapted from Talley et al. (2011); Aguiar-Gonzalez et al. (2016). SCTR = Seychelles Chagos Thermocline Ridge; MPA = Chagos Marine Protected Area; SECC = South Equatorial Countercurrent

Figure 3: Bathymetry of the study domain. Shaded regions denote the sub-areas related to other published studies. Black line: coast; grey line: 200 m contour; black dashed line: 1000 m contour; thick black line: MPA boundary.

Figure 4: Central spectra for the 10-band classifications from Level 2 (upper) and Level 3 (lower) datasets with two to nine classes, compared with the 6-band JSM classification (shaded).

Figure 5: Mapping of the N5, N8 and JSM classes onto published PFT algorithms. SLC=Synechococcus-like cyanobacteria; xDiat was assigned where the residual PHYSAT-like spectrum resembled the diatom criteria in Alvain et al. (2008) but with higher values; SynPro=spectrum matches SLC or Prochlorococcus except for one waveband, which fell in the other small-cell category. Algorithm acronyms are explained in section 3.1.

Figure 7: Number of classes held at each pixel over the 17 year study period, by month, for Level 3 data, $N=5$. The Chagos MPA outline is shown in black.

Figure 8: Monthly 17-year climatologies of key remote sensing variables. The Chagos MPA outline and 0 m contour are shown in black, and the 3000 m isobath in grey.

Figure 9: Variability of remotely sensed parameters in the MPA and z1000 domains: Top row = N5; Middle row = N8; Lower row = JSM; black=MPA; cyan=whole domain with depth > 1000 m (z1000).

Figure 10: Variability of remotely sensed parameters for each sub-domain using the N5 classification.

Figure 11: Averaged Mann-Kendall-Sen correlation coefficients between variables within each class for each domain: a) MPA, b) SCTR-W, c) SCTR-C, d) Wiggert-N, e) Wiggert-S, f) z1000. Only results with a p-value < 0.01, confirmed using bootstrapping to remove autocorrelation and subsampling effects, are shown.

Figure 12: OCO-2 lowest altitude pCO₂ distributions within the z1000 domain for the N5 classes applied to MODIS Level 2 data, January 2015.

Figure 13: a) to e) Density of patches of each N5 class (a=class 1, e=class 5); f) to j) Average patch area (f=class 1, j=class 5); k) to o) Average distance between patches (k=class 1, o=class 5) for the MPA, Wiggert-N and Wiggert-S domains. Note different y-axis limits are used to show detail.

Figure 14: IOTC fishing catch and catch per unit effort for the Indian Ocean surface (top row; a, b,c) and longline (lower row; d, e, f) fisheries.

Figure 15: Time-series for each N5 class within the MPA for a) Chl, b) FLH, c) FLH:Chl, d) SST, e) SSH and f) ∇SSH .

Figure 16: a) Summary of the temporal trends found within each domain ($p < 0.05$), including ranges of the Sen regression coefficient and rates of change. Trends for specific classes are denoted by class number; trends for the entire domain are indicated by block colour (red=positive trend; blue=negative trend). b) Summary of significant ($p < 0.05$) correlations between remotely sensed variables and the Indian Dipole Mode Index, including ranges of the Sen regression coefficient. Correlations for specific classes are denoted by class number; correlations for the entire domain are indicated by block colour (red=positive trend; blue=negative trend). Full correlation results, including 95% confidence intervals on the regression slopes, are given in Table S1.