2020-09-24

Choice of molecular assay determines ranavirus detection probability and inferences about prevalence and occurrence

Wynne, FJ

<http://hdl.handle.net/10026.1/16520>

10.3354/dao03518

Diseases of Aquatic Organisms

Inter-Research Science Center

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Choice of molecular assay determines ranavirus detection probability and inferences about incidence and impact

Running page head: Choice of assay determines ranavirus detection

Felicity J Wynne¹, Robert Puschendorf¹, Mairi E Knight¹, Stephen J Price^{2, 3*}

¹School of Biological and Marine Sciences, University of Plymouth, Drake Circus, Plymouth, Devon, PL4 8AA, United Kingdom

²UCL Genetics Institute, Gower Street, London, WC1E 6BT, United Kingdom

³Institute of Zoology, Zoological Society of London, Regent's Park, London, NW1 4RY, United Kingdom

*Corresponding author: Stephen Price s.price@ucl.ac.uk

Abstract

Infectious diseases are emerging at an increasing rate and reliable diagnostic strategies are essential for effective surveillance and research. Ranavirus is an emerging pathogen that can cause morbidity, mortality and population declines in ectothermic hosts. Despite the economic and conservation importance of these viruses, there is no standardized approach to diagnostics, with many molecular assays currently in use. Here, we compared the inter-assay variation and intra-assay precision among two commonly used quantitative and two conventional polymerase chain reaction assays, using laboratory propagated virus and field-collected amphibian and reptile samples. The assays varied in their diagnostic and analytical sensitivity and diagnostic specificity as well as the sequence similarity of oligonucleotides to a set of published ranavirus isolates. Some assays exhibited poor diagnostic sensitivity resulting in a high proportion of false negative results whilst false positive results were also generated. Amplicon sequences from the two conventional PCRs were used to confirm

24 results but were also shown to be useful for inferring viral lineage. Inaccurate results bias
25 estimates of the distribution and impacts of ranaviruses and together these findings emphasize
26 that molecular assays should be chosen carefully in the context of study aims and ideally
27 combined with other lines of evidence when diagnosing infection and disease.

28 **Key words**

29 Diagnostic methods, nested PCR, qPCR, ranavirus, sensitivity, specificity

1. INTRODUCTION

Disease emergence is occurring at an unprecedented rate (Jones et al. 2008) requiring surveillance systems which incorporate sensitive and reliable diagnostic assays that limit bias (Kuiken et al. 2005, Hyatt et al. 2007, Lachish et al. 2012, DiRenzo et al. 2017). Molecular methods – predominantly nucleic acid-based techniques such as conventional polymerase chain reaction (PCR), quantitative (q)PCR and sequencing – do not usually provide explicit information about pathology but are increasingly used for pathogen detection due to their perceived ease of use, reducing costs and rapid turnaround times (Boyle et al. 2004, Hyatt et al. 2007, Black et al. 2017). They also often have broader applications, greater sensitivity and specificity and require less training and skill than traditional techniques, such as histology (Hyatt et al. 2007, Lorch et al. 2010, Skerratt et al. 2011).

Ranaviruses (family *Iridoviridae*) have caused incidents of mortality and morbidity in amphibians, reptiles and bony fish around the world (Gray et al. 2009, Price et al. 2017). The distribution of *Ranavirus* and the drivers of outbreaks are often unclear. Ranaviruses have been suggested to be ubiquitous (Warne et al. 2016, Campbell et al. 2018) but severe disease outbreaks are patchy in distribution and many studies have linked disease emergence to human behaviors including translocations of infectious materials (Price et al. 2014, 2016, Rijks et al. 2016). Ranaviruses have been associated with population declines of a frog species in the United Kingdom and entire amphibian assemblages in Iberia (Teacher et al. 2010, Price et al. 2014, Rosa et al. 2017), underlining them as important pathogens of potentially significant conservation concern. Reliable, sensitive and specific diagnostic methods are essential for detecting low intensity infections, which will aid a more complete understanding of ranavirus disease ecology and epidemiology.

The majority of laboratories use a single molecular assay when screening samples for ranavirus infection, but approaches are not standardized despite the availability of published guidelines from the World Organisation for Animal Health (OIE 2017, Black et al. 2017). Eleven different PCR methods for ranavirus detection were used in the published literature between 2009-2014, but information about the sensitivity or specificity of these assays was usually lacking (Black et al. 2017). Therefore, results may not be reproducible, low intensity infections may be missed, prevalence estimations may be biased, and pathogens may be misidentified. Sequencing of PCR amplicons can be used to confirm the specificity of an assay and the sequences can also be used to investigate a pathogen's taxonomy and genetics (OIE 2017). Nevertheless, even when the primary goal is presence-absence detection, laboratories may favor qPCR and have frequently utilized unpublished methods (Black et al. 2017). Additionally, there appears to be both uncertainty and inconsistency about how to define a positive result from qPCR assays (e.g. Miller et al. 2015, Black et al. 2017), which can reduce the comparability and accuracy of results. This study compared the inter-assay variation and intra-assay precision of four molecular methods used in ranavirus detection. The Mao *et al* (1997) assay is one of the most commonly PCR methods used for ranavirus diagnostics (Black et al. 1997). Meng *et al.*(2013) is a nested method based on Mao *et al* (1997) which might therefore offer greater specificity and sensitivity. The Brunner *et al.*(2004) assay is one of the most commonly used qPCR methods in the literature and Leung *et al.* (2017) is a new qPCR assay which is likely to be attractive to users since it is paired with a host assay enabling precise quantification of viral loads.

2. MATERIALS AND METHODS

Four molecular assays were compared for their analytical and diagnostic (see Saah & Hoover 1997 for definitions) sensitivity and diagnostic specificity, including two conventional PCRs (from Mao *et al.*, 1997 and Meng *et al.*, 2013; Table 1) and two probe-based qPCRs

(Brunner *et al.*, 2004 and Leung *et al.*, 2017; Table 1), respectively referred to as “Mao PCR”, “Meng PCR”, “Brunner qPCR” and “Leung qPCR”. All assays targeted the viral major capsid protein (MCP) gene, the main viral coat protein, conserved among all members of the *Iridoviridae* (Eaton *et al.* 2007). Oligonucleotide sequences (primers and probes) were compared to an alignment of 21 ranaviruses with published whole genomes (Table S1) to obtain orientations and annealing positions and to check for polymorphisms in binding regions. Assay performance was compared using both cultured ranavirus isolates and field-collected samples.

DNA was extracted alongside extraction controls (nuclease free water as a negative control and cell-cultured ranavirus as a positive control) using an ammonium acetate protocol and purified using ethanol washes (Nicholls *et al.* 2000). Samples were stored at -20°C until use. Conventional PCRs were run in 8µl reactions and qPCRs in 20µl reactions. Two microliters of template DNA were used in every reaction, regardless of assay used. A PCR positive control consisting of DNA extracted from a cultured ranavirus isolated from a UK frog (RUK13; GenBank accession KJ538546; Price, 2014) and a no-template negative control consisting of nuclease-free water were run in duplicate on each PCR plate.

The Mao PCR reaction mixtures comprised forward and reverse primers MCP4 and MCP5 (Table 1; Mao *et al.*, 1997) at a final concentration of 0.5µM, 4µl of 2X DreamTaq Green PCR Master Mix (ThermoFisher Scientific, Massachusetts, USA), and nuclease-free water to take the total volume excluding template to 6µL. The nested Meng PCR used the Mao PCR as the first step but both primers were diluted to a final concentration of 0.05µM. The second step of the nested PCR used primers MCP-IF and MCP-IR from Meng *et al.* (2013) at a concentration of 0.4µM, but all other reagents and concentrations were the same as used for the Mao PCR. The Brunner qPCR used the rtMCP-for and rtMCP-rev primers (Picco *et al.* 2007) at a final concentration of 0.9µM with 10µl TaqMan Universal 2x PCR Master Mix

(ThermoFisher Scientific, Massachusetts, USA), rtMCP-probe at a final concentration of 0.25 μ M, and nuclease-free water to take the volume to 18 μ L (Brunner *et al.*, 2004 - cited in Picco and Collins, 2007; Whitfield *et al.*, 2012, 2013). Reaction mixtures for the Leung qPCR comprised forward and reverse primers (MCP_F and MCP_R) at a final concentration of 0.5 μ M, 10 μ L TaqMan Universal 2x PCR Master Mix (ThermoFisher Scientific, Massachusetts, USA), MCP_probe at a final concentration of 0.25 μ M, and nuclease-free water to take the volume to 18 μ L (Leung *et al.* 2017).

The Mao and Meng PCRs were run on a Techne PCRmax Alpha Cycler 1 (Fisher Scientific, Loughborough, UK) and thermocycler settings comprised a 10-minute strand dissociation step at 95°C, 23 cycles of 95°C for 30 seconds, 62°C (decreasing by 0.5°C per cycle) for 30s, and 72°C for 30s, 25 cycles of 95°C, 50°C and 72°C each for 30s, and a final elongation step of seven minutes at 72°C. The qPCRs were run on a StepOne Real-Time PCR system (Applied Biosystems, California, USA) and began with a two-minute step at 50°C to prevent carry-over contamination with Uracil-N glycosylase (UNG) followed by a 10-minute strand dissociation step at 95°C. The Brunner qPCR was run for 50 cycles comprising steps of 95°C for 20 seconds, 54°C for 20 seconds and 72°C for 30 seconds. The Leung PCR was run for 50 cycles comprising steps of 95°C for 15 seconds, 60°C for 30 seconds and 72°C for 30 seconds. Conventional PCR products were visualized on 100ml 2% agarose gels containing 5 μ L SYBR Safe DNA Gel Stain (Invitrogen, California, USA) alongside a 100-base pair ladder (Thermo Fisher Scientific, Massachusetts, USA). Positive results were defined in each of two ways: 1) using a “majority-rules” approach where a consensus was taken by considering the overall majority returned from replicates (Miller *et al.* 2015), 2) by taking amplification in any replicate as evidence of a positive sample.

2.1 Comparisson of assay sensitivity and specificity

Assay analytical sensitivity was compared using laboratory isolates grown in culture in epithelioma papulosum cyprini cells (Winton et al., (2010); ECACC 93120820). Isolates were used from two of the lineages of the amphibian-associated ranaviruses (frog virus (FV) 3-like and common midwife toad ranavirus (CMTV)-like; (Price et al. 2017). A ten-fold dilution series of DNA extractions of each isolate (1 to 1^{-10}), diluted in water, was used as template DNA.

To investigate the relative diagnostic sensitivities and specificities of the assays, the Meng PCR was considered a ‘gold standard’ on account of the high sensitivity of nested protocols (Hafz et al. 2005, Miller & Sterling 2007) and the capacity to confirm virus amplification by sequencing amplicons, as is recommended by the OIE (OIE 2017). DNA extractions of samples from an archive comprising toe-clips and liver tissue from a diverse assemblage of Costa Rican herpetofauna (Table S2) were screened using the Meng PCR and positive results confirmed by sequencing at Macrogen Inc. (F. Wynne unpubl. data). Forty “known positive” and forty “known negative” samples were selected at random from this larger set. Diagnostic sensitivities of the Mao PCR, Brunner qPCR and Leung qPCR were calculated relative to the Meng PCR (gold standard) as the proportion of the total known positive samples that amplified. Diagnostic specificities were also calculated relative to the gold standard as the proportion of known negatives that did not amplify.

2.2 Intra-assay precision

Intra-assay precision - defined as an assay’s capacity to generate consistent results among replicates within the same run - was investigated by using each assay to run six replicate reactions per sample. Fifteen known positive and fifteen known negative samples were

selected at random from the sets used above. The intra-assay precision was calculated as the proportion of replicates that returned the expected result.

2.3 Phylogenetic analyses of amplicon sequences from Mao and Meng PCRs

Phylogenetic trees were generated and used to compare the MCP regions amplified by the Meng and Mao PCRs. A multiple-sequence alignment of the complete coding sequence of the MCP gene of 21 ranavirus isolates with published whole genomes (Table S1) was trimmed to the maximum possible length of both the Meng and Mao amplicon sequences by mapping the respective primer pairs of each assay to the alignment. Phylogenetic trees were then constructed with Mr Bayes v3.2.2 (Huelsenbeck & Ronquist 2001) using a generalized time-reversible model of nucleotide substitution with rate variation modelled as a gamma distribution with four rate categories. Two runs of four MCMC chains were run for 500,000 generations and the default program settings were used for other parameters.

3. RESULTS

3.1 Polymorphisms within binding regions of primers and probes

The Mao PCR primers were 100% sequence identical to all 21 ranavirus isolates but there were mismatches (single nucleotide polymorphisms) between the oligonucleotide sets of the other assays and some of the 21 isolates (Table 2). The Meng_IR primer had between one and four mismatches with eight of the isolates and appeared to be poorly suited for use with the whole group of *Ambystoma tigrinum* virus (ATV)-like ranaviruses (Price 2016, Price et al. 2017).

3.2 Analytical sensitivity of assays in detection of cultured ranavirus isolates

The Leung qPCR was two orders of magnitude more analytically sensitive than other assays regardless of the viral lineage of the cultured isolate (FV3 or CMTV; Table 3). The Meng

PCR and Brunner qPCR performed similarly, exhibiting slightly greater analytical sensitivity than the Mao PCR (Table 3).

3.3 Assay performance against field-collected samples

The assays did not perform equally against the field-collected samples (Fisher's exact test, $P = 2.1 \times 10^{-19}$). The Leung qPCR showed the highest diagnostic sensitivity, detecting all of the known positives and therefore demonstrating 100% comparative diagnostic sensitivity with the results from the Meng PCR confirmed by sequencing (the proposed gold standard; Figure 1a). The Brunner qPCR showed lower diagnostic sensitivity, only correctly identifying 11 of 40 (27.5%) positive samples if a majority-rules approach was applied. If amplification in any replicate was used to define a positive result, then an additional nine known positives were correctly identified taking the total sensitivity to 20 of 40 (50%; Figure 1a). The Mao PCR performed extremely poorly, correctly identifying only one known positive (2.5%) using a majority-rules approach with just two additional samples giving an ambiguous result, and the remaining 37 samples showing no amplification in either replicate (Figure 1a).

There was no significant difference between the assays in terms of diagnostic specificity against known negative samples (Fisher's exact test, $P = 0.106$). Using a majority-rules approach, all assays showed 100% diagnostic specificity, correctly identifying all of the known negatives. However, taking any amplification (either replicate) as a positive result reduced the Brunner qPCR's comparative diagnostic specificity to 92.5% (three of the known negative samples showed amplification in one replicate; Figure 1b). Since the Brunner qPCR had a significantly lower diagnostic sensitivity than the Meng PCR or Leung qPCR in screens of known positive samples it is very unlikely that this amplification was explained by a greater capacity to detect ranavirus than exhibited by those assays. Instead, it is most likely

that this amplification was non-specific and that the Brunner qPCR generated false positive results for these three (out of 40) samples.

3.4 Intra-assay precision

The Leung qPCR performed most consistently of the assays tested in terms of the proportions of replicate reactions showing amplification with known positive samples: all samples (n = 15) were identified as positive when applying a majority-rules approach with only a single replicate in a total of 90 (15 samples, 6 replicates each) showing no amplification (Fig. 2a). The Brunner qPCR performed relatively poorly: none of the 15 positives would have been called positive using a majority-rules approach and just 38.9% of all replicates showed amplification (Fig 2a). The Mao PCR performed very poorly with just a single replicate out of 90 yielding a band on gels (Fig 2a). The Brunner qPCR had the lowest intra-assay precision for negative samples, however both qPCRs showed high intra-assay precision: 93 of 96 replicates (96.9%) of the Brunner qPCR and 94 of 96 replicates (97.9%) of the Leung qPCR showed no amplification (Fig. 2b). The Mao PCR showed 100% consistency among negative samples; no replicate produced bands on gels (Fig. 2b). When a majority-rules approach was applied to these results, no false positive results were generated by any assay.

3.5 Phylogenetic signal of PCR amplicons

The phylogenetic signal of the Meng PCR amplicon (320 base pairs [bp]) was compared to that of the longer Mao PCR amplicon (531 bp). The two amplicons returned well-supported trees with very similar topologies (Fig. 3). Neither amplicon contained sufficient signal to return the CMTV-like ranaviruses as the monophyletic group that is obtained when using much larger datasets (Price 2016). The amplicon sequences of both assays incorrectly placed ToRV1 in the FV3-like clade but this isolate is a mosaic of two highly divergent ranavirus types which requires whole genome data to resolve its true position (Price 2016). The shorter

Meng PCR amplicon exhibited some loss of resolution relative to the Mao PCR amplicon, especially in distinguishing CMTV-like viruses, but otherwise the two trees were highly concordant and both amplicons yielded accurate information that could be used to infer the ranaviral group (ATV-like, CMTV-like, FV3-like).

4. DISCUSSION

Ranaviruses are important pathogens which cause an emerging disease, spread by humans in some cases, that has had community-level impacts on amphibians (Picco & Collins 2008, Schloegel et al. 2009, Price et al. 2014, 2016, 2017). Sensitive and reliable diagnostics are required to yield accurate estimates of incidence and prevalence. We found considerable variation in the performance of four molecular assays that are routinely used for ranavirus diagnostics (both surveillance of wild populations and experimental laboratory studies).

Assays varied in their suitability for ranavirus detection based on the viral genotype: in particular, one of the Meng PCR primers contained multiple mismatches to a whole group of amphibian-associated ranaviruses, the ATV-like viruses, affecting the optimal temperature for primer annealing and increasing the probability of false negative results. Assays also varied in their diagnostic sensitivity and specificity as well as precision of results among replicates. The Leung qPCR showed 100% comparative diagnostic sensitivity and specificity, but the Mao PCR showed poor diagnostic sensitivity against our set of field-collected samples and the Brunner qPCR also resulted in false negatives as well as potential false positives. Although the Leung qPCR performed well in all experiments, neither qPCR assay allowed sequencing of amplicons as confirmation of results (OIE 2017) or as a basis for inferences about viral genotype (Black et al. 2017), meaning that alternative assays (or the use of a second assay) may be required when working with samples from regions or species where ranavirus has not been previously detected. In this respect, the Meng nested PCR has

high diagnostic sensitivity and yielded reasonable phylogenetic signal compared to the Mao PCR, enabling viral isolates to be roughly placed within the major groups of amphibian-associated ranaviruses (Black et al. 2017), although it appeared to be poorly suited for amplification of a whole group of ranaviruses due to mismatches in primer regions.

The use of an inappropriate assay that yields false negative results is a significant problem for estimates of the current ranavirus distribution. Low intensity infections of individuals as well as whole populations/communities where ranavirus occurs at low prevalence may be falsely reported as negative, biasing estimates of ranavirus distribution. This is important for three main reasons: 1) patchy distributions can be taken as evidence of long-distance dispersal and can favor the inclusion of agents outside of the straightforward host-pathogen interaction in conceptual models explaining pathogen distributions. 2) If, for example, assay sensitivity is only sufficient for detecting infections from visceral organs of individuals experiencing active disease, perceptions of the impact, host range and virulence of these viruses could be biased, which may influence the focus of both management approaches and research effort. 3) Assays with poor sensitivity could also bias understanding of the genetic diversity among ranaviruses (in the context of virulence and other key viral life history traits) if some genotypes have low transmission rates or increased latency for example. As with false negatives, false positives bias estimates of distribution and impact but might also have important consequences for conservation and commercial activities. Considering amplification of any replicate indicative of infection led to probable false positives during this study, whereas using a majority-rules approach reduced this, indicating it to likely be a more accurate approach, which should be adopted as standard. Given its OIE status, there are pressures to consider ranavirus presence when undertaking translocations of herpetofauna for both commercial and conservation purposes. False positive results could interfere with

important economic activities such as the culture and supply of amphibians for food or conservation actions.

In spite of the notifiable status of ranavirus disease and OIE guidelines for diagnostic tests, there is currently no standardized approach to diagnostics (OIE 2017, Black et al. 2017).

Many molecular assays have been used for ranavirus detection, often without checks on the appropriateness or power of the assays (Jaramillo et al. 2012, Black et al. 2017). Our study highlights the importance of selecting and optimizing methods that are compatible with study aims.

Acknowledgements

We thank Andrew Cunningham for use of the RUK13 ranavirus isolate. This work was supported by the Natural Environment Research Council (grant numbers: NE/M000338/1, NE/M000591/1, NE/M00080X/1). This work was carried out under the following CONAGEBIO permits R-029-2015-OT-CONAGEBIO, R-030-2015-OT-CONAGEBIO.

Literature cited

289 Black Y, Meredith A, Price SJ (2017) Detection and Reporting of Ranavirus in Amphibians:
 290 Evaluation of the roles of the World Organisation for Animal Health and the published
 291 literature. *J Wildl Dis* 53:509–520

292 Boyle DG, Boyle DB, Olsen V, Morgan J, Hyatt A (2004) Rapid quantitative detection of
 293 chytridiomycosis (*Batrachochytrium dendrobatidis*) in amphibian samples using real-
 294 time Taqman PCR assay. *Dis Aquat Organ* 60:141–148

295 Brunner JL, Schock DM, Davidson EW, Collins JP (2004) Intraspecific reservoirs: Complex
 296 life history and the persistence of a lethal ranavirus. *Ecology* 85:560–566

297 Campbell LJ, Hammond SA, Price, SJ, Sharma, MD, Garner, TWJ, Birol, I, Helbing, CC,
 298 Wilfert, L, Griffiths AGF (2018) A novel approach to wildlife transcriptomics provides
 299 evidence of disease-mediated differential expression and changes to the microbiome of
 300 amphibian populations. *Mol Ecol* 27:1413-1427

301 DiRenzo GV, Campbell Grant EH, Longo AV, Che-Castaldo C, Zamudio KR, Lips KR
 302 (2017) Imperfect pathogen detection from non-invasive skin swabs biases disease
 303 inference. *Methods Ecol Evol* 9:380–389

304 Eaton HE, Metcalf J, Penny E, Tcherepanov V, Upton C, Brunetti CR (2007) Comparative
 305 genomic analysis of the family Iridoviridae: re-annotating and defining the core set of
 306 iridovirus genes. *Virology* 4:11

307 Gray MJM, Miller DDL, Hoverman JTJ (2009) Ecology and pathology of amphibian
 308 ranaviruses. *Dis Aquat Organ* 87:243–266

309 Hafez HM, Hauck R, Lüschoff D, McDougald L (2005) Comparison of the Specificity and
 310 Sensitivity of PCR, Nested PCR, and Real-Time PCR for the Diagnosis of
 311 Histomoniasis. *Avian Dis* 49:366–370

312 Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogenetic trees.
313 Bioinformatics 17:754–755

314 Hyatt A, Boyle DD, Olsen V, Boyle DD, Berger L, Obendorf D, Dalton A, Kriger K, Hero
315 M, Hines H, Phillott R, Campbell R, Marantelli G, Gleason F, Colling A (2007)
316 Diagnostic assays and sampling protocols for the detection of *Batrachochytrium*
317 *dendrobatidis*. Dis Aquat Organ 73:175–192

318 Jaramillo D, Tweedie A, Becker JA, Hyatt A, Crameri S, Whittington RJ (2012) A validated
319 quantitative polymerase chain reaction assay for the detection of ranaviruses (Family
320 Iridoviridae) in fish tissue and cell cultures, using EHNV as a model. Aquaculture 356–
321 357:186–192

322 Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, Daszak P (2008) Global
323 trends in emerging infectious diseases. Nature 451:990–993

324 Kuiken T, Leighton FA, Fouchier RAMM, LeDuc JW, Peiris JSMM, Schudel A, Stöhr K,
325 Osterhaus ADMEME (2005) Pathogen surveillance in animals. Science 309:1680–1681

326 Lachish S, Gopalaswamy AM, Knowles SCL, Sheldon BC (2012) Site-occupancy modelling
327 as a novel framework for assessing test sensitivity and estimating wildlife disease
328 prevalence from imperfect diagnostic tests. Methods Ecol Evol 3:339–348

329 Leung WTM, Thomas-Walters L, Garner TWJ, Balloux F, Durrant C, Price SJ (2017) A
330 quantitative-PCR based method to estimate ranavirus viral load following normalisation
331 by reference to an ultraconserved vertebrate target. J Virol Methods 249:147–155

332 Lorch JM, Gargas A, Meteyer CU, Berlowski-Zier BM, Green DE, Shearn-Bochsler V,
333 Thomas NJ, Blehert DS (2010) Rapid Polymerase Chain Reaction Diagnosis of White-
334 Nose Syndrome in Bats. J Vet Diagnostic Investig 22:224–230

335 Mao J, Hedrick R. P, Chinchar V., Chinchar G V. (1997) Molecular characterization,
 336 sequence analysis, and taxonomic position of newly isolated fish iridoviruses. *Virology*
 337 229:212–220

338 Meng Y, Zhang H, Liang H, Zeng L, Xiao H, Xie C (2013) Development of a loop-mediated
 339 isothermal amplification assay for rapid detection of iridovirus in the Chinese giant
 340 salamander. *J Virol Methods* 194:211–216

341 Miller DL, Pessier AP, Hick P, Whittington RJ (2015) Comparative Pathology of
 342 Ranaviruses and Diagnostic Techniques. In: *Ranaviruses*. Springer International
 343 Publishing, Cham, p 171–208

344 Miller KM, Sterling CR (2007) Sensitivity of nested PCR in the detection of low numbers of
 345 *Giardia lamblia* cysts. *Appl Environ Microbiol* 73:5949–5950

346 Nicholls JA, Double MC, Rowell DM, Magrath RD (2000) The evolution of cooperative and
 347 pair breeding in thornbills *Acanthiza* (Pardalotidae). *J Avian Biol* 31:165–176

348 OIE (2016) Manual of Diagnostic Tests for Aquatic Animals.

349 OIE (2017) Chapter 2.1.2 Infection with Ranavirus Manual of diagnostic tests for aquatic
 350 animals In: OIE. Manual of Diagnostics for Aquatic Animals

351 Picco AM, Brunner JL, Collins JP (2007) Susceptibility of the Endangered California Tiger
 352 Salamander, *Ambystoma californiense*, to *Ranavirus* Infection. *J Wildl Dis* 43:286–290

353 Picco AM, Collins JP (2007) Fungal and Viral Pathogen Occurrence in Costa Rican
 354 Amphibians. *J Herpetol* 41:746–749

355 Picco AM, Collins JP (2008) Amphibian Commerce as a Likely Source of Pathogen
 356 Pollution. *Conserv Biol* 22:1582–1589

357 Price SJ (2014) Emergence of a virulent wildlife disease: using spatial epidemiology and
358 phylogenetic methods to reconstruct the spread of amphibian viruses. :158

359 Price SJ (2016) Comparative Genomics of Amphibian-like Ranaviruses, Nucleocytoplasmic
360 Large DNA Viruses of Poikilotherms. *Evol Bioinform Online* 11:71–82

361 Price SJ, Ariel E, MacLaine A, Rosa GM, Gray MJ, Brunner JL, Garner TWJ (2017) From
362 fish to frogs and beyond: Impact and host range of emergent ranaviruses. *Virology*
363 511:272–279

364 Price SJ, Garner TWJ, Cunningham AA, Langton TES, Nichols RA (2016) Reconstructing
365 the emergence of a lethal infectious disease of wildlife supports a key role for spread
366 through translocations by humans. *Proc R Soc B* 283:20160952

367 Price SJ, Garner TWJ, Nichols RA, Balloux F, Ayres C, Mora-Cabello De Alba A, Bosch J
368 (2014) Collapse of amphibian communities due to an introduced ranavirus. *Curr Biol*
369 24:2586–2591

370 Rijks JM, Saucedo B, Sluijs AS der, Wilkie GS, Asten AJAM van, Broek J van den,
371 Boonyarittichaijij RMS, Sterren F van der, Martel A, Pasmans F, Hughes J, Gröne A,
372 Beurden SJ van, Kik MJL (2016) Investigation of Amphibian Mortality Events in
373 Wildlife Reveals an On-Going Ranavirus Epidemic in the North of the Netherlands.
374 *PLoS One* 11:e0157473

375 Rosa GM, Sabino-Pinto J, Laurentino TG, Martel A, Pasmans F, Rebelo R, Griffiths RA,
376 Stöhr AC, Marschang RE, Price SJ, Garner TWJ, Bosch J (2017) Impact of
377 asynchronous emergence of two lethal pathogens on amphibian assemblages. *Sci Rep*
378 7:43260

379 Saah AJ, Hoover DR (1997) “Sensitivity” and “Specificity” Reconsidered: The Meaning of

380 These Terms in Analytical and Diagnostic Settings. *Ann Intern Med* 126:91

381 Schloegel LM, Picco AM, Kilpatrick AM, Davies AJ, Hyatt AD, Daszak P (2009) Magnitude
382 of the US trade in amphibians and presence of *Batrachochytrium dendrobatidis* and
383 ranavirus infection in imported North American bullfrogs (*Rana catesbeiana*). *Biol*
384 *Conserv* 142:1420–1426

385 Skerratt LF, Mendez D, McDonald KR, Garland S, Livingstone J, Berger L, Speare R (2011)
386 Validation of Diagnostic Tests in Wildlife: The Case of Chytridiomycosis in Wild
387 Amphibians. *J Herpetol* 45:444–450

388 Teacher AGFF, Cunningham AA, Garner TWJJ (2010) Assessing the long-term impact of
389 *Ranavirus* infection in wild common frog populations. *Anim Conserv* 13:514–522

390 Warne RW, LaBumard B, LaGrange S, Vredenburg VT, Catenazzi A (2016) Co-Infection
391 by Chytrid Fungus and Ranaviruses in Wild and Harvested Frogs in the Tropical Andes.
392 *PLoS One* 11:e0145864

393 Whitfield SM, Donnelly MA, Geerdes E, Kerby J (2012) Ranavirus infection in native
394 amphibians at La Selva Biological Station, Costa Rica: First report of ranavirus in
395 Central America. *Herpetol Rev* 43:425–427

396 Whitfield SM, Geerdes E, Chacon I, Ballesteros Rodriguez E, Jimenez RR, Donnelly MA,
397 Kerby JL (2013) Infection and co-infection by the amphibian chytrid fungus and
398 ranavirus in wild Costa Rican frogs. *Dis Aquat Organ* 104:173–178

399 Winton, J, Batts, W, LeBerre, M, Bremont, M, Fijan, N (2010) Current lineages of the
400 epithelioma papulosum cyprini (EPC) cell line are contaminated with fathead minnow,
401 *Pimephales promelas*, cells. *J Fish Dis* 33(8): 701-704.

Tables

Table 1. Primer and probe sequences of ranavirus diagnostic PCRs compared for sensitivity and specificity.

Assay	Amplicon size (bp)	Oligo ID	Sequence (5'→3')	Strand	Position in coding sequence
Mao PCR	531	MCP 4	GACTTGGCCACTTATGAC	Sense	43-60
		MCP 5	GTCTCTGGAGAAGAAGAA	Anti-sense	556-573
Meng (nested) PCR	320	MCP-IF	TCGCTGGTGTTCCTATCAT	Anti-sense	457-476
		MCP-IR	CTGCCAAGATGTCGGGTAAC	Sense	158-177
Brunner qPCR	70	rt-MCP-for	ACACCACCGCCCAAAGTAC	Sense	1298-1317
		rtMCP-rev	CCGTTCATGATGCGGATAATG	Anti-sense	1347-1367
		rtMCP-probe	CCTCATCGTTCTGGCCATCAACCAC	Sense	1320-1344
Leung qPCR	97	MCP_F	GTCCTTTAACACGGCATACT	Sense	381-401
		MCP_R	ATCGCTGGTGTTCCTATC	Anti-sense	459-477
		MCP_probe	TTATAGTAGCCTR*TGCGCTTG GCC	Anti-sense	432-455

*R is the IUPAC nucleotide ambiguity code for the bases Adenine or Guanine

Table 2. Single nucleotide polymorphisms (SNPs) within regions of the major capsid protein (MCP) gene complementary to primer and probe sequences. Full names and accession numbers for virus isolates are provided in Table S1.

Assay	Oligo ID	Isolate	SNP count
Mao PCR	MCP 4	Not applicable	100% identical to all isolates
	MCP 5	Not applicable	100% identical to all isolates
Meng PCR	MCP-IF	SERV	1
		SERV	2
	MCP-IR	Rmax	2
		CodIV	2
		TFV	1
		EHNV	2
		ESV	2
		ToRV1	4
		ATV	2
		SERV	1
Brunner qPCR	rt-MCP-for	PPIV	1

	rtMCP-rev	Not applicable	100% identical to all isolates
		SERV	1
	rtMCP-probe	GGRV	1
		PPIV	1
		SERV	3
	MCP_F	ESV	1
	MCP_R	SERV	1
Leung qPCR		SERV	1
		GGRV	1
	MCP_probe	ATV	1
		BIV	1

415

416

417

418

419

420

421

422

Table 3. Limit of detection of four PCR methods for detecting ranavirus isolates from two major groups of amphibian-associated ranaviruses (frog virus 3-like [FV3] or common midwife toad virus-like [CMTV]). DNA extracted from cultured isolates was diluted from 1 to $1e^{-10}$.

Isolate	Limit of Detection				
	Mao	Meng	Brunner	Leung	
FV3	$1e^{-6*}$	$1e^{-6*}$	$1e^{-6}$	$1e^{-8}$	
CMTV	$1e^{-6*}$	$1e^{-6}$	$1e^{-6}$	$1e^{-8}$	

* amplification in one of two replicates at the dilution given

445

446 **Figure 1. Relative performance of four molecular assays in detecting ranavirus from**

447 **non-lethal samples of Costa Rican amphibians.** Assay performance is relative to a nested

448 PCR assay (Meng et al. 2013) used as a ‘gold standard’. a) Relative sensitivity tested against

449 known positives, b) Relative specificity tested against known negatives. 95% confidence

450 intervals shown.

Figure 2. Comparison of the capacity of ranavirus molecular assays to generate consistent results among replicate reactions (intra-assay precision). a) Proportion of replicates that returned the expected result with known positive samples. b) Proportion of replicates that returned the expected result with known negative samples. 95% confidence intervals shown.

Figure 3. Comparison of the phylogenetic signal contained in sequences from PCR amplicons generated by two PCR methods (Mao PCR [531 bp] and Meng PCR [319 bp]). Both assays target the viral major capsid protein gene and an alignment of the full locus for amphibian-associated ranaviruses, with published complete genomes, was trimmed to the lengths of the respective PCR amplicons. Trees were constructed using default settings in Mr Bayes and drawn as cladograms (see main text). Support values at nodes are posterior probabilities. Full names of viruses, GenBank accession numbers and citations are provided in Table S1.