2015-02

Practice expectations for graduates of Australian critical care nurse education: Development of the standard of practice and evaluation of critical-care-nursing tool (SPECT)

Gill, F

<http://hdl.handle.net/10026.1/4183>

10.1016/j.aucc.2014.10.002

Australian Critical Care

Elsevier BV

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Papers and Poster Abstracts: The 39th Australian and New Zealand Scientific Meeting on Intensive Care and the 20th Annual Paediatric and Neonatal Intensive Care Conference, Melbourne, Australia, October 2014

Nursing Scholarship Papers

Practice expectations for graduates of Australian critical care nurse education: Development of the standard of practice and evaluation of critical-care-nursing tool (SPECT)

F. Gill^{1,2,*}, G. Leslie², C. Grech³, J. Latour^{2,4}

¹ Princess Margaret Hospital for Children, Australia

² Curtin University, Australia

³ University of South Australia, Australia

⁴ Plymouth University, United Kingdom

Winner: Nursing Scholarship award sponsored by

Introduction: To achieve optimal patient outcomes, nurses working in critical care should hold a specialty qualification. Currently no uniform practice outcomes for critical care programs exist. Varying interpretations of a “critical care qualification” have been used as there are no agreed practice standards to measure outcomes. Additionally there have been no reported attempts to include health service consumer input into the development of practice standards.

Study objectives: To develop national practice standards for graduates of critical care nurse education and a clinical assessment tool (SPECT) to measure practice.

Methods: A multi-step, mixed method design including a literature review, analysis of existing programs ($n = 22$) and identification of health consumer views via individual and group interviews ($n = 17$). The findings informed a Delphi study of 105 nurse stakeholders to determine critical care graduate practice standards. The SPECT was based upon the Delphi results and pilot-tested.

Results: The literature review, analysis of courses and health consumers views were used to develop 99 draft practice standards. The Delphi study resulted in 98 agreed standards (organised in 6 domains) that describe newly graduated critical care nurses should be able to manage the care of critically ill patients in most complex situations. They should not be expected to perform leadership

roles. The standards were further arranged into a practical tool to measure graduate practice. This involved elimination of some duplication and re-grouping of some statements as elements. The SPECT contained 86 statements – initial testing revealed strong content validity, good internal consistency, test-retest and intra-rater agreement, with at least moderate agreement for 94% of statements. The majority of panel responses (66%) were ranked as clinically feasible.

Conclusion: For the first time health consumers have contributed to the components of critical care nurse graduate practice. The standards and SPECT provide the means to clearly define the expected practice level for graduates of critical care education programs.

Acknowledgements: Thank you to: Australian College of Critical Care Nurses, Department of Health Western Australia, Helen Bailey Scholarship 2011, the participants for each phase of the study.

<http://dx.doi.org/10.1016/j.aucc.2014.10.002>

The transition from active treatment to initiation of end-of-life care within the intensive care unit: A clinician's perspective

L. Brooks^{1,*}, P. Nicholson², E. Manias³

¹ School of Nursing and Midwifery, Faculty of Health, Deakin University, Australia

² Nursing, School of Health Sciences, The University of Melbourne, Australia

³ SONM, Deakin University, Royal Melbourne Hospital, The University of Melbourne, Australia

Introduction: Advances in knowledge and technology present a reality that life in the ICU can be artificially sustained for a period, in patients with terminal conditions. Clinicians in the ICU are commonly faced with decisions involving withholding or withdrawing this life sustaining therapy and this presents many clinical and ethical challenges.

Objectives: The study was conducted in a 24-bed ICU located in a major tertiary referral hospital in Melbourne, Australia. We aimed to explore the perspectives and experiences of nurses and physicians providing end-of-life care in the ICU. In particular, perceived