2013-09-01

Cadmium contamination of agricultural soils and crops resulting from sphalerite weathering.

Robson, TC

<http://hdl.handle.net/10026.1/4355>

10.1016/j.envpol.2013.09.001

Environ Pollut

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1 Disclaimer: This is a pre-publication version. Readers are recommended to consult
2 the full published version for accuracy and citation. Published in Environmental
3 Pollution, 184, 283-289 (2014), doi: 10.1016/j.envpol.2013.09.001.
4

5 **Cadmium contamination of agricultural soils and crops**

6 **resulting from sphalerite weathering**

7 T.C. Robson^{a*}, C.B. Braungardt^a, J. Rieuwerts^a, P. Worsfold^a

8
9 ^a Biogeochemistry Research Centre, Plymouth University, Drake Circus. Plymouth, PL4
10 8AA, United Kingdom.

11
12 *thomas.robson@plymouth.ac.uk

13 charlotte.braungardt@plymouth.ac.uk

14 john.rieuwerts@plymouth.ac.uk

15 P.Worsfold@plymouth.ac.uk

16
17 *Corresponding author: Thomas C. Robson Tel: (+44)7712 599565; School of
18 Geography, Earth and Environmental Science, Plymouth University, Drake Circus,
19 Plymouth, Devon, PL4 8AA.

23 **Abstract**

24 The biogeochemistry and bioavailability of cadmium, released during sphalerite
25 weathering in soils, were investigated under contrasting agricultural scenarios to assess
26 health risks associated with sphalerite dust transport to productive soils from mining.

Laboratory experiments (365 d) on temperate and sub-tropical soils amended with sphalerite ($< 63 \mu\text{m}$, 0.92 wt.% Cd) showed continuous, slow dissolution ($0.6 - 1.2 \% \text{ y}^{-1}$). Wheat grown in spiked temperate soil accumulated $\approx 38 \%$ ($29 \mu\text{mol kg}^{-1}$) of the liberated Cd, exceeding food safety limits. In contrast, rice grown in flooded sub-tropical soil accumulated far less Cd ($0.60 \mu\text{mol kg}^{-1}$) due to neutral soil pH and Cd bioavailability was possibly also controlled by secondary sulfide formation. The results demonstrate long-term release of Cd to soil porewaters during sphalerite weathering. Under oxic conditions, Cd may be sufficiently bioavailable to contaminate crops destined for human consumption; however flooded rice production limits the impact of sphalerite contamination.

Capsule:

Sphalerite dissolves steadily in oxic agricultural soils and can release highly bioavailable Cd, which may contaminate food crops destined for human consumption.

Keywords: metals; sulfide weathering; human health; rice; risk assessment

Introduction

Cadmium (Cd) is considerably environmentally mobile, bioavailable and toxic to humans (Smolders and Mertens, 2013) and there are linkages between mineral exploitation, Cd soil contamination and human health hazards, for example the contamination of soils by Japan's Jinzu River and its association with the debilitating 'itai-itai' disease (Ishihara et al., 2001). In the region bordering Guangdong and Hunan provinces (China), decades of metal production were shown to have contaminated river sediments and agricultural soils (e.g. Chenzhou) as far as 60 km from the source ($> 9 \mu\text{mol Cd kg}^{-1}$) (Limei et al., 2008) and inhabitants are considered at risk of chronic health effects from consuming locally grown rice and vegetables (H. Zhao et al., 2012; Zhuang et al., 2009). Crop safety is a concern because the primary human intake routes for Cd are tobacco smoking and diet (Järup, 2003), both of which link human exposure to soil contamination. Chronic

toxicity may arise because Cd has a long biological half-life (15 – 30 years) and accumulates in the body, resulting in kidney disease, osteoporosis, lung and prostate cancer and endocrine disruption (Henson and Chedrese, 2004; Järup, 2003).

Mining and ore processing produces fine mineral particles enriched with potentially toxic metals and metalloids (e.g. As, Cd, Hg). These particles are vulnerable to fluvial (Miller et al., 2004; Simmons et al., 2005) and aeolian (Castillo et al., 2013; Zota et al., 2009) transport, for example through erosion from exposed tailings, so they can behave as vectors for toxic elements. Several investigators have reported the spatial distribution and concentration of toxic metals in soils affected by mineral exploitation, but less is understood about how transported mineral particles influence soil quality in terms of the biogeochemical cycling of toxic metals and the risk they pose to human health through crop contamination.

Commonly exploited sulfide ores (e.g. sphalerite) are vulnerable to oxidative and acid-promoted dissolution under moist, oxic surface conditions and may release metals into soil porewater and surface waters, providing a source of potentially toxic metals for plant uptake. This situation is most likely to affect communities in developing and rapidly industrialising countries, where environmental regulations may be either weak or poorly enforced and soils impacted by mining may be used for agriculture (Miller et al., 2004; Zhuang et al., 2009).

Sphalerite (ZnS), the primary geologic source of zinc (Zn) and Cd, occurs commonly around the world. Sphalerite is notable for its tendency for isomorphic substitution of Zn by other metals and Cd is generally present in solid solution at 0.2–1 % (Smolders and Mertens, 2013). The alteration mechanisms proposed for sphalerite include oxidative dissolution, either by molecular oxygen (Eq. 1) or Fe(III) (Eq. 2), and acid-promoted dissolution (Eq. 3) (Heidel et al., 2011). Cd is released from solid solution during sphalerite dissolution (Stanton et al., 2008).

Laboratory experiments in aqueous media (Acero et al., 2007; Stanton et al., 2008) showed that sphalerite dissolution follows a first order reaction with respect to $[\text{H}^+]$ (pH 1 – 4.2); the rate increases with temperature (25 – 70 °C) and is independent of dissolved oxygen concentrations (6.3 – 270 μM dO_2), suggesting that the process described by Eq. 1 is of minimal importance. Apart from differences in experimental design, the iron (Fe) content of the sphalerite is also proposed to influence the dissolution rate (Weisener et al., 2003).

The aim of this study was to determine the rate of sphalerite dissolution and Cd release in soils of contrasting geologic and climatic provenance, and the bioavailability of the Cd to key crops under relevant agricultural scenarios.

Methods

Investigative approach

This study comprised of: (1) Laboratory batch incubations of sphalerite-spiked (0.1 % m/m) soils to determine sphalerite dissolution behaviour and (2) Phytoavailability experiments where *Triticum aestivum* (spring wheat) and *Oryza sativa* (rice) were grown in samples of the temperate and flooded sub-tropical soils, respectively, to evaluate the bioavailability of Cd released from sphalerite.

Reagents and materials

Reagents were of analytical grade or higher (ROMIL, Sigma-Aldrich, Fisher) and ultra-high purity water (UHP, $\geq 18.2 \text{ M}\Omega \text{ cm}^{-1}$) was used for all experiments. Specimen sphalerite was obtained from a private collection (Richard Tayler Minerals, Cobham, UK).

Soil and mineral sampling, preparation and characterisation

A temperate soil (inceptisol from the Tamar Valley, Cornwall, United Kingdom) was sampled from low-intensity grassland, used only for haymaking for the past decade. The sub-tropical soil (The University of Hong Kong Kadoorie Centre) comprised sub-surface oxisol from secondary forest and horticultural soil, in equal parts by volume.

Experimental soils (< 2 mm, dried at 40°C) were fertilised with dried (50°C , ≥ 72 h) well-rotted animal dung (milled and sieved to < 2 mm) at 10 % m/m, bringing the organic matter content to the upper range for productive soils. Sphalerite was finely ground, sieved (ball mill, $< 63\ \mu\text{m}$) and stored in a desiccating, N_2 -purged atmosphere (see Laboratory batch incubation experiments). The $< 63\ \mu\text{m}$ fraction represents clay and silt size particles, which are thought to account for the majority of fugitive dust mass flux (Kon et al., 2007).

Experimental soils were characterised using standard methods (Carter and Gregorich, 2007): total sulfur (S), nitrogen, organic/inorganic carbon (NC2500 elemental analyser, Carlo Erba), eCEC, organic matter, texture and pH (United States Environmental Protection Agency, 2004). For elemental analyses, soils and sphalerite were microwave-digested (MarsXpress, CEM) in 50 % v/v 1:3 HNO_3 :HCl. Bulk mineralogy was evaluated using powder X-ray diffraction (XRD) ($\text{Cu-K}\alpha$, $2\text{-}70^{\circ} 2\theta$, $0.02^{\circ}/\text{S}$, D5000, Siemens). The sphalerite was also examined using scanning electron microscopy with energy dispersive spectroscopy (SEM-EDS) (JSM-7100F, JEOL/Aztec EDS, Oxford Instruments).

Laboratory batch incubation experiments

Batch incubations (≤ 12 months) were conducted in polypropylene beakers using 100 g aliquots of the experimental soils. The soils were either spiked with 0.1 g ground sphalerite or left as controls (no ZnS) and then their moisture was maintained gravimetrically at 75 % field capacity. Triplicate incubation batches were sacrificed and analysed after 0 hours, 7, 30, 90, 180, 270 and 365 days under laboratory conditions. Sacrificed soils were freeze-dried, homogenized and sub-sampled for analyses. Soluble major ions (NO_3^- , SO_4^{2-}), cation-exchangeable Cd/Zn and EDTA-extractable Cd/Zn were extracted in UHP water (1:5), 0.01 M CaCl_2 (1:5) and 0.1 M EDTA (1:30, pH 7.5),

respectively (2 h agitation, reciprocating shaker). Extracts were centrifuged, filtered (0.45 μm) and preserved until analysis (freezing, acidification or refrigeration).

An abiotic control experiment, analogous to the first 30 days of the batch incubation experiments, was performed using sterile soils. Soils were fractionally sterilised to ensure overkill of both microorganisms and endospores, using three cycles of steam heating ($97 \pm 2^\circ\text{C}$, 2 h) and overnight incubated ($37 \pm 1^\circ\text{C}$). The incubations were performed in autoclaved (121°C , 1 h), foam-bunged (tortuous path filter) glass conical flasks. Sterility was maintained by using only heat-sterilised ($\geq 250^\circ\text{C}$) implements, filter-sterilised (0.22 μm) water and observing best practise for sterile handling.

Phytoavailability experiments

Larger (2 kg), analogous, soil incubations were established in polypropylene pots and maintained in parallel with those of the batch experiment. After 180 days, 10-day wheat seedlings (350 seeds m^{-2}) were transplanted into the temperate soils. The plants were matured to ripeness (112 days) under glasshouse conditions ($24 \pm 3^\circ\text{C}$, $66 \pm 3\%$ RH) and then the stems and ears were rinsed (≥ 5 times) with water and freeze-dried. The tissues were comminuted (food processor) and $0.5 \pm 0.01\text{ g}$ ($n = 3$) of tissue was microwave-digested in 10 mL HNO_3 (50 % v/v conc.).

The sub-tropical soils received 14-day rice seedlings, were flooded with 6 cm standing water and plants matured (152 days) inside a growth chamber (Fitotron PG660, Sanyo; 16 h light, 8 h dark at 27°C , followed by 12 h light, 12 h dark at 24°C) and then treated as per the wheat. The drained, saturated soils were core-sampled ($\varnothing 2.5\text{ cm}$, $n = 5$) inside an anoxic chamber (Coy laboratory products). Bulk cores were sub-sampled for aqueous extraction, capped and sealed with Parafilm and then extracted as previously described (see Laboratory batch incubation experiments). Dissolved sulfide was determined in soil water extracts (in an anoxic chamber) using the methylene blue method (Cline, 1969). Solid-phase Fe speciation and acid-volatile sulfide were determined after Lovley and Phillips (1986) and Allen et al. (1993), respectively. Porewater pH and Eh were determined in the water that collected in the core voids.

Analytical techniques

Anion, Fe (II) and dissolved sulfide (S^{2-}) concentrations were determined using ion chromatography (DX-500, Dionex Corporation) or spectrophotometry (8453 UV-Vis, Agilent), respectively. Aqueous metal concentrations were determined using ICP-OES (725-ES, Varian) and ICP-MS (X-series 2 + Collision Cell, Thermo Scientific). Half-cell redox potentials (Hanna HI9025, BDR Gelplas ORP) were corrected against ZoBell's solution. Certified reference materials (BCR 320R channel sediment, IRMM 804 rice flour and PACS-1 marine sediment) were used to verify the satisfactory accuracy and performance of the methods (Table S1). Statistical analyses were performed using the Sigmaplot 12 software package (Systat Software).

Results and discussion

Experimental soil and sphalerite characterisation

Both experimental soils were of circum-neutral pH, rich in organic matter, with a similar moderate eCEC (Table 1). Both soils were Fe rich (5.5 – 6.3 % Fe m/m) but the sub-tropical soil contained 7 times less manganese, more aluminium (6.2 % vs. 3.9 %) and a lower proportion of poorly crystalline Fe and aluminium oxides/hydroxides (factor of 2 – 4). Both soils had similar S, Zn and Cd concentrations, which fell within (Cd) or just above (Zn) the range expected for uncontaminated soils (Mertens and Smolders, 2013). The main crystalline phases were clinocllore ($(Mg,Fe^{2+})_5Si_3Al_2O_{10}(OH)_8$), muscovite ($KAl_2(AlSi_3O_{10})(OH)_2$), illite ($(K,H_3O)(Al,Mg,Fe)_2(Si,Al)_4O_{10}[(OH)_2,(H_2O)]$) and quartz (SiO_2) in the temperate soil (sandy loam) and kaolinite ($Al_2Si_2O_5(OH)_4$), orthoclase ($KAlSi_3O_8$), microcline ($KAlSi_3O_8$), gibbsite ($Al(OH)_3$) and quartz in the sub-tropical soil (silt loam).

Elemental (wet) analyses and SEM-EDS examination showed that the experimental mineral consisted of ZnS ($Zn_{1.01}S_{0.99}$) with 0.3 % m/m Fe and 0.9 % m/m Cd. An XRD analysis and reference to mineral databases also confirmed the crystalline structure to be that of sphalerite and not the sphalerite polymorph, wurtzite (Figure S1).

Soil biogeochemical conditions throughout oxic incubations

The pH was not controlled during the experiment and a notable decrease was observed in the pH of both temperate (- 1.05 pH units) and sub-tropical (- 0.43 pH units) soils during the initial 30 days of incubation. After 30 days, the temperate and sub-tropical soils fluctuated around pH 5.53 ± 0.13 and pH 6.40 ± 0.08 , respectively. Acid buffering experiments, performed after Magdoff and Bartlett (1985), demonstrated that the sub-tropical soil had significantly greater buffering capacity ($\approx 43\%$ at neutral-acid pH) than the temperate soil, which was devoid of carbonate, partly explaining the disparate pH decline in the early stages of the experiment (Figure S2).

Data from an abiotic control experiment covering the initial 30 days incubation of the temperate soil (Figure 1) indicate that the pH decline was mediated by soil microbiota, and was coupled with sharp increases in nitrate and sulfate concentrations, reflecting ammonium and sulfur-oxidising bacterial activity. Both can contribute to soil acidity especially if percolation is prevented, as was the case in these experiments.

[Approximate location for Figure 1]

Sphalerite dissolution had a negligible effect on soil pH, since control and spiked incubation soil pH generally differed by < 0.1 pH unit throughout the batch incubations. Redox potential (Eh) was determined in soil extracts, which were performed as for the pH determinations (see Soil and mineral sampling, preparation and characterisation). The data indicated consistently oxic conditions (350 – 400 mV) throughout the experiment.

Geochemical conditions in sub-tropical soil during rice cultivation

After rice cultivation (180 days oxic, 152 days flooded) the sub-tropical soils had attained neutral pH (7.01 ± 0.08) and moderately reducing conditions (Eh - 23 ± 6 mV) (Figure 2 b). Flooded paddy soils often attain neutral pH as many important reduction reactions, e.g. $\text{Fe(III)} > \text{Fe(II)}$, consume free protons (Ponnamperuma, 1972).

[Approximate location for Figure 2.]

Redox indicators and CaCl_2 -extractable metal concentrations suggest that the availability of Cd in porewater, and therefore to the rice plants, was limited by the formation of secondary Fe/Cd/Zn sulfide phases. Depleted soluble nitrate ($> 99\%$) and sulfate ($\geq 94\%$) (Figure 2 a), together with a significant proportion of acid-extractable Fe(II) ($66 \pm 10\%$), indicated the influence of nitrate, Fe and sulfate-reducing anaerobes (Inglett et al., 2005). No dissolved sulfide was detected in soil extracts but acid-volatile sulfide (AVS) was found in the reduced control soils ($200 \pm 16 \mu\text{mol S}^{2-} \text{ kg}^{-1}$), providing evidence for the formation of amorphous secondary sulfides (e.g. greigite, mackinawite). An AVS determination was not possible in the spiked soils because sphalerite itself is acid-volatile. Concurrent with sulfide formation, net (control-corrected) CaCl_2 -extractable Cd and Zn concentrations were considerably lower in the reduced soils, compared with their oxic equivalents (Figure 2 c), and net EDTA-extractable Cd (Cd_{net}) concentrations also fell from 0.23 ± 0.02 to $0.12 \pm 0.02 \mu\text{mol kg}^{-1}$. Much of the depleted sulfate was unaccounted for by AVS formation, suggesting other contributory mechanisms. There was no olfactory evidence for $\text{H}_2\text{S}_{(\text{g})}$ evolution and sulfate adsorption is minimal in pH neutral soils (Scherer, 2009), excluding adsorption effects. Other potential mechanisms for the observed sulfate and extractable Cd depletion were the formation of non-acid-volatile sulfides (e.g. pyrite, greenockite) and plant uptake, respectively, and these are given further consideration in the section 'Uptake by paddy rice grown in sub-tropical soil'.

Sphalerite dissolution

Cd and Zn extraction protocols

The 0.01 M CaCl_2 protocol was applied to provide a 'snapshot' reflecting plant-available concentrations at a given time (Meers et al., 2007). The 0.1 M EDTA extraction protocol was selected because it effectively scavenges metal cations from solid soil phases (Lo and Yang, 1999; Schecher, 2001), providing an indication of the total Cd and Zn release from sphalerite and total plant-available concentrations over a longer timescale.

256 Cd and Zn release trends

257 The clear distinction between Cd concentrations obtained from control and spiked
258 incubations (Figure 3a/a_i & b/b_i) evidences the release of Cd from sphalerite
259 dissolution. This divergence can be seen after 7 days incubation of both the temperate
260 and sub-tropical soils. Cd concentrations extracted from control incubations were
261 relatively constant throughout the incubation duration, which demonstrates that Cd
262 extractability was not affected by changes in soil pH during the oxic incubations.

263

264 [Approximate location for Figure 3.]

265

266 Extractable Zn concentrations (Figure 3c/c_i & d/d_i) were considerably higher than Cd
267 concentrations, reflecting the molar Zn:Cd ratio in the sphalerite. In the temperate soil
268 (Figure 3c & d), comparison with the control shows that Zn was released from sphalerite
269 with increasing incubation time and, as with the Cd, the release curve did not exhibit a
270 change in slope over the last 180 days of the experiment. In the temperate control soils,
271 Zn concentrations from the beginning and the end of the experiment did not
272 significantly differ ($p = 0.05$).

273

274 The CaCl₂-extractable Zn concentrations in spiked sub-tropical soils were higher than in
275 the control soil (Figure 3c_i), most notably during the last 180 days of the experiment;
276 however the CaCl₂-extractable concentrations in control soils varied significantly ($p >$
277 0.05 , ANOVA) during incubation. The EDTA-extractable Zn concentrations in spiked and
278 control sub-tropical soils (Figure 3d_i) fluctuated throughout the incubation time and
279 there was no significant difference between concentrations at the beginning and end of
280 the experiment ($p < 0.05$, ANOVA). Comparison of the Zn release trends with those of Cd,
281 which were linear with incubation time, suggest that Zn released from the sphalerite
282 was in equilibrium with another solid phase and recalcitrant to EDTA complexation. The
283 net release indicated by CaCl₂-extractable concentrations (Figure 3c_i) was masked in the
284 EDTA-extractable data, as the EDTA extraction was not sensitive to minor variations in
285 Zn lability.

Dissolution rate, trends and limits

EDTA-extractable Cd concentrations were the most suitable indicator of the extent and rate of sphalerite dissolution, in this case meaning alteration from the original sulfide species. Unlike Zn, Cd release trends were clear and consistent (concentration vs. time) in both experimental soils. Additionally, Cd release is directly correlated with sphalerite dissolution rate (Stanton et al., 2008) and EDTA dissolves sphalerite oxidation products but not the sulfide itself (Rumball and Richmond, 1996), as confirmed by preliminary experiments (data not shown).

Net Cd release (Cd_{net}) from the sphalerite was calculated by subtracting EDTA-extractable concentrations obtained from control incubations from those obtained from the respective spiked incubations. The Cd_{net} data were used to estimate the percentage sphalerite dissolved at each incubation time point (Table 2). The relationship between Cd_{net} and incubation duration was linear ($R^2 \geq 0.96$) for both temperate and sub-tropical soils, indicating a constant rate of Cd release, and therefore sphalerite dissolution, throughout the experiments. Several studies on sphalerite dissolution in aqueous solution showed that dissolution rates decline during the initial few hundred hours of exposure and then attain an apparent steady state (Acero et al., 2007; Stanton et al., 2008; Weisener et al., 2003). It was proposed that this change is concurrent with the formation of Zn-deficient, polysulfide and elemental S product layers on sphalerite particles, and a shift from reaction rate-limited dissolution to dissolution limited by reagent diffusion (i.e. H_3O^+ , O_2 , Zn^{2+} , Cd^{2+} and/or S) through those product layers (Weisener et al., 2003). Acero et al. (2007) argued that, because steady state was attained, the layers were not passivating and initially high dissolution rates probably resulted from micro-crystals and oxidised phases on the pre-exposure sphalerite surfaces. In this study the slower dissolution rate remained constant over long durations (hundreds of days), regardless of whether these product layers are porous, and therefore not diffusion limiting, or whether they are in equilibrium with bulk solution, and therefore do not accumulate.

The data from this study are consistent with slow steady-state dissolution. The constant Cd release excludes the significant formation of stable secondary Cd phases, which

would have produced declining CaCl_2 and EDTA-extractable concentrations with increasing incubation duration by sequestering Cd^{2+} from the porewater. The absence of secondary phases was evidenced by SEM-EDS examination of sphalerite platelets exposed to field conditions for 2 years. Full details of this method are provided in Robson et al. (2013).

Dissolution rates observed at 365 days are indicative of the annual average, approximately 1 and $0.5 \mu\text{mol Cd g}^{-1} \text{ZnS a}^{-1}$ for the temperate and sub-tropical soils, respectively. Accounting for the reducing surface area of the given mass of dissolving sphalerite and assuming proportionality between dissolution rate and surface area (shrinking particle model) (Pradhan et al., 2010; Safari et al., 2009), the half-life of the sphalerite was estimated to be 50 and 94 years in the temperate and sub-tropical soils, respectively.

The slower dissolution rates observed in the sub-tropical soil were attributed to the prevailing soil pH. Based on kinetics data from Acero et al. (2007), a change in porewater pH from 5.53 (temperate soil) to 6.40 (sub-tropical soil) would result in dissolution rates being reduced by 66 %; therefore pH is likely to be the most significant factor affecting sphalerite dissolution in oxic soils. The shift to neutral pH observed in flooded sub-tropical soils (from pH 6.40) is associated with a further 53 % reduction in the dissolution rate, based on pH alone.

Cd uptake by crops

Uptake by wheat grown in temperate soil

Grain and stem Cd concentrations in the spring wheat grown in the sphalerite-spiked temperate soil were considerably higher (by a factor of ≥ 75) than in plants grown in the control soil (Figure 4). The wheat grain contained $29.0 \pm 3.3 \mu\text{mol Cd kg}^{-1}$, around 8 times higher than the international food safety limit of $3.6 \mu\text{mol kg}^{-1}$ (FAO/WHO, 2006).

[Approximate location for Figure 4.]

The data suggest that high Cd concentrations in grain produced from the spiked soil were proportional to the magnitude of the bioavailable Cd pool in that soil. Stem-to-grain transfer factors (TF) were the same for plants grown in spiked and control soil; therefore the translocation rate was independent of the phytoaccessible Cd concentration in the soil and the stem Cd concentration (Figure 4).

Stem bioconcentration factors (BCF), based on Cd_{net} values, for plants grown in spiked soils were 25 times higher than for those grown in control soils. The probable explanation for this observation is soil 'ageing'. Cd is generally regarded as exhibiting minimal ageing effect (Smolders and Mertens, 2013) but Hamon et al. (1998) demonstrated that around 1 % of soil Cd could be rendered unavailable for plant uptake per year of soil residence time. Ageing may have rendered the antecedent Cd in the soils far less phytoavailable than the Cd recently introduced by sphalerite dissolution.

Uptake by paddy rice grown in sub-tropical soil

Rice stem and grain Cd concentrations of plants grown in spiked soils were 3 – 4 times higher than in control soil plants (Figure 4). Although the plants were contaminated by the sphalerite, the edible tissue concentration ($0.597 \pm 0.019 \mu\text{mol Cd kg}^{-1}$) was well below applicable Chinese ($1.78 \mu\text{mol Cd kg}^{-1}$) and international ($3.56 \mu\text{mol Cd kg}^{-1}$) food safety limits (FAO/WHO, 2006; USDA Foreign Agriculture Service, 2010). For comparison, Cd concentrations in the wheat (spiked soil) were higher than those for rice by a factor of 49 in seeds and 24 in stems. Given that the rate of sphalerite dissolution in the experimental soils only varied by a factor of 2, the tissue concentrations illustrate significant differences in the Cd bioavailability and/or uptake behaviour in the rice and wheat soil-plant systems.

The data suggest that, all factors being equal, the rice had a propensity for Cd uptake similar to or greater than the wheat. In control soils, the wheat and rice stem concentrations were similar ($0.74 - 0.85 \mu\text{mol kg}^{-1} \text{Cd}$) and the rice stem BCF was much higher than for the wheat (Figure 4). Also, the rice TF increased in spiked soils (+ 53 %), indicating that the plants responded to higher Cd availability by enhancing stem-to-grain translocation. In light of this apparent propensity for uptake, the relatively low

rice tissue Cd concentrations suggest that decreased Cd availability in the paddy soil porewater limited uptake. This proposition is supported by CaCl_2 -extractable Cd concentrations that were below the detection limit (Figure 2 c), Cd_{net} concentrations that were reduced by 49 % (versus oxic incubation) and stem BCF values that were the same (6.36 – 6.40) in spiked and control soils (i.e. equal bioavailability).

Soil-to-rice Cd transfer was examined to determine if this could explain the depleted extractable Cd concentrations obtained from the sub-tropical soils (see Geochemical conditions in sub-tropical soil during rice cultivation). The plant roots were not analysed but their biomass is always much smaller than the stem biomass and therefore assuming equal contribution by the root and stems provided a conservative estimate (Kibria and Ahmed, 2006). Although the neutral soil pH and plant uptake might explain the depletion of CaCl_2 -extractable Cd, these factors cannot entirely explain the decreased Cd_{net} . Firstly the EDTA extraction, from which Cd_{net} is derived, would have been insensitive to the shift towards neutral soil pH. Secondly, after considering the estimated total Cd uptake by rice, a 33 % decrease in Cd_{net} still remained unaccounted for. Therefore it is likely that the formation of non-acid-volatile secondary sulfides (see Geochemical conditions in sub-tropical soil during rice cultivation) contributed to the low bioavailability and rice uptake of Cd in this study (de Livera et al., 2011).

Lowland rice is traditionally grown under near-constant standing water; however increasing global population and freshwater demand have catalysed the adoption of new agricultural practises such as 'system of rice intensification' (SRI), a set of management principles that discourage flooded agriculture (Africare et al., 2010; L. Zhao et al., 2010). A shift towards more oxic soil management will remove the protective biogeochemical conditions afforded by soil flooding and enhance the bioavailability of Cd in soils.

Conclusions

Sphalerite exhibits slow, steady dissolution behaviour in oxic agricultural soils developed under contrasting geoclimatic conditions and is accompanied by the release of the guest element Cd. Sphalerite contamination impacts soil quality for decades to

centuries, long after its introduction to soils ceases. The liberated Cd is highly bioavailable under oxic conditions, as indicated by *Triticum aestivum*, and has the potential to contaminate crops and pose a human health hazard. Data from *Oryza sativa* indicate that flooded rice production can limit these impacts by neutralising soil pH and possibly by providing sulfate-reducing conditions, under which secondary Cd sulfides can form. The recently publicised advantages of ending a reliance upon flooded agriculture (increased yields, reduced water consumption) suggests that growing rice under more oxic conditions will increase in popularity. Adopters of these new practices working Cd or sphalerite-impacted soil will sacrifice the protective biogeochemical conditions afforded by flooding, increasing the risk of producing contaminated food.

Acknowledgements

This research was funded by a Plymouth University PhD studentship and the Seale Hayne Educational Trust Fund. The authors would like to acknowledge OEA Laboratories Ltd, Kelly Bray, UK, for assistance with S analysis, and thank The University of Hong Kong Kadoorie Centre for supplying soil samples. We also thank the three anonymous reviewers, whose feedback helped us improve the manuscript.

References

- Acero, P., Cama, J., Ayora, C., 2007. Sphalerite dissolution kinetics in acidic environment. *Appl. Geochem.* 22, 1872–1883.
- Africare, Oxfam America, WWF-ICRISAT Project, 2010. More Rice for People, More Water for the Planet. WWF-ICRISAT Project, Hyderabad, India.
- Allen, H.E., Fu, G., Deng, B., 1993. Analysis of acid-volatile sulfide (AVS) and simultaneously extracted metals (SEM) for the estimation of potential toxicity in aquatic sediments. *Environ. Toxicol. Chem.* 12, 1441–1453.
- Carter, M.R., Gregorich, E.G., 2007. *Soil Sampling and Methods of Analysis*, 2nd ed. CRC Press, Boca Raton, USA.
- Castillo, S., de la Rosa, J.D., de la Campa, A.M.S., González-Castanedo, Y., Fernández-Caliani, J.C., Gonzalez, I., Romero, A., 2013. Contribution of mine wastes to atmospheric metal deposition in the surrounding area of an abandoned heavily polluted mining district (Rio Tinto Mines, Spain). *Sci. Total. Environ.* 449, 363–372.
- Cline, J.D., 1969. Spectrophotometric determination of hydrogen sulfide in natural waters. *Limnol. Oceanogr.* 14, 454–458.
- de Livera, J., McLaughlin, M.J., Hettiarachchi, G.M., Kirby, J.K., Beak, D.G., 2011. Cadmium solubility in paddy soils: effects of soil oxidation, metal sulfides and competitive

- ions. *Sci. Total. Environ.* 409, 1489–1497.
- FAO/WHO, 2006. Report of the 38th session of the Codex Committee on Food Additives and Contaminants. Food and Agriculture Organization of the United Nations & World Health Organization, The Hague, Netherlands.
- FAO/WHO, 2010. Summary Report of the Seventy-Third Meeting of Joint FAO/WHO Expert Committee on Food Additives. Food and Agriculture Organization of the United Nations & World Health Organization, The Hague, Netherlands.
- Hamon, R.E., McLaughlin, M.J., Naidu, R., Correll, R., 1998. Long-term changes in cadmium bioavailability in soil. *Environ. Sci. Technol.* 32, 3699–3703.
- Heidel, C., Tichomirowa, M., Breitkopf, C., 2011. Sphalerite oxidation pathways detected by oxygen and sulfur isotope studies. *Appl. Geochem.* 26, 2247–2259.
- Henson, M.C., Chedrese, P.J., 2004. Endocrine disruption by cadmium, a common environmental toxicant with paradoxical effects on reproduction. *Exp. Biol. Med.* (Maywood) 229, 383–392.
- Inglott, P.W., Reddy, K.R., Corstanje, R., 2005. Anaerobic soils, in: Hillel, D. (Ed.), *Encyclopedia of Soils in the Environment*. Academic Press Inc, Waltham MA, USA, pp. 71–78.
- Ishihara, T., Kobayashi, E., Okubo, Y., Suwazono, Y., Kido, T., Nishijyo, M., Nakagawa, H., Nogawa, K., 2001. Association between cadmium concentration in rice and mortality in the Jinzu River Basin, Japan. *Toxicology* 163, 23–28.
- Järup, L., 2003. Hazards of heavy metal contamination. *Br. Med. Bull.* 68, 167–182.
- Kibria, M.G., Ahmed, M.J., 2006. Cadmium and lead uptake by rice (*Oryza Sativa* L.) grown in three different textured soils. *Soil & Environ.* 25, 70–77.
- Kon, L.C., Durucan, S., Korre, A., 2007. The development and application of a wind erosion model for the assessment of fugitive dust emissions from mine tailings dumps. *Int. J. Min. Reclam. Environ.* 21, 198–218.
- Limei, Z., Xiaoyong, L., Tongbin, C., Xiulan, Y., Hua, X., Bin, W., Lixia, W., 2008. Regional assessment of cadmium pollution in agricultural lands and the potential health risk related to intensive mining activities: a case study in Chenzhou City, China. *J. Environ. Sci. (China)* 20, 696–703.
- Lo, I.M., Yang, X.Y., 1999. EDTA extraction of heavy metals from different soil fractions and synthetic soils. *Water Air Soil Pollut.* 109, 219–236.
- Lovley, D.R., Phillips, E.J.P., 1986. Availability of ferric iron for microbial reduction in bottom sediments of the freshwater tidal Potomac River. *Appl. Environ. Microbiol.* 52, 751–757.
- Magdoff, F.R., Bartlett, R.J., 1985. Soil and pH buffering revisited. *Soil Sci. Soc. Am. J.* 49, 145–148.
- Meers, E., Laing, Du, G., Unamuno, V., Ruttens, A., Vangronsveld, J., Tack, F.M.G., Verloo, M.G., 2007. Comparison of cadmium extractability from soils by commonly used single extraction protocols. *Geoderma* 141, 247–259.
- Mertens, J., Smolders, E., 2013. Zinc, in: Alloway, B.J. (Ed.), *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and Their Bioavailability*. Springer, Dordrecht, Netherlands, pp. 465–493.
- Miller, J.R., Hudson-Edwards, K.A., Lechler, P.J., Preston, D., Macklin, M.G., 2004. Heavy metal contamination of water, soil and produce within riverine communities of the Rio Pilcomayo Basin, Bolivia. *Sci. Total Environ.* 320, 189–209.
- Ponnamperuma, F.N., 1972. The chemistry of submerged soils. *Adv. Agron.* 24, 29–88.
- Pradhan, D., Kim, D.J., Chaudhury, G.R., Sohn, J.S., Lee, S.W., 2010. Dissolution kinetics of complex sulfides using acidophilic microorganisms. *Mater. Trans.* 51, 413–419.

- Robson, T.C., Braungardt, C.B., Keith-Roach, M.J., Rieuwerts, J.S., Worsfold, P.J., 2013. Impact of arsenopyrite contamination on agricultural soils and crops. *J. Geochem. Explor.* 125, 102–109.
- Rumball, J.A., Richmond, G.D., 1996. Measurement of oxidation in a base metal flotation circuit by selective leaching with EDTA. *Int. J. Miner. Process.* 48, 1–20.
- Safari, V., Arzpeyma, G., Rashchi, F., Mostoufi, N., 2009. A shrinking particle - shrinking core model for leaching of a zinc ore containing silica. *Int. J. Miner. Process.* 93, 79–83.
- Schecher, W., 2001. Thermochemical Data Used in MINEQL+ version 4.5. Environmental Research Software, Hallowell, UK.
- Scherer, W.H., 2009. Sulfur in soils. *J. Plant Nutr. Soil Sci.* 172, 326–335.
- Simmons, R.W., Pongsakul, P., Saiyasitpanich, D., Klinphoklap, S., 2005. Elevated levels of cadmium and zinc in paddy soils and elevated levels of cadmium in rice grain downstream of a zinc mineralized area in Thailand: implications for public health. *Environ. Geochem. Health* 27, 501–511.
- Smolders, E., Mertens, J., 2013. Cadmium, in: Alloway, B.J. (Ed.), *Heavy Metals in Soils: Trace Metals and Metalloids in Soils and Their Bioavailability*. Springer, Dordrecht, Netherlands, pp. 283–311.
- Stanton, M.R., Gemery-Hill, P.A., Shanks Iii, W.C., Taylor, C.D., 2008. Rates of zinc and trace metal release from dissolving sphalerite at pH 2.0–4.0. *Appl. Geochem.* 23, 136–147.
- United States Environmental Protection Agency, 2004. Test methods for evaluating solid waste, physical/chemical methods [online]. URL <http://www.epa.gov/epawaste/hazard/testmethods/sw846/online/index.htm> [accessed 10 Mar 2012].
- USDA Foreign Agriculture Service, 2010. GAIN report: National food safety standard - maximum levels of contaminants in food [online]. URL http://gain.fas.usda.gov/Recent%20GAIN%20Publications/National%20Food%20Safety%20Standard-Maximum%20Levels%20of%20Contaminants%20in%20Food_Beijing_China%20-%20Peoples%20Republic%20of_8-19-2010.pdf [accessed 24 Jan 2013].
- Weisener, C.G., Smart, R., Gerson, A.R., 2003. Kinetics and mechanisms of the leaching of low Fe sphalerite. *Geochim. Cosmochim. Acta* 67, 823–830.
- Zhao, H., Xia, B., Fan, C., Zhao, P., Shen, S., 2012. Human health risk from soil heavy metal contamination under different land uses near Dabaoshan mine, southern China. *Sci. Total. Environ.* 417–418, 45–54.
- Zhao, L., Wu, L., Li, Y., Animesh, S., Zhu, D., Uphoff, N., 2010. Comparisons of yield, water use efficiency, and soil microbial biomass as affected by the system of rice intensification. *Commun. Soil Sci. Plant Anal.* 41, 1–12.
- Zhuang, P., McBride, M., Xia, H., Li, N., Li, Z., 2009. Health risk from heavy metals via consumption of food crops in the vicinity of Dabaoshan mine, south China. *Sci. Total. Environ.* 407, 1551–1561.
- Zota, A.R., Willis, R., Jim, R., Norris, G.A., Shine, J.P., Duvall, R.M., Schaider, L.A., Spengler, J.D., 2009. Impact of mine waste on airborne respirable particulates in northeastern Oklahoma, United States. *J. Air Waste Manag. Assoc.* 59, 1347–1357.

542 Table 1: **Characterisation data for the temperate and sub-tropical experimental**
543 **soils.** Uncertainties reported as ± 1 standard deviation ($n = 5$). eCEC = Effective cation
544 exchange capacity; LOI = Organic matter content, determined by loss on ignition.

	Temperate soil	Sub-tropical soil
pH	6.58 ± 0.07	6.83 ± 0.12
eCEC (cmol ⁺ kg ⁻¹)	14.6 ± 0.3	13.1 ± 0.2
C _{organic} (% m/m)	5.57 ± 0.20	4.62 ± 0.06
C _{inorganic} (% m/m)	< LOD	0.47 ± 0.37
LOI (% m/m)	12.3 ± 0.5	13.1 ± 0.7
Al (mol kg ⁻¹)	1.44 ± 0.11	2.32 ± 0.11
Al _{oxalate} (mol kg ⁻¹)	0.121 ± 0.001	0.0975 ± 0.0018
Fe (mol kg ⁻¹)	0.973 ± 0.018	1.13 ± 0.03
Fe _{oxalate} (mol kg ⁻¹)	0.202 ± 0.002	0.0602 ± 0.0014
Mn (mmol kg ⁻¹)	37.6 ± 3.2	5.25 ± 0.31
S (mmol kg ⁻¹)	17.0 ± 2.2	17.1 ± 1.7
Cd (μ mol kg ⁻¹)	2.70 ± 0.55	2.67 ± 0.27
Zn (mmol kg ⁻¹)	2.05 ± 0.14	2.13 ± 0.17

545

546

Table 2: **Net Cd release (Cd_{net}) and percentage sphalerite dissolution** determined after 7 – 365 days incubation in both temperate and sub-tropical soils. Uncertainties are reported as ± 1 standard deviation ($n = 3$). Asterisks indicate insignificant differences between the spiked and control incubation values.

Days	Temperate soil		Sub-tropical soil	
	Cd_{net} (nmol Cd g⁻¹ ZnS)	% dissolution	Cd_{net} (nmol Cd g⁻¹ ZnS)	% dissolution
7	18.2 $\pm$ 15.4	0.02 $\pm$ 0.02	*	*
30	76.9 $\pm$ 21.1	0.09 $\pm$ 0.03	73.6 $\pm$ 57	0.09 $\pm$ 0.07
90	261 $\pm$ 38	0.32 $\pm$ 0.05	148 $\pm$ 24	0.18 $\pm$ 0.03
180	475 $\pm$ 223	0.58 $\pm$ 0.27	228 $\pm$ 24	0.28 $\pm$ 0.03
270	756 $\pm$ 60	0.93 $\pm$ 0.07	425 $\pm$ 18	0.52 $\pm$ 0.02
365	998 $\pm$ 212	1.23 $\pm$ 0.26	464 $\pm$ 27	0.57 $\pm$ 0.03

Figure 1: **Influence of microbiota upon soil pH:** Sulfate, pH (a) and nitrate (b) in biotic and abiotic control incubations of the temperate soil. Uncertainties are reported as ± 1 standard deviation ($n = 3$).

Figure 2: **Redox indicators and metal availability in flooded paddy soils:** (a) SO_4^{2-} and NO_3^- , (b) pH and Eh and (c) CaCl_2 -extractable Zn/Cd in the sub-tropical soil (180-365 days), under both oxic (filled symbols) and anoxic (hollow symbols) conditions. Uncertainties are reported as ± 1 standard deviation ($n = 3$).

Figure 3: **Cd and Zn release during sphalerite weathering:** CaCl_2 -extractable and EDTA-extractable Cd (a/b_{I-II}) and Zn (a/b_{III-IV}) concentrations in temperate (a_{I-IV}) and sub-tropical (b_{I-IV}) experimental soils. Uncertainties are reported as ± 1 standard deviation ($n = 3$). Note that all Cd concentrations were below the detection limit ($0.002 \mu\text{mol Cd kg}^{-1}$) until day 180 of sub-tropical soil incubation (b_I). Asterisks denote statistically significant ($p > 0.05$, ANOVA) differences between spiked and control soils.

Figure 4: **Plant uptake of cadmium:** total Cd tissue concentrations, stem bioconcentration factors (BCF) based upon Cd_{net} concentrations and stem-to-grain transfer factors (TF) for spring wheat grown in the temperate experimental soil and rice in the flooded sub-tropical experimental soil. Uncertainties based on ± 1 standard deviation.

571 Table S1: Certified and determined concentrations obtained for certified reference
572 materials. Uncertainties are reported as ± 1 standard deviation ($n = 5$).

Χερτιφιεδ ματεριαλ	Παραμετεροσ	Χερτιφιεδ	Δετερμινεδ
IRMM 804 Rice flour	Cd ($\mu\text{mol kg}^{-1}$)	14.3 ± 0.6	14.3 ± 1.1
	Zn ($\mu\text{mol kg}^{-1}$)	353 ± 29	360 ± 32
BCR 320R Channel Sediment	Cd ($\mu\text{mol kg}^{-1}$)	23.5 ± 1.6	21.1 ± 0.2
	Zn (mmol kg^{-1})	4.88 ± 0.31	4.85 ± 0.19

573

574

575

576 Figure S1: **XRD characterisation:** X-ray diffractogram obtained for the experimental
577 sphalerite used in this study, plotted together with an exemplar pattern for wurtzite.

578

579

580 Figure S2: **Soil pH buffering curves for temperate and sub-tropical experimental**
581 **soils:** The curves were produced by adding variable concentrations of H_2SO_4 (x-axis) to
582 the soils and determining slurry pH after overnight equilibration (y-axis).

583

590