2016-04-01

'Something normal in a very, very abnormal environment' Nursing work to honour the life of dying infants and children in neonatal and paediatric intensive care in Australia

Bloomer, MJ

<http://hdl.handle.net/10026.1/4966>

10.1016/j.iccn.2015.09.001

Intensive and Critical Care Nursing

Elsevier BV

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1 'Something normal in a very, very abnormal environment' – nursing work to honour the life of dying
2 infants and children in neonatal and paediatric ICU

3 Melissa J Bloomer ^a (Primary/Corresponding Author) Melissa.bloomer@monash.edu

4 Ruth Endacott ^b Ruth.endacott@monash.edu

5 Beverley Copnell ^b Beverley.copnell@monash.edu

6 Margaret O'Connor ^a Margaret.oconnor@monash.edu

7
8 ^a Monash University, School of Nursing and Midwifery, Building E, PO Box 527, Frankston , VIC 3199,
9 Australia

10 ^b Monash University, School of Nursing and Midwifery, 10 Chancellors Walk, Clayton Campus,
11 Clayton, VIC, 3800, Australia

12
13 This is an accepted manuscript of an article published by Elsevier in Intensive and Critical Care
14 Nursing in April 2016 available at: <http://dx.doi.org/10.1016/j.iccn.2015.09.001>

Abstract

The majority of deaths of children and infants occur in paediatric and neonatal intensive care settings. For nurses, managing an infant/child's deterioration and death can be very challenging. Nurses play a vital role in how the death occurs, how families are supported leading up to and after the infant/child's death. This paper describes the nurses' endeavours to create normality amidst the sadness and grief of the death of a child in paediatric and neonatal ICU. Focus groups and individual interviews with registered nurses from NICU and PICU settings gathered data on how neonatal and paediatric intensive care nurses care for families when a child dies and how they perceived their ability and preparedness to provide family care. Four themes emerged from thematic analysis: (1) respecting the child as a person; (2) creating opportunities for family involvement/connection; (3) collecting mementos; and (4) planning for death. Many of the activities described in this study empowered parents to participate in the care of their child as death approached. Further work is required to ensure these principles are translated into practice.

Keywords

Neonatal intensive care unit (NICU)
Paediatric intensive care unit (PICU)
Paediatric nursing
End of life care
Death
Dying
Rituals

Implications for Clinical Practice*

- Pediatric and neonatal intensive care nurses need specialist skills to be able to support families and honour the dying child.

- Creating opportunities for families to interact with the dying child, and inform the death are essential.
- There is scope for neonatal and pediatric intensive care nurses to go beyond the ‘routine’ collection of mementos as a way of supporting grieving families.

Background

The primary goal of therapy in intensive care units (ICUs) is to assist the recovery of seriously ill patients through technologically sophisticated care (Copnell, 2005; Hamric & Blackhall, 2007).

Despite this goal, mortality is higher than most other health care settings (Yu & Chan, 2010); this is no less for paediatric and neonatal intensive care (PICU and NICU), as the majority of in-hospital deaths of children and infants take place in these settings (Chang et al., 2013; Stark et al., 2008; Yorke, 2011).

For parents, the death of an infant or child is regarded as one of the most devastating events endured (Davies, 2004). Not only is parental grief intense, but it can also be overwhelming because they lose a part of themselves and the hopes and dreams they had for the child (Davies, 2001; Davies, 2004). Given that the majority of children who die are hospitalised at the time of their death (Chang, et al., 2013), the experience and grief of parents associated with the loss of their child is influenced by interactions with, and actions of, various clinicians involved in their child’s care (Butler et al., 2015; Meert et al., 2001).

Deterioration in a child’s condition and/or their death can also result in personal grief and a sense of failure for nurses, especially if they are primary carer /nurse within the care team (Hamric & Blackhall, 2007; Yam et al., 2001). Consequently, nurses often attempt to rationalise a death as a way of coping. But rationalising the death of an infant or child is significantly harder than an adult

because society regards the death of a young person as an 'undeserved death' (O'Connor et al., 2010).

When death occurs in NICU and PICU, nursing care of the family is equally as challenging as caring for the dying infant or child in the time leading up to and after death. Nurses play a role in what happens as death approaches, how the death occurs and how the family are engaged after death. Nurses express concerns about knowing what to say to parents, how to prepare them for imminent death and how to support parental grief (Ahern, 2013).

Numerous studies have identified other factors that impact upon the parental experience. Caring actions, sensitive and caring staff and understandable explanations were cited in one study as being most helpful to parents (Brooten et al., 2012). Similarly, other studies have stressed the importance of communication between clinicians and parents (Darbyshire et al., 2013; Meert, et al., 2001; Yorke, 2011), with another paper stating that families relied more heavily of nursing professionals than other family members (Beckstrand et al., 2009). Importantly however, when clinicians demonstrate caring in other ways, which go above and beyond the professional role (Meyer et al., 2006), the parental experience is also improved. A recent PICU study found that measures such as making the PICU environment more comfortable and welcoming, minimising the imposing nature of the technology, and facilitating parents' time at the bedside by giving them access to a telephone and hospital meals were identified by bereaved parents as strategies to improve their experience (Yorke, 2011).

Parents also appreciate the rituals and practices designed to mark the death of an infant/child and create memories (Butler, et al., in press). The making of memories is increasingly recognised as an aid in parental coping and grieving (Schott et al., 2007). Often nurse initiated, they include keepsakes such as a lock of hair, hand and foot prints, memory boxes and hospital ID bands (Abib El Halal et al., 2013; Brooten, et al., 2013; Meert et al., 2009; Meyer, et al., 2006). These keepsakes, also referred to in the literature as mementos, were described as helpful for parents in their grief (Brooten, et al.,

2013). Photographs of the infant/child before and after death are also common and are said to provide both memory and meaning to the parents' experience of their child after the loss (Heartfelt, n.d.; Michelson et al., 2013; Yorke, 2011).

Other studies report that rather than physical keepsakes, parents and families instead appreciated other actions/gestures by clinicians that demonstrated that the infant/child was important and special and respectful of the parent and family's needs (McGuinness et al., 2014; Meert, et al., 2009). This included having time alone with the infant/child and being encouraged and supported to provide care to the infant/child leading up to the death (Brooten, et al., 2013; Meert, et al., 2009; Yorke, 2011).

Some studies of parental experiences indicated that activities which they valued were not universally performed, for reasons which are unclear. Parents reported not being allowed to be with their child either during or after death (Neidig & Dalgas-Pelish, 1991), a lack of privacy while their child was dying (Abib El Halal, et al., 2013), and not receiving desired mementos, an outcome justified against miscommunication between families and clinicians (Widger & Picot, 2008).

This paper reports on one aspect of a larger study that examined nurses' experiences of caring for families when a child dies, and their preparedness to provide family centred end-of-life care. We describe the nurses' endeavours to create memories for families and facilitating families' interactions with the dying infant/child amidst the sadness and grief of the death of a child in paediatric and neonatal ICU.

Method

Focus groups and individual interviews were utilised to gather data on how NICU and PICU nurses care for families when a child dies and how they perceived their ability and preparedness to provide family care. Thematic analysis provided a framework to elicit themes which highlighted the actions taken by nurses to create a sense of normality around the death.

Setting and Sample

The study was conducted in two NICUs and two PICUs located in two Australian metropolitan hospitals. Convenience sampling was used in this study. Details of the study were forwarded to NICU and PICU nursing staff via email, following contact with Nurse Unit Managers. Potential participants were asked to determine their own eligibility according to the inclusion criteria. Twenty-one registered nurses from NICU (N=8) and PICU (N=13) settings participated. Participants were eligible for inclusion if they were employed as a Registered Nurse (RN), had at least one year of nursing experience in either NICU or PICU and had cared for at least one child who had died in the NICU or PICU.

Interested nursing staff were asked to make contact with the researchers to register their interest and interview times and locations were negotiated following dissemination of an Explanatory Statement. At the time of meeting with the potential participants, the research and the Explanatory Statement were discussed and all questions answered. Written consent was then gained prior to commencement of individual and focus group interviews.

Data Collection

Focus groups (defined in this study as an interview of two or more participants) and individual interviews both utilised a semi-structured conversational approach, facilitated by the researcher to encourage communication and interaction amongst participants (Liamputtong, 2009). For those unavailable for a focus group, individual interviews were conducted using the same semi-structured conversational approach used in the focus groups.

From the four participating units, six focus group interviews were conducted, with between two and four participants in each focus group, and four nurses were interviewed individually, totalling 21 participants. The type of interview was determined mostly by the availability of willing participants.

Where possible, participants were grouped together to enable a focus group interview, but where scheduling did not allow this, participants were instead offered individual interviews. All interviews (focus group and individual) were conducted between November 2012 and January 2013 in a private, quiet space located in the hospitals. Interviews lasted between 17 and 50 minutes. They were audio-taped and transcribed verbatim for analysis. No field notes were made during the interviews.

Data Analysis

Inductive content analysis was used to analyse the data because it is well-suited to analysing multifaceted and sensitive phenomena that are characteristic in nursing (Elo & Kyngas, 2008; Moretti et al., 2011). Inductive content analysis, in which coding categories are derived directly and inductively from the raw data, is an approach to qualitative analysis that attempts to limit, as much as possible the researchers' subjective interpretations (Moretti, et al., 2011). The process involved two researchers independently reading and re-reading the interview manuscripts to gain a sense of the whole and highlighting phrases that appear to capture the essence of the interviews; taking notes of the content; 112 grouping content together into categories; revising the categories and organising and/or linking the categories and subcategories until the themes were determined (Elo & Kyngas, 2008; Moretti, et al., 2011).

This process of analysis was made more robust by the development of a study protocol, in which all procedures were described in detail and established prior to data analysis, assuring consistency and transparency and that the themes developed out of the raw data rather than any pre-conceived themes (Moretti, et al., 2011). Once both researchers had determined the final themes, the findings were shared amongst the team and the analysis further refined until the final themes were settled.

Ethical Considerations

Whilst this study gained ethical approval as 'low risk', given the highly emotive nature of the work of neonatal and paediatric intensive care nurses in providing end of life care, the authors considered it important to safeguard the welfare of nurse participants. An employee counselling service was available to support any participants who became distressed as a result of their participation in this study, however this was not necessary. No patient or family data was collected as part of this research. Furthermore, In order to protect the anonymity of participants, once eligibility of individuals was determined, no other demographic data was collected. The intent was to explore the experience and emotion as reflected by the participants at that time, irrespective of these other factors.

Results

Six focus groups and four individual interviews were conducted across the four units (NICU x2 and PICU x 2). For the purposes of reporting participant voices, quotes have been anonymised and instead are labelled according to the type of interview (individual or focus group), interview number/transcript page number and whether the interview involved PICU or NICU staff.

Participants expressed that their role and responsibilities change as death neared. When it was evident that death was likely, participants said that the focus of their role shifted to place more emphasis on the family: *'It's still caring for the patient, but maybe a little – yes, a little bit more split between for the family and the patient'* (FG Int 5/10, PICU).

Inductive analysis identified four key themes that demonstrated how participants worked to ensure families were left with heart-warming memories of their child following their death. The key themes derived from the analysis were: (1) respecting the child as a person; (2) creating opportunities for family involvement/connection; (3) collecting mementos; and (4) planning for death.

1. Respecting the child as a person

193 Participants expressed the need to acknowledge the child as a person and demonstrate respect for
194 their personhood. This was achieved largely by consideration of their physical appearance.
195 Participants expressed that rather than just using their technical expertise and skill, it was important
196 to consider how the child was presented to the parents and family:

197 *'We are very pedantic sometimes in the way you sort of – well, not just care for them,*
198 *but their physical appearance as well...we had braided her hair and things like*
199 *that...and they're so excited that somebody has taken some time and some care'* (FG
200 Int 155 3/5, PICU).

201

202 Participants saw this as a way of keeping 156 the parental and family focus on the child, rather than
203 the technological interventions surrounding them:

204 *'...to actually have, to look at how their child is presented and not just the physical*
205 *[interventions] that we're doing, but you know, their outward appearance. So it's not*
206 *just the machines...'* (FG Int 3/6, PICU)

207 *'lots of them [parents] have never held them without tubes, never seen their faces*
208 *without stuff stuck on them...'* (FG Int 2/8, NICU).

209 Another participant explained their role not just in preparing the baby, but also in presenting a
210 respectful environment:

211 *'...we can get the baby into some sort of fit state of presentation....Because*
212 *sometimes the babies really do look like they've been in a war zone by the time*
213 *they're dying, and so it's pretty important to make sure the baby is clean and that the*
214 *environment is clean, and all the extraneous clutter is no longer there....and*
215 *redressing the baby in clothing that they may choose...'* (FG Int 3/10, PICU).

216

In circumstances where changing and grooming the child was not possible due to the fragile nature of the child's critical illness, participants worked hard to emphasise the positives in the infant/child in communication with their parents and family:

'I talk about the positive features of the nice look of the baby. I find whatever features are looking sort of attractive...whether it's his skin or hair – sometimes they have a nice head of hair, or whatever it is. I sort of like to talk about what's positive about the baby...it's an honouring of their life, it's more of a recognition. Like this was a human being that was – you know, had lots of perfection about it too' (FG Int 10/4, NICU).

2. Creating Opportunities

A second theme arising from the interview data was the imperative to create opportunities for parents and families to be with and spend time with their child leading up to and after their death, in ways that that they previously may not have been able to:

'...I want to make this experience the best it can be, so you can have the nice memories and that side of things...because it's their baby and their child and their dreams for the future' (FG Int 2/5, NICU).

'[One parent said] "I've never had a chance to lay down with my baby until it died". You can put a baby on a bed and you can put a mother or father on that bed. It doesn't matter, but let them be – so something normal in a very, very abnormal environment' (FG Int 3/10, PICU).

Despite the fragility of the critically ill infant/child, the importance of needing to make opportunities for parents to have time with them was not forgotten, and often required significant commitment by the staff:

'I like to make sure that the parents have a lot of time holding their baby – so even if the child has been really critically ill and not moved much, I may call colleagues in to

244 *help me, make sure I can safely move that baby off its bed ... and into the parents'*
245 *arms, and that can just take hours'* (FG Int 10/2, NICU).

246

247 Another participant spoke of how she listened to families and the needs they express, through
248 including extended family such as siblings and grandparents into the experiences she facilitated:

249

250 *'A lot of the families, you sort of ... 199 get led by them. If they want their children to*
251 *be involved, then you obviously deal with that and sort of, you know, try and make it*
252 *as normal - I know it's not a normal situation, but try and make it as normal for them*
253 *as possible; still sort of refer to the baby as their brother or sister, or you know, try*
254 *and get them involved if we're giving the baby a wash and things like that'* (Individual
255 Int 9/13, NICU).

256

257 **3. Collecting mementos**

258 Participants discussed at length, the importance of collecting mementos for families prior to and
259 after a death. 'Memory boxes' were used as a way of enabling the collection of mementos and
260 trinkets that may be appreciated by the family after death:- *'every time I find something that I*
261 *think they might like, I just put it all in a book for them'* (FG Int 10/16, NICU).

262

263 Collecting handprints and taking photos seemed a common way to connect family members to their
264 infant. Participants suggested that these were ways to prepare families for impending death:

265 *'...because they're helping families and the siblings put the handprints together, and*
266 *put all the fingerprints together...I was really sort of helping and getting the family a*
267 *little bit more prepared for what was going to happen..."* (Individual Int 4/5, PICU)

268

269 *'Photographs, like it's important to try and get [photographs]. I try and get [volunteer*

photography service] *in if we've got time and preparation*' (FG Int 10/16, NICU).

4. Providing an appropriate environment for the death

As well as creating opportunities for families to spend time with their critically ill infant or child when death was approaching, participants recounted stories of the lengths taken to give the parents an opportunity to have an input into planning for an eventual death. Participants stressed that any infant/child death is a life cut too short, and thus any way in which they could accommodate the wishes of the family were prioritised. For some, this involved creating an environment that was less clinical and technical than a standard ICU environment:

'We have a room...there's a bed in there...they took the baby in with them, and everybody stayed in there overnight for quite a few hours...then in the next day there was some back and forwards with bringing the baby back up from the mortuary so anyone else who need to come to say goodbye did so...' (FG Int 10/2, NICU).

In other cases, in fulfilling the parents' wishes, death occurred outside of the hospital, facilitated and supported by ICU staff:

'We did have a baby years ago who died travelling down the river on a boat cruise with her twin, and it was expected she would die, so that was where she died, and her parents were really happy with that...' (Individual Int 9/9, NICU).

Another participant described another out of hospital death, as requested by the family and facilitated by ICU staff:

'we recently had a death at the [animal park]. Our consultant went out of the way to provide what the parents wanted, and they were very focussed on animals and, yes, they took him to see them' (FG Int 10/21, NICU).

296

297 This same child's death was also brought up by another participant, who recalled the significance of
298 the experience, not just for the family, but also the staff:

299 *'Withdrawal of care happened in a quiet place in the [animal park]...we were*
300 *absolutely amazed that it was possible to do that. I think the [animal park] was*
301 *something important to them. It was something that meant a lot to them, and*
302 *therefore, they wanted their child to see the animals, to go to the [animal park] with*
303 *them, and he did'* (Individual Int 9/9, NICU).

304

305 Data analysis demonstrated the lengths to which NICU and PICU nurses will go, to facilitate family
306 care as their child approached the end of their life, which goes beyond what might be evident or
307 feasible in other ICU environments.

308

309 **Discussion**

310 In this study, it became clear that the focus group interviews provided much richer data than the
311 individual interviews. This is thought to be attributed to the synergistic nature of focus groups, in
312 which a stronger social context is created (Finch & Lewis, 2003), resulting in nurse participants
313 engaging more deeply with each other, most likely because of their shared experiences. When this
314 occurs, they are less influenced by the external researcher.

315

316 Several studies have detailed the needs of parents and families inside the ICU including consistent
317 staff presence and expressions of sympathy and caring (Latour, 2008; Meert, et al., 2001). Through
318 respecting the child as a person and emphasising the child as an individual, the nurses in our study
319 were demonstrating the child's importance to the family. Other studies also highlight the need for
320 adequate information and coordinated care (Meert, et al., 2001; Meyer, et al., 2006) and
321 preservation of the preservation of the parent-child relationship leading up to death (Meyer, et al.,

2006). Similarly this study reveals the lengths to which NICU and PICU nurses went to to create opportunities for parental and extended family involvement in care.

Existing research literature has highlighted the potential of early recognition of a life-limiting illness and advance care planning as a way of improving the care of paediatric and neonatal patients and their families (Mitchell & Dale, 2015). Other research has demonstrated that, where available, specialist paediatric palliative care services can allow for end of life care to be provided at home, rather than in hospital (Chang, et al., 2013). Transferring the child home however is not always an option, but what is important is privacy leading up to a child's death (Latour, 2008).

The themes that emerged from this study highlight the steps taken by NICU and PICU nurses to work with parents to honour the life of the dying child. Activities described within all the themes have been identified by parents as being helpful to them if performed well (Butler, et al., 2015). But to our knowledge, no other studies have detailed instances where families and clinicians have worked together to create special memories through the place of death. This study highlighted one example where families and clinicians were able to facilitate a death at a specially chosen place outside of the ICU and greater hospital, in this case an animal park; a venue chosen by the family for its special significance in relation to the dying infant.

The nurses in our study did not detail the conversations behind facilitating a death outside of the ICU; that is, it was not clear whether such requests were always initiated by parents, on the suggestion of clinicians, or as a collaborative idea that evolved between parents and staff. They also did not indicate whether parental requests were ever refused. Going to such lengths to facilitate a child's death at a specific location, with the express purpose of creating memories for the families, significantly contribute to the complexity of end of life care in PICU and NICU. What it also demonstrated, similar to the work of Meert et al (2001), was the desire to preserve the parent-child relationship by allowing parents to make at least some of the decisions in their child's last moments.

Despite the positive impact of such actions as demonstrated in this study, there are scant other examples in the research literature. We recommend that actions like this designed to facilitate family coping and create memories be included in future studies, particularly prospective studies of management of dying infants and children, which have hitherto taken a mechanistic approach to the topic (Copnell, 2005).

Many of the activities described in this study empowered parents to participate in the care of their child as death approached. These findings resonate with the body of work undertaken by Latour and colleagues in the Netherlands (Latour, 2008; Latour et al., 2012; Latour, 2011).

The empowerment of parents has developed over time in NICU (Davis et al., 2003; Latour, et al., 2012) and PICU (Sturdivant & Warren, 2009). However, whilst family-centred care in NICU has been reported to have benefits for parental stress, comfort and confidence (Cooper et al., 2007), the principles are not always translated into practice (Petersen et al., 2004). Similarly, studies in PICU indicate deficits in the provision of family centred care (Butler, et al., 2015).

Limitations

This study was undertaken in two hospitals in one city in Australia. The findings cannot be assumed to be generalizable to all PICU and NICU settings. Demographic data, such as age, sex, qualification and experience of participants might have provided further insights in this study; however, the four themes were evident across all interviews (paediatric and neonatal), indicating that the demography of the participants may have had little effect.

Conclusion

This paper has described nursing work to create memories for families amidst the sadness and grief of the death of a child in PICU and NICU. The themes illustrated nurses' concerns for family members in preparing for the death, how to present the child to their parents in the best way, and the enabling of opportunities for family involvement including the creation of mementos. Many activities derived from the literature and described in this study were seen to empower parents' participation in caring for their child both at the end of their life and after the death. Further work is required to ensure these principles are translated into practice.

Funding

This work was supported by a Nurses Board of Victoria Legacy Limited Major Research Grant.

References

- Abib El Halal, G., Piva, J., Lago, P., El Halal, M., Cabral, F., Nilson, C., et al. (2013). Parents' perspectives on the deaths of their children in two Brazilian paediatric intensive care units. *International Journal of Palliative Nursing*, 19(10), 495-502.
- Ahern, K. (2013). What neonatal intensive care nurses need to know about neonatal palliative care. *Advances in Neonatal Care* 13(2), 108-114.
- Beckstrand, R., Rawle, N., LCallister, L., & Mandleco, B. (2009). Pediatric nurses' perceptions of obstacles and supportive behaviours in end-of-life care. *American Journal of Critical Care*, 19(6), 543-552.
- Bloomer, M., O'Connor, M., Copnell, B., & Endacott, B. (2015). Nursing care for the families of the dying child/infant in paediatric and neonatal ICU: Nurses' emotional talk and sources of discomfort. A mixed methods study. *Australian Critical Care*. doi: 10.1016/j.aucc.2015.01.002
- Brooten, D., Youngblut, J. M., Seagrave, L., Caicedo, C., Hawthorne, D., Hidalgo, I., et al. (2013). Parent's Perceptions of Health Care Providers Actions Around Child ICU Death: What Helped, What Did Not. *American Journal of Hospice and Palliative Medicine*, 30(1), 40-49. doi: 10.1177/1049909112444301
- Butler, A., Hall, H., Willetts, G., & Copnell, B. (in press). Parents' experiences of healthcare provider actions when their child dies: An integrative review of the literature. *Journal for Specialists in Pediatric Nursing*. doi: 10.1111/jspn.12097
- Chang, E., Macleod, R., & Drake, R. (2013). Characteristics influencing location of death for children with life-limiting illness. *Archives of Disease in Childhood*, 98(6), 419-424.

411 Cooper, L., Gooding, J., Gallagher, J., Sternesky, L., Lesdsky, R., 343 & Berns, S. (2007). Impact of a
 412 family-centred care initiative on NICU care, staff and families. *Journal of Perinatology*,
 413 27(S32-S37).

414
 415 Copnell, B. (2005). Death in the pediatric ICU: Caring for the children and families at the end of life.
 416 *Critical Care Nursing Clinics of North America* 17(4), 349-360.

417 Darbyshire, P., Cleghorn, A., Downes, M., Elford, J., Gannoni, A., McCullagh, C., et al. (2013).
 418 Supporting bereaved parents: A phenomenological study of a telephone intervention
 419 programme in a paediatric oncology unit. *Journal of Clinical Nursing*, 22(3-4), 540-549.

420 Davies, A. (2001). Children's nurisng. Death of adolescents: Parental grief and coping strategies.
 421 *British Journal of Nursing*, 10(20), 1332.

422 Davies, R. (2004). New understandings of parental grief: Literature review. *Journal of Advanced*
 423 *Nursing*, 46(5), 506-513.

424 Davis, L., Mohay, H., & Edwards, H. (2003). Mothers' involvement in caring for their premature
 425 infants: An historical overview. *Journal of Advanced Nursing*, 42(6), 578-586.

426

427 Elo, S., & Kyngas, H. (2008). The qualitative content analysis process. *JAN*, 62(1), 107-115.

428

429 Finch, H., & Lewis, J. (2003). Focus Groups. In R. J & J. Lewis (Eds.), *Qualitative Research Practice:*
 430 *A Guide for Social Science Students and Researchers*. London: Sage Publications.

431

432 Hamric, A., & Blackhall, L. (2007). Nurse-physician perspectives on the care of dying patients in
 433 intensive care units: Collaboration, moral distress and ethical climate. *Critical Care Medicine*,
 434 35(2), 422-429.

435

436 Heartfelt. (n.d.). Heartfelt: Giving the gift of photographic memories Retrieved 13 Jan, 2015, from
 437 <https://www.heartfelt.org.au/>

438 Latour, J. (2008). Parental needs during end-of-life care: one room for improvement. *Pediatric*
 439 *Critical Care Medicine*, 9(6), 661-662.

440 Latour, J., Duivenvoorden, H., Hazelet, J., & van Goudoever, J. (2012). Development and validation of
 441 a neonatal intensive care parent satisfaction instrument. *Pediatric Critical Care Medicine*,
 442 13(5), 554-559.

443 Latour, J. M. (2011). *Empowerment of parents in the intensive care*. Doctor of Philosophy,
 444 Erasums University Rotterdam, Rotterdam.

445 Liamputtong, P. (2009). *Qualitative research methods* (3rd ed.). South Melbourne: Oxford University
 446 Press.

447 McGuinness, D., Coughlan, B., & Power, S. (2014). Empty arms: Supporting bereaved mothers during
 448 the immediate postnatal period. *British Journal of Midwifery*, 22(4), 246-252.

449 Meert, K., Briller, S., Schim, S., Thurston, C., & Kabel, A. (2009). Examining the needs of bereaved
 450 parents in the pediatric intensive care unit: A qualitative study. *Death Studies*, 33(8), 712-
 451 740.

452 Meert, K., Thurston, C., & Thomas, R. (2001). Parental coping and bereavement outcome after the
 453 death of a child in the pediatric intensive care unit. *Pediatric Critical Care Medicine*, 2(2),
 454 324-328.

455 Meyer, E., Ritholz, M., Burns, J., & Truog, R. (2006). Improving the quality of end-of-life care in the
 456 pediatric intensive care unit: Parents' priorities and recommendations. *Pediatrics*, 117(3),
 457 649-657.

458 Michelson, K., Blehart, K., Hochberg, T., & James, K. (2013). Bereavement photography for children:
 459 Program development and health care professionals' response. *Death Studies*, 37(6), 513-
 460 528.

461 O'Connor, M., Davis, M., & Abernathy, A. (2010). Language, discourse and meaning in palliative
 462 medicine. *Progress in Palliative Care*, 18(2), 66-71.

463 Schott, J., Henley, A., & Kohner, N. (2007). *Pregnancy loss and the death of a baby: guidelines for*
 464 *professionals* (3rd ed.). London: SANDS.

465 Stark, Z., Hynson, J., & Forrester, M. (2008). Discussing withholding and withdrawing life-sustaining
 466 medical treatment in paediatric inpatients: Audit of current practice. *Paediatric Child Health*,
 467 44(7-8), 399-403.

468 Stolberg, J. (2011). Leaving footprints on our hearts - how can midwives provide meaningful
 469 emotional support after a perinatal death? Part 2. *Midwifery Digest*, 21(2), 235-241.

470 Truog, R., Campbell, M., Curtis, J., Haas, C., JM, L., Rubenfeld, G., et al. (2008). Recommendations for
 471 end-of-life care in the intensive care unit: A consensus statement by the American College of
 472 Critical Care Medicine. *Critical Care Medicine*, 36(3), 953-963.

473 Yorke, D. (2011). Parents' memories of haivng a child die in the PICU. *The World of Critical Care*
 474 *Nursing*, 8(3), 97-102.

475 Yu, H., & Chan, S. (2010). Nurses' Response to Death and Dying in an Intensive Care Unit: A
 476 Qualitative Study. *Journal of Clinical Nursing*, 19, 1167-1169.

477