2016-05

The Use of a Dynamic Elastomeric Fabric Orthosis in Supporting the Management of Athletic Pelvic and Groin Injury

Sawle, L

<http://hdl.handle.net/10026.1/5486>

10.1123/jsr.2014-0266

Journal of Sport Rehabilitation

Human Kinetics

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

This is an accepted article published by the Journal of Sport Rehabilitation available at

<http://journals.humankinetics.com/jsr-current-issue/jsr-volume-25-issue-2-may-/the-use-of-a-dynamic-elastomeric-fabric-orthosis-in-supporting-the-management-of-athletic-pelvic-and-groin-injury>

DOI [10-1123/jsr.2014-0266](https://doi.org/10.1123/jsr.2014-0266)

Title Evaluating a dynamic elastomeric fabric orthosis developed to aid in the management of athletic pelvic pain

Author names and affiliations Leanne Sawle^{ab}, Jennifer Freeman^b, Jonathan Marsden^b, Martin Matthews^a

^aDM Orthotics. Unit 2 Cardrew Way, Cardrew Industrial Estate, Redruth, Cornwall TR15 1SH

^bPlymouth University. Faculty of Health, Education and Society. Peninsula Allied Health Centre, Derriford Road, Plymouth, Devon PL6 8BH

Corresponding author Leanne Sawle. DM Orthotics. Unit 2 Cardrew Way, Cardrew Industrial Estate, Redruth, Cornwall TR15 1SH. 01209219205 l.sawle@dmorthotics.com

Word count 2999

22

23

24

25

26

27 **Objectives:** To evaluate a dynamic elastomeric fabric orthosis (DEFO) developed to aid the
28 management of athletic pelvic pain.

29 **Design:** A series of single case studies, with randomised onset of intervention. Daily
30 assessments were undertaken over 15 days (a minimum baseline and intervention period of 6
31 days). A follow up session was undertaken after 1 month.

32 **Setting:** Because of the demands of daily testing over a three week period, participants were
33 tested in locations convenient to them. This included Plymouth University, DM Orthotics and
34 the homes of participants.

35 **Participants:** Eight athletes, confirmed through clinical screening, as presenting with pain
36 affecting the pelvic girdle.

37 **Interventions:** During the intervention period the athletes wore a bespoke DEFO (in the form
38 of shorts).

39 **Main outcome measures:** Force produced on bilateral resisted hip adduction (squeeze test),
40 and self-scored pain (using a numerical rating scale 0-10) at rest and on completing an active
41 straight leg raise (ASLR) and a broad jump. A questionnaire on DEFO usage was administered
42 on completing the study.

43 **Results:** Athletes responded differently, but all but one case showed significant improvement
44 on at least one outcome measure.. People presenting with decreased force output on the
45 squeeze test, and a very asymmetric ASLR tended to show the greatest improvement. All
46 athletes reported they would continue to wear the DEFO.

47 **Conclusions:** This DEFO may support the physiotherapeutic management of athletic pelvic
48 pain.

49

50 Word count: 241

51 Key words: DEFO, Athletes, pelvic, groin, pain, squeeze test

Introduction

Athletic pelvic pain is a common problem; sports like football, cricket and ice hockey have notably high rates of groin injury (1). With different aetiologies affecting, and referring pain to the pelvic region, athletic pelvic pain can be a difficult to manage phenomena made harder by the common occurrence of more than one site of injury, and issues with accurate assessment (2).

Weakness and/or delayed activation in local core stabilisers, particularly transversus abdominus, has been associated with certain types of pelvic pain(1). Pelvic belts have been used in the management of pelvic pain (3;4). Pelvic belts may act by addressing issues in “force closure”; the stabilisation of joints through active and passive musculo-tendinous and ligamentous actions. Pain affecting the lumbopelvic region has also been hypothesised to result from impaired proprioceptive feedback (5) and pelvic belts may also have an effect on proprioception ., In people with longstanding adduction pain pelvic belts reduce pain during active hip adduction(the “ squeeze test”) and active straight leg raise (ASLR) in 50% and 25% of participants respectively (6). Clinical experience suggests that in practice belts are difficult to keep in position, especially during sporting activities, and therefore tend to be limited to use in controlled rehabilitation and training sessions.

Dynamic elastomeric fabric orthoses (DEFO’s) are a recent advancement in orthotics; increasingly used in the management of neurological (7), and musculoskeletal conditions (8). These bespoke orthoses apply forces (through selectively positioned elastomeric panelling) which act to stabilise and align body segments, in order to promote normal function and reduce unwanted movements. The compression that these orthoses can provide, may offer an opportunity to replicate the action of pelvic belts in a more dynamic form.

Based upon the use of pelvic belts and how they may aid management of pelvic pain by improving force closure and / or proprioception (4), a study was undertaken to establish whether diagonally applied force to the pelvis can be as / or more effective in reducing pain during clinical tests compared to those delivered transversely, the usual line of action of a pelvic belt (9). This work highlighted that diagonal forces towards the site of pain may have additional benefits in improving pain and function, compared to no belt or a transverse application. These results informed the development of a DEFO (figure 1) aimed at aiding the management of athletic pelvic pain.

The aim of the current study is to evaluate the effectiveness of this novel, newly developed DEFO on athletes pelvic pain and function using a series of single case studies.

87 **Methods**

88 *Study design*

89 An AB single case study design was used with a randomised onset of intervention. The study
90 was approved by the local ethics committee (Plymouth University) and athletes participated
91 after informed written consent was obtained.

92 *Sample*

93 A mixed sex sample of eight athletes with pelvic pain, as determined by the screening
94 procedure below.

95 *Eligibility Criteria*

96 Potential athletes were over 18 years old; with a history of pelvic pain presenting
97 during sport or rest. There was no time minimum or maximum on duration of pain, to allow
98 for inclusion of both acute and chronic conditions.

99 *Inclusion criteria*

100 In line with European Guidelines a battery of tests was used for screening purposes
101 (11), ensuring the inclusion of tests to identify sacroiliac joint, symphysis pubis and adductor
102 pain:-

103 *ASLR* – This has shown to be a reliable measure of impaired load transfer through the
104 lumbopelvic region (5), and valid for use in athletes with groin pain (12). Athletes in supine
105 with their legs 20 cm apart, raised each leg 20 cm above the plinth. Athletes self-scored how
106 difficult they found this task using a rating of 0-5 (where 5 indicates most difficult). The final
107 score was determined by adding scores from both legs (range 0 – 10). Scores 1-10 were
108 defined as positive.

Faber's test- From supine the athlete was asked to position their leg into flexion, abduction and external rotation (placing one foot across the knee of their opposite leg). pressure was applied to the externally rotated knee, whilst stabilising the opposite ASIS; a positive test was determined by pain being provoked by pressure (13).

Resisted hip adduction – In line with other studies examining groin pain (4;14), from a supine position the athlete was asked to adduct their leg and maintain adduction against an external resistance. The test was positive if pelvic pain was reproduced.

Thigh thrust- This was regarded as the most sensitive test of sacroiliac pain (15). The athlete lay in supine, whilst the therapist flexed their knee and hip, before applying a downwards force through the knee towards the pelvis. The test was positive if pain was provoked by pressure (16).

Gaenslens – From supine, the therapist flexed the knee and hip, whilst extending the opposite leg to lie off the plinth. Overpressure was applied to the flexed knee towards the pelvis, and to the iliac crest of the abducted/extended leg. A positive test was determined by pain provoked

When used in isolation these tests are limited in terms of sensitivity and specificity of diagnosis. However, these increase when they are used as a battery of tests. Therefore, for inclusion into the study people had to have a positive test on at least two of these five tests (17)(16).

Exclusion criteria

People were excluded if they had:

- i. Osteoporosis – to exclude risk of a pelvic fracture being responsible for pelvic pain

- 132 ii. Anorexia - to exclude those at risk of osteoporosis
- 133 iii. Neurological signs, determined by clinical examination, which may influence pain
- 134 perception, and to exclude lumbosacral radicular syndrome
- 135 iv. Pregnancy - to ensure the safety of the mother and foetus
- 136 v. Co-morbidities (rheumatological, neurological or systemic disease) which may
- 137 impact upon the outcome measures
- 138 vi. Suspected fracture based on clinical examination (e.g pain as measured on a numerical
- 139 rating scale (NRS) >8/10, deformity, acute swelling, significant leg-length discrepancy,
- 140 and mechanism of injury)
- 141 vii. Trochanteric bursitis
- 142 viii. Muscle / tendon rupture
- 143 ix. Inguinal herniation
- 144 x. Previous pelvic fracture

145

146 *Procedures*

147 Fifteen daily assessments were undertaken, with at least six during each phase
148 9(baseline and intervention).The onset of the intervention was randomised for each subject
149 using random codes generated in Matlab (Mathworks UK). The intervention could either
150 commence on day 7, 8, or 9. In the intervention phase participants wore the DEFO for
151 activities of their choice and completed a training diary.

152

153 *Outcome Measures*

154 Outcome measures were self-reported pain on a numerical rating scale (NRS) at rest,
155 during a squeeze test, ASLR, and a 1m broad jump. Additionally the force applied during a

156 squeeze test was measured via a load cell. With the legs extended participants were asked to
157 squeeze the load cell as hard as possible. To avoid aggravating the pain with repeated
158 measures participants were instructed to stop squeezing if the pain exceeded 5/10 in the NRS.
159 This was not seen in any of the participants.

160 In each phase measures were repeated after 10 minutes rest. At baseline regular sport
161 shorts were worn during both tests; during the intervention phase participants first wore the
162 DEFO and then ordinary shorts. This allowed an evaluation of the “orthotic effects “ of the
163 DEFO as well as any carry over effects that may be seen when not wearing the orthosis. At
164 one month post the initial testing period, participants were retested as for the intervention
165 period, and completed a questionnaire on their usage of the DEFO.

166

167

Analysis

Visual analysis of trend, level and slope was undertaken on all data (18). The mean (+/- 2 SD) (19) was plotted for the force data and pain scores. Celeration lines were fitted (19;20) and the point of non-overlapping data (PND) statistic (21) was calculated where appropriate.

AB multiple-baseline randomisation tests (22) were undertaken using MatLab (MathWorks, UK) for each outcome measure (with and without the DEFO) (22). This enabled the data from the single case studies to be combined, and examined to see whether any improvement in pain and/or function was due to the intervention or chance. Significance level was $p = 0.05$.

Results

Athletes from various sports, presenting with acute and chronic conditions were assessed (table 1). The effects of the DEFO for each participant are summarised in table 2 and examples of the visual analyses undertaken are demonstrated in figures 2-4. Visual analyses of the single case studies, with the exception of number 5, demonstrated significant effects from wearing the DEFO upon pain and/or function according to at least one measure of significant change.

Group Analysis

The results of the group analysis, combining all 8 participants using a randomisation test are shown in table 3. As a group there was no significant effect of the DEFO or control conditions on the outcome measures tested.

Trends

Examination of the results for each outcome measures identified trends in patient responses which may help build a profile of those benefitting the most from this DEFO.

Force output: Cases studies 1, 3, 4 and 7 demonstrated a significant improvement (above mean + 2SD on 2 consecutive occasions) in force output on the squeeze test wearing the DEFO, and all but study 5 for the intervention period wearing ordinary shorts. This was not related to the site of pain (table 1). . When the forces produced in the squeeze test were normalised by body weight, cases studies 1,3,4 and 7 had a lower baseline force (0.8; 1.3; 1.6; and 1.5 N per kg respectively) compared to the other cases. One exception was case study 6 who also showed a low force output to body weight value (1.2 N per kg). Case 6 did show a trend towards improved force output according to the celeration line; an extended intervention period may have made any improvement clearer.

All of the cases (1, 3, 4 and 7) that showed an improvement in force production with the intervention thus demonstrated a force output of less than 2 N per kg of body weight; values above 2N were seen in the cases where the DEFO had no significant effect upon force output. It is evident that these case studies presented with a fluctuating baseline on the squeeze test; possibly indicating that daily variations in pain affected function.

ASLR

All case studies except number 6 demonstrated bilateral pain responses during the ASLR. When the pain score differences between the right and left ASLR were calculated, cases 1, 3 and 7 demonstrated the biggest differences highlighting a one-sided presentation in pain. When resting pain score was subtracted from the ASLR score causing the most pain (to examine functional impact upon pain), these same cases studies were observed as having the biggest differences. That is, those cases that showed an improvement in force production during the squeeze test, tended to have pain that was considerably aggravated by an ASLR, mainly on the side of their symptoms.

The ASLR is used as a functional indicator of the pelvis' ability to effectively transfer loads from upper to lower limbs (23). Mens et al (4) suggests that those who have a positive ASLR will improve with the introduction of a belt; this has been supported by the work of Lee (24) who has demonstrated this using hands to stabilise the pelvis. This suggests that cases 1, 3, and 7 may have load transfer deficits.

Baseline and intervention resting pain levels

A comparison of baseline and intervention resting pain levels showed cases 1 and 2 had the most benefit from the DEFO.

Case number 2 not only experienced the biggest reduction in pain for pain at rest, but also for ASLR right, and pre-jump pain scores. They also showed significant decreases in ASLR left and post-jump pain scores. This was a case which had a clinically significant improvement in pain, not force. This may be explained in that this participant demonstrated SIJ hypomobility. SIJ hypomobility was diagnosed through thorough clinical assessment (including tests of lumbo-sacral mobility), as this was a historical patient prior to being involved in the case study. This finding suggests that the DEFO may have another mechanism other than force closure. Case study 2 may provide support for the notion that the DEFO is influencing proprioception.

Considering the various pelvic presentations (pain location; duration, history etc.), it may be expected that some participants would show a stronger effect in terms of function (4); whilst in others an effect upon pain is more apparent. More insight into the mechanism(s), and/or pathology associated with pelvic pain may explain this, but this is an issue already acknowledged as being challenging (2).

Subjective responses

Athletes reported that they would continue to wear the DEFO. DEFO usage varied considerably; from participant 1 only wearing his DEFO for activities which caused pain (football and squash), to participant 2 who wore the DEFO when playing rugby, and for 8-9 hours a day during periods of acute pain. Participant 8 chose to wear their DEFO to aid their learning of a new skill, ski-ing, because they felt that it aided their stability. Participants 2, 3, 4, 6, 7 and 8 reported that wearing the DEFO improved their balance and/or posture; case study 6 also reported that the DEFO improved their power training.

Excluding participant 1, who only intended to wear the DEFO during rehabilitation, the remaining participants reported that they would continue to wear the DEFO

for purposes including pain control/ injury prevention, improved posture, and feelings of stability.

Potential Explanations

An athletic DEFO and its effect upon pelvic pain is a novel concept. However, findings from pelvic belt literature may offer some explanations. Belts have been shown to significantly reduce pain and improve force output (as measured by the squeeze test) on athletes with adductor pain (4), and reduce pain in those with posterior pelvic pain (25).

Adductor pain and osteitis pubis have been discussed as examples where pain may arise from instability (6). This DEFO may address force closure deficit (26), by providing cylindrical pressure to improve loading and enhance stability. Decreased force closure may be associated with some pelvic presentations, but other mechanisms need to be considered. Mens et al (5) explained how proprioceptive deficit and impaired muscle function may also be a cause of lumbopelvic pain. Case study 2, where there was SIJ hypomobility, supports the notion that other mechanisms may be involved.

In this case consideration should be given as to whether this DEFO addresses other causes of pelvic dysfunction. Kraemer et al (27), for example, demonstrated the ability of compression shorts to improve proprioception at the hip, suggesting that enhancement of cutaneous receptors may have resulted in improved joint position sense. A similar finding was observed by McNair and Heine (28), finding lumbar bracing improves trunk proprioception; more so in those with a proprioceptive deficit. Even elastic bandage compression has been shown to enhance knee proprioception (29). Therefore enhancing proprioception may improve pain response.

Group Comparisons: Randomisation tests

Randomisation tests showed no significant effect of intervention on any outcome measure. Pre-jump pain score (for the control) demonstrated a trend towards decreased pain scores ($p = 0.054$); this may reflect some degree of carryover from wearing the DEFO, but is mentioned with caution as trends were not seen in other scores.

Results support the conclusions drawn from the visual analyses, which highlighted variation between the athletes in responding to the DEFO. Some reacted by significantly increasing their squeeze test force, others responded with significantly decreased pain scores, others showed little effect e.g. case 5. This mixed picture may be explained by numerous factors. Firstly by the differing pelvic conditions that presented and therefore the existence of varying aetiologies. Secondly in that some patients had more than one site of pain, and thus can be expected to respond differently. Finally, the low pain scores invoked by many of these tests may have influenced the non-significant findings; more stressful tests, as discussed by Verrall et al (30) may be useful.

These explanations may help inform the development of patient profiles to categorise patient subgroups who may respond to this DEFO with reduction in pain and/or functional improvements.

Limitations

Outcome measures

The selected outcome measures were based on standardised, validated tests used clinically, and/or in pelvic belt research. However, some measures e.g. ASLR did not challenge this group of athletes as much as expected, and more stressful tests may have been appropriate for the mixture of pelvic presentations tested (30).

Analyses methods

The mean $\pm$ 2SD often represented a value less than 0. In terms of participants' rating of their pain on the scale of 0-10, less than 0 is an invalid value. Therefore using this form of analysis did not always show a significant result even if a participant's pain was consistently lower during the intervention period. Russo (31) describes how these "floor effects" demonstrate that this is a poor measure of performance. This supports the selection of more stressful measures of performance, in order to avoid this.

A The PND line was of limited use with low pain levels; one very low pain score in the baseline period results in the PND line being set at this level e.g. 0. This criticism has been noted in literature (32).

Future Work

Considering the subjective responses which related to performance benefits, examining the impact of the DEFO on a functional measure of balance and a field test of power is a good starting point for evaluating performance.

Conclusion

The nature of these single case studies and the time-scale involved enabled detailed data to be collected. These results have indicated that this DEFO may have a positive effect upon the pain and / or function of some athletes with pelvic pain. Furthermore, the studies provide preliminary information which is helpful in identifying a patient profile who appear to respond positively to wearing this DEFO. There is evidence indicating that a low force output, and/or asymmetric ASLR could be a useful predictor in terms of responding to the DEFO.

In conclusion this DEFO may have a role in supporting the physiotherapeutic management of athletic pelvic pain. Further work examining its impact upon performance measures may help in understanding of the mechanisms behind its function, and to ascertain if there are effects on performance markers. This may assist in understanding those patients who experience the most benefit from wearing this DEFO.

Ethical Approval: Granted by Plymouth University (08/2009)

Funding: The project was funded by a Knowledge Transfer Partnership between Plymouth University and DM Orthotics . Funding was provided by the Department of Health and DM Orthotics.

Conflict of interest: The DEFO evaluated was developed and patented by DM Orthotics.

Reference List

- (1) Maffey L, Emery C. What are the risk factors for groin strain injury in sport? A systematic review of the literature. *Sports Medicine* 2007;37(10):881-94.
- (2) Ficek K, Rzepka R, Orawczyk T, Zotnierz Z. Groin pain in athletes - clinical experience. *Journal of Human Kinetics* 2008;19:141-8.
- (3) Lee D. *The Pelvic Girdle*. 3rd ed. Oxford: Churchill Livingstone; 2004.
- (4) Mens J, Inklaar H, Koes BW, Stam HJ. A new view on adduction-related groin pain. *Clinical Journal of Sport Medicine* 2006;16(1):15-9.
- (5) Mens JMA, Vleeming A, Snijders CJ, Koes BW, Stam HJ. Reliability and Validity of the Active Straight Leg Raise Test in Posterior Pelvic Pain Since Pregnancy. *Spine (Phila Pa 1976)* 2001;26:1167-71.
- (6) Jansen JACG. Longstanding adduction-related groin pain in athletes Erasmus; 2010.
- (7) Rennie DJ, Attfield SF, Morton RE, Polak FJ, Nicholson J. An evaluation of lycra garments in the lower limb using 3-D gait analysis and functional assessment (PEDI). *Gait and Posture* 2000;12:1-6.

- 352 (8) Matthews M, Crawford R. The use of dynamic Lycra orthosis in the treatment of scoliosis: A
353 case study. *Prosthetics and Orthotics International* 2006;30(2):171-4.
- 354 (9) Sawle L, Freeman J, Marsden J, Matthews MJ. Exploring the effect of pelvic belt
355 configurations upon athletic lumbopelvic pain. *Prosthetics and Orthotics International*
356 2012;Online first ; July 2.
- 357 (10) Mens JMA, Damen L, Snijder CJ, Stam HJ. The mechanical effect of a pelvic belt in patients
358 with pregnancy-related pelvic pain. *Clinical Biomechanics* 2006;21:122-7.
- 359 (11) Vleeming A, Albert HB, Ostgaard HC, Sturesson B, Stuge B. European guidelines for the
360 diagnosis and treatment of pelvic girdle pain. *European Spine Journal* 17[6], 794-819. 2008.
361 Ref Type: Journal (Full)
- 362 (12) Cowan SM, Schache AG, Brukner P, Bennell KL, Hodges PW, Coburn P, et al. Delayed Onset of
363 Transversus Abdominus in Long-Standing Groin Pain. *Medicine and Science in Sports and*
364 *Exercise* 2004;36(12).
- 365 (13) Merriman LM, Turner W. Assessment of the lower limb. 2nd ed. London: Elsevier Health
366 Sciences; 2002.
- 367 (14) Holmich P, Uhrskou P, Ulnits L, Kanstrup I, Bachmann Nielsen M, Munch Bjerg AaKK.
368 Effectiveness of active physical training as treatment for longstanding adductor-related groin
369 pain in athletes: randomized trial. *Lancet* 1999;353:439-43.
- 370 (15) Laslett M, Aprill CN, McDonald B, Young SB. Diagnosis of sacroiliac joint pain: Validity of
371 individual provocation tests and composites of tests. *Manual Therapy* 2005;10:207-18.
- 372 (16) Cattely P, Winyard J, Trevaskis J, Eaton S. Validity and reliability of clinical tests for the
373 sacroiliac joint: A review of the literature. *Australasian Chiropractic and Osteopathy*
374 2005;10(2):73-80.
- 375 (17) Stuber KJ. Specificity, sensitivity, and predictive values of clinical tests of the sacroiliac joint: A
376 systematic review of the literature. *The Journal of the Canadian Chiropractic Association*
377 2007;51(1):30-44.
- 378 (18) Wolery M, Harris SR. Interpreting results of single-subject research designs. *Physical Therapy*
379 1982;62(445):452.
- 380 (19) Reza Nourbakhsh M, Ottenbacher KJ. The statistical analysis of single-subject data: A
381 comparative examination. *Physical Therapy* 1994;74(768):776.
- 382 (20) Hojem MA, Ottenbacher KJ. Empirical investigation of visual-inspection versus trend-line
383 analysis of single-subject data. *Physical Therapy* 1988;68(983):988.
- 384 (21) Kazdin AE. Statistical analysis for single-case experimental designs. In: Barlow DaHM, editor.
385 Single case experimental designs: Strategies for studying behavior change. New York:
386 Pergamon; 1984. p. 285-324.
- 387 (22) Todman JB, Dugard P. Single-Case and Small-n Experimental Designs a Practical Guide to
388 Randomization Tests. New York; London: Lawrence Erlbaum Associates; 2001.
- 389 (23) Vleeming A, Albert HB, Ostgaard HC, Sturesson B, Stuge B. European guidelines for the
390 diagnosis and treatment of pelvic girdle pain. *European Spine Journal* 2008;17(6):794-819.

- 391 (24) Lee D. The one-leg standing test and the active straight leg raise test: A clinical interpretation
392 of two tests of load transfer through the pelvic girdle. Orthopaedic Review 2005.
- 393 (25) Ostgaard HC, Zetherstrom G, Roos-Hansson E, Svanberg B. Reduction of back and posterior
394 pelvic pain in pregnancy. Spine (Phila Pa 1976) 1994 Apr;19(8):894-900.
- 395 (26) Lee D, Vleeming A. Impaired load transfer through the pelvic girdle. A new model of altered
396 neutral zone function. Proceedings of the Third interdisciplinary world congress on low back
397 and pelvic pain 1998.
- 398 (27) Kraemer WJ, Bush JA, Newton RU, Duncan ND, Volek JS, Denegar CR, et al. Influence of a
399 compression garment on repetitive power output before and after different types of muscle
400 fatigue. Sports Medicine, Training and Rehabilitation 1998;8(2):163-84.
- 401 (28) McNair PJ, Heine PJ. Trunk proprioception: Enhancement through lumbar bracing. Archives of
402 Physical and Medical Rehabilitation 1999;80:96-9.
- 403 (29) Perlau R, Frank C, Fick G. The effect of elastic bandages on human knee proprioception in the
404 uninjured population. American Journal of Sports Medicine 1995;23(2):251-5.
- 405 (30) Verrall GM, Slavotinek JP, Barnes PG, Fon GT. Description of pain provocation tests used for
406 the diagnosis of sports related chronic groin pain: relationship of tests to defined clinical (pain
407 and tenderness) and MRI (pubic bone marrow oedema) criteria. Scandinavian Journal of
408 Medicine and Science in Sports 2005;15:36-42.
- 409 (31) Russo R. Statistics for the Behavioural Sciences: An Introduction. Sussex: Psychology Press.
410 Taylor and Francis Group.; 2003.
- 411 (32) Morgan DL, Morgan RK. Single-Case Research Methods for the Behavioral and Health
412 Sciences. London, California, New Delhi and Singapore: Sage; 2009.
413
414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

Single Case Study Number	Sex	Sport	Age (yrs)	Height (m)	Weight (Kg)	Site of pain	Duration of Pain
1	Male	Football, running, cycling, squash	37	1.86	73	Adductor	3 months
2	Male	Rugby, running, cycling	31	1.81	84.4	SIJ	3-4 years
3	Female	Power walking	62	1.67	63.4	Adductor and SIJ	6 months
4	Female	Yoga, aerobic/ power training programmes	29	1.72	87.5	SIJ	3 years; worse last 18months
5	Female	Boxing training	53	1.53	39.4	SIJ	2 years
6	Female	Aerobic/ power training programmes	26	1.65	58.1	SIJ	2 years
7	Female	Ski-ing	42	1.60	68.2	SIJ	17years

8	Female	Cycling, swimming	34	1.68	54	SIJ	20 years
---	--------	----------------------	----	------	----	-----	----------

432

433 Table 1.Demographics

434

435

436

437

438

439

440

Subject/ Outcome measures	1	2	3	4	5	6	7	8
Resisted Adduction Force‡	CI↑ Msd↑ PND =58.3%	↑ CI PND = 12.5%	↑ CI Msd↑ PND =62.5%	↑ CI Msd↑ PND =100%	↓ CI Msd↓ PND = 0%	PND =0% CI =variable	↑CI Msd↑ PND =33.3%	↑CI PND =25%
Pain on resisted adduction	CI↓ PND = 8.3%	↓ CI PND = 25%	PND = 12.5%	PND = 0%	PND =0%	* PND =0%	↓CI PND = 88.9%	↓CI Msd↓ PND =100%
ASLR	CI↓	↓ CL	↓CI	↓CI	PND	PND	↓CI	↓CI

Right	PND= 0%	PND = 0%	PND =0%	PND =14.3%	=0%	=0%	PND =0%	PND =0%
ASLR	CI ↓	↓CI	↓CI	↓CI	PND	PND	↓CI	↓CI
Left	PND= 0%	PND =0%	PND =0%	PND = 14.3%	=0%	=0%	PND =0%	PND =0%
PAIN	CI ↓	↓CI	PND	↓CI	↓CI	PND	↓CI	↓CI
REST	PND= 0%	PND =25%	=0%	PND =0%	PND =0%	=0%	PND =0%	PND =12.5%
PRE JUMP	CI ↓ PND = 25%	↓CI PND = 37.5%	PND = 0%	↓CI PND= 14.29%	PND =0%	PND =0%	↓CI PND =0%	↓CI PND =0%
POST	CI ↓	↓CI	PND	↓CI	PND	PND	↓CI	↓CI
JUMP	PND = 8.3%	PND =75%	=37.5%	PND =14.29%	=0%	=0%	PND =0%	PND =0%

441

442 CI indicates the change relative to the celeration line whilst msd indicates the change relative to the
443 mean +/- 2 standard deviation line. Figures indicate the PND score, ↑ indicates that in phase B
444 measures were above the celeration. ↓ indicates that in phase B measures were below the celeration.
445 *Indicates that the pain dropped to zero. Increases above the celeration line (↑) for the resisted
446 adduction test indicates improvement in force production, whilst for the measures of pain a decrease
447 (↓) indicates a decrease in pain

448

449 Table 2: Summarising the change in l outcome measures.

450

451

452

453

454

455

456

457

458

459

460

Outcome measure	Intervention (DEFO)	Control (shorts)
Pain at rest	0.15	0.40
Force on squeeze test	0.26	0.38
Pain on squeeze test	0.17	0.54
Pain on right ASLR	0.25	0.23
Pain on left ASLR	0.19	0.16
Pain pre broad jump	0.15	0.054
Pain post broad jump	0.17	0.51

461

462 Table 3 Randomisation test p values for each outcome measure

463

464

465

466

467

468

469

470

471

472

473 Figure 1: The DEFO

Figure 2: An example of improved force output

Figure 3: An example of decreased pain scores

Figure 4: An example of no significant change observed