2017-01-30

Combined sea-level and climate controls on limestone formation, hiatuses and ammonite preservation in the Blue Lias Formation, South Britain (uppermost Triassic Lower Jurassic)

WEEDON, GRAHAMP

<http://hdl.handle.net/10026.1/8505>

10.1017/S001675681600128X

Geological Magazine

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Combined sea-level and climate controls on limestone formation, hiatuses and ammonite preservation in the Blue Lias Formation, South Britain (uppermost Triassic–Lower Jurassic)

Short title/running head: **Sea level and climate controls of the Blue Lias**

Graham P. Weedon^{1*}, Hugh C. Jenkyns² and Kevin N. Page³

* Corresponding author

1 graham.weedon@metoffice.gov.uk, Met Office, Maclean Building, Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK.

2 Department of Earth Sciences, University of Oxford, South Parks Road, Oxford, OX1 3AN, UK.

3 School of Geography, Earth and Environmental Sciences, Plymouth University, Drake Circus, Plymouth, PL4 8AA, UK.

In press: *Geological Magazine*

January 2017

Abstract

Lithostratigraphic and magnetic-susceptibility logs for four sections in the Blue Lias Formation are combined with a re-assessment of the ammonite biostratigraphy. A Shaw plot correlating the West Somerset coast with the Devon/Dorset coast at Lyme Regis, based on 63 common biohorizon picks, together with field evidence, demonstrate that intra-formational hiatuses are common. Compared to laminated shale deposition, the climate associated with light marl is interpreted as both drier and stormier. Storm-related non-deposition favoured initiation of limestone formation near the sediment-water interface. Areas and time intervals with reduced water depths had lower net accumulation rates and developed a greater proportion of limestone.

Many homogeneous limestone beds have no ammonites preserved, whereas others contain abundant fossils. Non-deposition encouraged shallow sub-seafloor cementation which, if occurring after aragonite dissolution, generated limestones lacking ammonites. Abundant ammonite preservation in limestones required both rapid burial by light marl during storms as well as later storm-related non-deposition and near-surface carbonate cementation that occurred prior to aragonite dissolution.

The limestones are dominated by a mixture of early framework-supporting cement that minimized compaction of fossils, plus a later micrograde cement infill. At Lyme Regis, the relatively low net accumulation rate ensured that final cementation of the limestones took place at relatively shallow burial depths. On the West Somerset coast, however, much higher accumulation rates led to deeper burial before final limestone cementation. Consequently, the oxygen-isotope ratios of the limestones on the West Somerset coast, recording precipitation of the later diagenetic calcite at higher temperatures, are lower than those at Lyme Regis.

Keywords:

Sedimentary cycles; Sea level; Climatic cycles; Blue Lias Formation; Diagenesis; Hiatuses

1. Introduction

The origin and significance of the limestone beds and nodules in the uppermost Rhaetian (Triassic) to Sinemurian (Jurassic) Blue Lias Formation of Britain has long been a subject of interest (Day, 1865; Richardson, 1923; Kent, 1936). Hallam (1960; 1964) argued that the limestones owed their characteristics to both diagenetic and primary (i.e. depositional) factors. Shukri (1942) speculated on the possible role of climate but ruled out forcing by Milankovitch orbital-precession cycles because the limestone beds at Lyme Regis are more widely spaced in the lower Sinemurian compared to the Hettangian. Following the demonstration by Hays, Imbrie & Shackleton (1976) of orbital forcing of Pleistocene to Recent climate, House (1985; 1986) and Weedon (1985; 1986) revived the idea of orbital-climatic (Milankovitch cycle) forcing to explain the interbedded lithologies.

One of the reasons the Blue Lias Formation has attracted so much research interest is that it exhibits two styles of sedimentary cyclicity. Visually most obvious are the alternations of limestones and non-limestones, but also present are alternations of homogeneous, organic-carbon-poor strata (grey limestone plus light grey marl) with laminated organic-carbon-rich strata (black to very dark grey laminated shale plus laminated limestone). Dark grey marls represent intermediate compositions. The homogeneous limestone beds and layers of nodules are considered to have formed diagenetically within light marl beds while the much rarer laminated limestone beds and laminated limestone nodules formed within primary laminated shale beds (Hallam, 1964; Weedon, 1986; 1987a; Arzani, 2004).

Weedon (1986; 1987a) showed, using time-series analysis, that the alternation of the limestone with the non-limestones can encode a similar signal of regular cycles to that of the alternating homogeneous and laminated rock types. However, in thicker sections such as those

exposed on the West Somerset Coast, multiple limestone beds and/or nodule horizons occur within thick (tens of centimetre- to metre-scale) light marl beds. A recent analysis of orbital forcing in the relatively thick sections on the West Somerset Coast (Ruhl *et al.*, 2010) specifically avoided sampling the limestones that nonetheless represent a critical part of the sequence. This paper is designed to clarify the nature of the information conveyed by the limestones, both laminated and homogeneous.

----- Figure 1 near this position -----

New, high-resolution lithostratigraphic and magnetic-susceptibility logs for four sections (Fig. 1) are presented together with refinements of the ammonite biostratigraphy. The sections chosen within ‘typical offshore’ Blue Lias, span all the ammonite zones of the Hettangian Stage and represent a wide range of net accumulation rates. The aims are to explain why: a) limestone bed thicknesses are apparently independent of the net accumulation rate; b) the spacing of the limestone beds within ammonite zones can be related to Milankovitch orbital cycles although there are large variations in spacing from zone to zone and from place to place; c) laminated limestone beds and laminated limestone nodules are comparatively rare; and d) some limestone beds preserve abundant ammonite fossils, but others preserve none. We present new evidence for intra-formational hiatuses and a synthesis of limestone formation in terms of combined climatic and sea-level controls plus an improved understanding of the preservation of the ammonites.

2. Bio- and chronostratigraphy

2.a. Ammonite biochronology and correlation

The Jurassic System is divided into a sequence of 11 globally applicable stages that are subdivided, without gaps or overlap, into regional sequences of ammonite-correlated ‘zones’ (Ogg & Hinnov, 2012). Since ammonite zones completely fill each stage, and are now usually

explicitly defined with a basal stratotype, they can also be considered to be chronostratigraphical units (Callomon, 1985; 1995; Page, 1995; 2003; in press). The great majority of zones do not conform in any way to a classical biozone since they do not correspond to the range of any ammonite species or assemblage.

In the present work, we refer neutrally to the ammonite-correlated stratigraphical units as ‘Standard’ Zones and Subzones (*sensu* Callomon, 1985), rather than using an epithet to describe their character (i.e. ‘chronozone’ and ‘subchronozone’ as in Page, 2010a; 2010b; 2010c). However, as chronozones, these units can be explicitly correlated using proxies other than ammonites, such as local lithological changes, other faunal or floral elements and isotopic ‘events’ (Jenkyns *et al.*, 2002).

Throughout the Jurassic System, infra-subzonal units that are often referred to as biohorizons provide a very high-resolution biochronology. Biohorizons typically correspond to the stratigraphic range of a specific indicator species. Normally, the bases of ammonite Standard Zones are explicitly, or effectively, defined by the bases of specific biohorizons, and hence correlated, as potential timelines, by the stratigraphically lowest occurrences of the indicator species.

The initial biohorizontal framework for the Lower Sinemurian Stage in the UK of Page (1992) led to the schemes applied by Page (2002; 2010b) to the Devon and Dorset coastal sections. The scheme for the Hettangian Stage of the Devon/Dorset coast (Page, 2010b; 2010c) was based largely on the version for West Somerset by Page (2005), but recent re-sampling of the latter area has revealed even greater biostratigraphical detail, which is used here. This new scheme provides a sequence of 55 biohorizons for the Hettangian, as opposed to the 27 used by Page (2010b; 2010c).

Zonal and subzonal boundaries throughout the Hettangian have been reviewed, based on new sampling by K.N.P. on coastal sections (Fig. 1a) in East Devon (Lyme Regis), West

Somerset (Doniford, St Audries Bay, East Quantocks Head to Kilve) and South Glamorgan, South Wales (St. Mary's Well Bay, Lavernock). The biohorizon boundaries, where constrained to within a few centimetres stratigraphically, are indicated in Table 1. Within the Lower Sinemurian of West Somerset, the placement of biohorizon boundaries are as previously published (e.g. Bloos & Page, 2000b; Page 2010b; 2010c).

-----Table 1 near this position-----

2.b. The base of the Jurassic

The Global Stratigraphic Section and Point (GSSP) for the base of the Jurassic has been defined in the Kuhjoch section in the Austrian Calcareous Alps (Hillebrandt, Krystyn & Kuerschner, 2007; Hillebrandt & Krystyn, 2009). Subsequently Page (2010b; 2010c) revised the zonal framework for the base of the Hettangian Stage in the UK by inserting the former authors' Tilmanni Zone below the Planorbis Zone (which was previously considered to be the lowest zone of the Jurassic System).

Page (2010b; 2010c) concluded that only the top of the Tilmanni Zone in the UK had yielded ammonites, specifically *Psiloceras erugatum* (Phillips) as illustrated by Bloos & Page (2000a). Nevertheless, in the UK the base of the Zone and hence the base of the Jurassic System can still be approximated using the most positive part of a positive $\delta^{13}\text{C}_{\text{org}}$ interval, as demonstrated by Clémence *et al.* (2010). At St Audries Bay on the West Somerset coast (Fig. 1), the base of the Hettangian corresponds to a level less than 1.5 m above the base of the Blue Lias Formation (Clémence *et al.*, 2010). At St Mary's Well Bay near Lavernock, South Wales (Fig. 1), the $\delta^{13}\text{C}_{\text{carb}}$ curve of Korte *et al.* (2009), based on measurements of *Liostrea* sp. samples, suggests that the base of the Jurassic System lies about 1.9 m above the top of the Langport Member (Penarth Group, Upper Triassic). The $\delta^{13}\text{C}_{\text{carb}}$ curve produced from bulk

sediment samples from the coast at Lyme Regis on the Devon/Dorset border by Korte *et al.* (2009) is, however, of limited use for correlation because it does not show the same signature as elsewhere. Nevertheless, it is likely that the base of the Jurassic System in the area also lies in the lowest part of the Blue Lias Formation.

2.c. The base of the Planorbis Zone

The base of the Planorbis Zone, as correlated by the first appearance of *Neophyllites* at the base of the Hn3 *imitans* Biohorizon (Page, 2010a), corresponds to the base of bed 9 of Whittaker & Green (1983) on the West Somerset Coast (Page & Bloos, 1995; Bloos & Page 2000a). In South Glamorgan, in St. Mary's Well Bay, Lavernock, the base of the Planorbis Zone is about 12 cm below the base of bed 30 of Waters and Lawrence (1987): i.e. the level indicated by Hodges (1994) within his bed 38. At Lyme Regis, this level correlates with the base of bed H25 of Lang (1924); see Page (2002).

2.d. Southam Quarry near Long Itchington

At Southam Quarry, Long Itchington, Warwickshire, below bed 10 of Clements *et al.* (1975; 1977) ammonites are rare (Radley, 2008) and the lowest part of the Blue Lias Formation is inferred to belong to the Liasicus Zone (i.e. the Tilmanni and Planorbis zones are missing). This interpretation follows from the location of Long Itchington in relation to the reconstructed age of onlap of the Blue Lias Formation onto the London-Brabant Platform by Donovan, Horton & Ivimey-Cook (1979). Furthermore, it is consistent with the records of Old, Sumbler, & Ambrose (1984, p. 34), who cite *Laqueoceras* and *Waehneroceras* 'close above the Langport Member' at Southam Quarry. *Laqueoceras* certainly indicates the lower part of the Laqueus Subzone Hn18 Biohorizon of Page (2010b; 2010c). However, specimens of *Waehneroceras* from Southam Quarry include species from the upper part of the Portlocki Subzone, including

abundant examples of *W. portlocki* (Wright), in limestone nodules and rare *W. cf. shroederi* Lange in limestone beds. These observations suggest that at least biohorizons Hn16 and Hn17 of Page (2010b; 2010c) are also present.

3. The sections studied and the lithostratigraphic and magnetic-susceptibility logs

Lithological logging, whilst measuring magnetic susceptibility (MS) in the field, utilized the five microfacies or rock-types of Weedon (1986; 1987a), as adopted by others (e.g. Bottrell & Raiswell 1989, Arzani 2004; 2006; Paul, Allison & Brett 2008) i.e. homogeneous grey limestone, laminated grey limestone, light grey marl, dark grey marl and millimetre-laminated black shale. Limestone nodules occur within light marl beds and locally pass laterally into continuous limestone beds (e.g. at Lyme Regis bed 37 or ‘Rattle’: Lang (1924) and Hesselbo & Jenkyns (1995) provide all the limestone bed names for the Devon/Dorset coast). Laminated limestone nodules occur within the laminated black shales. Large differences in net accumulation rates in the sections studied are suggested by the thicknesses of zones shown in Fig. 1b, and the average composition of the rock-types is indicated in Fig.1c and discussed further in Section 4.

3.a. Measurements of magnetic susceptibility in the field

Volume magnetic susceptibility (vol MS) logging of sections used a Bartington Instruments MS2 meter combined with an F-probe in direct contact with fresh faces of rock at right angles to the bedding. Instrument drift, related to temperature changes, was removed by taking measurements in the air more than 1 metre from the cliff face between each rock-surface measurement. At the levels of limestone nodules, MS was measured in the limestone rather than light marl in cases where limestone constituted more than half of that stratigraphic level (Weedon *et al.*, 1999). As far as possible, calcite ‘beef’ veins and lenses (Richardson, 1923;

Marshall, 1982), most prevalent at the base of laminated shale beds at Lyme Regis, were avoided during MS measurement.

----- Figure 2 near this position -----

Measurements of weight-specific MS (wt MS) using a Bartington Instruments MS2B sensor for samples from Lyme Regis, and the West Somerset coast show the expected strong, linear correlation with vol MS (Pearson's $r = +0.967$, $N = 74$, $P < 0.001$, Fig. 2a). This relationship and the similarity of both the vol MS and wt MS logs to %CaCO₃ in Fig. 2b confirm that the field measurements are neither significantly affected by changes in magnetic properties due to weathering, nor by incomplete rock sensing due to imperfect surface contact and/or undetected voids.

The lithological columns shown in Figs 3–6, with captions indicating the references used in bed numbering, illustrate large variations in the thickness of the marls and shales and highly variable spacing and overall proportions of limestone beds. In the logs, the laminated black shales are shown as the most recessed lithology on the profiles as a convention to aid their identification rather than as an indication of the lowest resistance to weathering. The fixed spacing of the MS measurements at each section was designed to resolve the smallest variations in bulk composition. The measurements were obtained at 2 cm spacing at Lyme Regis; 3 cm at Southam Quarry, Long Itchington and 4 cm spacing on both the West Somerset coast and at St. Mary's Well Bay, Lavernock. In Southam Quarry, the faces were measured for magnetic susceptibility in 1994 when the section in the south west between the A423 and the A426 was fully accessible down to the disconformity with the Upper Triassic Langport Member. The vol MS log for Lyme Regis was reported previously (Weedon *et al.*, 1999) but Fig. 4, primarily of the Hettangian portion, shows revisions to the zonal and subzonal boundaries.

----- Figure 3 near this position -----

The log for the West Somerset coast (Fig. 5a and b; Palmer, 1972; Whittaker & Green, 1983) is composite. The section at St Audries Bay was measured from the base of the Blue Lias Formation to the top of bed 101 of Whittaker & Green (1983), overlapping the MS log at Quantock's Head by about 3.5 m. The Quantock's Head to Kilve section was measured for MS from bed 97 to bed 163. The splice level of the composite record in bed 96, at 56.90 m above the base of the Blue Lias Formation, represents the base of the Quantock's Head magnetic-susceptibility log. Note that in Table 1 the biohorizon levels of the Quantock's Head section are listed both relative to the base of the log (i.e. at the splice level) and relative to the base of the Formation according to the heights of the St Audries Bay section within the composite column.

----- Figure 4 near this position -----

3.b. Origin of the variations in vol MS

It was reported previously for the Blue Lias Formation that MS is inversely correlated with %CaCO₃ (Hounslow, 1985; Weedon *et al.*, 1999; Deconinck *et al.*, 2003; Ruhl *et al.*, 2010). This relationship is supported by current measurements of wt MS using a Bartington Instruments MS2B cavity sensor and vol MS (Fig. 2a). The right-hand linear regression line of Fig. 2a allows estimation of carbonate contents from the field measurement of vol MS (using: %CaCO₃ = 99.06 – (13.79 x vol MS), $r = -0.892$, $N = 74$, $P < 0.001$). Calcium carbonate measurements have been shown as open circles on top of the MS logs in Figs 4 and 5 by scaling %CaCO₃ (bottom scale) relative to the vol MS (top scale) according to this regression. Despite the good agreement between the estimated and measured %CaCO₃ shown on the right of Fig 2b, the uncertainty (95% confidence interval) in individual estimates of %CaCO₃ from vol MS using the regression is about ±20.0% – ranging between ±19.6 and ±20.4%, depending on the difference between the individual vol MS and the mean vol MS (Williams, 1984, p. 313).

----- Figure 5a near this position -----

----- Figure 5b near this position -----

The average MS values for these mudrocks are fairly low, thereby contrasting with younger British Jurassic cyclic mudrocks such as the Kimmeridge Clay Formation (Weedon *et al.*, 1999). It was shown for the Blue Lias Formation (Hounslow, 1985; Bixler, Elmore & Engel, 1998; Deconinck *et al.*, 2003; Hounslow, Posen & Warrington, 2004) that the stratigraphic variations in MS are consistent with variable dilution of paramagnetic clays by non-ferroan, and thus diamagnetic, calcite (wt MS = $+0.84 \times 10^{-8}$ SI, Bleil & Petersen, 1987). Note that despite the presence of locally rather large percentages of pyrite (maximum 12% in the laminated shales, Weedon, 1987a) in a pure, unweathered, state this mineral is also diamagnetic (wt MS = -0.48×10^{-8} SI, Collinson, 1983) and consequently contributes very little to the whole-rock MS signal. Hence, stratigraphic logs of MS in the Blue Lias Formation provide an accurate picture of the occurrence of the limestones (Figs 2a, 2b and 3).

The marls and shales of the Liasicus Zone at all four localities have consistently higher average MS than the underlying Tilmanni and Planorbis and overlying Angulata and Bucklandi zones (Figs 3–6, Weedon *et al.*, 1999; Deconinck *et al.*, 2003). This observation suggests that there were long-term (zonal-scale) variations in the composition of the paramagnetic (mainly clay mineral) components (Deconinck *et al.*, 2003). The relatively high average MS of the lower part of the section in Southam Quarry (Fig. 6) is consistent with the interpretation (Section 2.d) that the lowermost strata that lie disconformably on the Langport Member belong to the Liasicus Zone and not to the Planorbis Zone.

At about the 31 m level in the mid-Liasicus Zone (upper Portlocki Subzone) of the St Audries Bay section (Fig. 5a), the average vol MS is unusually high with values extending outside the range of the %CaCO₃ versus vol MS regression of Fig. 2a. Lower stratigraphic resolution sampling by Deconinck *et al.* (2003) indicates that, at this level on the West

Somerset coast, the clay minerals switch to higher kaolinite/illite ratios, but surprisingly, not to iron-rich clays. The unusually high MS might indicate an association of increased kaolinite with a source of different paramagnetic components and/or small amounts of ferromagnetic components such as magnetite-like minerals that, unusually for the Blue Lias Formation, survived dissolution by H₂S during diagenesis (Deconinck *et al.*, 2003). A similar interval of unusually high values of vol MS, of around 8.0×10^{-8} SI, is found at about 8.2 m in Southam Quarry in the Liasicus Zone (Fig. 6). This interval of high average MS in the marls and shales is located at the top of the Portlocki subzone, Liasicus Zone at both St Audries and Lyme Regis and may provide a correlatable change in paramagnetic mineralogy (Figs 4 and 5a).

----- Figure 6 near this position -----

4. Variations in whole-rock composition

The typical ‘offshore’ facies of the Hettangian Blue Lias Formation of interest here shows considerable lateral and stratigraphic variations in bulk composition as discussed in Sections 4.b and 4.c. To provide a context for these descriptions we start with consideration of the ‘marginal facies’.

4.a. Marginal facies and ‘near-shore’ Blue Lias

The Blue Lias was deposited at the same times as palaeo-coastline deposits that include the Sutton Stone and Southerndown Beds of South Glamorgan (Trueman, 1920; 1922; Hallam, 1960; Wobber, 1965; Fletcher, 1988; Sheppard, 2006) and the Brockley Down Stone of the Radstock Plateau near Bristol (D. L. Loughman, unpub. Ph.D. thesis, University of Birmingham, 1982; Donovan & Kellaway, 1984). These so-called marginal facies, formed at and close to palaeo-coastlines, lie unconformably on Carboniferous Limestone and comprise

conglomeratic limestones with grainstone textures including Carboniferous Limestone and chert lithoclasts, derived bioclasts, ooids and bioclasts (Wobber, 1965; 1966; Fletcher, 1988).

On the coast near Southerndown, Glamorgan, 31 km west north west of the exposure at Lavernock (Fig. 1a), the unconformity surface is exposed and consists of multiple wave-cut platforms overlain by associated successive breccia and conglomeratic and grainstone limestones of the Sutton Stone (Fletcher, 1988; Sheppard, 2006). Both the unconformity surface and Carboniferous Limestone lithoclasts in the overlying Sutton Stone have *Trypanites* borings and are encrusted by oysters and colonial corals (Johnson & McKerrow, 1995; Simms, Little & Rosen, 2002; Sheppard, 2006). Some lithoclasts in the Sutton Stone are imbricated (Wobber, 1963) and elongate bioclasts have been used as palaeocurrent indicators that indicate long-shore drift and complex current movements influenced by the local islands of Carboniferous rocks (Wobber, 1966). The matrix-supported and rounded boulder- and cobble-sized lithoclasts in the Sutton Stone are succeeded stratigraphically by pebble-grade lithoclasts, micrites and thin shales of the Southerndown Beds, which grade laterally into the near-shore facies of Blue Lias (Trueman, 1922; Wobber, 1965; Wilson et al., 1990; Sheppard, 2006).

Although the platform erosion probably started in the latest Triassic, ammonites have been recovered in stratigraphic order from the Sutton Stone and Southerndown Beds from the Planorbis, Liasicus and Angulata Zones (Hodges, 1986; Sheppard, 2006). These observations are not consistent with nearly instantaneous deposition of the Sutton Stone by a single hurricane (Ager, 1986), but rather with the latest Triassic and Early Jurassic erosion of the wave-cut platforms and deposition of the marginal facies under storm influence over millions of years during episodic rises in relative sea level (Trueman, 1922; Fletcher, 1988; Sheppard, 2006).

Just six kilometres south of the palaeo-coastline of Southerndown at Nash Point, the Blue Lias of the Bucklandi Zone represents an atypical limestone-dominated facies often full of bioclast fragments in both the marls and limestones (Hallam, 1960; Weedon, 1987a;

Trueman, 1930). These strata representing Units B and C of the Porthkerry Formation have much higher limestone proportions than found in more 'typical' Blue Lias (an average of 60 to 85 % limestone by thickness in sections tens of metres thick, Waters & Lawrence, 1987; Wilson et al., 1990; Warrington & Ivimey-Cook, 1995). Units B and C consist of metre-scale bedding units formed of closely spaced nodular limestones and 'anastomosing mudstone beds' that probably result from pressure dissolution (Waters & Lawrence, 1987). Local decimetre-scale mudstone beds are present, permitting correlation of limestone-shale bed groups, but organic-carbon contents are low and neither laminated shales nor laminated limestones are present (Sheppard, Houghton & Swan, 2006). Although much of the bedding was formed purely during diagenesis, the presence of hummocky cross-stratification at Nash Point proves strong storm influences (Sheppard, Houghton & Swan, 2006).

Thirty kilometres from the palaeo-coastline of South Glamorgan, typical 'offshore' Blue Lias facies is present at Lavernock in the Tilmanni to Liasicus zones as logged here (Fig. 3). In the lower part of the Angulata Zone Unit A of the Porthkerry Formation consists of about 55 % by thickness of normal offshore Blue Lias with nodular limestones and marls. However, in the same area near Cardiff, the upper part of the Angulata Zone, 'Unit B' of the Porthkerry Formation of Waters and Lawrence (1987), is of a similar facies to the Bucklandi Zone at Nash Point. The Porthkerry Formation is overlain gradationally by uppermost Angulata Zone and Bucklandi Zone oolitic 'marginal facies' (Waters & Lawrence, 1987). Thus, Units B and C of the Porthkerry Formation near Cardiff and Nash Point represent a 'near-shore' facies of the Blue Lias, which has been envisaged as a storm-derived aragonitic lime mud deposited on a marine shelf (Sheppard, Houghton & Swan, 2006).

4.b. Rock-type compositions

The different rock-types of the ‘typical’ or ‘offshore’ Blue Lias consist mostly of various proportions of bioclasts, calcite microspar, clay minerals, organic matter, pyrite and minor quartz silt (Hallam, 1960; Weedon, 1986; 1987a). Whole rock compositions quoted here are based on the samples from both Lyme Regis and the Somerset Coast (Weedon, 1987a). Numbers of samples by rock type and locality are indicated in the caption for Fig. 1c and sample locations are indicated in Figs 4 and 5. The limestones (total organic carbon, TOC, mean = 0.54% $\pm$ 0.09 ($\pm$ 95% confidence interval), range = 0.14 to 1.64%, N = 48; CaCO₃, mean = 79.3% $\pm$ 2.2, range = 51.8 to 88.7%, N = 48) form beds with contacts ranging from planar to nodular. Limestone nodules are typically entirely enclosed by light marl but can be ‘welded’ onto limestone beds. On the Devon/Dorset and West Somerset coasts, when seen in plan on the foreshore, limestone nodules are often found to be linked laterally at some levels. The limestones generally have a homogeneous bioclastic wackestone to bioclastic carbonate mudstone fabric reflecting the thoroughly bioturbated nature of the precursor light marls. The light marls (TOC mean = 1.42% $\pm$ 0.26, range = 0.38 to 4.41%, N = 40; CaCO₃, mean = 43.8% $\pm$ 2.7, range = 25.7 to 69.9%, N = 40) are friable and homogeneous with planar contacts against dark marls and laminated shale beds.

Normally the dark marls (TOC mean = 2.35% $\pm$ 0.41, range = 0.51 to 6.51%, N = 39; CaCO₃, mean = 40.1% $\pm$ 3.7, range = 19.9 to 61.0%, N = 39) are homogeneous like the light marls and friable, but can become fissile on weathering. However, Fig. 7a shows a laminated dark marl fabric, not visible in the field, consisting of abundant microspar lenses in places forming nearly continuous laminae that alternate with other laminae comprising clay and organic matter (Weedon, 1987a). The figure shows a *Chondrites* burrow with a homogeneous light marl filling that penetrates the dark marl fabric. Burrow mottles at the contacts of light and dark marl beds are common at Lyme Regis as both light marl burrow-fills within dark marl and *vice versa* (Hallam, 1964).

The laminated shales (TOC mean = $5.46\% \pm 0.77$, range = 1.53 to 12.80%, $N = 51$; CaCO_3 , mean = $35.5\% \pm 3.4$, range = 11.6 to 68.4%, $N = 51$), usually with sharp basal bedding contacts against dark marl or light marl, produce a brown streak when scratched and weather into fissile, sometimes paper thin, laminae. Sub-microscopically they have scattered calcite microspar lenses dispersed within wavy sub-millimetre laminae of clay and organic matter (Weedon, 1987a). Laminated limestone (TOC mean = $1.42\% \pm 0.67$, range = 0.90 to 3.00%, $N = 7$; CaCO_3 , mean = $77.2\% \pm 6.2$, range = 62.4 to 87.2%, $N = 9$) forms beds that have planar contacts with laminated shale beds or nodules that are enclosed entirely by laminated shale (Weedon, 1987a; Arzani, 2004).

----- Figure 7 near this position -----

Hallam (1987) and Arzani (2006) doubted the presence of rock-forming quantities of nannofossils as a possible precursor to the microspar. However, calcareous nannofossils, both as true coccoliths as well as *Schizosphaerella*, are well documented from throughout the Tilmanni to Bucklandi zone interval at Lyme Regis and on the West Somerset coast (Hamilton, 1982; Bown, 1987; Bown & Cooper, 1998; Van de Schootbrugge *et al.*, 2007; Clémence *et al.*, 2010). Indeed, Weedon (1986; 1987a; 1987b) argued that the microspar lenses in the dark marls and laminated shales represent neomorphosed zooplankton faecal pellets containing calcareous nannofossils. Very similar fabrics to Fig. 7a are known in the Kimmeridge Clay Formation of Dorset (Kimmeridgian to Tithonian, Fig. 2d, f and g of Pearson, Marshall & Kemp, 2004), which has abundant nannofossils (e.g. Young & Bown, 1991; Lees, Bown & Young, 2006). Aggregates of nannofossils surrounded by clay minerals and organic matter in the Blue Lias Formation (Figs 8b, 8c and 8d; figure 3 of Weedon, 1986; figures 6B and 6C of Arzani, 2006) have similar dimensions to the microspar lenses, supporting their interpretation as neomorphosed zooplankton faecal pellets (Pearson, Marshall & Kemp, 2004).

Weedon (1986; 1987a) argued that nannofossil preservation was best in the more organic carbon-rich laminated shales and dark marls, as confirmed recently by Clémence *et al.* (2010). The high organic-carbon contents may have helped the preservation of primary organic coatings on the coccoliths, which in turn prevented calcite dissolution by acidic pore waters during diagenesis (Bukri, Dent Glasser & Smith, 1982). In the less organic-carbon-rich sediments, partial or total microspar aggradation of nannofossils (e.g. Fig. 8b) was inferred to be the normal situation in the Blue Lias (Weedon, 1986; 1987a; 1987b). The homogeneity of the light marls and most of the dark marls was explained as resulting from burrowers mixing the organic-matter- and clay-rich laminae with faecal pellets. In the limestones and light marls, however, very rare isolated elliptical structures, representing partially corroded or overgrown coccoliths provide the only *in situ* evidence for calcareous nannofossil precursors to the microspar (Fig. 8e; Fig. 2.4B of Weedon, 1987a).

If most carbonate mud in the ‘offshore’ Blue Lias Formation was derived from very shallow coastal waters rather than from nannofossil material, different localities would be expected to have different rock-type compositions. However, the average %CaCO₃ and %TOC of the limestone, light marl, dark marl and laminated shales are statistically indistinguishable between the Devon/Dorset coast and the West Somerset coast sections (i.e. overlapping 95% confidence intervals in Fig. 1c). Therefore, considering the large difference in zonal and average bed thicknesses between these locations, the lithological composition of the offshore Blue Lias facies was not influenced by either the local net accumulation rates or by local differences in water depth. The consistency of facies between localities and across the range of accumulation rates represented is consistent with a primarily hemipelagic, rather than shoreline-influenced origin for the carbonate mud (Weedon, 1986).

The majority of the laminated limestone beds were formed by the coalescence of laminated limestone nodules (Weedon, 1987a; Arzani, 2004). However, at Lyme Regis, bed

H30 categorized as laminated limestone does not have internal lamination but is homogeneous and without macrofossils. As for laminated limestone beds H32 and H36, it is underlain and overlain by laminated shale, has sharp planar top and bottom contacts, and has exceptionally low, slightly negative magnetic susceptibility values. The lowest measured vol MS of -0.13×10^{-8} SI implies, by extrapolating the regression of Fig. 2a (Section 3.b), 97.3 %CaCO₃ and hence these limestones represent cementation of a primary sediment almost devoid of organic matter and clay. Partly due to the sharp, planar base Hesselbo & Jenkyns (1995) inferred a dilute turbidity-current origin for bed H30 (which they named 'Intruder'). However, in the absence of direct evidence for basal erosion, grading or cross-lamination alternative explanations for the deposition of almost pure carbonate mud should be considered.

4.c. Lateral and stratigraphic variations in limestone proportions

Unlike the marls and laminated shale beds, average limestone bed thicknesses are consistently close to 10-15 cm at all the logged localities and in every zone of the Hettangian (Fig. 1b). Hallam (1964; 1986) regarded the consistency in limestone bed thickness of the Blue Lias, independent of locality and zonal thickness, as indicative of a diagenetic rather than a primary control on limestone formation. Although average limestone bed thicknesses remain nearly constant, the proportion of limestone beds by thickness per ammonite zone varies substantially from place to place and stratigraphically (Fig. 1b). Areas with thinner biozones (lower net accumulation rate or 'condensed sections') have higher average proportions of limestone (Fig. 1b, figure 2 of Page, 1995). For example, all four ammonite zones of the Hettangian of Lyme Regis have a far higher proportion of limestone than those on the West Somerset coast (Fig. 1b).

At Lyme Regis, representing an area of low net accumulation rate, all the zones of the Hettangian are of similar thickness and the average bed thicknesses for all rock types are also

similar in the different zones (Fig. 1b). By contrast, on the West Somerset coast the oldest zones (Tilmanni and Planorbis) are much thinner than the succeeding (Liasicus and Angulata) zones; significantly, the large change in zonal thickness there is mirrored by changes in average bed thickness of the light and dark marls and the laminated shales (Fig. 1b).

The association of changes in average non-limestone bed thickness with changing zonal thickness is interpreted here as indicating a large change in net accumulation rates at the end of the Planorbis Zone times or start of the Liasicus Zone for sections on the West Somerset coast. In this study, only the base of the Liasicus Zone was logged at Lavernock (Fig. 3). However, the much greater thickness of the Liasicus Zone interval at Lavernock compared to the Tilmanni and Planorbis zones (Waters & Lawrence, 1987; Warrington & Ivimey-Cook, 1995) indicates a similar change in net accumulation rate to that of the West Somerset coast.

The Liasicus Zone is consistently associated with lower proportions of limestone than preceding and succeeding Hettangian zones in different localities (Fig. 1b, Hallam, 1960; Palmer, 1972). This difference in character has led to member-scale subdivisions of the Blue Lias Formation with locally applied names i.e. the Lavernock Shales (Richardson, 1905; Waters & Lawrence, 1987); the St Audries Shales (Palmer, 1972) and the Saltford Shales (Donovan, 1956; Ambrose, 2001).

In the lower Liasicus Zone, higher velocities on downhole sonic logs related to relatively higher clay content, and higher gamma-ray counts related both to thorium and potassium in the marls and shales and to uranium within organic matter, allow ready correlation of this portion of the Blue Lias across Britain (Whittaker, Holliday & Penn, 1985). The base of this mudrock-dominated interval is diachronous because, in the English Midlands, there is biostratigraphical evidence that it occurs within the Planorbis Zone (Old, Sumblar & Ambrose 1987; Ambrose 2001), whereas in West Somerset and Lavernock the base is close to the base of the Liasicus Zone (Figs 3 and 5).

Mapping of the oldest strata of the Blue Lias Formation on the London-Brabant Platform and the Radstock Plateau near Bristol shows that the Liasicus Zone in the Hettangian and Semicostatum Zone of the Sinemurian were times of accelerated onlap (Donovan, Horton & Ivimey-Cook, 1979; Donovan & Kellaway, 1984). An inferred sea-level rise during these zones was suggested by Hallam (1981) and broadly supported by Hesselbo & Jenkyns (1998) and Hesselbo (2008).

At Lyme Regis, in the Liasicus Zone, the zonal thickness and the average thicknesses of the various rock-types are similar in the overlying and underlying zones, but there is nevertheless a lower proportion of limestone (Fig. 1b). Therefore, rising sea levels were apparently associated with reduced probability of limestone formation. The near-constancy of net accumulation rate at Lyme Regis, as indicated by the near-uniform average bed thicknesses of the marl and laminated shale, presumably resulted from an underlying tectonic block maintaining a relatively level sea floor within turbulent-water depths (due to storm winnowing) despite sea-level rise (Sellwood & Jenkyns, 1975). Apparently, rising sea levels during the Liasicus Zone times at Lyme Regis led to slight increases in water depth and less probability of limestone formation, but the increase in accommodation space was too small to allow substantially increased net sedimentation rates.

On the West Somerset coast and at Lavernock, increased accommodation space and higher sedimentation rates during the Liasicus Zone account for locally greater thicknesses of mudrocks (i.e. greater zonal thicknesses and greater average thickness of marl and laminated shale beds) compared to the Tilmanni and Planorbis zones (Figs 1b, 3, 5a). In these localities, the rapid increase in net accumulation rates apparently resulted from faster, probably fault-related, subsidence that was coincident with the sea-level rise (i.e. the increased subsidence rates exaggerated the effect of sea-level rise on local water depths). Fault-related activity is documented for the Hettangian in the Wessex Basin (Jenkyns & Senior, 1991) and South

Glamorgan in South Wales (Wilson et al., 1990) and fault movement occurred at some point during the Jurassic associated with the Bristol Channel area (Waters & Lawrence, 1987; Bixler, Elmore & Engel, 1998). Furthermore, fossil methane seeps dating from the Bucklandi Zone in the Sinemurian might be related to fault movement on the West Somerset coast (Allison, Hesselbo & Brett, 2008; Price, Vowles-Sheridan & Anderson, 2008).

At Southam Quarry near Long Itchington, the majority of the Liasicus Zone strata consist of laminated shale (Figs 1b and 6). This observation might be thought to indicate that greater water depth due to sea-level rise in the Liasicus Zone was associated with the conditions required for the preservation of lamination (especially bottom-water dysoxia and anoxia, cf. Hallam & Bradshaw, 1979). However, in both West Somerset and Lyme Regis, although there is a reduction in the proportions of limestone in the Liasicus Zone, unlike Long Itchington the proportions of laminated shale are lower than in the succeeding Angulata Zone (Figs 1b, 4 and 5a and b; Weedon, 1987a). Hence, it is the formation of a lower proportion of limestones, not the development of laminated shales, which can be consistently related to rising sea level/greater water depth.

5. Intra-formational hiatuses

Weedon (1986; 1987a; 1987b) reasoned that differences between localities in the numbers of regular sedimentary cycles, thought to result from Milankovitch orbital forcing of climate, could be explained by the presence of intra-formational hiatuses. Although Hallam (1960; 1987) disagreed, there are numerous lines of evidence, summarized in the sub-sections below, for both intermittent sea-floor erosion and for missing stratigraphic intervals within the offshore Blue Lias Formation of southern Britain.

----- Figure 8 near this position -----

5.a. Shaw plot

In Table 1 there are 63 levels in the Hettangian and Lower Sinemurian, out of 130 available, where the same biohorizon level (i.e. biohorizon top or biohorizon base) has been located to within a few centimetres both at Lyme Regis and on the West Somerset coast. This stratigraphic refinement allowed construction of the Shaw plot in Fig. 8 that illustrates the correlation of these two sites. At Lyme Regis, the base of the Angulata Zone lies somewhere between the lowest level with recorded *Schlotheimia* at 11.06 m (base of bed H84) and the highest recorded level with *Waehneroceras* at 9.66 m (within bed H71), as indicated by short horizontal lines in Fig. 8. Using the line of correlation between St Audries and Lyme Regis, its location has been inferred to be close to 10.54 m (within bed H77 Fig. 4, Table 1) as indicated using the long grey horizontal arrow in Fig. 8. Alternatively, the line of correlation from Quantock's Head rather than St Audries Bay, with fewer biohorizon constraints, would imply that the base of the Angulata Zone lies close to 10.13 m at Lyme Regis (within bed H73).

Subzones on the West Somerset coast sections are thicker than those in the corresponding section at Lyme Regis, as indicated in figures at the bottom of Fig. 8. Treating the biohorizon picks as time lines, the amount of strata within each interval results from the combination of the sedimentation rates and the effects of intra-formational hiatuses: i.e. the net accumulation rate. Shallower slopes on the line of correlation indicate higher net accumulation rates on the West Somerset Coast relative to the Devon/Dorset coast.

In contrast to the Lyme Regis section, on the West Somerset Coast the much thicker Liasicus and Angulata zones compared to the Tilmanni and Planorbis zones are mirrored by the increased average thicknesses of the corresponding marl and shale beds (Fig. 1b, Section 4.c). Hence, the major decrease in the average slope of the line of correlation in Fig. 8 at the base of the Liasicus Zone is interpreted as due to substantially increased net accumulation rates in West Somerset. In detail, the breaks in slope (nearly horizontal or nearly vertical segments)

of the line of correlation are interpreted as due to hiatuses or intervals of exceptionally low accumulation rate, and have been labelled with the corresponding bed numbers. These breaks in slope of the line of correlation are discussed in Section 5.c. The breaks in slope are defined by multiple points, significantly reducing the likelihood that they simply result from misplacing the tie-levels due to collection failure for key ammonites.

5.b. Minor erosion

Several lines of field evidence for minor sea-floor erosion were used by Weedon (1986) to argue for storm-induced bottom-water turbulence. Protrusive *Diplocraterion* burrow mottles, which are found only within the limestones and light marls, have been described at all four localities studied here as well as at Saltford, Avon (Fig. 9c, Sellwood, 1970; Donovan & Kellaway, 1984; Weedon, 1987a; Mogadam & Paul, 2000; Barras & Twitchett, 2007). Tiering analysis at Lyme Regis (Mogadam & Paul, 2000) showed that *Diplocraterion* cross-cuts all other trace fossils except *Chondrites* which is well known to have been produced by late-stage deposit feeders within anoxic sediments (Bromley & Ekdale, 1984). The observations of Mogadam & Paul (2000) suggest re-excavation of the vertical U-shaped burrow following rapid centimetre-scale sediment removal (Seilacher, 2007).

At Lyme Regis, dark marl beds that are only a few centimetres thick are locally missing for several metres laterally (Weedon, 1986). Within limestone bed 23 ('Mongrel') locally a thin dark marl bed is replaced laterally by centimetre-scale, metre-wide scours filled with bioclastic packstone. The former presence of a thin bed of dark marl is in some places recorded by a layer of isolated dark marl burrow mottles within what otherwise appears to be a single limestone bed (e.g. in bed 19, Fig. 9d). In West Somerset, this phenomenon is occasionally observed in thick light marl beds but more typically an opposite arrangement occurs with isolated bands of light burrow-fills within dark marls representing relics of thin light marl beds.

Paul, Allison & Brett (2008) also noted that occasional ‘exotic’ sediment fills within uncrushed ammonites at Lyme Regis could indicate material subsequently removed by erosion.

Radley (2008) argued that at Southam Quarry, Long Itchington, erosion within laminated shale was linked to distal storm influences below normal storm wave-base. He described siltstone scour-fills within Liasicus Zone laminated shales as well as highly elongated limestone nodules containing imbricated ammonites, which were interpreted as gutter casts. The scour fills are associated with a trace fossil indicative of a crustacean escape structure and therefore very rapid deposition (Radley, 2008; O’Brien, Braddy & Radley, 2009). Similar presumed scour fills, full of ammonites and shell fragments, within light marl beds are also known in West Somerset, albeit mainly in the Planorbis Subzone (e.g. bed 24).

A scour fill at Lyme Regis in the top of a limestone bed, probably from the Rotiforme Subzone, Bucklandi Zone, consists of a shallow depression around 50 cm across filled with coarse shelly debris and abundant small articulated echinoids (?*Diademopsis*) and common articulated ophiuroids (Page collection, Bristol City Museum and Art Gallery). These echinoderms retained their articulation due to rapid burial, presumably as waning storm-currents deposited previously suspended sediment. A second level with common, but more scattered *Diademopsis* in laminated mudstone bed 17 (Planorbis Zone and Subzone) near Watchet, West Somerset, may indicate a similar event, although there is no equivalent concentration of shelly debris (K. N. P. pers. obs. 2015).

Found commonly throughout the sections on the Devon/Dorset and West Somerset coast are centimetre-scale horizons with: a) concentrated bivalve fragments; b) lenses of echinoid debris; c) abundant large ammonites on the surface or base of limestone beds, and/or d) scattered large ammonites and nautiloids with encrusting oysters and/or crinoid debris on the tops of limestone beds. Some of these features have been previously recorded by Paul, Allison & Brett (2008) and they indicate either periods of increased sea-floor turbulence that

led to winnowing of fines, fragmentation of shells and the concentration of bioclasts or periods of very slow, or halted deposition. In the latter case, epifauna such as oysters and crinoids had time to attach to the ‘benthic islands’ created by large shells. Such observations can coincide with additional evidence for significant intervals of missing strata.

5.c. Major hiatuses

The following sub-sections summarize the Shaw plot and field evidence for significant hiatuses at many levels within the Hettangian and Lower Sinemurian offshore facies of the Blue Lias of southern Britain. Note that in some cases these major hiatuses can be potentially linked to sea-level rises and in others to sea-level falls.

5.c.1. Tilmanni and Planorbis zones, Hettangian

A widespread, diachronous erosion surface across southern Britain at the base of the Blue Lias Formation has been explained in terms of latest Triassic sea-level fall followed by a rise in the earliest Jurassic (Tilmanni Zone, Donovan, Horton & Ivimey-Cook, 1979; Hallam, 1981; 1997; Wignall, 2001; Hesselbo, Robinson & Surlyk, 2004). In the Bristol area at several localities (Filton railway cutting, Chipping Sodbury, Henleaze, Wick and Doynton), Donovan & Kellaway (1984) recorded Johnstoni Subzone (i.e. upper Planorbis Zone) strata directly on top of the ‘Pre-planorbis Beds’ (i.e. the Tilmanni Zone). Widespread condensation by winnowing of mud-grade sediment during the Tilmanni and Planorbis zones apparently led to numerous centimetre-scale horizons of small bivalve fragments and closely spaced limestone beds at Lyme Regis, West Somerset Coast, and Lavernock (Kent, 1937; Hallam, 1960; Hesselbo & Jenkyns, 1995; 1998). Furthermore, Kent (1937) noted limestone intraclasts encrusted by serpulids as well as borings, indicative of sea-floor erosion and non-deposition,

within the Tilmanni and/or Planorbis zones of Nottinghamshire (the precise stratigraphic levels not being reported).

The apparent thinness of the *Neophyllites*-bearing levels at the base of the Planorbis Subzone (biohorizons Hn3 and Hn4, bed 9 of Whittaker & Green, 1983), suggests that there is condensation at this level in West Somerset (Page, 1995; Bloos & Page, 2000a). This supposition is confirmed by a corresponding break in slope in Fig. 8. A similar break in slope, implying a gap in the West Somerset coast section, occurs at the base of the Johnstoni Subzone (bed 25). A break in slope associated with bed 25 is also found on a Shaw plot of the West Somerset coast versus Lavernock (Table 1 data, not illustrated). Currently, there is no independent biostratigraphical evidence for a gap at this level on the West Somerset coast but, as noted in Section 5.b, presumed scour fills are present in bed 24.

5.c.2. Liasicus Zone, Hettangian

Liasicus Zone strata have not provided much field evidence for stratigraphic gaps. Radley (2008) noted in Southam Quarry, near Long Itchington, the presence of limestone nodules encrusted by serpulids and oysters (*Liostrea*) with scratches and grooves attributed to crustaceans. This evidence certainly indicates sea-floor exhumation of the nodules and a period of non-deposition (though this might have been short-lived). He inferred an Angulata Zone age, assuming causal factors such as lowered sea level rather than sediment starvation. However, comparison of the stratigraphic level shown (figure 5 of Radley, 2008) for the encrusted nodules with Fig. 3, combined with the biostratigraphy of Clements *et al.* (1975; 1977), indicates that this erosion and non-deposition occurred during the Liasicus Zone.

Breaks in slope in Fig. 8 indicate possible hiatuses within the Liasicus Zone at Lyme Regis associated with unnamed limestone beds H58 and H68. Consistent with increased bottom-water turbulence, if not direct evidence for non-deposition, is the presence of abundant

macroconch *Waehneroceras* commonly with encrusting *Liostrea* on the top surface of bed H58, as well as lenses of echinoid debris, especially spines in a few centimetres of the overlying marl. Figure 8 also indicates a stratigraphic gap at the base of bed 67 at the base of the Laqueus Subzone on the West Somerset Coast. Two metres below bed 67, the top of bed 65 (Fig. 5a) also yields common macroconch *Waehneroceras*.

Hence, there was apparently increased bottom-water turbulence at the end of the Portlocki Subzone and beginning of the Laqueus Subzone in both relatively slowly and relatively quickly subsiding areas. Since bed H68 at Lyme Regis and beds 65 to 67 on the West Somerset coast occur at the levels of increased MS (Section 3.b), the mid-Liasicus increase in bottom-water turbulence was apparently associated with a change in sediment composition. Sequence stratigraphic analysis of the marginal facies in Glamorgan (Section 4.a) led Sheppard (2006) to infer a major flooding surface in the mid Liasicus Zone – consistent with the analysis by Hesselbo & Jenkyns (1998) of coeval marine strata from across Britain.

5.c.3. Angulata Zone, Hettangian

Correlation of a downhole log from the borehole at Burton Row in north Somerset, 20 km north east of Quantock's Head (Fig. 1a), with the lithological log of Weedon (1987a) from the Devon coast suggested to Smith (1989) that a gap exists in the Angulata Zone at Lyme Regis. Deconinck *et al.* (2003) inferred, using kaolinite/illite ratios from the West Somerset coast, that a stratigraphic gap occurs at Lyme Regis at the top of the Angulata Zone. Hesselbo & Jenkyns (1995; 1998) also argued for condensation and thus reduced accumulation rates in the middle Angulata Zone at Lyme Regis, as indicated by the unusually close spacing of limestone beds and nodule-bearing horizons (e.g. bed 1 or 'Brick Ledge').

Figure 8 confirms the presence of stratigraphic gaps in the Angulata Zone at Lyme Regis with at least at one level in the Complanata Subzone (bed 1c within 'Brick Ledge') and

at two levels within the Depressa Subzone (bed 7c or ‘Lower Skulls’, and bed 17 or ‘Upper White’). The Shaw plot also indicates a gap at bed 134 in the Depressa Subzone on the West Somerset coast (Fig. 8), a level characterized by abundant large macroconch ammonites (*Schlotheimia*), again suggesting low net accumulation rates.

----- Figure 9 near this position -----

5.c.4. Bucklandi Zone, Sinemurian

Direct evidence for erosion at Lyme Regis during the Conybeari Subzone, Bucklandi Zone, is provided by a limestone intraclast with the size and shape reminiscent of a small limestone nodule, but found *in situ* within limestone bed 25 (‘Top Copper’, Fig. 9a, Weedon, 1987a). The intraclast surface has the characteristics of a hardground as it is bored both by bivalves (possible *Lithophaga* crypts, Fig. 2.5e of Weedon, 1987a) and by *Talpina ramosa* Von Hagenow (150–250 µm diameter, possibly made by phoronids). The intraclast surface is encrusted by at least two generations of *Liostrea*, which also have *Talpina ramosa* borings (Figs 9ai and aii). The intraclast surface is directly overlain by an unusual packstone of small gastropods, benthic foraminifera and larger bivalve fragments and succeeded by normally bioturbated wackestone. Thus, it appears that a period of increased turbulence led to exhumation of a limestone nodule, exposure for sufficient time for repeated colonization and boring (over at least two years based on the two or three generations of bivalves), followed by burial whilst the turbulence was sufficient to lead to generation of the packstone (i.e. grain-supported fabric) before a return to typical carbonate mudstone and wackestone deposition.

There is no biostratigraphic evidence for missing strata at the level of bed 25 at Lyme Regis containing the limestone intraclast (Fig. 9a). However, large ammonites are common in the limestone bed 153 on the West Somerset coast, which correlates biostratigraphically with bed 24 at Lyme Regis (i.e. Biohorizon Sn3b with *Metophioceras* ex *grp rouvillei* (Reynès),

etc). In Fig. 8, the kink in the line of correlation at the level of bed 153 (at about 81.5 m) indicates condensation of the West Somerset section relative to Lyme Regis. Hence, erosion/non-deposition was apparently simultaneous at Lyme Regis and on the West Somerset coast during the middle part of the Conybeari Subzone. The break in slope of the line of correlation was determined more by sediment condensation at the level of bed 153 in West Somerset than by erosion associated with beds 24 and 25 at Lyme Regis.

In bed 29 at Lyme Regis ('Top Tape', Fig. 4), also within the Conybeari Subzone, Hallam (pp8–9, 1960) noted the presence of glauconite, both as discrete 'granules' and in microfossil infills, this mineral being an indicator of reduced accumulation rate (Odin & Matter, 1981). At beach level on either side of Seven Rock Point, west of Lyme Regis, bed 29 forms a prominent ammonite 'pavement' covered with large specimens of *Metophioceras* ex gr. *conybeari* (J. Sowerby) (Sn5b *conybeari* Biohorizon: Page 2002, 2010b; 2010c), and this is matched by its correlative bed 161 in West Somerset, which has the same group of species in abundance on its base. The equivalent level is also recognizable palaeontologically near Pilton in north Somerset (pers. obs. by K.N.P., 1992) and correlates with the Calcaria Bed in the Bristol-Bath area where the widespread evidence for truncation, condensation and phosphatization of fossils has been specifically attributed to shallowing (Kellaway & Donovan, 1984). Figure 2 of Page (1995) clearly illustrates, for the Hettangian/Sinemurian boundary interval of the West Somerset Coast, Lyme Regis area and Saltford Cutting near Bristol, the way that lateral loss and condensation of biohorizons is associated with an increased proportion of limestone compared to marl and shale.

A widespread late Angulata–early Bucklandi erosive phase is also indicated by the non-sequences and condensation at this level described in eastern France, southern Germany and Austria, contrasting with the relative completeness of the Sinemurian GSSP section on the West Somerset coast (Page *et al.*, 2000; Bloos & Page, 2002). An indication for shallowing at

this time is also provided in South Wales near Cardiff where typical offshore Blue Lias of Unit A of the Porthkerry Formation in the Angulata Zone is overlain by the near-shore Unit B facies (i.e. limestone dominated and lacking laminated shales and laminated limestones, Section 4.a). Unit B and C facies are, in turn, overlain by uppermost Angulata Zone and Bucklandi Zone oolitic marginal facies suggesting further shallowing (Waters & Lawrence, 1987; Wilson et al, 1990).

In the Cardiff area, the Porthkerry Formation with Bucklandi and Semicostatum Zone ammonites is found on top of a bored surface on the oolitic marginal facies (Waters & Lawrence, 1987). Similarly, near Southerndown Porthkerry Formation Unit D of the same zones (normal offshore Blue Lias with about 60 % nodular limestones and marls) occurs on top of marginal facies (Wilson et al., 1990). The nature of these successions is consistent with a rise in relative sea level at the end of the Bucklandi Zone.

5.c.5. Early Semicostatum Zone, Sinemurian

As reviewed by Donovan & Kellaway (1984, e.g. their figures 5 and 6), a non-conformity in the Bristol area is overlain by phosphatised sediments and fossils with truncation of the Bucklandi Zone and successive overstepping of all the Hettangian biozones and the Rhaetian onto the Radstock Plateau to the south (i.e. approaching the Carboniferous Limestone massif of the Mendip Hills).

As evidence for coeval erosion and condensation at Lyme Regis, Hallam (1960) illustrated the truncation of a *Diplocraterion* burrow, and noted glauconite and phosphate associated with bed 49 ('Grey Ledge'). This level is now known to be associated with up to four missing biohorizons from the top of the Bucklandi Subzone to the base of the Scipionianum Subzone, within the lower part of the Semicostatum Zone (Page, 2003, 2010b; 2010c). Additionally, on the Devon/Dorset coast, Gallois & Paul (2009) showed that there are

lateral variations in the amount of strata removed at this level from the uppermost Blue Lias Formation and that locally there is a bored surface on bed 49 and, throughout the area, deep *Diplocraterion* burrows are present, commonly at more than one level in beds 47–49. It should be noted, however, that the ammonite faunas recorded through this interval indicate that these levels may be of slightly different ages in different places and hence what are termed beds 47, 48 or 49 in different places may not be exactly the same lithostratigraphical entities. Hallam (1981) linked the change in facies above bed 49 to both the accelerated onlap on the London Platform described by Donovan, Horton & Ivimey-Cook (1979) and to sea-level rise.

There is no evidence for a break in sedimentation in the late Bucklandi to basal Semicostatum zones on the West Somerset coast where the biohorizon succession appears to be complete. Interestingly, however, there is a concentration of ammonites at the level of bed 247 of Whittaker & Green (1983) in the Lyra Subzone (Page 1992; 2009, biohorizon Sn 15b of Page, 2010b), which may indicate at least one phase of reduced accumulation rates in the early Semicostatum Zone. There are several levels in the 5–6 m of shale-marl/limestone alternations, below which are concentrations of shelly debris, with common large ammonites, and one prominent level of *Diplocraterion* burrows in bed 242 (K.N.P. pers. obs. 2013–2015). Notably, the ammonite fauna of bed 247 (e.g. *Paracoronicerias* ex gr. *charlesi* Donovan) is very close in terms of biohorizonal assignment to that recorded from the top c. 6–10 cm of bed 49 at the ‘Slabs’, between Axmouth and Lyme Regis (Page, 2002), immediately below the non-sequence and with Scipionianum Subzone strata immediately above. The lower part of bed 49, however, still appears to belong the Bucklandi Subzone (and Zone) because large *Arietites* are common.

6. Preservation of ammonites

It has long been believed that since the macrofossils, especially ammonites and nautiloids, are largely uncrushed, and horizontal sections of burrow mottles are nearly circular in limestone beds of the Blue Lias Formation, cementation must have started prior to significant compaction, and hence was both early and at shallow depth (Kent, 1936; Hallam, 1964; Weedon 1987a; Arzani, 2006; Paul, Allison & Brett, 2008). Since the uncrushed ammonite and gastropod shells in the limestones would have originally been aragonite, the cementation must also have pre-dated dissolution or inversion to calcite.

At Lyme Regis, ammonites completely enclosed within limestone beds typically have their originally aragonitic shells replaced by sparry calcite and the inner and outer whorls are uncrushed. By contrast, ammonites diagenetically ‘welded’ onto the surfaces of such beds usually present an uncrushed internal mould of the outer whorls with the shell entirely absent. Importantly, their inner, septate whorls are represented predominantly by an external mould on the surface of the limestone bed.

This differential preservation according to the location of the ammonite specimens can be explained by the carbonate cementation generating a framework-supporting fabric that formed first at the centres of (i.e. mid-way through) what became the limestone beds. When and where the cementation and framework support occurred before aragonite dissolution, the full three-dimensional shape of the aragonitic ammonite shell was preserved, allowing for a later sparry calcite replacement and infill. However, on the outer margins of limestone beds, cementation and framework support appears to have been later than aragonite dissolution, albeit before significant compaction. Consequently, although this later cementation could preserve the three-dimensional shape of the shell where already filled with marl (in the body chamber and, in some cases, perforated sections of the phragmocone), it could not preserve the thickness of the shell or support inner whorls that lacked internal sediment. Arzani’s (2006)

study of the limestone micro-fabrics also showed that, despite the typically uncrushed nature of the macrofossils, cementation of the edges of the limestones did occur during compaction.

Although a cursory examination of the Devon–Dorset and West Somerset sections may give an impression that ammonites do not occur in every limestone bed, they have been recorded at far more levels than noted by Paul, Allison & Brett (2008) at Lyme Regis (Page, 2002). Ammonites are common only in certain beds but prolonged examination over many years by K.N.P. has led to the recovery of very rare isolated specimens from many levels throughout the Devon–Dorset and West Somerset coastal successions. The simplest explanation for the apparent lack of ammonites within some limestone beds at Lyme Regis is preservation failure caused by cementation following a combination of aragonite dissolution, burrowing and compaction that destroyed all trace of the shells. Which leads to the critical question: why do some limestones record cementation before aragonite dissolution and others do not? This issue is re-visited in Section 8.

Paul, Allison & Brett (2008) suggested that the apparent lack of dissolution and encrustation of uncrushed ammonite and nautiloid shells within limestone beds, given the average bed thickness, implies much more rapid sedimentation of these lithologies than the net rate for the formation as a whole. As evidence for rapid sedimentation, they cited the presence of ammonite shells preserved in a near-vertical orientation within limestone beds, as well as cases of ammonite imbrication. However, their observations were primarily based on bed 41 (‘Best Bed’, Bucklandi Zone), which has abundant small ammonites (including *Vermiceras scylla* (Reynès)) that would have been readily re-oriented by large burrowers. Just two additional beds in the Johnstoni Subzone in the Planorbis Zone have common small ammonites. Otherwise the limestone beds typically have a large-bodied ammonite fauna.

In addition, although encrustation on ammonites appears to be mainly observed on the surfaces of limestone beds, its reported absence from specimens *within* limestone beds (Paul,

818 Allison & Brett 2008) is not necessarily evidence for rapid sedimentation because biologically
819 relevant factors may have been influential. In fact, the benthic macrofauna of the limestone
820 beds is commonly characterized by low diversity and locally high density, indicative of either
821 a ‘high-stress’ environment or opportunistic colonization during brief periods of ‘improved’
822 conditions. For example, rhynchonellid brachiopods dominate bed 11 and the tops of beds 19
823 and 23 at Lyme Regis, beds 147 and 151 on the Somerset Coast and bed 21c at Long Itchington,
824 whereas *Gryphaea* dominates at the tops of bed 13 and beds 14 to 17 and 31 at Lyme Regis
825 and at the tops of beds 136 and 138 on the Somerset Coast. It is likely that these controls on
826 benthic colonization were due to intermittent bottom-water dysoxia which would have also
827 inhibited encrustation of ammonites. There are some exceptions to these general observations,
828 however, as there is evidence that some of the rhynchonellid brachiopods may have inhabited
829 *Thalassinoides* burrows preferentially. Additionally, occasional winnowing may have
830 concentrated some of the shelly material, for instance in bed 31 at Lyme Regis. At this latter
831 level, *Gryphaea* apparently show current sorting as they are dominated by the larger, heavier
832 lower valve in random orientations throughout the bed – possibly indicative of rapid deposition
833 after a storm (C. R. C. Paul, pers. comm. 2016).

834 In the light and dark marls and the black laminated shales, without the early cementation
835 of the limestones, the aragonitic fossils were much less likely to be preserved and hence subject
836 to compaction and early diagenetic dissolution. At Lyme Regis in particular, ammonites are
837 absent from most of the light and dark marls, but in some of the laminated shales very poorly
838 preserved shell-less impressions can sometimes be found. Scattered oyster xenomorphs in the
839 marls, however, prove that ammonites were once present (Paul, Allison & Brett 2008).
840 Dissolution of aragonite and bioturbation and hence destruction of any remaining trace of shell
841 can be assumed to have removed most of the ammonites from such levels, especially when no

842 traces of other aragonitic shells, such as bivalves, are observed (Wright, Cherns & Hodges,
843 2003).

844 In West Somerset, however, the situation is more complex. White aragonitic shells
845 persist in laminated shales through much of the succession, although they are only well
846 preserved and iridescent at certain levels in the Planorbis and middle Liasicus zones. By
847 analogy with calcareous nannofossil preservation (Section 4.b), the preservation of iridescent
848 aragonite is likely to be connected to the high organic-carbon contents of the laminated shales,
849 as has proven to be the case in other Jurassic shale sequences from the British Isles (e.g. Hall
850 & Kennedy, 1967; Hudson & Martill, 1991).

851 Commonly, on the West Somerset coast in the laminated shales and especially in the
852 marls, only an impression of the shell remains, picked out by a residual brown organic film:
853 the remains of the organic content of the shell, in particular the periostracum. This pattern of
854 preservation indicates that the aragonite dissolution post-dated the end of bioturbation in the
855 marls. In the upper Bucklandi and Lyra subzones, levels with white aragonite preservation can
856 pass laterally into areas with brown calcitic shell preservation as major faults are approached.
857 In such cases, the inversion of aragonite to calcite in the marls and shales is clearly associated
858 with the circulation of fluids along such faults, for instance during the tectonic inversion of the
859 basin, as described by Bixler, Elmore & Engel (1998). At some levels, both brown calcitic and
860 white aragonitic shells may also contain pyrite, commonly associated with specific layers in
861 the shell.

862 In the Burton Row borehole in North Somerset, pristine iridescent aragonite is
863 characteristic of the laminated shales of most of the Hettangian and Sinemurian succession. On
864 the coast, inversion to brown calcite (e.g. in parts of the Liasicus and Semicostatum zones) was
865 due to the effects of fluids and/or raised temperatures generated during tectonic inversion
866 (Water & Lawrence, 1983; Bixler, Elmore & Engel, 1998), whereas conversion of iridescent

aragonite in the laminated shales to white and powdery aragonite was probably due to near-surface weathering and/or the effects of ground-water penetration.

----- Figure 10 near this position -----

7. Oxygen and carbon isotopes and timing of limestone formation

7.a. Lyme Regis

Figures 2b and 10a show that, at Lyme Regis, oxygen isotopes in the centres of the limestone beds are as high as -1.5 ‰ VPDB with carbon-isotope values close to 0.0 ‰ VPDB. These values are not much lighter than the isotopes of unaltered benthic bivalve (*Gryphaea*) calcite (Weedon, 1987a). Hence, carbon and oxygen isotopes of the Blue Lias, combined with the near-absence of evidence for compaction of macrofossils, have long been considered consistent with early cementation (Campos & Hallam, 1979; Gluyas, 1984; Weedon, 1987a; Arzani, 2006; Paul, Allison & Brett, 2008). The decreases of oxygen-isotope values towards the edges of limestone beds, where calcium carbonate contents are lower, are consistent with cementation occurring progressively during the early stages of compaction (Gluyas, 1984; Weedon, 1987a; Arzani, 2006; Paul, Allison & Brett, 2008). Disseminated framboidal and euhedral pyrite found throughout the limestones and the non-ferroan nature of the calcite microspar indicate at least some carbonate generation during sulphate reduction (Gluyas, 1984).

Raiswell (1988) postulated that cementation of the Blue Lias limestone beds and nodules was associated with anaerobic methane oxidation by sulphate within the sulphate reduction zone. Both normal sulphate reduction and anaerobic methane oxidation generate carbonate that neutralizes the acidity produced during the precipitation of iron monosulphide and pyrite oxidation (Raiswell, 1988; Bottrell & Raiswell, 1989). Since there is a restricted depth interval involved, anaerobic methane oxidation was used by Raiswell (1988) to explain

Hallam's (1964) observations of the limited range of limestone bed thickness. In particular, Raiswell's model requires that shallow cementation (<1 m burial) occurred during a pause in sedimentation.

Bottrell & Raiswell (1989) further demonstrated that pyrite formation occurred over a longer period in the limestones than in the light marl and dark marls. To explain the sulphur-isotope composition of the pyrite in the limestones they invoked incorporation of sulphur derived from the bioturbational oxidation of previously formed pyrite. Such a process implies limestone cementation close to the zone of burrowing and hence not greatly below the sediment-water interface. $\delta^{13}\text{C}$ of carbonate from sulphate reduction or aerobic methane oxidation is typically less than -10.0 ‰ VPDB. Bottrell & Raiswell (1989) explained the relatively heavy carbon-isotope values of around 0.0 ‰ VPDB in the limestones as indicating that the bulk of the carbonate was derived from dissolution of aragonite and high-Mg bioclasts. Later authors concurred with this suggested source of carbonate (Wright, Cherns & Hodges, 2003; Arzani, 2004; 2006; Paul, Allison & Brett, 2008). Some limestone beds at Lyme Regis contain centimetre-scale cracks partly filled with bioclastic sediment, indicating that cementation and fissuring occurred close enough to the sediment-water interface to allow access to unconsolidated material, presumably following a phase of sea-floor erosion (Hesselbo and Jenkyns, 1995).

Plots of isotope ratios in the Blue Lias Formation show measurements lying near a trend-line that slopes from near the origin down towards more negative oxygen- and carbon-isotope ratios. Trends in isotope composition from heavier oxygen isotopes at the centres of nodules to lighter oxygen isotopes at the edges (Fig. 10, Weedon, 1987a, Arzani, 2006) apparently confirm the supposition that oxygen isotopes became lighter through time, most likely due to increasing pore-water temperatures. However, the carbon isotopes in limestone nodules can increase rather than decrease towards the outside of nodules and consequently do

not lie along the main trend of measurements at Lyme Regis (Fig 10a) and West Somerset (Fig. 10c).

Assuming that oxygen-isotope values only decreased during the diagenesis of the Blue Lias Formation, an explanation is needed for why the limestone beds can have average carbonate with $\delta^{18}\text{O} = -4.5$ and $\delta^{13}\text{C} = -1.5$ ‰ VPDB while limestone nodules can have average carbonate with the same $\delta^{18}\text{O}$, but $\delta^{13}\text{C} = -3.0$ ‰ VPDB (Fig. 10a). The key to this problem is to regard the main trend in the isotopes in the limestone beds as indicating a mixing line between the composition of early and late carbonate rather than as a time series.

It is now recognized that early meniscus calcite cement associated with carbonate nodule formation could have provided a supporting framework that allowed the bulk of cementation to occur much later during compaction (Curtis *et al.*, 2000; Raiswell & Fisher, 2000). This model helps explain why it is so rare to find shallow limestone nodules forming in modern organic-rich marine sediments (Raiswell & Fisher, 2000). In the Blue Lias Formation, the very early cement in the limestones with oxygen and carbon isotopes relatively close to 0.0 ‰ VPDB was apparently sufficient to prevent the compaction of large macrofossils. At a later phase of diagenesis, once compaction had started in earnest, the remaining pore spaces within the limestone beds could then be partly filled with carbonate that had much lighter oxygen and, typically also lighter carbon isotopes. This model means that, if cemented relatively quickly, limestone nodules can show progressive changes in isotopes that, unlike the limestone beds, record a snapshot of the history of changing pore-water isotopic composition in concentric zones (including pore-water carbon isotopes becoming heavier while oxygen isotopes became lighter). Although Arzani (2006) demonstrated that microspar growth was displacive, the cementation cannot have continued much beyond the end of occlusion of the pore spaces because the limestone beds and nodules have oxygen isotopes not lighter than about -6.0 ‰.

At Lyme Regis, the oxygen isotopes of the calcite microspar in the light and dark marls and the black laminated shale (-3.5 to -6.0 ‰ VPDB) are much lighter on average than the limestone beds (Fig. 10a). Hence, the marls and shales appear to have been cemented later than the limestone beds. However, one sample of light marl from mid-way through bed 15c (BL307, Table 3B of Weedon, 1987a, Figs 2b and 10a) with %CaCO₃ = 38.2 and %TOC = 0.61 has an oxygen-isotope value that is far heavier ($\delta^{18}\text{O}$ = -1.99 ‰ VPDB) than the other marl and shale samples. This sample is interpreted to represent light marl that was cemented by early meniscus cement, as though destined to become the centre of a limestone bed or nodule, but the amount of cement was apparently insufficient to provide enough framework support to prevent compaction during burial.

Ferroan and non-ferroan calcite beef occurring near the bases of some laminated shales at Lyme Regis with $\delta^{18}\text{O}$ less than -6 ‰ (Campos & Hallam, 1979), not shown in Fig. 10a, probably formed as a result of over-pressuring during deep burial (Marshall, 1982).

7.b. West Somerset Coast

Unaltered carbonate of *Liostrea* from the Tilmanni Zone of the West Somerset Coast have generally much heavier carbon isotopes (+1.6 to +3.9 ‰, Fig. 10c, van de Schootbrugge *et al.*, 2007) than those found in *Gryphaea* from the Angulata and Bucklandi Zones at Lyme Regis (+0.5 to +2.5 ‰ VPDB, Figs 2b and 10b, Weedon, 1987a). Korte *et al.* (2009) measured $\delta^{13}\text{C}$ of between +1.5 and +5.0 ‰ VPDB for *Liostrea* in the Tilmanni Zone of Lavernock (not shown) with progressively lighter average values in the upper part of the Tilmanni, Planorbis and lower Liasicus zones. The different ranges of values in the Tilmanni and the Angulata–Bucklandi zones have been related to secular trends in carbon isotopes within the Hettangian rather than diagenetic factors (Korte *et al.*, 2009). However, all three groups of measurements of unaltered benthic bivalve calcite indicate $\delta^{18}\text{O}$ between about 0.0 and -2.0 ‰ VPDB.

The limestones on the West Somerset Coast with $\delta^{18}\text{O} = -4.0$ to -12.5 ‰ VPDB have far lighter oxygen isotopes than those at Lyme Regis (compare Fig. 10a with Figs 10b and 10c). The isotopically lightest values from a limestone bed (Fig. 10b) are derived from beds associated with mud volcanoes developed in the Bucklandi Zone strata near Kilve (Cornford, 2003; Allison, Hesselbo & Brett, 2008; Price, Vowles-Sheridan & Anderson, 2008). Isotope values from material described as comprising the mud volcano mounds themselves (i.e. breccia, 'tufa', crust etc) have been excluded from Fig. 10b.

Oxygen isotopes from limestone beds from the Tilmanni and Planorbis Zones on the West Somerset Coast are not only lighter than the limestone beds at Lyme Regis, but also generally lighter, rather than heavier, than their associated marls and black laminated shales (Fig. 10c, Arzani, 2004). The much lighter oxygen isotopes suggest that the limestones were cemented later than the marls and shales. Nevertheless, large macrofossils within the limestone beds of the West Somerset Coast are preserved uncrushed in the same manner as observed at Lyme Regis. Accordingly, cementation started prior to compaction. These apparently conflicting interpretations can be reconciled if it is hypothesised that the West Somerset Coast limestone beds had an early framework cement with oxygen isotopes close to 0.0 ‰ VPDB that was joined by carbonate formed much later that had much lighter oxygen-isotope values. Thus, the marls and laminated shales became fully compacted and fully cemented at a time when there remained framework-supported pores within the limestone beds still available for later cementation by calcite with very light oxygen-isotope values.

Campos & Hallam (1979) explained their observations of much heavier average oxygen isotopes in the limestones of the Lower Jurassic of Devon/Dorset compared to the Lower Jurassic of the Yorkshire Coast in terms of net sedimentation rate. Their explanation is followed here with respect to the comparison of the West Somerset with the Devon/Dorset coast, assuming that the total time for cementation was approximately the same in both cases. In both

991 areas, cementation appears to have started very early, but the limestones at Lyme Regis were
992 buried more slowly and so their final carbonate was probably precipitated at relatively low
993 temperatures (overall oxygen isotopes heavier than the marls and shales). The more rapidly
994 buried limestone beds on the West Somerset Coast apparently ceased full cementation at much
995 greater depths and thus were formed from pore waters with higher temperatures (overall
996 oxygen isotopes lighter than the marls and shales).

997 A prolonged cementation history on the West Somerset Coast is indicated by very
998 negative $\delta^{18}\text{O}$ in vein calcite (-9 to -12 ‰ VPDB - not shown in Fig. 10), probably associated
999 with Late Jurassic to Early Cretaceous extensional and Cenozoic compressional faulting
1000 (Bixler, Elmore & Engel, 1998).

1002 8. Synthesis

1003 In the offshore facies, at the scale of ammonite zones, lateral and stratigraphic variations
1004 in the characteristics of the Blue Lias Formation are not due to variations in the composition
1005 of specific rock types. Neither the water depth nor the net accumulation rate influenced the
1006 average composition of the rock type (Fig. 1c), a conclusion that follows from the hemipelagic
1007 origin of the sediment (Section 4.b). The key variables determining the characteristics of the
1008 Formation are: a) the average thicknesses of beds of light marl, dark marl and black laminated
1009 shale; and b) the proportion of limestone.

1010 Variations in the average thickness of the marls and shales are associated with
1011 variations in zonal thickness and were determined by the net accumulation rate (Section 4.c).
1012 The uniformly and relatively thin zones of the Hettangian at Lyme Regis are associated with
1013 thin beds of marls and shales. On the West Somerset Coast, the younger stratigraphically
1014 thicker zones are associated with thicker beds of marls and shales whose deposition post-dated

a major increase in accumulation rate at the start of the Liasicus Zone (Figs 1b and 8, Section 4.c).

On the scale of ammonite zones, the proportion of limestones was linked to the net accumulation rate as determined by rates of subsidence and sea-level variations. The abundant evidence for intra-formational hiatuses and evidence for storm-related scours and winnowing episodes (Section 5) are consistent with the overall deposition of fines limited by the storm wave base. Raiswell's (1988) model for the formation of limestone nodules and beds within a restricted interval of anaerobic methane oxidation by sulphate during pauses in sedimentation accords with the field and geochemical evidence for cementation close to the sediment-water interface (Sections 6 and 7). However, although initiation of limestone formation started early, final cementation appears to have occurred well below the sediment-water interface (Section 7). Such a history explains the overall rarity of limestone intraclasts despite the prevalence of sea-floor erosion and re-deposition of sediments in storm events.

The restriction of protrusive *Diplocraterion* traces to light marls and limestone beds supports the idea that deposition of light marl, rather than dark marl or black laminated-shale, was often associated with significant bottom-water turbulence. At Lyme Regis, increased proportions of angular- compared to rounded-wood particles, interpreted as indicating relatively high bottom-water turbulence, occur near the tops of many limestone beds and light marls (Waterhouse, 1999a). This observation has apparently not been tested for the West Somerset Coast (e.g. Bonis, Ruhl & Kürschner, 2010) probably because woody material is much rarer.

The explanation of orbital-climatic (Milankovitch) control of the alternations of homogenized organic-carbon poor sediments with laminated organic carbon-rich sediments in the offshore facies of the Blue Lias and correlatives remains popular (House, 1985; 1986; Weedon, 1986; 1987a; 1987b, Waterhouse 1999a; 1999b; Weedon *et al.*, 1999; Hanzo *et al.*,

2000; Bonis, Ruhl & Kürschner, 2010; Ruhl *et al.*, 2010). Intervals of increased fluvial drainage during wetter climates are likely to have led to salinity stratification, enhanced stability of the water column and increased fluxes of clay minerals from land. These factors were probably significant in promoting both higher productivity and preservation of marine organic matter in the darker laminated and more clay-rich sediments (Weedon, 1986; Fleet *et al.*, 1987).

The precursors to the limestone beds, the light marls, were probably not only associated with a relatively dry climate compared to the dark marl and black laminated shale deposition (Weedon, 1986), but also stormier conditions (Waterhouse, 1999b). The increased storminess causing erosion and/or winnowing during the deposition of light marl would have frequently led to pauses in sedimentation (non-deposition and/or hiatuses) so that limestone beds could be formed. However, increases in water depth and accommodation space, such as during the Liasicus Zone, would have reduced the probability that storms would promote the formation of limestones during deposition of light marl.

The palaeo-latitude of southern Britain was about 35° N in the Early Jurassic (Smith, Smith & Funnell, 1994). The ‘stormier’ climate associated with deposition of light marl is envisaged as resulting from rare intense tropical cyclones, rather from than frequent mid-latitude depressions. Increased storm influence during deposition of light marl (more frequent turbulent bottom water) is consistent with a drier climate overall (less rainfall and lower flux of clay). Such storm events led to episodic increases in bottom-water turbulence well below the normal storm wave-base. Individual storm events could have led to sea-floor erosion, but multiple storms would have been necessary for the prolonged phases of non-deposition required for the initiation of limestone formation by carbonate cementation of light marls near the sediment-water interface.

In the Blue Lias Formation as a whole, beds of dark grey to black laminated limestone and layers of laminated limestone nodules are much rarer than the homogeneous limestones.

1065 This distribution of lithologies was apparently because, at the associated water depths and in
1066 the wetter climatic regime associated with the laminated shales (with high clay fluxes), storms
1067 were usually too rare and/or too weak to cause pauses in sedimentation. In the field, laminated
1068 limestone nodules at Lavernock and Southam Quarry show thicker laminae in the centres of
1069 nodules that pinch laterally into much thinner laminae within the enclosing laminated shales
1070 (Weedon, 1987a). Hence, laminated limestones apparently formed prior to significant
1071 compaction just like the homogeneous grey limestone beds (Weedon, 1986; 1987a; Arzani,
1072 2004). However, exceptionally ‘clean’ laminated limestone beds such as bed H30 (Intruder) at
1073 Lyme Regis may represent deposition from low-density turbidity currents (Hesselbo &
1074 Jenkyns, 1995).

1075 The observations of gutter casts and crustacean escape structures in Southam Quarry at
1076 Long Itchington prove that storms manifestly caused sea-floor erosion and re-deposition during
1077 times of laminated-shale deposition (Radley, 2008; O’Brien, Braddy & Radley, 2009). In
1078 general, laminated limestones are most abundant within the Tilmanni and Planorbis zones.
1079 Hence during the early Hettangian, water depths were apparently shallow enough for even
1080 rare/weak storms to cause the pauses in sedimentation required to generate laminated limestone
1081 beds and laminated limestone nodule layers.

1082 According to the Raiswell (1988) model of anaerobic methane oxidation by sulphate,
1083 limestone formation started a metre so below the sediment-water interface. The range of
1084 preservation of ammonites (Section 6) indicates that for different limestone beds the
1085 cementation formed a framework-supporting fabric: a) before aragonite dissolution and before
1086 compaction (ammonites fully preserved), b) after aragonite dissolution, but before significant
1087 compaction (only external moulds of sediment-free inner whorls preserved, especially on the
1088 edges of limestone beds) and c) after aragonite dissolution and after compaction started (no
1089 ammonites shells preserved, although calcitic oyster xenomorphs may be present).

Paul, Allison & Brett (2008) argued that the ammonites within the limestones at Lyme Regis were preserved due to rapid deposition, which seems plausible, given the evidence for storm-related deposition (Section 5, Weedon, 1986; 1987a; Waterhouse, 1999b; Radley, 2008). However, not mentioned by Paul, Allison & Brett (2008), was that additionally an interval of non-deposition after the rapid burial by light marl seems also to have been necessary to initiate limestone formation. Given the homogenization of the light marl by burrowers prior to limestone formation, the non-deposition and cementation apparently did not immediately follow rapid sedimentation. However, unless non-deposition occurred before aragonite dissolution (i.e. within a few thousand years), cementation would not have been early enough to preserve the ammonites.

The Blue Lias Formation is well known for fully articulated ichthyosaur and plesiosaur fossils (e.g. Milner & Walsh, 2010). Skeletons enclosed by laminated shale were probably protected from scavengers by bottom-water anoxia. However, given that many fully articulated skeletons that somehow avoided disarticulation by scavengers are found also in the marls and homogeneous limestones, rapid burial of carcasses during storms could also explain their preservation.

Normally, the aragonite of ammonites buried within accumulating light marl would have dissolved and left no trace unless pyritized (very rare in the Blue Lias Formation) or preserved as external moulds in oyster xenomorphs. Limestone beds that do not preserve ammonites were apparently formed during phases of non-deposition that did not quickly follow an episode of rapid light marl deposition.

9. Conclusions

In the offshore hemipelagic facies of the Blue Lias Formation, the spacing of individual limestones within zones and the varying limestone proportions of limestones from zone-to-

1115 zone and from site-to-site, can all be related to the bottom-water turbulence. The model adopted
1116 represents a synthesis of several key papers accompanied by the new biostratigraphic and
1117 sedimentological data presented here.

1118 There is abundant evidence for winnowing, non-deposition and sea-floor erosion within
1119 the formation, as indicated by the Shaw plot (Fig. 8) and the field evidence (Section 5).
1120 Initiation of limestone formation is explained by periods of non-deposition causing
1121 cementation of the zone of anaerobic methane oxidation by sulphate close to the sediment-
1122 water interface (Raiswell, 1988, Bottrell & Raiswell, 1989). This mechanism explains the
1123 decoupling of limestone bed thickness from lateral and stratigraphic variations in net
1124 accumulation rate (Hallam, 1964; Raiswell, 1988). The periods of increased sea-floor erosion
1125 or non-deposition that initiated the formation of individual limestone beds and horizons of
1126 limestone nodules can be attributed to storms (Weedon, 1986; 1987a; Waterhouse 1999b;
1127 Radley, 2008).

1128 Deposition of light marl was associated with a drier climate characterized by much
1129 stronger and/or more frequent, though possibly still rare, major storms whereas more tranquil
1130 conditions were typical for the deposition of dark marl and laminated shale (Waterhouse,
1131 1999b). Thus, limestone beds and nodule horizons were much more likely to form within light
1132 marl beds. However, occasionally during deposition of laminated shale, storms did lead to non-
1133 deposition and formation of laminated limestone beds and laminated limestone nodule horizons
1134 (Radley, 2008). Relatively shallow water depths during the Tilmanni and Planorbis Zones, and
1135 thus greater influence of storms on bottom-water turbulence during laminated-shale deposition,
1136 explains the concentration of laminated limestones within this interval. Conversely, relatively
1137 greater water depths, for example during the Liasicus Zone, meant that storm-related bottom-
1138 water turbulence was reduced and thus less likely to cause non-deposition and the formation
1139 of homogeneous limestone beds and nodule horizons within the light marls.

A strong storm influence on deposition in the offshore Blue Lias Formation is not surprising, given the well-known storm-related deposition of the ‘marginal facies’ (Johnson & McKerrow, 1995; Simms, 2004; Sheppard, 2006) and of the near-shore facies of the Blue Lias in South Wales (Section 4.a, Sheppard, Houghton & Swan, 2006). Storm influences have also been documented from Hettangian and Lower Sinemurian strata, of Blue Lias character, in the Paris Basin (Hanzo *et al.*, 2000).

In the Lower Pliensbachian of Yorkshire, Van Buchem, McCave & Weedon (1994) showed that the orbitally forced cycles of grain size were related to varying storm intensity. The coeval Belemnite Marls of Dorset (or Stonebarrow Marl Member of the Charmouth Mudstone Formation of Page, 2010b) exhibit orbitally controlled alternations of light and dark marls (Weedon & Jenkyns, 1999). Similar orbitally controlled cyclicity has been documented in the Pliensbachian strata of the Mochras borehole at Llanbedr, Wales (Ruhl *et al.*, 2016). Sellwood (1970) believed the rhythms of the Belemnite Marls indicated cycles in bottom-water turbulence but invoked short-term changes in sea level as a primary forcing mechanism. However, similarly to the Blue Lias, the changes in sediment composition in the Belemnite Marls can be linked to orbital climatic cycles (Weedon & Jenkyns, 1999), with indications of varying bottom-water turbulence (variations in characteristics of and types of burrow traces, Sellwood, 1970) linked to cycles in storminess. Deeper water conditions compared to the depositional environment of the Blue Lias may explain the lack of non-sequences and consequent lack of limestones except at the base and top of the Belemnite Marls (Hesselbo & Jenkyns, 1995; Weedon & Jenkyns, 1999).

Normally, ammonites in the Blue Lias are not preserved in the light marls unless represented by crushed specimens, commonly with associated organic films; or their former presence may be indicated by oyster xenomorphs. Ammonites preserved within limestone beds required rapid burial under light marl (Paul, Allison & Brett, 2008), most likely caused by

storm-related re-deposition (Section 5, Radley, 2008). However, preservation would only normally occur when rapid burial was followed fairly quickly by a period of non-deposition that initiated limestone formation. Non-deposition was probably also linked to storm activity and occurred after homogenization of the sediment by burrowers, but before aragonite dissolution. If non-deposition or hiatus formation did not quickly follow rapid deposition, formation of limestone beds devoid of ammonites took place, as a combination of aragonite dissolution, bioturbation and early compaction removed all trace of the shells. Ammonites within the laminated shales on the west coast of Somerset are preserved as: a) shell-less impressions, b) impressions with residual brown films, c) as white aragonite, d) brown calcite near faults or e) as pristine (i.e. iridescent) aragonite that is likely to have owed its survival to the protective effects of organic matter.

To account for the stable-isotope signatures of the limestone beds, it is suggested that initially early carbonate cementation produced framework-supporting fabrics that were usually strong enough to resist compaction of large macrofossils such as ammonites (if not previously dissolved). However, the final oxygen- and carbon-isotope signatures within the limestones represent a mixture of the early plus much later cement that filled the micrograde framework pores. At Lyme Regis, due to the low net accumulation rate and the relatively shallow depths at which cementation finished, the oxygen isotopes in the limestone beds are heavier than those of marls and shales. By contrast, on the West Somerset coast, the much higher net accumulation rates ensured that final cementation of the limestone beds occurred much deeper in the sediment pile so that the overall oxygen-isotope values are generally lighter than the associated marls and shales and also lighter than the limestones at Lyme Regis.

Acknowledgements

The Royal Society provided a grant to G.P.W. in the early 1990s for the purchase of a

Bartington MS meter. Stuart Robinson kindly supplied 38 ‘SAB’ samples (Hesselbo, Robinson & Surlyk, 2004) from the Tilmanni and Planorbis biozones from St Audries Bay for MS analysis to supplement the collection obtained by Weedon (1987a). Simon Robinson provided helpful advice on interpretations of MS. Our thanks go to Chris Paul for his constructive review.

K.N.P. thanks the Orchard-Wyndham estate, including the late Dr Catherine Wyndham, and the East Quantoxhead Estate, including the late Colonel Sir Walter Luttrell, for ongoing permission to sample on the Estate’s foreshore areas in West Somerset. Bob Cornes and Tom Sunderland of Natural England provided help and advice regarding permission for sampling in West Somerset and within the Axmouth-Lyme Regis National Nature Reserve, west of Lyme Regis, respectively. Raymond Roberts of Natural Resources Wales (formerly Countryside Council for Wales) advised on the St. Mary’s Well Bay section at Lavernock. K.N.P. would also like to thank María-José Bello-Villalba and Joseph Bello-Page for assistance in the field.

References

- Ager, D. 1986. A reinterpretation of the basal ‘Littoral Lias’ of the Vale of Glamorgan. *Proceedings of the Geologists’ Association* **97**, 29–35.
- Allison, P. A., Hesselbo, S. P., & Brett, C. A. 2008. Methane seeps on an Early Jurassic seafloor. *Palaeogeography, Palaeoclimatology, Palaeoecology* **270**, 230–238.
- Ambrose, K. 2001. The lithostratigraphy of the Blue Lias Formation (Late Rhaetian-Early Sinemurian) in the southern part of the English Midlands. *Proceedings of the Geologists’ Association* **112**, 97–110.
- Arzani, N. 2004. Diagenetic evolution of mudstones: black shales to laminated limestones, an example from the Lower Jurassic of SW Britain. *Journal of Sciences Islamic Republic of Iran* **15**, 257–267.

- 1215 Arzani, N. 2006. Primary versus diagenetic bedding in the limestone-marl/shale alternations of
1216 the epeiric seas, an example from the Lower Lias (Early Jurassic) of SW Britain.
1217 *Carbonates and Evaporites* **21**, 94–109.
- 1218 Barras, C. G. & Twitchett, R. J., 2007. Response of the marine infauna to Triassic-Jurassic
1219 environmental change: ichnological data from southern England. *Palaeogeography,*
1220 *Palaeoclimatology, Palaeoecology* **244**, 223–241.
- 1221 Bixler, W. G., Elmore, R. D., & Engel, M. H. 1998. The origin of magnetization and
1222 geochemical alteration in a fault zone, Kilve, England. *Geological Journal* **33**, 89–105.
- 1223 Bleil, S. & Petersen, N. 1987. Magnetic properties of natural minerals. In *Physical properties*
1224 *of rocks Volume 1 subvolume B* (ed G. Angenheister) pp. 308–365. Springer.
- 1225 Bloos, G. & Page, K. N. 2000a. The basal Jurassic ammonite succession in the North-West
1226 European Province - Review and new results. In: *Advances in Jurassic Research 2000.*
1227 *Proceedings of the Fifth International Symposium on the Jurassic System* (eds R. L.
1228 Hall & P. Smith), pp. 27–40. GeoResearch Forum **6**, Trans Tech Publications.
- 1229 Bloos, G. & Page, K. N. 2000b. The proposed GSSP for the base of the Sinemurian Stage near
1230 East Quantoxhead/West Somerset (SW England) - The ammonite sequence. In:
1231 *Advances in Jurassic Research 2000. Proceedings of the Fifth International Symposium*
1232 *on the Jurassic System* (eds R. L. Hall & P. L. Smith) pp. 13–26. GeoResearch Forum
1233 **6**, Trans Tech Publications.
- 1234 Bloos, G. & Page, K. N. 2002. The Global Stratotype Section and Point for base of the
1235 Sinemurian Stage (lower Jurassic). *Episodes* **25**, 22–28.
- 1236 Bonis, N. A., Ruhl, M. & Kürschner, W.M. 2010. Milankovitch-scale palynological turnover
1237 across the Triassic–Jurassic transition at St. Audrie’s Bay, SW UK. *Journal of the*
1238 *Geological Society, London* **167**, 877–888.

- 1239 Bottrell, S. & Raiswell, R. 1989. Primary versus diagenetic origin of Blue Lias rhythms
1240 (Dorset, UK): evidence from sulphur geochemistry. *Terra Nova* **1**, 457–456.
- 1241 Bown, P.R. 1987. Taxonomy, evolution, and biostratigraphy of late Triassic-early Jurassic
1242 calcareous nannofossils. *Special Papers in Palaeontology* **38**, Palaeontological
1243 Association, 118 pp.
- 1244 Bown, P. R. & Cooper, M. K. E. 1998. Jurassic. In *Calcareous Nannofossil Biostratigraphy*,
1245 (ed. P. R. Bown), pp. 34–85. British Micropalaeontological Society Publications Series,
1246 Chapman and Hall.
- 1247 Bromley, R. G. & Ekdale, A. A. 1984. *Chondrites*: a trace fossil indicator of anoxia in
1248 sediments. *Science* **224**, 872–874.
- 1249 Bukri, P. M., Dent Glasser, L. S. & Smith, D. N. 1982. Surface coatings on ancient coccoliths.
1250 *Nature* **297**, 145–147.
- 1251 Callomon, J. 1985. Biostratigraphy, chronostratigraphy and all that - again! In: *International*
1252 *Symposium on Jurassic Stratigraphy, Erlangen 1984* (eds O. Michelsen, O. & A. A.
1253 Zeiss), pp. 611-624. Geological Survey of Denmark, Copenhagen 3.
- 1254 Callomon, J. 1995. Time from fossils: S.S. Buckman and Jurassic high-resolution
1255 geochronology. *Milestones in Geology* (ed. M. L. Le Bas), pp. 127–150. Geological
1256 Society Memoir 16.
- 1257 Campos, J. G. & Hallam, A. 1979. Diagenesis of English Lower Jurassic limestones as inferred
1258 from oxygen and carbon isotope analysis. *Earth and Planetary Science Letters* **45**, 23–
1259 31.
- 1260 Clémence, M-E., Bartolini, A., Gardin, S., Paris, G., Beaumont, V. & Page, K. N. 2010. Early
1261 Hettangian benthic-planktonic coupling at Doniford (SW England).
1262 Palaeoenvironmental implications from the aftermath of the end-Triassic crisis.
1263 *Palaeogeography, Palaeoclimatology, Paleoecology* **295**, 102–115.

- 1264 Clements, R. G. & members of the field studies in local geology class 1975. *The geology of the*
 1265 *Long Itchington Quarry (Rugby Portland Cement Co. Ltd., Southam Works)*. Percival
 1266 Guildhall, Rugby and Dept. of Geology, University of Leicester, 30 pp.
- 1267 Clements, R. G. & members of the field studies in local geology class, Percival Guildhouse,
 1268 Rugby 1977. *The Geology of the Parkfield Quarry, Rugby (Rugby Portland Cement Co.*
 1269 *Ltd., Rugby Works, Rugby, Warwickshire)*. Dept. of Geology, University of Leicester,
 1270 54 pp.
- 1271 Collinson, D. W., 1983. *Methods in rock magnetism and palaeomagnetism: Techniques and*
 1272 *Instrumentation*. Chapman and Hall, London, 503 pp.
- 1273 Cornford, C. 2003. Triassic palaeopressure and Liassic mud volcanoes near Kilve, west
 1274 Somerset. *Geoscience in South-West England* **10**, 430–434.
- 1275 Curtis, C. D., Cope, J. C. W., Plant, D. & Macquaker, J. H. S. 2000. ‘Instantaneous’
 1276 sedimentation, early microbial sediment strengthening and a lengthy record of chemical
 1277 diagenesis preserved in Lower Jurassic ammonitiferous concretions from Dorset.
 1278 *Journal of the Geological Society, London* **157**, 165–177.
- 1279 Day, E. C. H. 1865. On the Lower Lias of Lyme Regis. *Geological Magazine* **2**, 518–519.
- 1280 Deconinck, J-F., Hesselbo, S. P., Debuisser, N., Averbuch, O., Baudin, F. and Bessa, J. 2003.
 1281 Environmental controls on clay mineralogy of an Early Jurassic mudrock (Blue Lias
 1282 Formation, southern England). *International Journal of Earth Sciences* **92**, 255–266.
- 1283 Donovan, D. T. 1956. The zonal stratigraphy of the Blue Lias around Keynsham, Somerset.
 1284 *Proceedings of the Geologists’ Association* **66**, 182–212.
- 1285 Donovan, D. T., Horton, A., & Ivimey-Cook, H. C. 1979. The transgression of the Lower Lias
 1286 over the northern flank of the London Platform. *Journal of the Geological Society,*
 1287 *London* **136**, 165–173.

- 1288 Donovan, D. T. & Kellaway, G. A. 1984. *Geology of the Bristol District; the Lower Jurassic*
1289 *rocks*. Memoir of the British Geological Survey Special Sheet. HMSO, 69 pp.
- 1290 Fleet, A. J., Clayton, C. J., Jenkyns, H. C. & Parkinson, D. N. (1987). Liassic source rock
1291 deposition in western Europe (eds J. Brooks & K. Glennie) pp 59–70. In *Petroleum*
1292 *Geology of north-west Europe, 1*, Graham & Trotman, London.
- 1293 Fletcher, C. J. N. 1988. Tidal erosion, solution cavities and exhalative mineralization associated
1294 with the Jurassic unconformity at Ogmore, South Glamorgan. *Proceedings of the*
1295 *Geologists' Association* **99**, 1–14.
- 1296 Gallois, R. W. & Paul, C. R. C. 2009. Lateral variations in the topmost part of the Blue Lias
1297 and basal Charmouth Mudstone Formations (Lower Jurassic) on the Devon and Dorset
1298 Coast. *Geoscience in South-West England* **12**, 125–133.
- 1299 Gluyas, J. G. 1984. Early carbonate diagenesis within Phanerozoic shales and sandstones of
1300 the NW European shelf. *Clay Minerals* **19**, 309–321.
- 1301 Hall, A. & Kennedy, W. J. 1967. Aragonite in fossils. *Philosophical Transactions of the Royal*
1302 *Society, London* **B168**, 377–412.
- 1303 Hallam, A. 1960. A sedimentary and faunal study of the Blue Lias of Dorset and Glamorgan.
1304 *Philosophical Transactions of the Royal Society, London* **B243**, 1–44.
- 1305 Hallam, A. 1964. Origin of limestone-shale rhythms in the Blue Lias of England: a composite
1306 theory. *Journal of Geology* **72**, 157–169.
- 1307 Hallam, A. 1981. A revised sea-level curve for the early Jurassic. *Journal of the Geological*
1308 *Society, London* **138**, 735–743.
- 1309 Hallam, A., 1986. Origin of minor limestone-shale cycles: climatically induced or diagenetic?
1310 *Geology* **14**, 609–612.
- 1311 Hallam, A., 1987. Reply to Weedon, 1987 Comment on: 'Origin of limestone-shale cycles:
1312 climatically induced or diagenetic?' *Geology* **15**, 93–94.

- 1313 Hallam, A. 1997. Estimates of the amount and rate of sea-level change across the Rhaetian–
 1314 Hettangian and Pliensbachian–Toarcian boundaries (latest Triassic to early Jurassic).
 1315 *Journal of the Geological Society, London* **154**, 773–779.
- 1316 Hallam, A. & Bradshaw, M. J. 1979. Bituminous shales and oolitic ironstones as indicators of
 1317 transgression and regression. *Journal of the Geological Society, London* **136**, 157–164.
- 1318 Hamilton, G. B. 1982. Triassic and Jurassic calcareous nannofossils. In *A stratigraphical index*
 1319 *of calcareous nannofossils* (ed. A. R. Lord), pp. 17–39. British Micropalaeontological
 1320 Society Series, Ellis Horwood, Chichester.
- 1321 Hanzo, M., Lathuilière, B., Alméras, Y., Dagallier, G., Guérin-Franiette, S., Guillocheau, F.,
 1322 Huault, V., Nori, L. & Rauscher, R. 2000. Paléoenvironnements dans le Calcaire à
 1323 gryphées du Lias de Lorraine, de la carrière de Xeuilley au Bassin parisien. *Eclogae*
 1324 *Geologicae Helveticae* 93, 183–206.
- 1325 Hays, J. D., Imbrie, J. & Shackleton, N. J. 1976. Variations in the Earth’s orbit: pacemaker of
 1326 the ice ages. *Science* **194**, 1121–1132.
- 1327 Hesselbo, S. P. 2008. Sequence stratigraphy and inferred relative sea-level change from the
 1328 onshore British Jurassic. *Proceedings of the Geologists’ Association* **119**, 19–34.
- 1329 Hesselbo, S. P. & Jenkyns, H. C. 1995. A comparison of the Hettangian to Bajocian successions
 1330 of Dorset and Yorkshire. In *Field Geology of the British Jurassic* (ed P. D. Taylor) pp.
 1331 105–150. Geological Society, London.
- 1332 Hesselbo, S. P. & Jenkyns, H. C. 1998. British Lower Jurassic sequence stratigraphy. In
 1333 *Mesozoic and Cenozoic Sequence Stratigraphy of European Basins* (eds J. Hardenbol,
 1334 J. Thierry, M. B. Farley, T. Jacquin, P-C. de Graciansky & P. Vail) pp. 562–581. SEPM
 1335 Special Publication 60.

- 1336 Hesselbo, S. P., Robinson, S. A. & Surlyk, F. 2004. Sea-level change and facies development
1337 across potential Triassic-Jurassic boundary horizons, SW Britain. *Journal of the*
1338 *Geological Society, London* **161**, 365–379.
- 1339 Hillebrandt, A. V., Krystyn, L., & Kuerschner, W. M. 2007. A candidate GSSP for the base of
1340 the Jurassic in the Northern Calcareous Alps (Kuhjoch section, Karwendel Mountains,
1341 Tyrol, Austria). *International Sub-commission on Jurassic Stratigraphy Newsletter* **34**,
1342 2–20.
- 1343 Hillebrandt, A. V. & Krystyn, L. 2009. On the oldest Jurassic ammonites of Europe (Northern
1344 Calcareous Alps, Austria) and their global significance. *Neues Jahrbuch für Geologie*
1345 *und Paläontologie, Abhandlungen* **253**, 163–195.
- 1346 Hodges, P. 1986. The Lower Lias (Lower Jurassic) of the Bridgend area, South Wales.
1347 *Proceedings of the Geologists' Association* **97**, 237–242.
- 1348 Hodges, P. 1994. The base of the Jurassic System: new data on the first appearance of
1349 *Psiloceras planorbis* in southwest Britain. *Geological Magazine* **131**, 841–844.
- 1350 Hounslow, M. W. 1985. Magnetic fabric arising from paramagnetic phyllosilicate minerals in
1351 mudrocks. *Journal of the Geological Society, London* **142**, 995–1106.
- 1352 Hounslow, M. W., Posen, P. E. & Warrington, G. 2004. Magnetostratigraphy and
1353 biostratigraphy of the Upper Triassic and lowermost Jurassic succession, St. Audrie's
1354 Bay, UK. *Palaeogeography, Palaeoclimatology, Palaeoecology* **213**, 331–358.
- 1355 House M. R. 1985. A new approach to an absolute time scale from measurements of orbital
1356 cycles and sedimentary microrhythms. *Nature* **313**, 17–22.
- 1357 House, M. R. 1986. Are Jurassic microrhythms due to orbital forcing? *Proceedings of the*
1358 *Ussher Society* **6**, 299–311.
- 1359 Hudson, J. D. & Martill, D. M. 1991. The Lower Oxford Clay: production and preservation of
1360 organic matter in the Callovian (Jurassic) of central England. In *Modern and ancient*

- 1361 *continental Shelf Anoxia* (eds R. V. Tyson & T. H. Pearson), pp. 363–379. Geological
 1362 Society, London Special Publication 58.
- 1363 Jenkyns, H. C., Jones, C. E., Gröcke, D. R., Hesselbo, S. P. & Parkinson, D. N. 2002.
 1364 Chemostratigraphy of the Jurassic System: applications, limitations and implications
 1365 for palaeoceanography. *Journal of the Geological Society, London* **159**, 351–378.
- 1366 Jenkyns, H. C. & Senior, J. R. 1991. Geological evidence for intra-Jurassic faulting in the
 1367 Wessex Basin and its margins. *Journal of the Geological Society, London* **148**, 245–
 1368 260.
- 1369 Johnson, M. E. & McKerrow, W. S. 1995. The Sutton Stone: an early Jurassic rocky shoreline
 1370 deposit in South Wales. *Palaeontology* **38**, 529–541.
- 1371 Kent, P. E. 1936. The formation of hydraulic limestones of the Lower Lias. *Geological*
 1372 *Magazine* **73**, 476–478.
- 1373 Kent, P. E. 1937. The Lower Lias of South Nottinghamshire. *Proceedings of the Geologist's*
 1374 *Association* **48**, 163–174.
- 1375 Korte, C., Hesselbo, S. P., Jenkyns, H. C., Rickaby, R. E., & Spotl, C. 2009.
 1376 Palaeoenvironmental significance of carbon- and oxygen-isotope stratigraphy of
 1377 marine Triassic–Jurassic boundary sections in SW Britain. *Journal of the Geological*
 1378 *Society, London* **166**, 432–445.
- 1379 Lang, W. D. 1924. The Blue Lias of the Devon and Dorset Coasts. *Proceedings of the*
 1380 *Geologists' Association* **35**, 169–184.
- 1381 Lees, J. A., Bown, P. R. & Young, J. R. 2006. Photic zone palaeoenvironments of the
 1382 Kimmeridge Clay Formation (Upper Jurassic, UK) suggested by calcareous
 1383 nannoplankton palaeoecology. *Palaeogeography, Palaeoclimatology, Palaeoecology*
 1384 **235**, 110–134.

- 1385 Loughman, D. L. 1982. *A facies analysis of the Triassic/Jurassic boundary beds of the world:*
 1386 *with special reference to North-West Europe and the Americas*. Ph.D. Thesis University
 1387 of Birmingham (unpublished).
- 1388 Marshall, J. D. 1982. Isotopic composition of displacive fibrous calcite veins: reversals in pore-
 1389 water composition trends during burial diagenesis. *Journal of Sedimentary Petrology*
 1390 **52**, 615–630.
- 1391 Milner, A.C. and Walsh, S., 2010. Reptiles. In *Fossils from the Lower Lias of the Dorset Coast*,
 1392 (eds A. R. Lord & P. G. Davis), pp. 372–394. Palaeontological Association, London.
- 1393 Moghadam, H. V. & Paul, C. R. C. 2000. Trace fossils of the Jurassic, Blue Lias, Lyme Regis,
 1394 Southern England. *Ichnos* **7**, 283–306.
- 1395 O'Brien, L. J., Braddy, S. J., & Radley, J. D. 2009. A new arthropod resting trace and associated
 1396 suite of trace fossils from the Lower Jurassic of Warwickshire, England. *Palaeontology*
 1397 **52**, 1099–1112.
- 1398 Odin, G. S. & Matter, A. 1981. De glauconiarum origine. *Sedimentology* **28**, 611–641.
- 1399 Ogg, J. G. & Hinnov, L. A. 2012. Jurassic. In *The Geologic Time Scale 2012* (eds F. M.
 1400 Gradstein, J. G. Ogg, M. Schmitz & G. Ogg), pp. 731–791. Elsevier.
- 1401 Old, R. A., Sumbler, M. G., & Ambrose, K. 1987. *Geology of the Country around Warwick*.
 1402 Memoir of the British Geological Survey Sheet 84 (England and Wales).
- 1403 Page, K. N. 1992. The sequence of ammonite-correlated horizons in the British Sinemurian
 1404 (Lower Jurassic). *Newsletters on Stratigraphy* **27**, 129–156.
- 1405 Page, K. N. 1995. East Quantoxhead, Somerset, England: a potential Global Stratigraphic
 1406 Section and Point (GSSP) for the base of the Sinemurian Stage (Lower Jurassic).
 1407 *Proceedings of the Ussher Society* **9**, 446–450.

- 1408 Page, K. N. 2002. A review of the ammonite faunas and standard zonation of the Hettangian
1409 and Lower Sinemurian succession (Lower Jurassic) of the East Devon Coast (South
1410 west England). *Geoscience in South-West England* **10**, 293–303.
- 1411 Page, K. N. 2003. The Lower Jurassic of Europe - its subdivision and correlation. In *The*
1412 *Jurassic of Denmark and adjacent areas* (eds J. Ineson, & F. Surlyk), pp. 23–59.
1413 Geological Survey of Denmark and Greenland, Bulletin 1.
- 1414 Page, K. N. 2005. The Hettangian ammonite faunas of the west Somerset Coast (South West
1415 England) and their significance for the correlation of the candidate GSSP (Global
1416 Stratotype and Point) for the base of the Jurassic System at St. Audries Bay. *Còlloque*
1417 *l'Hettangien a Hettange de la science au patrimoine*, Université Henri Poincaré, pp.
1418 15–19.
- 1419 Page, K. N. 2010a. High resolution ammonite stratigraphy of the Charmouth Mudstone
1420 Formation (Lower Jurassic: Sinemurian-Lower Pliensbachian) in south-west England,
1421 UK. *Volumina Jurassica* **7**, 19–29.
- 1422 Page, K. N. 2010b. Stratigraphical framework. In *Fossils from the Lower Lias of the Dorset*
1423 *Coast* (eds A. R. Lord & P. G. Davis), pp. 33–53. Palaeontological Association Field
1424 Guides to Fossils 13.
- 1425 Page, K. N. 2010c. Ammonites. In *Fossils from the Lower Lias of the Dorset Coast* (eds A. R.
1426 Lord & P. G. Davis), pp. 169–261. Palaeontological Association Field Guides to Fossils
1427 13.
- 1428 Page, K. N. in press. From Oppel to Callomon (and beyond!): building a high resolution
1429 ammonite-based biochronology for the Jurassic System. *Lethaia*.
- 1430 Page, K. N. & Bloos, G. 1995. The base of the Jurassic System in West Somerset, South-West
1431 England – new observations on the succession of ammonite faunas of the lowermost
1432 Hettangian Stage. *Geoscience in South-West England* **9**, 231–235.

- 1433 Page, K. N., Bloos, G., Bessa, J. L., Fitzpatrick, M., Hart, M. B., Hesselbo, S., Hylyon, M.,
 1434 Morris, A. & Randall, D. E. 2000. East Quantoxhead, Somerset: A candidate Global
 1435 Stratotype Section and Point for the base of the Sinemurian Stage (Lower Jurassic). In
 1436 *Advances in Jurassic Research 2000* (eds R. L. Hall & P. L. Smith), pp. 163–72. Proc.
 1437 Fifth International Symposium on the Jurassic System, GeoResearch Forum 6, Trans
 1438 Tech Publications.
- 1439 Palmer, C. P. 1972. The Lower Lias (Lower Jurassic) between Watchet and Lillstock in North
 1440 Somerset (United Kingdom). *Newsletters on Stratigraphy* **2**, 1–30.
- 1441 Paul, C. R. C., Allison, P.A. & Brett, C. E. 2008. The occurrence and preservation of ammonites
 1442 in the Blue Lias Formation (lower Jurassic) of Devon and Dorset, England and their
 1443 palaeoecological, sedimentological and diagenetic significance. *Palaeogeography,*
 1444 *Palaeoclimatology, Palaeoecology* **270**, 258–272.
- 1445 Paris, G., Beaumont, V., Bartolini, A., Clémence, M-E. & Gardin, S. 2010. Nitrogen isotope
 1446 record of a perturbed paleoecosystem in the aftermath of the end-Triassic crisis,
 1447 Doniford section, SW England. *Geochemistry, Geophysics, Geosystems* **11**, Q08021,
 1448 doi:10.1029/2010GC003161.
- 1449 Pearson, S. J., Marshall, J. E. A. & Kemp, A. E. S. 2004. The White Stone Band of the
 1450 Kimmeridge Clay Formation, an integrated high-resolution approach to understanding
 1451 environmental change. *Journal of the Geological Society, London* **161**, 675–683.
- 1452 Price, G. D., Vowles-Sheridan, N. & Anderson, M. W. 2008. Lower Jurassic mud volcanoes
 1453 and methane, Kilve, Somerset, UK. *Proceedings of the Geologists' Association* **119**,
 1454 193–201.
- 1455 Radley, J. D. 2008. Seafloor erosion and sea-level: Early Jurassic Blue Lias Formation of
 1456 central England. *Palaeogeography, Palaeoclimatology, Palaeoecology* **270**, 287–294.

- 1457 Raiswell, R. 1988. Chemical model for the origin of minor limestone-shale cycles by anaerobic
1458 methane oxidation. *Geology* **16**, 641–644.
- 1459 Raiswell, R. & Fisher, Q. J. 2000. Mudrock-hosted carbonate concretions: a review of growth
1460 mechanisms and their influence on chemical and isotopic composition. *Journal of the*
1461 *Geological Society, London* **157**, 239–251.
- 1462 Richardson, L. 1905. The Rhaetic and contiguous deposits of Glamorganshire. *Quarterly*
1463 *Journal of the Geological Society, London* **61**, 385–424.
- 1464 Richardson, W. A. 1923. Petrology of the Shales-with-‘Beef’. *Quarterly Journal of the*
1465 *Geological Society, London* **79**, 88–98.
- 1466 Ruhl, M., Deenen, M. H. L., Abels, H. A., Bonis, N. R., Krijgsman, W. & Kürschner, W. M.
1467 2010. Astronomical constraints on the duration of the early Jurassic Hettangian stage
1468 and recovery rates following the end-Triassic mass extinction (St. Audrie’s Bay/East
1469 Quantoxhead, UK). *Earth and Planetary Science Letters* **296**, 262–276.
- 1470 Ruhl, M., Hesselbo, S. P., Hinnov, L., Jenkyns, H. C., Xu, W., Riding, J. B., Storm, M.,
1471 Minisini, D., Ullmann, C. V. & Leng, M. J. 2016. Astronomical constraints on the
1472 duration of the Early Jurassic Pliensbachian Stage and global climatic fluctuations.
1473 *Earth and Planetary Science Letters* **455**, 149–165.
- 1474 Seilacher, A. 2007. *Trace Fossil Analysis*. Springer, 226 pp.
- 1475 Sellwood, B. W. 1970. The relation of trace fossils to small-scale sedimentary cycles in the
1476 British Lias. In *Trace Fossils* (eds T. R. Crimes & J. C. Harper), pp. 489–504.
1477 *Geological Journal Special Issue 3*.
- 1478 Sellwood, B. W. & Jenkyns, H. C. 1975. Basins and swells and the evolution of an epeiric sea
1479 (Pliensbachian–Bajocian of Great Britain). *Journal of the Geological Society, London*
1480 **131**, 373–388.

- 1481 Sheppard, H. T. 2006. Sequence architecture of ancient rocky shorelines and their response to
 1482 sea-level change: an Early Jurassic example from South Wales, UK. *Journal of the*
 1483 *Geological Society, London* **163**, 595–606.
- 1484 Sheppard, H. T., Houghton, R. D. & Swan, A. R. H. 2006. Bedding and pseudo-bedding in the
 1485 Early Jurassic of Glamorgan: deposition and diagenesis of the Blue Lias in South
 1486 Wales. *Proceedings of the Geologists' Association* 117, 249–264.
- 1487 Shukri, N. M. 1942. Rhythmic bedding in the Lower Lias of England. *Faculty of Science Fouad*
 1488 *I University Cairo Bulletin* **24**, 66–73.
- 1489 Simms, M. J. 2004. British Lower Jurassic Stratigraphy: an Introduction. In *British Lower*
 1490 *Jurassic Stratigraphy* (eds M. J. Simms, N. Chidlaw, N. Morton, & Page, K. N.) pp. 3–
 1491 51. Geological Conservation Review Series 30.
- 1492 Simms, M. J., Little, C. T. S. & Rosen, B. R. 2002. Corals not serpulids: mineralized colonial
 1493 fossils in the Lower Jurassic marginal facies of South Wales. *Proceedings of the*
 1494 *Geologists' Association* **113**, 31–36.
- 1495 Smith, A. G. Smith, D. G. & Funnell, B. M. 1994. *Atlas of Mesozoic and Cenozoic Coastlines*.
 1496 Cambridge University Press, 99 pp.
- 1497 Smith, D. G. 1989. Stratigraphic correlation of presumed Milankovitch cycles in the Blue Lias
 1498 (Hettangian to earliest Sinemurian), England. *Terra Nova* **1**, 457–460.
- 1499 Trueman, A. E., 1920. The Liassic rocks of the Cardiff District. *Proceedings of the Geologists'*
 1500 *Association* **31**, 93–107.
- 1501 Trueman, A. E. 1922. The Liassic rocks of Glamorgan. *Proceedings of the Geologists'*
 1502 *Association* **33**, 245–284.
- 1503 Trueman, A. E. 1930. The Lower Lias (Bucklandi Zone) of Nash Point, Glamorgan.
 1504 *Proceedings of the Geologists' Association* **61**, 149–159.

- 1505 Van Buchem, F. S. P., McCave, I. N. & Weedon, G. P. 1994. Orbitally induced small-scale
1506 cyclicity in a siliciclastic epicontinental setting (Lower Lias, Yorkshire, UK). In *Orbital*
1507 *Forcing and Cyclic Sequences* (eds P.L. de Boer & D.G. Smith), pp. 345–366.
1508 International Association of Sedimentologists Special Publication 19.
- 1509 van de Schootbrugge, B., Tremolada, F., Rosenthal, Y., Bailey, T. R., Feist-Burkhardt, S.,
1510 Brinkhuis, H., Pross, J., Kent, D. V. & Falkowski, P.G. 2007. End-Triassic calcification
1511 crisis and blooms of organic-walled ‘disaster species’. *Palaeogeography,*
1512 *Palaeoclimatology, Palaeoecology* **244**, 126–141.
- 1513 Warrington, G. & Ivimey-Cook, H. C. 1995. The Late Triassic and Early Jurassic of coastal
1514 sections in west Somerset and South and Mid-Glamorgan. In *Field Geology of the*
1515 *British Jurassic* (ed P. D. Taylor) pp. 9–30. Geological Society, London.
- 1516 Waterhouse, H. K. 1999a. Orbital forcing of palynofacies in the Jurassic of France and the
1517 United Kingdom. *Geology* **27**, 511–514.
- 1518 Waterhouse, H. K. 1999b. Regular terrestrially derived palynofacies cycles in irregular marine
1519 sedimentary cycles, Lower Lias, Dorset, UK. *Journal of the Geological Society,*
1520 *London* **156**, 1113–1124.
- 1521 Waters, R. A. & Lawrence, D. J. D. 1987. *Geology of the South Wales Coalfield, Part III, the*
1522 *country around Cardiff*. Memoir 1:50,000 Geological sheet 263 (England and Wales),
1523 HMSO.
- 1524 Weedon, G. P. 1985. Palaeoclimatic information from ancient pelagic and hemipelagic cyclic
1525 sediments. *Terra Cognita* **5**, 110.
- 1526 Weedon, G. P. 1986. Hemipelagic shelf sedimentation and climatic cycles: the basal Jurassic
1527 (Blue Lias) of South Britain. *Earth and Planetary Science Letters* **76**, 321–335.

- 1528 Weedon, G. P. 1987a. *Palaeoclimatic significance of open-marine cyclic sequences*. D. Phil.
 1529 Thesis University of Oxford, 2 volumes available at:
 1530 <http://ora.ox.ac.uk/objects/uuid:aa009e6b-d429-4340-b3c5-30f5227f0148>.
- 1531 Weedon, G. P. 1987b. Comment on: ‘Origin of limestone-shale cycles: climatically induced or
 1532 diagenetic?’ by Hallam, A., 1986. *Geology* **15**, 92–94.
- 1533 Weedon, G. P., & Jenkyns, H. C. 1999. Cyclostratigraphy and the Early Jurassic time scale:
 1534 data from the Belemnite Marls, Dorset, southern England. *Geological Society of*
 1535 *America Bulletin* **111**, 1823–1840.
- 1536 Weedon, G. P., Jenkyns, H. C., Coe, A. L. & Hesselbo, S. P. 1999. Astronomical calibration
 1537 of the Jurassic time-scale from cyclostratigraphy in British mudrock formations.
 1538 *Philosophical Transactions of the Royal Society, London* **357**, 1787–1813.
- 1539 Whittaker, A. & Green, G. W. 1983. Geology of the country around Weston-super-Mare.
 1540 Memoire 1:50,000 Geological sheet 279, New series with parts of sheets 263 and 295,
 1541 HMSO.
- 1542 Whittaker, A., Holliday, D. W. & Penn, I. E. 1985. *Geophysical logs in British Stratigraphy*.
 1543 Geological Society Special Report 18, Geological Society, London, 74 pp.
- 1544 Wignall, P. B. 2001. Sedimentology of the Triassic-Jurassic boundary beds in Pinhay Bay
 1545 (Devon, SW England). *Proceedings of the Geologists’ Association* **112**, 349–360.
- 1546 Williams, R. B. G. 1984. *Introduction to Statistics for Geographers and Earth Scientists*.
 1547 MacMillan, 349 pp.
- 1548 Wilson, D., Davies, J. R., Fletcher, C. J. N. & Smith, M. 1990. *Geology of the South Wales*
 1549 *Coalfield, Part VI, the country around Bridgend*. Memoir 1:50,000 Geological sheets
 1550 261 and 262 (England and Wales), HMSO.
- 1551 Wobber, F. J. 1963. A directional structure from the Lower Jurassic of Great Britain. *Journal*
 1552 *of Sedimentary Petrology* **34**, 692–693.

- 1553 Wobber, F. J. 1965. Sedimentology of the Lias (Lower Jurassic) of South Wales. *Journal of*
1554 *Sedimentary Petrology* **35**, 683–703.
- 1555 Wobber, F. J. 1966. A study of the depositional area of the Glamorgan Lias. *Proceedings of*
1556 *the Geologists' Association* **77**, 127–137.
- 1557 Wright, V. P., Cherns, L. & Hodges, P. 2003. Missing molluscs: field testing taphonomic loss
1558 in the Mesozoic through large-scale aragonite dissolution. *Geology* **31**, 211–214.
- 1559 Young, J. R. & Bown, P. R. 1991. An ontogenetic sequence of coccoliths from the Late Jurassic
1560 Kimmeridge Clay of England. *Palaeontology* **4**, 843–850.

1561

1562 **Figure captions**

1563 Figure 1.

1564 (a) Location of the sections illustrated in Figs 3–6.

1565 (b) Formation characteristics according to Hettangian ammonite zone and location. Data from
1566 logs shown in Fig 3–6. The symbols in Fig. 1a indicate the localities. The percentage of
1567 limestone refers to the thickness of limestone logged as limestone or laminated limestone beds,
1568 or if more than 50% of a particular level, limestone or laminated limestone nodules. Ave. =
1569 average, L/D = Light/Dark, Lam. = laminated.

1570 (c) Average rock-type composition at Lyme Regis and on the West Somerset coast (St Audries
1571 Bay and Quantock's Head sections) in terms of weight per cent calcium carbonate (%CaCO₃),
1572 total organic carbon (TOC) and TOC re-expressed on a carbonate-free basis (TOCcf).
1573 Horizontal bars denote 95% confidence intervals around the mean. Data and analytical methods
1574 from Weedon (1987a): 90 samples from the Angulata and Bucklandi zones at Lyme Regis and
1575 57 samples from the Tilmanni to Bucklandi zones on the West Somerset coast. These data are
1576 supplemented using a further 38 SAB samples (Hesselbo *et al.*, 2008) from the Tilmanni and
1577 Planorbis zones at St Audries Bay. Numbers of samples analyzed by rock-type for Lyme Regis

1578 vs West Somerset coast respectively are: limestone and limestone nodules 31 vs 17; light marl
1579 23 vs 17; dark marl 20 vs 19; laminated shale 16 vs 35; laminated limestone samples from west
1580 Somerset coast alone, 7. The average %TOC_{cf} values and their uncertainties in the different
1581 rock types are consistent with the limestones formed by carbonate cementation of light marls
1582 and with the laminated limestones formed by carbonate cementation of laminated shales.

1583

1584 Figure 2.

1585 (a) Scatter plots of field measurements of vol MS, and laboratory measurements of wt MS and
1586 %CaCO₃. The dashed lines indicate the 95% confidence intervals of the regressions. *N* =
1587 number of samples, *r* = Pearson's *r* (degree of correlation). ○ = sample from Lyme Regis, + =
1588 sample from West Somerset coast. The stratigraphic locations of the samples are indicated in
1589 Figs 4 and 5.

1590 (b) Profiles of vol MS and %CaCO₃ measurements at Lyme Regis in beds 13 to 23 (bed
1591 numbers from Lang, 1924). + = measurement of vol MS, wt MS or %CaCO₃, ○ = estimated
1592 %CaCO₃ based on regression between vol MS and %CaCO₃ (Fig. 2a) – see text Section 3.b.

1593 (c) Profiles of measurements of δ¹⁸O and δ¹³C and %TOC at Lyme Regis in beds 13 to 23. + =
1594 bulk sample δ¹⁸O and δ¹³C from Weedon (1987a), □ = bulk sample δ¹⁸O and δ¹³C from Paul,
1595 Allison, & Brett (2008), ○ = *Gryphaea* sp. sample δ¹⁸O and δ¹³C from Weedon (1987a). Key
1596 to rock-types in Fig. 3.

1597

1598 Figure 3. Lithostratigraphic log, vol MS measurements and ammonite biostratigraphy at St.
1599 Mary's Well Bay, Lavernock, Wales. Homogeneous limestone nodules are indicated by black
1600 ellipses. Laminated limestone nodules are indicated by white ellipses. At the levels where vol
1601 MS was measured within limestone nodules rather than light marl, the nodules are indicated
1602 on the right of the lithostratigraphic log. Bed numbers from Waters & Lawrence (1987).

1603

1604 Figure 4. Lithostratigraphic log, vol MS and %CaCO₃ measurements and ammonite
1605 biostratigraphy to the west of Lyme Regis, Devon. The location of the base of the Angulata
1606 Zone is discussed in Section 5.a. For key to lithologies see Fig. 3. The concentration of samples
1607 at the level of bed 37 relates to the profile shown in Figs 2b and 2c. ○ = sample measured for
1608 %CaCO₃ plotted with reference to lower horizontal axis, B = level with abundant calcite beef,
1609 Lport Mbr = Langport Member, Angul. = Angulata, Buc. = Bucklandi, Extra. = Extranodosa,
1610 Ext. = Extranodosa. Bed numbers from Lang (1924). Lang (1924) and Hesselbo & Jenkyns
1611 (1995) provide the names of limestone beds on the Devon/Dorset coast.

1612

1613 Figure 5. (a) Lithostratigraphic log, vol MS and %CaCO₃ measurements and ammonite
1614 biostratigraphy at St Audries Bay, Somerset. For key to lithologies see Fig. 3. Black circles
1615 indicate samples measured for %CaCO₃ plotted with reference to lower horizontal axis. ○ =
1616 sample measured for %CaCO₃ plotted with reference to lower horizontal axis, B = level with
1617 abundant calcite beef. Bed numbering from Whittaker and Green (1983). (b) As for 5a but a
1618 composite section from St Audries Bay and Quantock's Head. The join (splice) of the St
1619 Audries Bay section (St A.) with the Quantocks Head section (Q. H.) is indicated within bed
1620 96 (Section 3.a). GSSP = Global Stratigraphic Section and Point.

1621

1622 Figure 6. Lithostratigraphic log, vol MS measurements and ammonite biostratigraphy at
1623 Southam Quarry, near Southam and Long Itchington, Warwickshire. For key to lithologies see
1624 Fig. 3. L. M. = Langport Member. Bed numbers from Clements *et al.* (1975).

1625

1626 Figure 7.

1627 (a) Photomicrograph of dark marl thin-section (plane polarized light, width of view 3 mm).
1628 Sample BL120 and Fig. 2.2M of Weedon, (1987a) bed 32a, Rotiforme Subzone, Bucklandi
1629 Zone, Lyme Regis.

1630 (b) SEM photograph showing coccoliths *in situ* partly engulfed by euhedral microspar
1631 overgrowth (width of view 14 μ m). Dark marl sample BL120 and Fig. 2.4G of Weedon
1632 (1987a), bed 32a, Bucklandi Zone, Lyme Regis.

1633 (c) SEM photograph showing an aggregate of coccoliths surrounded by a matrix of organic
1634 matter and clay (width of view 56 μ m). Dark marl sample BL120 and Fig. 2.4F of Weedon
1635 (1987a), bed 32a, Bucklandi Zone, Lyme Regis.

1636 (d) SEM photograph showing an aggregate of coccoliths next to *Schizosphaerella punctulata*
1637 (centre) both surrounded by a matrix of organic matter, clay and calcite microspar (width of
1638 view 53 μ m). Laminated shale sample BL117, and Fig. 2.4H of Weedon (1987a), topmost
1639 laminated shale of bed 32, Bucklandi Zone, Lyme Regis.

1640 (e) SEM photograph showing a corroded coccolith in limestone (width of view ~8 μ m).
1641 Limestone sample BL103 and Fig. 2.4A of Weedon (1987a), bed 29, Conybeari Subzone,
1642 Bucklandi Zone, Lyme Regis.

1643

1644 Figure 8. Shaw plot constructed using the joint locations of biohorizon bases and biohorizon
1645 tops at Lyme Regis and the West Somerset Coast (St Audries Bay and Quantock's Head
1646 sections) as listed in Table 1. The location of the base of the Angulata Zone at Lyme Regis has
1647 been estimated using the West Somerset data via the line of correlation data (long grey arrows,
1648 Section 5.a). The ratios of the thicknesses of subzones in the West Somerset composite section
1649 to subzones in the Lyme Regis section are indicated at the bottom of the plot. Vertical or
1650 horizontal arrows with bed numbers indicate the levels at which breaks in slope of the line of
1651 correlation imply relative condensation or intra-formational hiatuses. + = St Audries Bay

1652 section versus Lyme Regis section, × Quantock's Head section versus Lyme Regis section.
1653 Tilm. = Tilmanni, Planorb. = Planorbis, Pl. = Planorbis, Jo. = Johnsoni, Laq. = Laqueus, Extran.
1654 = Extranodosa, Depr. = Depressa, Conyb. = Conybeari, Roti. = Rotiforme.

1655

1656 Figure 9.

1657 (a) Photograph of polished limestone section showing bored and encrusted limestone intraclast
1658 with a hardground-like surface (diagrammatic description below) collected *in situ* from within
1659 bed 25 (Top Copper) below Devonshire Head, Lyme Regis. *i*) Photographic negative image of
1660 acetate peel showing *Liostrea* encrusting the intraclast surface and overlain by bioclastic
1661 packstone. Two *Liostrea* individuals (left and right) encrust a third individual – demonstrating
1662 at least two generations of encrustation. Short arrows indicate the *Talpina ramosa* Von
1663 Hagenow borings into the intraclast surface and within the encrusting *Liostrea*. *ii*) As for *i*) but
1664 a different section of the intraclast surface (see diagram for location).

1665 (b) Limestone intraclasts on the surface of a fallen block of limestone, Monmouth Beach, Lyme
1666 Regis. The lens cap is 5 cm in diameter.

1667 (c) Protrusive *Diplocraterion* forming dark marl burrow-fill within limestone bed 19
1668 (Specketty), Seven Rock Point, Lyme Regis.

1669 (d) Horizon of isolated dark marl burrow fills indicating the former presence of a bed of dark
1670 marl that has been removed locally by seafloor erosion, bed 19, Seven Rock Point, Lyme Regis.

1671

1672 Figure 10.

1673 (a) Compilation of oxygen- and carbon-isotope measurements for Lyme Regis from the
1674 Angulata and Bucklandi Zones. Note several data points from limestone beds (black squares)
1675 near $\delta^{18}\text{O} = -4.5$ and $\delta^{13}\text{C} = -1.5$ ‰ VPDB are obscured by the data for marl or laminated shale
1676 (unfilled triangles). Sources of measurements for whole-rock: Campos and Hallam (1979);

1677 Gluyas (1984); Weedon (1987a); Arzani (2004; 2006); Paul, Allison & Brett (2008), for
 1678 *Gryphaea* sp.: Weedon (1987a).
 1679 (b) Compilation of oxygen- and carbon-isotope measurements for West Somerset Coast (Kilve)
 1680 for the Bucklandi Zone. Sources of measurements for whole rock: Allison, Hesselbo & Brett
 1681 (2008); Price, Vowles-Sheridan & Anderson (2008). Measurements plotted exclude atypical
 1682 rock-types associated with the Bucklandi Zone mud mounds.
 1683 (c) Compilation of oxygen- and carbon-isotope measurements for West Somerset Coast (St
 1684 Audries and Doniford) for the Tilmanni and Planorbis Zones. Sources of measurements for
 1685 whole rock: Arzani (2004; 2006); Clémence *et al.* (2010 as listed in Supplementary information
 1686 of Paris *et al.*, 2010); for *Liostrea* sp.: van Schootbrugge *et al.*, (2007, their ‘unaltered’ samples
 1687 only).

1688 Table 1. Ammonite biohorizon limits on the West Somerset (St Audries Bay and Quantock's
1689 Head), Devon/Dorset (Lyne Regis) and South Wales (Lavernock) coast sections.

1690

1691

Zone	Subzone	Biohorizon	Pos.	St A.	Qu. H.	CWS	Lav.	L.R.
Bucklandi	Rotiforme	Sn7 <i>rotiforme</i>	Base	-	92.69	92.69	-	18.06
Bucklandi	Rotiforme	Sn6 cf. <i>defneri</i>	Top	-	90.66	90.66	-	17.94
Bucklandi	Rotiforme	Sn6 cf. <i>defneri</i>	Base	-	90.38	90.38	-	17.90
Bucklandi	Rotiforme	Sn5c <i>silvestri</i>	Top	-	87.76	87.76	-	17.76
Bucklandi	Rotiforme	Sn5c <i>silvestri</i>	Base	-	87.70	87.70	-	17.70
Bucklandi	Conybeari	Sn5b <i>conybeari</i>	Top	-	86.72	86.72	-	17.56
Bucklandi	Conybeari	Sn5b <i>conybeari</i>	Base	-	86.52	86.52	-	17.43
Bucklandi	Conybeari	Sn5a <i>elegans</i>	Top	-	86.48	86.48	-	-
Bucklandi	Conybeari	Sn5a <i>elegans</i>	Base	-	86.16	86.16	-	-
Bucklandi	Conybeari	Sn4 <i>rotator</i>	Top	-	82.40	82.40	-	16.78
Bucklandi	Conybeari	Sn4 <i>rotator</i>	Base	-	82.32	82.32	-	16.76
Bucklandi	Conybeari	Sn3b <i>rouvillei</i>	Top	-	81.64	81.64	-	16.64
Bucklandi	Conybeari	Sn3b <i>rouvillei</i>	Base	-	81.46	81.46	-	16.40
Bucklandi	Conybeari	Sn3a <i>rotarius</i>	Top	-	81.28	81.28	-	16.14
Bucklandi	Conybeari	Sn3a <i>rotarius</i>	Base	-	81.18	81.18	-	16.12
Bucklandi	Conybeari	Sn2b <i>conybearoides</i>	Top	-	79.98	79.98	-	-
Bucklandi	Conybeari	Sn2b <i>conybearoides</i>	Base	-	79.78	79.78	-	-
Bucklandi	Conybeari	Sn2a <i>Metophioceras</i> sp. A	Top	-	79.34	79.34	-	-
Bucklandi	Conybeari	Sn2a <i>Metophioceras</i> sp. A	Base	-	78.64	78.64	-	15.22
Bucklandi	Conybeari	Sn1 <i>quantoxense</i>	Top	-	78.14	78.14	-	15.21
Bucklandi	Conybeari	Sn1 <i>quantoxense</i>	Base	-	78.06	78.06	-	15.16
Angulata	Depressa	Hn27b <i>quadrata</i> 2	Top	-	77.50	77.50	-	14.96
Angulata	Depressa	Hn27b <i>quadrata</i> 2	Base	-	77.42	77.42	-	14.94
Angulata	Depressa	Hn27a <i>quadrata</i> 1	Top	-	73.42	73.42	-	-
Angulata	Depressa	Hn27a <i>quadrata</i> 1	Base	-	73.38	73.38	-	14.56
Angulata	Depressa	Hn26b <i>princeps</i>	Top	-	72.14	72.14	-	13.66
Angulata	Depressa	Hn26b <i>princeps</i>	Base	-	72.04	72.04	-	13.50
Angulata	Depressa	Hn26a <i>depressa</i> 1	Top	-	69.94	69.94	-	-
Angulata	Depressa	Hn26a <i>depressa</i> 1	Base	-	69.52	69.52	-	13.36
Angulata	Complanata	Hn25 <i>striatissima</i>	Top	-	67.10	67.10	-	-
Angulata	Complanata	Hn25 <i>striatissima</i>	Base	-	66.90	66.90	-	-
Angulata	Complanata	Hn24d grp. <i>vaihingensis</i>	Top	-	65.68	65.68	-	-
Angulata	Complanata	Hn24d grp. <i>vaihingensis</i>	Base	-	65.26	65.26	-	-
Angulata	Complanata	Hn24c aff. <i>complanata</i>	Top	-	63.78	63.78	-	12.58
Angulata	Complanata	Hn24c aff. <i>complanata</i>	Base	-	63.60	63.60	-	12.46
Angulata	Complanata	Hn24b <i>phoebetica</i>	Top	-	63.14	63.14	-	-
Angulata	Complanata	Hn24b <i>phoebetica</i>	Base	-	63.04	63.04	-	-
Angulata	Complanata	Hn24a <i>complanata</i>	Top	-	57.40	57.40	-	12.22
Angulata	Complanata	Hn24a <i>complanata</i>	Base	-	57.26	57.26	-	12.12
Angulata	Complanata	Hn23c cf. <i>polyeides</i>	Top	54.74	-	54.74	-	-
Angulata	Complanata	Hn23c cf. <i>polyeides</i>	Base	54.60	-	54.60	-	-
Angulata	Complanata	Hn23b <i>similis</i>	Top	52.66	52.09	52.66	-	12.10
Angulata	Complanata	Hn23b <i>similis</i>	Base	52.58	51.95	52.58	-	12.06
Angulata	Complanata	Hn23a grp. <i>stenorhyncha</i>	Top	49.48	-	49.48	-	12.00
Angulata	Complanata	Hn23a grp. <i>stenorhyncha</i>	Base	49.26	-	49.26	-	11.90
Angulata	Extranodosa	Hn22 cf. <i>germanica</i>	Top	47.42	51.24	47.42	-	-
Angulata	Extranodosa	Hn22 cf. <i>germanica</i>	Base	47.22	51.02	47.22	-	-
Angulata	Extranodosa	Hn21c <i>amblygonia</i> 3	Top	45.40	-	45.40	-	11.36
Angulata	Extranodosa	Hn21c <i>amblygonia</i> 3	Base	45.00	-	45.00	-	11.28
Angulata	Extranodosa	Hn21b cf. <i>pyncnotycha</i>	Top	43.26	-	43.26	-	-
Angulata	Extranodosa	Hn21b cf. <i>pyncnotycha</i>	Base	43.12	-	43.12	-	-
Angulata	Extranodosa	Hn21a <i>atrox</i>	Top	42.24	-	42.24	-	-

Angulata	Extranodosa	Hn21a <i>atrox</i>	Base	42.14	-	42.14	-	-
Angulata	Extranodosa	Hn20c <i>hadrotychus</i>	Top	40.86	-	40.86	-	-
Angulata	Extranodosa	Hn20c <i>hadrotychus</i>	Base	40.76	-	40.76	-	-
Angulata	Extranodosa	Hn20b <i>Schlotheimia</i> sp. 1b	Top	40.64	-	40.64	-	-
Angulata	Extranodosa	Hn20b <i>Schlotheimia</i> sp. 1b	Base	40.58	-	40.58	-	-
Angulata	Extranodosa	Hn20a <i>Schlotheimia</i> sp. 1a	Top	40.08	-	40.08	-	-
Angulata	Extranodosa	Hn20a <i>Schlotheimia</i> sp. 1a	Base	39.96	40.10	39.96	-	10.54
Liasicus	Laqueus	Hn19d aff. <i>bloomfieldense</i>	Top	39.72	-	39.72	-	-
Liasicus	Laqueus	Hn19d aff. <i>bloomfieldense</i>	Base	39.60	-	39.60	-	-
Liasicus	Laqueus	Hn19c <i>bloomfieldense</i>	Top	38.80	-	38.80	-	-
Liasicus	Laqueus	Hn19c <i>bloomfieldense</i>	Base	38.66	-	38.66	-	-
Liasicus	Laqueus	Hn19b cf. <i>subliassicus</i>	Top	37.22	-	37.22	-	-
Liasicus	Laqueus	Hn19b cf. <i>subliassicus</i>	Base	36.98	-	36.98	-	-
Liasicus	Laqueus	Hn19a cf. <i>laqueolus</i>	Top	36.80	-	36.80	-	-
Liasicus	Laqueus	Hn19a cf. <i>laqueolus</i>	Base	36.32	-	36.32	-	-
Liasicus	Laqueus	Hn18d cf. <i>polyspeirum</i>	Top	35.80	-	35.80	-	-
Liasicus	Laqueus	Hn18d cf. <i>polyspeirum</i>	Base	35.68	-	35.68	-	-
Liasicus	Laqueus	Hn18c cf. <i>costatum</i>	Top	33.96	-	33.96	-	9.66
Liasicus	Laqueus	Hn18c cf. <i>costatum</i>	Base	33.46	-	33.46	-	9.53
Liasicus	Laqueus	Hn18b cf. <i>gallbergensis</i>	Top	33.08	34.74	33.08	-	9.25
Liasicus	Laqueus	Hn18b cf. <i>gallbergensis</i>	Base	33.00	34.21	33.00	-	9.18
Liasicus	Laqueus	Hn 18a <i>laqueus</i>	Top	33.00	33.81	33.00	-	9.03
Liasicus	Laqueus	Hn 18a <i>laqueus</i>	Base	32.88	33.53	32.88	-	8.97
Liasicus	Portlocki	Hn17c cf. <i>latimontanum</i>	Top	32.48	33.10	32.48	-	8.94
Liasicus	Portlocki	Hn17c cf. <i>latimonanum</i>	Base	32.28	32.92	32.28	-	-
Liasicus	Portlocki	Hn17b aff. <i>benecke</i>	Top	29.94	-	29.94	-	-
Liasicus	Portlocki	Hn17b aff. <i>benecke</i>	Base	29.86	-	29.86	-	-
Liasicus	Portlocki	Hn17a cf. <i>gottingense</i>	Top	29.14	-	29.14	-	-
Liasicus	Portlocki	Hn17a cf. <i>gottingense</i>	Base	28.94	-	28.94	-	8.84
Liasicus	Portlocki	Hn16b grp. <i>portlocki</i>	Top	28.46	-	28.46	-	8.80
Liasicus	Portlocki	Hn16b grp. <i>portlocki</i>	Base	28.16	-	28.16	-	8.70
Liasicus	Portlocki	Hn16a cf. <i>crassicosta</i>	Top	27.20	-	27.20	-	-
Liasicus	Portlocki	Hn16a cf. <i>crassicosta</i>	Base	25.92	-	25.92	-	-
Liasicus	Portlocki	Hn15 <i>hagenowi</i>	Top	25.62	-	25.62	-	8.21
Liasicus	Portlocki	Hn15 <i>hagenowi</i>	Base	25.12	-	25.12	-	8.16
Liasicus	Portlocki	Hn14d <i>harpotychum</i>	Top	19.16	-	19.16	-	-
Liasicus	Portlocki	Hn14d <i>harpotychum</i>	Base	19.11	-	19.11	-	-
Liasicus	Portlocki	Hn14c <i>Waehneroceras</i> sp. nov.	Top	17.68	-	17.68	-	-
Liasicus	Portlocki	Hn14c <i>Waehneroceras</i> sp. nov.	Base	17.44	-	17.44	-	-
Liasicus	Portlocki	Hn14b <i>iapetus</i>	Top	15.60	-	15.60	-	6.66
Liasicus	Portlocki	Hn14b <i>iapetus</i>	Base	13.92	-	13.92	-	6.61
Liasicus	Portlocki	Hn14a aff. <i>franconium</i>	Top	13.62	-	13.62	14.67	-
Liasicus	Portlocki	Hn14a aff. <i>franconium</i>	Base	13.54	-	13.54	14.58	6.58
Planorbis	Johnstoni	Hn13c 'post'- <i>intermedium</i>	Top	13.50	-	13.50	14.46	-
Planorbis	Johnstoni	Hn13c 'post'- <i>intermedium</i>	Base	13.42	-	13.42	14.19	-
Planorbis	Johnstoni	Hn13b grp. <i>intermedium</i>	Top	13.10	-	13.10	12.93	6.42
Planorbis	Johnstoni	Hn13b grp. <i>intermedium</i>	Base	12.82	-	12.82	12.84	-
Planorbis	Johnstoni	Hn13a aff. <i>torus</i>	Top	-	-	-	12.51	-
Planorbis	Johnstoni	Hn13a aff. <i>torus</i>	Base	-	-	-	12.39	6.08
Planorbis	Johnstoni	Hn12 <i>johnstoni</i>	Top	12.48	-	12.48	12.12	5.90
Planorbis	Johnstoni	Hn12 <i>johnstoni</i>	Base	12.10	-	12.10	11.43	5.86
Planorbis	Johnstoni	Hn11d <i>Caloceras</i> sp. 5	Top	-	-	-	10.77	-
Planorbis	Johnstoni	Hn11d <i>Caloceras</i> sp. 5	Base	-	-	-	10.65	-
Planorbis	Johnstoni	Hn11c <i>Caloceras</i> sp. 4	Top	11.30	-	11.30	10.53	-
Planorbis	Johnstoni	Hn11c <i>Caloceras</i> sp. 4	Base	11.22	-	11.22	10.41	-
Planorbis	Johnstoni	Hn11b aff. <i>tortlie</i>	Top	10.94	-	10.94	10.17	5.20
Planorbis	Johnstoni	Hn11b aff. <i>tortile</i>	Base	10.78	-	10.78	10.11	5.14
Planorbis	Johnstoni	Hn11a <i>Caloceras</i> sp. 2	Top	10.02	-	10.02	-	-
Planorbis	Johnstoni	Hn11a <i>Caloceras</i> sp. 2	Base	9.98	-	9.98	-	5.06
Planorbis	Johnstoni	Hn10 aff. <i>aries</i>	Top	9.82	-	9.82	9.87	-

Planorbis	Johnstoni	Hn10 aff. <i>aries</i>	Base	9.66	-	9.66	9.72	4.88
Planorbis	Planorbis	Hn9 <i>bristoviense</i>	Top	-	-	-	9.45	4.79
Planorbis	Planorbis	Hn9 <i>bristoviense</i>	Base	-	-	-	9.39	4.76
Planorbis	Planorbis	Hn8 <i>sampsoni</i>	Top	-	-	-	8.25	-
Planorbis	Planorbis	Hn8 <i>sampsoni</i>	Base	-	-	-	8.10	-
Planorbis	Planorbis	Hn7 <i>plicatulum</i>	Top	9.08	-	9.08	7.44	3.78
Planorbis	Planorbis	Hn7 <i>plicatulum</i>	Base	8.78	-	8.78	7.35	3.72
Planorbis	Planorbis	Hn6 <i>planorbis</i> β	Top	8.58	-	8.58	6.93	3.70
Planorbis	Planorbis	Hn6 <i>planorbis</i> β	Base	8.18	-	8.18	6.72	3.66
Planorbis	Planorbis	Hn5 <i>planorbis</i> α	Top	8.18	-	8.18	6.72	3.66
Planorbis	Planorbis	Hn5 <i>planorbis</i> α	Base	5.98	-	5.98	6.33	3.42
Planorbis	Planorbis	Hn4 <i>antecedens</i>	Top	5.58	-	5.58	6.15	3.02
Planorbis	Planorbis	Hn4 <i>antecedens</i>	Base	5.42	-	5.42	6.06	2.94
Planorbis	Planorbis	Hn3 <i>imitans</i>	Top	-	-	-	5.49	-
Planorbis	Planorbis	Hn3 <i>imitans</i>	Base	5.42	-	5.42	4.71	2.82
Tilmanii		Hn2 <i>erugatum</i>	Top	5.38	-	5.38	4.65	-
Tilmanii		Hn2 <i>erugatum</i>	Base	5.26	-	5.26	4.56	-
Tilmanii		Hn1 (no ammonites)	Top	5.26	-	5.26	4.56	-
Tilmanii		Hn1 (no ammonites)	Base	?1.50	-	?1.50	?1.86	?0.60

1692

1693 Height in metres of biohorizon boundaries above base of the Blue Lias Formation. The value
1694 in ***bold italics*** for the base Angulata Zone at Lyme Regis was estimated using the correlation
1695 with the St. Audries Bay section (see Fig. 8, Section 5.a.). Pos. = Position, St A. = St Audries
1696 Bay section, Somerset, Qu. H. = Quantock's Head section, Somerset, CWS = Composite West
1697 Somerset coastal sections using common splice level at 56.90 m (below this level the heights
1698 for Quantock's Head differ from the CWS heights), Lav. = St. Mary's Well Bay section near
1699 Lavernock, South Wales, L.R. = Pinhay Bay to Devonshire Head section, Lyme Regis, Dorset.

1700

Fig. 1

1701

1702

1703

Fig. 2

1704
1705

Fig. 3

Fig. 4

Lyme Regis (Pinhay Bay – Devonshire Head), Devon

Fig. 5a

Fig. 5b

Fig. 6

Southam Quarry, Long Itchington, Warwickshire

Fig. 7

1712

1713

Fig. 8

Fig. 9

Fig. 10