2017-08-21

Mesoscale and Submesoscale Effects on Mixed Layer Depth in the Southern Ocean

Bachman, S

<http://hdl.handle.net/10026.1/9609>

10.1175/JPO-D-17-0034.1

Journal of Physical Oceanography

American Meteorological Society

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

AMERICAN METEOROLOGICAL SOCIETY

Journal of Physical Oceanography

EARLY ONLINE RELEASE

This is a preliminary PDF of the author-produced manuscript that has been peer-reviewed and accepted for publication. Since it is being posted so soon after acceptance, it has not yet been copyedited, formatted, or processed by AMS Publications. This preliminary version of the manuscript may be downloaded, distributed, and cited, but please be aware that there will be visual differences and possibly some content differences between this version and the final published version.

The DOI for this manuscript is doi: 10.1175/JPO-D-17-0034.1

The final published version of this manuscript will replace the preliminary version at the above DOI once it is available.

If you would like to cite this EOR in a separate work, please use the following full citation:

Bachman, S., J. Taylor, K. Adams, and P. Hosegood, 2017: Mesoscale and Submesoscale Effects on Mixed Layer Depth in the Southern Ocean. *J. Phys. Oceanogr.* doi:10.1175/JPO-D-17-0034.1, in press.

© 2017 American Meteorological Society

1 Mesoscale and Submesoscale Effects on Mixed Layer Depth in the Southern 2 Ocean

3 S.D. Bachman and J.R. Taylor*

4 *Department of Applied Mathematics and Theoretical Physics, University of Cambridge, UK*

5 K.A. Adams and P.J. Hosegood

6 *School of Biological and Marine Sciences, Plymouth University, UK*

7 *Corresponding author address: DAMTP, Centre for Mathematical Sciences, Wilberforce Road,
8 Cambridge CB3 0WA, United Kingdom.

9 E-mail: J.R.Taylor@damtp.cam.ac.uk

ABSTRACT

10 Submesoscale dynamics play a key role in setting the stratification of the
11 ocean surface mixed layer and mediating air-sea exchange, making them es-
12 pecially relevant to anthropogenic carbon uptake and primary productivity in
13 the Southern Ocean. In this paper a series of offline-nested numerical simu-
14 lations is used to study submesoscale flow in the Drake Passage and Scotia
15 Sea regions of the Southern Ocean. These simulations are initialized from
16 an ocean state estimate for late-April 2015, with the intent to simulate fea-
17 tures observed during the Surface Mixed Layer at Submesoscales (SMILES)
18 research cruise which occurred at that time and location. The nested models
19 are downscaled from the original state estimate resolution of $1/12^\circ$ and grid
20 spacing of about 8 km, culminating in a submesoscale-resolving model with a
21 resolution of $1/192^\circ$ and grid spacing of about 500 m. The submesoscale eddy
22 field is found to be highly spatially variable, with pronounced “hotspots” of
23 submesoscale activity. These areas of high submesoscale activity correspond
24 to a significant difference in the 30-day average mixed layer depth, $\overline{\Delta H_{ML}}$, be-
25 tween the $1/12^\circ$ and $1/192^\circ$ simulations. Regions of large vertical velocities
26 in the mixed layer correspond with high mesoscale strain rather than large
27 $\overline{\Delta H_{ML}}$. It is found that $\overline{\Delta H_{ML}}$ is well-correlated with the mesoscale density
28 gradient but weakly correlated with both the mesoscale kinetic energy and
29 strain. This has implications for the development of submesoscale eddy pa-
30 rameterizations which are sensitive to the character of the large-scale flow.

31 1. Introduction

32 Submesoscale processes play a crucial role in the evolution of the oceanic surface boundary
33 layer. Recent work has highlighted the importance of near-surface submesoscales both as a means
34 of transporting heat and tracers into the oceanic interior via strong vertical circulations (Pollard
35 and Regier 1990; Rudnick 1996; Lapeyre and Klein 2006; Mahadevan and Tandon 2006), and as a
36 mechanism for fluxing large-scale energy downscale via unbalanced instabilities (e.g. McWilliams
37 et al. 2001; Molemaker et al. 2005; Taylor and Ferrari 2009, 2010; Thomas and Taylor 2010;
38 D’Asaro et al. 2011). The vertical transport associated with submesoscale motions has also been
39 shown to significantly affect primary production by redistributing phytoplankton, grazers, and
40 nutrients throughout the water column (Spall and Richards 2000; Mahadevan and Archer 2000;
41 Flierl and McGillicuddy 2002; Gargett and Marra 2002; Lévy et al. 2001, 2012; Lévy and Martin
42 2013; Omand et al. 2015).

43 The growing appreciation for the importance of submesoscales has spurred intensive research
44 into a wide variety of processes which occur at these scales within the ocean surface boundary
45 layer. There exists a rich set of instabilities and dynamics which constitute the broad class of
46 submesoscale flows, here defined in the dynamical sense to be motions with $\mathcal{O}(1)$ Rossby and
47 Richardson numbers and horizontal scales of 0.1 - 10 km (Thomas et al. 2008). Oceanic sub-
48 mesoscale motions are often associated with the presence of lateral density gradients, or fronts.
49 These fronts arise via mesoscale frontogenesis (Lapeyre and Klein 2006) and precondition the
50 mixed layer to a variety of submesoscale instabilities such as ageostrophic baroclinic instability
51 (Boccaletti et al. 2007), symmetric instability (Taylor and Ferrari 2009), and centrifugal instability
52 (Jiao and Dewar 2015), which in turn can be enhanced or suppressed through buoyancy forcing
53 and wind stress (Thomas 2005; Taylor and Ferrari 2010).

54 Because submesoscale turbulence is highly sensitive to atmospheric forcing, frontal strength,
55 and mixed layer depth, it can be expected to vary in strength on both fast and slow timescales.
56 Mixed layer baroclinic instability and forced symmetric instability both have growth timescales on
57 the order of hours to days (Stone 1966; Taylor and Ferrari 2009) and are capable of restratifying the
58 mixed layer (e.g. Boccaletti et al. 2007). Observations (Callies et al. 2015; Buckingham et al. 2016;
59 Thompson et al. 2016) and high-resolution modelling studies (e.g. Capet et al. 2008a; Mensa et al.
60 2013; Sasaki et al. 2014; Brannigan et al. 2015) suggest strong seasonal variation in the strength
61 of submesoscale turbulence, where deep wintertime mixed layers increase the available potential
62 energy that can be released by these instabilities.

63 Submesoscales are also expected to be energised through a downscale transfer from mesoscale
64 eddies, which are highly spatially variable (e.g. Klocker and Abernathey 2014). However, it is
65 unclear how submesoscale activity might vary with the energy of the mesoscale eddy field and
66 complex bottom topography. Rosso et al. (2014, 2015) used a $1/80^\circ$ regional model of the South-
67 ern Ocean to investigate the role of submesoscales in a region of complex bottom topography near
68 the Kerguelen Plateau, and identified submesoscales using a high-pass spatial filter with a $1/5^\circ$
69 cutoff. Using this method they found a strong correlation between upper-ocean vertical veloc-
70 ities, which was used as a proxy for submesoscale activity, and mesoscale eddy kinetic energy
71 and strain. No direct influence of topography on submesoscale features was observed, though
72 it was argued that topographic control over the mesoscale eddy field might indirectly affect the
73 submesoscales.

74 In this paper we use a series of nested high-resolution models to analyze submesoscale activ-
75 ity in a different location within the Southern Ocean, as part of SMILES (Surface Mixed Layer
76 Evolution at Submesoscales; <http://www.smiles-project.org/>). The simulations coincide
77 with observations collected on the SMILES project research cruise to the Scotia Sea, just east of

78 Drake Passage, in April-May 2015 (Adams et al. 2017). This region is characterized by an en-
79 ergetic mesoscale eddy field (Frenger et al. 2015) and strong fronts associated with the Antarctic
80 Circumpolar Current (ACC). Although mode water transformation and subduction occurs here
81 (Sallée et al. 2010; Cerovečki et al. 2013), the role of submesoscale processes is unknown. Sub-
82 mesoscale motions have the potential to modulate water mass properties across the mixed layer
83 and, therefore, may affect the oceanic uptake of tracers, such as atmospheric gases and heat.

84 The goal of this analysis is to investigate how and where submesoscale eddies affect the mixed
85 layer depth by comparing the output of the nested models. To do so, we will compare output
86 from the highest-resolution member of the series of models, which at $1/192^\circ$ (less than 500 m)
87 horizontal resolution is sufficient to resolve submesoscales, against the coarsest member, a $1/12^\circ$
88 mesoscale-permitting model. In this comparison, we intend to focus special attention on how
89 mixed layer submesoscales should be identified in high-resolution models like these, and to assess
90 how they are spatially correlated with larger, mesoscale features. The numerical model configu-
91 ration is described in Section 2. Analysis of the meso- and submesoscale influence on the mixed
92 layer depth and vertical transport is presented in Section 3. Concluding remarks appear in Section
93 4.

94 **2. Model description**

95 In this study the MITgcm (Marshall et al. 1997a) is used to conduct a series of offline-nested
96 simulations of the Drake Passage and Scotia Sea regions of the Southern Ocean. Each simula-
97 tion is run on a curvilinear, latitude-longitude grid, and uses open boundary conditions whose
98 configuration is described below.

99 The initial state and boundary conditions for the lowest resolution ($1/12^\circ$) MITgcm simulation
100 are provided by the Copernicus Marine Environment Monitoring Service Global Ocean $1/12^\circ$

101 Physics Analysis (hereafter CMEMS), which is produced by Mercator Ocean (<http://marine.copernicus.eu>). The domain of the $1/12^\circ$ simulation extends from 65° S to 45° S, and from
102 110° W to 40° W (Figure 1). The flow is initialized from the CMEMS ocean state estimate for this
103 region on 23 April 2015. The open boundary conditions are one-way nested, updated once per day,
104 and relaxed to the CMEMS state estimate for each subsequent day over a sponge region 2° wide
105 on all edges of the domain. The timescale of this relaxation increases linearly as one approaches
106 the edge of the domain, ranging from 30 days at the inner edge of the sponge region to one day at
107 the boundary.
108

109 The vertical grid spacing is 5 meters over the top 100 m of the water column and increases
110 by a factor of 1.1 for each level below that, up to a maximum of 50 m. The vertical grid
111 consists of 125 levels, thus extending down to 4600 m. Model bathymetry is provided by the
112 General Bathymetric Chart of the Oceans (GEBCO) 2014 global 30-arc-second (~ 1 km) product
113 (<http://www.gebco.net>), and is interpolated appropriately to match the resolution of each simula-
114 tion. Wind stress and surface heat forcing are provided by daily snapshots of the European Centre
115 for Medium-Range Weather Forecasts (ECMWF) atmospheric analysis for the time period from
116 April to July 2015, which are interpolated from $1/4^\circ$ to the appropriate resolution. Lastly, each
117 simulation uses a vertical viscosity $\nu_v = 10^{-4} \text{ m}^2 \text{ s}^{-1}$, a vertical temperature and salt diffusivity
118 $\kappa_v = 10^{-5} \text{ m}^2 \text{ s}^{-1}$, and a combination of modified harmonic and biharmonic Leith horizontal vis-
119 cosity (Leith 1996; Fox-Kemper and Menemenlis 2008) with tuning coefficients of 1.5 and 2.0,
120 respectively. Added to this is a biharmonic horizontal viscosity which varies in strength according
121 to the grid resolution according to $\nu_{bh} = 0.1 \times (\Delta x \Delta y)^{3/2} \text{ m}^4 \text{ s}^{-1}$ (e.g. Chassignet and Garraffo
122 2001). The K-Profile Parameterization (Large et al. 1994) is used to represent the vertical mixing
123 of momentum and tracers in the surface boundary layer.

124 The $1/12^\circ$ simulation is run from 23 April to 31 July 2015, with daily-averaged output. The next
 125 simulation in the nesting hierarchy, at $1/24^\circ$ resolution, uses the same domain extent as the $1/12^\circ$
 126 simulation, and is also run until 31 July 2015. The open boundary conditions for this simulation
 127 are also provided by the interpolated CMEMS state estimate. Due to computational expense, the
 128 final three simulations in the hierarchy, at $1/48^\circ$, $1/96^\circ$ and $1/192^\circ$ resolution, are run until 31
 129 June 2015 on a smaller domain, 60° S to 48° S and 80° W to 40° W. Detailed analysis is performed
 130 by time-averaging over the month of June 2015 (see below), giving an effective spin-up time of just
 131 over one month. Because the mesoscale eddy field in the $1/12^\circ$ CMEMS state estimate is already
 132 fully spun-up and the growth timescale of mixed layer submesoscale eddies is $\mathcal{O}(1)$ day (e.g.
 133 Fox-Kemper et al. 2008), this is sufficient spin-up time for both the submesoscale and mesoscale
 134 kinetic energy fields to saturate (not shown).

135 The open boundary conditions for these simulations are provided from the daily snapshots of the
 136 $1/24^\circ$ simulation. Each simulation in the nesting hierarchy is initialized using the model state of
 137 the simulation one level coarser and after one day of simulated time, e.g. the $1/24^\circ$ is initialized
 138 on 24 April using the solution of the $1/12^\circ$ simulation, and so on. This allows the model to
 139 adjust to each new resolution and reduces spurious numerical artifacts which may arise from the
 140 interpolation. The choice to double the grid resolution at each level of the nesting procedure
 141 was made to minimize the risk that these numerical artifacts would crash the model. While it is
 142 possible that larger jumps in resolution could have been taken without inducing a model crash,
 143 limited computing resources prevented exploration of more aggressive downscaling procedures.

144 The analysis in this manuscript will primarily use output from the highest-resolution, $1/192^\circ$,
 145 simulation and will focus on dynamics in the surface boundary layer. Surface fields from this
 146 simulation are saved as hourly averages, and full 3D fields are saved as daily averages. The
 147 horizontal resolution is anisotropic and varies with latitude, but remains between 290 and 380 m

in the zonal direction in this simulation. The meridional resolution is fixed at around 590 m. This simulation is four times higher resolution than the simulations of Rocha et al. (2016), which were also run for the Drake Passage region, and thus permits more small-scale variability, though unlike Rocha et al. (2016) these simulations do not include tidal forcing. The resolution of this simulation is expected to fully resolve submesoscale mixed layer baroclinic eddies (hereafter MLE).

The $1/192^\circ$ simulation is able to successfully capture key features of the circulation in and around the Scotia Sea region (e.g. Sokolov and Rintoul 2009). Flow along the ACC has a strong barotropic component, is predominantly zonal, and consists of several jets with speeds $> 1 \text{ m s}^{-1}$. The Subantarctic and Polar Fronts are located close together in the Drake Passage constriction. These fronts separate just east of Burdwood Bank (54° W) where the Subantarctic Front is redirected north to connect with the Malvinas Current (Figure 1). Mesoscale meanders and eddies develop south of the Scotia Ridge in the Scotia Sea, a region characterized as an eddy “hot spot” (Frenger et al. 2015). The time-averaged eddy kinetic energy from the model ranges from 10^{-2} to $10^{-1} \text{ m}^2 \text{ s}^{-2}$ (Figure 4), in agreement with EKE estimates calculated from altimetry-derived geostrophic surface currents (AVISO; 1993-2015).

3. Results

Due to the variability in the $1/192^\circ$ simulation on small spatial and fast time scales, further averaging is performed as part of the analysis. Following the notation of Rosso et al. (2015), temporal means will be denoted by an overbar ($\bar{\cdot}$) and are performed over the month of June 2015, and angle brackets $\langle \cdot \rangle$ indicate a spatial average. The fluctuating part of the flow is defined as the departure from the time mean. The mesoscale component, denoted with subscript M , is obtained by applying a 2D convolution filter of width $32\Delta x$, or $1/6^\circ$, to the fluctuations. This filter width, which is about 16 km, is chosen because it lies at the approximate cutoff between mesoscales,

171 whose characteristic horizontal length scales are 10-100 km, and submesoscales, which occupy the
 172 range of 1-10 km (e.g. Thomas et al. 2008). The submesoscale component, denoted with subscript
 173 s , is the residual between the unfiltered fluctuations and the mesoscale fields, and includes all
 174 dynamics smaller than the filter width.

175 *a. Change in mixed layer depth and vertical velocity*

176 The effects of downscaling from mesoscale-permitting to submesoscale-permitting resolution
 177 have been explored in previous studies comparing model dynamics at multiple scales (e.g. Capet
 178 et al. 2008a,b,c; Rosso et al. 2014, 2015, 2016), which is most readily seen in the appearance of
 179 MLE. MLE are energised by converting potential energy into kinetic energy, and in doing so tilt
 180 density surfaces toward the horizontal and increase the mixed layer stratification (e.g. Boccaletti
 181 et al. 2007; Fox-Kemper et al. 2008). Here we will define the mixed layer depth H_{ML} to be the shall-
 182 lowest depth where the change in density $\Delta\rho = \rho|_z - \rho|_{z=0} > 0.03 \text{ kg m}^{-3}$ (de Boyer Montégut
 183 et al. 2004). Because the effects of MLE lead to a higher rate of change in the density with depth,
 184 they can also result in a shallower mixed layer depth.

185 Figure 2a shows $\overline{H_{ML}}$ from the $1/192^\circ$ simulation, which exhibits significant variability in both
 186 magnitude and spatial distribution. The range of $\overline{H_{ML}}$ observed in the Drake Passage tends to
 187 remain between 75 and 250 m, broadly in agreement with Argo climatology of mixed layer depths
 188 for this region at the onset of the Southern Hemisphere winter (e.g. Dong et al. 2008; Holte and
 189 Talley 2009). The model $\overline{H_{ML}}$ field exhibits sharp meridional gradients in comparison with the
 190 Argo climatology, likely due both to the high resolution of the model and the coarse mapping of
 191 float profiles in the climatology (2 degrees in latitude and 5 degrees in longitude).

192 The change in $\overline{H_{ML}}$ between the $1/12^\circ$ and $1/192^\circ$ simulations, $\Delta\overline{H_{ML}}$, is shown in Figure 2b,
 193 where positive values indicate a shallowing of the mixed layer depth with increasing resolution.

As anticipated, $\overline{H_{ML}}$ indeed becomes shallower as the model resolution increases, but the change is greater in some regions than in others. In particular, in the westernmost region from 76° W to 72° W, $\Delta\overline{H_{ML}}$ exceeds 100 m in places, as well as in a conspicuous jet-like feature extending from the tip of the continent at 55° S. In contrast, the region east of 48° W shows almost no change in $\overline{H_{ML}}$ with increased resolution.

Submesoscale motions are also associated with a loss of balance and a corresponding increase in the strength of vertical circulations (e.g. Mahadevan and Tandon 2006; Capet et al. 2008b; Thomas et al. 2008; Klein and Lapeyre 2009). Modelling at higher resolution is expected to result in an increase in the root-mean square vertical velocity, $\overline{w_{rms}} = \sqrt{\overline{w^2}}$, as smaller-scale processes become better resolved. Indeed, the $\overline{w_{rms}}$ field from the $1/192^\circ$ simulation, shown in Figure 2c, is significantly intensified in comparison with the lower resolution simulations (see also Figure 9 for numerical values). If submesoscale dynamics are indeed assumed to be the principal driver of the change in $\overline{H_{ML}}$ and increase in $\overline{w_{rms}}$ between these models, this suggests some spatial inhomogeneity in the strength of the submesoscale eddy field. The nature of this inhomogeneity, and its implications for modelling of the ocean boundary layer, are investigated further on.

Also outlined in Figure 2 are the 400-m isobath (white line, panel (c)), and two regions, R1 and R2, which will be analysed in Section c. These regions are chosen because they exhibit the most extreme contrasts between their respective mesoscale and submesoscale motions, and the dynamical consequences of each. The 400-m isobath is chosen as a demarcation between the continental shelf, featuring $\mathcal{O}(1)$ km eddies whose size is limited by the shallow depth (Figure 3a), and deep water. To enable a fair comparison between different regions, the analysis in this paper will only consider locations where the depth is greater than 400 m. The 400-m isobath and analysis regions are outlined on all subsequent Figures as a visual aid.

217 *b. Submesoscale intensity varies spatially*

218 Submesoscale processes are associated with $\mathcal{O}(1)$ Rossby number (Thomas et al. 2008). One
219 metric for the local submesoscale intensity could be the Rossby number $Ro = |\zeta/f|$ based on the
220 vertical component of the relative vorticity $\zeta = \partial v/\partial x - \partial u/\partial y$, where (u, v) is the horizontal
221 velocity and f is the Coriolis parameter. While this definition of Ro can be straightforwardly
222 calculated from the simulation data, this metric does not distinguish submesoscale features from
223 strongly rotating mesoscale eddies or intense jets. Figure 3a shows a snapshot of ζ taken from
224 June 30, 2015, where strongly rotating mesoscale eddies can easily be identified east of 56° W.

225 To isolate submesoscale features from these larger structures we define the “mixed layer baro-
226 clinic” Rossby number, $Ro_b = |\zeta_b/f|$, where $\zeta_b = \zeta|_{z=0} - \zeta|_{z=-400m}$ is the difference in relative
227 vorticity between the surface and a depth of 400 m. This depth is chosen because it is well below
228 the maximum $\overline{H_{ML}}$ of 221 m within the domain (Figure 2) and deeper than the continental shelf, so
229 that statistics measured at this depth will be considered representative of the interior ocean in deep
230 water. The expectation is that submesoscale features which are confined to the mixed layer will
231 have large surface relative vorticity but small relative vorticity below the mixed layer. In contrast,
232 features such as jets and mesoscale eddies which extend well below the mixed layer are expected
233 to have similar relative vorticity at both depths, so ζ_b for these features will be small. Therefore,
234 this definition is intended to distinguish mixed layer submesoscales from these other features. Ro_b
235 is not calculated in regions where the ocean depth is less than 400 m.

236 Figure 3b shows $\overline{Ro_b}$, where it is apparent that the mesoscale structures on the eastern side of
237 the domain have been filtered out by the differencing operation. Values of $\overline{Ro_b}$ near $\mathcal{O}(1)$ suggest
238 higher activity of mixed layer submesoscales, whose location corresponds to the small vortical

features seen on the southwest corner of Figure 3a. Regions where the depth is shallower than 400 m have been grayed out, and are excluded from the detailed analysis in Section c.

c. Correlation between mesoscales, submesoscales, $\overline{w_{rms}}$, and $\overline{\Delta H_{ML}}$

Recent work by Rosso et al. (2015) employed a spatial filtering method to explore the relationship between vertical velocity and mesoscale eddy kinetic energy and strain in the Kerguelen Plateau region of the Southern Ocean. Following their approach, the kinetic energy associated with the mesoscale and submesoscale velocities can be defined $\frac{1}{2}\overline{|\mathbf{u}_M|^2}$ and $\frac{1}{2}\overline{|\mathbf{u}_S|^2}$, respectively. The mesoscale strain field can be diagnosed using the filtered velocity field as

$$\overline{S_M} = \left[\left(\frac{\partial u_M}{\partial x} - \frac{\partial v_M}{\partial y} \right)^2 + \left(\frac{\partial v_M}{\partial x} + \frac{\partial u_M}{\partial y} \right)^2 \right]^{1/2}. \quad (1)$$

Figure 4 shows the surface mesoscale and submesoscale kinetic energies and mesoscale strain.

The maps of $\overline{\Delta H_{ML}}$, $\overline{w_{rms}}$, $\overline{Ro_b}$, and the mesoscale fields in Figures 2 - 4 reveal an interesting spatial correlation between these quantities, where the largest vertical velocities are co-located with regions of high mesoscale KE and strain, and the largest values of $\overline{\Delta H_{ML}}$ occur where $\overline{Ro_b}$ is largest. Both results taken individually are unsurprising. Strong vertical circulations can occur at mesoscale fronts (e.g. Nagai et al., 2006) and filaments (e.g. Lapeyre and Klein, 2006, McWilliams et al., 2014) in addition to being often associated with submesoscale dynamics. A large change in mixed layer depth can occur in regions of intense submesoscale activity due to the influence of MLE in restratifying the boundary layer. A surprising feature of these maps is the appearance of regions with large $\overline{w_{rms}}$ and weak submesoscales with small $\overline{Ro_b}$, the most notable of which are in and around R2, and regions of strong submesoscale activity with large $\overline{Ro_b}$ and comparatively small $\overline{w_{rms}}$, such as the area in and north of R1.

1) CORRELATIONS WITHIN R1 AND R2

In the previous figures two regions, R1 and R2 (Figure 2b), have been outlined which will be analysed further here. R1, which extends from 78° W to 72° W and 58° S to 55° S, exhibits strong surface submesoscale activity as indicated by the maps of $\overline{Ro_b}$, $\overline{\Delta H_{ML}}$, and submesoscale kinetic energy (Figure 4b), but relatively weak mesoscale flow (Figure 4a, c). R2 extends from 59° W to 48° W and 58° S to 55.5° S and features large $\overline{w_{rms}}$, mesoscale kinetic energy, and mesoscale strain, but small $\overline{Ro_b}$ and $\overline{\Delta H_{ML}}$. Note that the mean $\overline{H_{ML}}$ in both regions is similar (Figure 2a), despite significant local variations in R1.

The vertical profiles of $\overline{w_{rms}}$ are consistent with the above interpretation of each region (Figure 5). For this analysis the vertical velocity field is filtered into mesoscale and submesoscale components before being squared and time-averaged, yielding $\overline{(w_{rms})_M} = \sqrt{\overline{w_M^2}}$ and $\overline{(w_{rms})_S} = \sqrt{\overline{w_S^2}}$. Vertical profiles of these fields are obtained by spatially averaging over R1 and R2, and are shown in Figure 5a and Figure 5b, respectively. The submesoscale component in R1 (red line, Figure 5a) features a local maximum in the mixed layer which extends down to 150 m, the approximate mean mixed layer depth for this region (Figure 2a), suggesting the presence of intensified vertical motions from submesoscales in the mixed layer. The submesoscale component in R2 (red line, Figure 5b) has less surface intensification. Both mesoscale and submesoscale components increase with depth, with the submesoscale component being larger than the mesoscale component at nearly all depths. These results are consistent with those of Rosso et al. (2015, Figure 3), who attributed part of the submesoscale component at the surface and the bottom intensification to internal lee wave activity. To further justify this point, histograms of bathymetry (Figure 5; gray bars) show that the largest vertical velocities occur at or slightly above the bottom depths in both R1 and R2.

281 The especially large velocities in R2 could also be partly due to the generation of lee waves from
 282 Drake Passage (e.g. Naveira Garabato et al. 2004; St. Laurent et al. 2012).

283 2) CORRELATIONS OVER THE FULL DOMAIN

284 Scatter plots can also be used to illustrate correlations between different variables in this analy-
 285 sis. Figure 6 shows how $\langle \overline{w_{rms}} \rangle$ and $\langle \overline{\Delta H_{ML}} \rangle$ trend with $\langle \overline{Ro_b} \rangle$, the mesoscale KE, and mesoscale
 286 strain over the full domain. In this analysis each field is averaged over 1° boxes and includes
 287 only locations where the mean depth over these boxes exceeds 400 m. Error bars are shown for
 288 each data point and represent one standard deviation above and below the mean for that box. The
 289 locations for each data point are indicated by color: blue dots indicate locations in R1, red dots
 290 indicate locations in R2, and gray dots indicate locations throughout the rest of the domain. A sys-
 291 tematic increase in $\langle \overline{w_{rms}} \rangle$ is observed at larger values of both mesoscale KE (panel (c), correlation
 292 coefficient $r = 0.80$) and strain (panel (e), $r = 0.73$). $\langle \overline{w_{rms}} \rangle$ also trends positively with $\langle \overline{Ro_b} \rangle$, con-
 293 sistent with submesoscale-driven vertical velocities. However, a second, sharper upward trend is
 294 evident near $\langle \overline{Ro_b} \rangle = 10^{-1}$, with vertical velocities approaching 100 m day^{-1} . These large vertical
 295 velocities and values of $\overline{Ro_b} \sim 10^{-1}$ correspond to locations with large mesoscale KE and strain
 296 in R2. Due to the two competing trends, an overall weak correlation exists between $\langle \overline{w_{rms}} \rangle$ and
 297 $\langle \overline{Ro_b} \rangle$ (panel (a), $r = 0.05$) across the domain. Conversely, $\langle \overline{\Delta H_{ML}} \rangle$ increases with $\langle \overline{Ro_b} \rangle$ (panel
 298 (b), $r = 0.64$) but shows no clear trend with either the mesoscale KE (panel (d), $r = -0.12$) or
 299 strain (panel (f), $r = 0.20$).

300 A full list of the correlation coefficients between $\langle \overline{w_{rms}} \rangle$, $\langle \overline{\Delta H_{ML}} \rangle$, and each of these variables
 301 appears in Table 1. In this Table different regions are indicated by font style, with boldface font
 302 indicating values over the whole domain, standard font values for R1, and italic font values for
 303 R2. The correlation coefficients tend to be consistent from region to region for strongly corre-

lated variables, whereas the coefficients for weakly correlated variables tend to have much more variation.

Due to the occurrence of many submesoscale instabilities at mixed layer fronts, extant submesoscale parameterizations have been designed to be sensitive to the frontal strength, $|\nabla_h b|$ (e.g. Fox-Kemper et al. 2008; Canuto and Dubovikov 2010; Bachman et al. 2017), where ∇_h is the horizontal gradient operator and b is the buoyancy. Maps of the frontal strength from both the $1/12^\circ$ and $1/192^\circ$ simulations are shown in Figure 7 (top row). The spatial pattern of the frontal strength qualitatively matches that of the change in mixed layer depth, $\overline{\Delta H_{ML}}$, between simulations (Figure 2b). The higher resolution model permits tighter fronts to form, reflected in a tendency for $|\nabla_h b|$ to be larger almost everywhere in the $1/192^\circ$ simulation. When $|\nabla_h b|$ from these simulations is coarse-grained over 1° boxes a positive correlation is evident between $\langle \overline{\Delta H_{ML}} \rangle$ and $\langle |\nabla_h b| \rangle$ in both the $1/12^\circ$ ($r = 0.58$) and $1/192^\circ$ ($r = 0.66$) models (Figure 7, bottom row). The correlation between $\langle |\nabla_h b| \rangle$ and $\langle \overline{w_{rms}} \rangle$ is weak ($r = -0.18$ for both models), as is the direct correlation between $\langle \overline{\Delta H_{ML}} \rangle$ and $\langle \overline{w_{rms}} \rangle$ ($r = 0.08$ for the $1/12^\circ$ model; $r = 0.07$ for the $1/192^\circ$ model; not shown).

319 *d. A possible mechanism for large $\overline{w_{rms}}$*

A question remains about how to physically interpret the large $\overline{w_{rms}}$ in R2 if it is not associated with submesoscale circulations. Bottom intensification of the vertical velocity due to topography can explain the large velocities below 3000 m, and the region is known to be a hotspot for lee wave generation (Watson et al. 2013). Rosso et al. (2015) found that such bottom-generated internal waves only occasionally reached the mid- to upper ocean, however, and that the dominant temporal frequency of the submesoscale vertical velocity was much slower than could be explained by

internal wave activity. A local maximum in $\overline{w_{rms}}$ shallower than 500 m depth in R2 also suggests a surface-intensified generation mechanism (Figure 5b).

Rocha et al. (2016) calculated horizontal wavenumber spectra in Drake Passage and found that ageostrophic motions in this region are likely dominated by internal waves, which imprint strongly on the near-surface kinetic energy at scales between 10 and 40 km and might explain the strong velocities in R2. A possible source of these waves was explored by Shakespeare and Hogg (2017), who highlighted the process of wave generation through frontogenesis in the Southern Ocean. Recent studies by Shakespeare and Taylor (2014, 2015, 2016) focused on wave generation and dynamics of the ageostrophic secondary circulation which develops at fronts undergoing large strain (up to $\mathcal{O}(f)$), and have led to a theoretical scaling for the vertical velocity associated with these fronts (Shakespeare 2015; Shakespeare and Taylor 2016),

$$W \sim H\zeta \left(1 + \frac{\zeta}{f}\right) \frac{S}{f^2} (f^2 + S^2)^{1/2}. \quad (2)$$

This scaling is a function of a depth scale, H , Coriolis parameter, f , large-scale relative vorticity, ζ , and large-scale strain, S . Here we compare this scaling to the simulated flow by using the mixed layer depth $\overline{H_{ML}}$ as the depth scale, a low-pass filtered $\overline{\zeta_M}$ as the large-scale relative vorticity, and $\overline{S_M}$ as the large scale strain. The map of W using these diagnosed parameters and a proportionality coefficient of 1.5 is shown in Figure 8a. Comparing against the map of $\overline{w_{rms}}$ in Figure 8b, the scaling is a good approximation to the diagnosed $\overline{w_{rms}}$ throughout the domain. The scaling is less skillful in the boundary current around the edge of the continent and on the continental shelf, but it is unclear whether H_{ML} and mesoscale parameters are appropriate in these shallow regions. These areas lie within the 400 m isobath (white line) and will not be considered further. Figure 8c shows

346 a scatter plot of the 1° -averaged $\langle \overline{w_{rms}} \rangle$ against $\langle W \rangle$. The scaling shows good agreement ($r = 0.78$)
347 with the diagnosed $\langle \overline{w_{rms}} \rangle$ across over an order of magnitude.

348 *e. Sensitivity of $\langle \overline{w_{rms}} \rangle$ to grid resolution*

349 The simulation results and comparison against theory suggest frontogenesis and complex bottom
350 topography as two mechanisms responsible for large $\langle \overline{w_{rms}} \rangle$ in the Scotia Sea region. Because both
351 the mesoscale strain field (Figure 4c) and bottom topography are highly variable in this region, it
352 is likely that the magnitude of $\langle \overline{w_{rms}} \rangle$ would vary significantly over the rest of the Southern Ocean
353 as well.

354 Very few modelling studies have been conducted at sufficient resolution to capture mesoscale,
355 submesoscale, and topographic interactions, particularly with regard to wave-driven vertical mo-
356 tions. Due to the important role waves play in exchanging energy with the large-scale flow at rough
357 topography (e.g. Nikurashin and Ferrari 2010a,b, 2011) and driving mixing in the deep ocean
358 (Wunsch and Ferrari 2004), such studies are needed to fill gaps in our understanding of how the
359 energy of the general circulation is dissipated. From an ocean modelling perspective, these studies
360 are needed to assess and accurately estimate dissipation due to unresolved wave generation and
361 breaking. The simulations used here offer a unique opportunity to explore these multi-scale inter-
362 actions because they are run at five different horizontal resolutions, spanning from a mesoscale-
363 permitting regime with no submesoscales in the $1/12^\circ$ model, to a submesoscale-resolving regime
364 with significant wave activity in the $1/192^\circ$ model.

365 Previous studies using high-resolution numerical simulations have found varying sensitivity of
366 $\langle \overline{w_{rms}} \rangle$ to changing the horizontal resolution. This sensitivity can be straightforwardly quantified
367 by defining an enhancement factor,

$$s = \frac{\text{Fractional change in } \langle \overline{w_{rms}} \rangle}{\text{Fractional change in resolution}}. \quad (3)$$

The realistic simulations of Rosso et al. (2014) and Capet et al. (2008a) found $s = 2.75$ and $s = 2.5$, respectively, which were much higher than $s = 0.57$ and $s = 0.2$ found by Lévy et al. (2001) and Lévy et al. (2012). The latter two simulations were run using an idealised, flat-bottom domain, however, implicating bottom topography as the reason for the pronounced difference in s between these studies.

Because each model in our nesting hierarchy is exactly twice the resolution of the previous model, we are able to calculate s as a function of resolution as well. Figure 9 shows how $\langle \overline{w_{rms}} \rangle$ is enhanced by increased resolution, where the $\overline{w_{rms}}$ fields are averaged vertically over the top 400 m and horizontally over (a) R1, (b) R2, and (c) the whole domain. As expected, the values of $\langle \overline{w_{rms}} \rangle$ monotonically increase with resolution, although s is dependent on both resolution and location. The values of s stay relatively consistent in R1, remaining between 1.1 and 1.4 each time the resolution is doubled. This is the same magnitude of increase seen in R2 and over the whole domain when the resolution is increased to $1/24^\circ$ and $1/48^\circ$. However, s increases noticeably each time when downscaling to $1/96^\circ$ and $1/192^\circ$.

We hypothesize that the lower values of s up to $1/48^\circ$ occur because the resolved mesoscale dynamics are relatively unchanged by downscaling between the $1/12^\circ$ and $1/48^\circ$ models. That is, the eddy flow up to this resolution is driven primarily by baroclinic turbulence, while smaller submesoscale instabilities, convection, and waves remain unresolved. The emergence of submesoscale dynamics and some internal wave activity causes a jump in s at $1/96^\circ$, which is further accentuated by a vigorous internal wave field appearing at $1/192^\circ$, particularly in R2. Interestingly, spatial inhomogeneity also begins to emerge at these high resolutions, as reflected by the

sharp increase in s in R2 compared with R1. Counterintuitively, it is R2 that is responsible for the largest value, $s = 2.4$, at $1/192^\circ$.

4. Conclusions

In this study a series of numerical simulations of the Scotia Sea region have been used to investigate the effects of mesoscale and submesoscale processes on the oceanic surface boundary layer. The highest-resolution member of the series has a grid spacing of about 500 m and is capable of resolving submesoscale dynamics, enabling an analysis of the oceanic boundary layer which is not possible using coarser models. The “baroclinic Rossby number”, Ro_b , defined as the difference in relative vorticity between the surface and the interior, has been used to identify regions of mixed layer submesoscale activity. A comparison of the highest-resolution model against the lowest-resolution model, which has a resolution of about 8 km and therefore is unable to resolve any submesoscales, shows significant differences in many key metrics, including relative vorticity, frontal strength, mixed layer depth, RMS vertical velocity, and kinetic energy.

Here we have highlighted differences in the time-averaged mixed layer depth, $\overline{\Delta H_{ML}}$, and RMS vertical velocity, $\overline{w_{rms}}$, between the low- and high-resolution models because these metrics are especially significant to the ocean’s role in affecting climate. Ocean-atmosphere exchange is modulated by the character of the mixed layer, with the mixed layer depth affecting the ocean’s ability to uptake and store heat and trace gases on short timescales. These air-sea interactions are especially important in the Southern Ocean, which is a key region for anthropogenic carbon uptake (Khatiwala et al. 2009; Sallée et al. 2012; Frölicher et al. 2015) through the subduction of mode and intermediate waters. Large, persistent vertical velocity can transport tracers between the mixed layer and the ocean interior where it can be stored on long timescales, and is also an indicator of nutrient supply for phytoplankton growth (e.g. Lévy et al. 2001). These metrics are expected to be

particularly sensitive to model resolution between the meso- and submesoscales, where dynamics become less constrained by the Earth’s rotation and vertical transport is enhanced. Understanding how the mixed layer responds to dynamics at multiple scales is therefore crucial to our ability to predict the future climate, making the models in this study especially useful in this regard.

Previous work by Rosso et al. (2015) used a submesoscale-resolving model to establish a relationship between regions of large submesoscale vertical velocity, $|\overline{w_S}|$, and mesoscale kinetic energy and strain, treating $|\overline{w_S}|$ as a proxy for near-surface submesoscale activity. However, $|\overline{w_S}|$ does not distinguish between small-scale processes like internal waves which can drive strong vertical motions, and the more climatically relevant mixed layer submesoscales which modulate air-sea exchange. In this work we take a slightly different approach, which is to first identify regions of mixed layer submesoscale activity using maps of $\overline{Ro_b}$ before performing analysis of $\overline{w_{rms}}$. In agreement with Rosso et al. (2015), we find that submesoscales are associated with enhanced $\overline{w_{rms}}$, but also find an even larger enhancement of $\overline{w_{rms}}$ which may be due to mesoscale frontogenesis (e.g. Shakespeare and Taylor 2014). We also find a close link between regions of enhanced $\overline{Ro_b}$ and large $\overline{\Delta H_{ML}}$, the latter of which is likely caused by resolving mixed layer baroclinic instability.

These results suggest a similar but nuanced interpretation relative to that of Rosso et al. (2015). Submesoscales are coincident with strong vertical velocities, but regions of strong vertical velocity should not necessarily be used as an indicator of enhanced mixed layer submesoscale activity. Mesoscale frontogenesis is suggested as a mechanism leading to large vertical velocity in certain regions where submesoscales are not necessarily present, and the magnitude of this velocity can exceed that associated with submesoscales. However, the regions of large vertical velocity are not always associated with a shallowing of the mixed layer depth. The interpretation of these results has significant consequences for the development of deterministic submesoscale eddy parameterizations, whose effects are sensitive to the mesoscale flow. Our results indicate no systematic rela-

436 tionship between mesoscale kinetic energy and strain and $\Delta\overline{H_{ML}}$, raising questions about whether
 437 these fields are appropriate to inform an eddy parameterization (e.g. Rosso et al. 2015). We find
 438 a stronger correlation between $\Delta\overline{H_{ML}}$ and the coarse-resolution lateral density gradient, the latter
 439 of which is already used as the basis for multiple submesoscale eddy closures (Fox-Kemper et al.
 440 2008; Bachman et al. 2017).

441 Internal waves act as a primary pathway toward energy dissipation and play a key role in driving
 442 mixing in the deep ocean (Wunsch and Ferrari 2004; Ferrari and Wunsch 2009). Much of this
 443 mixing and dissipation is due to wave breaking, a process which is parameterized in hydrostatic
 444 models by the use of vertical eddy viscosity but could be explicitly resolved upon moving to non-
 445 hydrostatic modelling. The richness of the internal wave field in the $1/192^\circ$ simulation suggests
 446 that it lies close to the resolution threshold where a nonhydrostatic model would be appropriate.
 447 The nonhydrostatic parameter (Marshall et al. 1997b), $\eta = h^2/(L^2 Ri)$, can be used as a gauge
 448 of whether a nonhydrostatic model is necessary, where h and L are characteristic depth and hor-
 449 izontal length scales, and the Richardson number, $Ri = N^2 h^2 / U^2$, is a function of the buoyancy
 450 frequency, N , and characteristic velocity scale, U . This parameter is likely to be largest in the
 451 mixed layer where Ri is small and the aspect ratio, h/L , is large. Using values from the $1/192^\circ$
 452 simulation, where $h = 100$ m is an approximate mixed layer depth, $L = \Delta x = 500$ m is an average
 453 grid spacing, and $Ri = 1$ for the mixed layer (e.g. Young 1994; Thomas et al. 2008; Bachman and
 454 Taylor 2016), we have $\eta = 1/25 \ll 1$, so that motion is approximately hydrostatic. It is possible
 455 that another downscaling to $1/384^\circ$ would require a nonhydrostatic model; however, because the
 456 computational burden of nonhydrostatic models is significantly higher, this realm of modelling
 457 tends to remain out of reach for regional studies such as those presented here.

458 The Southern Ocean has several characteristics, such as weak vertical stratification in the up-
 459 per ocean, strong mesoscale kinetic energy, and significant eddy-mean flow interaction (e.g.

Naveira Garabato et al. 2011), and further research is required to understand whether the correlations and localized submesoscale activity we find in the Scotia Sea region occur in the rest of the ocean as well. Our simulations indicate that submesoscales are spatially variable and can be highly active immediately adjacent to a region where they are nearly absent. It is unclear what causes this spatial inhomogeneity, especially given that the regions of highest mesoscale strain, where we would expect submesoscale-generating mechanisms like frontogenesis (Thomas and Ferrari 2008) and frontal instabilities (Mahadevan and Tandon 2006; Thomas et al. 2008) to be prevalent, are not always associated with elevated submesoscale activity. Further research is necessary to determine the causes and consequences of this observation, and is ongoing.

Acknowledgments. The authors gratefully acknowledge support from the Natural Environment Research Council, awards NE/J010472/1 and NE/J009857/1. This work used the ARCHER UK National Supercomputing Service (<http://www.archer.ac.uk>). Discussions and advice from Dave Munday about how to configure the simulations were extremely helpful. The altimeter products were produced by Ssalto/Duacs and distributed by Aviso, with support from Cnes (<http://www.aviso.altimetry.fr/duacs/>).

References

- Adams, K., P. Hosegood, J. Taylor, J.-B. Sallée, S. Bachman, and M. Stamper, 2017: Frontal circulation and submesoscale variability during the formation of a Southern Ocean mesoscale eddy. *Journal of Physical Oceanography*, **Accepted for publication**.
- Bachman, S., B. Fox-Kemper, J. Taylor, and L. Thomas, 2017: Parameterization of frontal symmetric instabilities. I: Theory for resolved fronts. *Ocean Modelling*, **109(1)**, 72–95, doi: 10.1016/j.ocemod.2016.12.003.

482 Bachman, S., and J. Taylor, 2016: Numerical simulations of the equilibrium between eddy-induced
 483 restratification and vertical mixing. *Journal of Physical Oceanography*, **46 (3)**, 919–935, doi:
 484 10.1175/JPO-D-15-0110.1.

485 Boccaletti, G., R. Ferrari, and B. Fox-Kemper, 2007: Mixed layer instabilities and restratification.
 486 *Journal of Physical Oceanography*, **37(9)**, 2228–2250, doi:10.1175/JPO3101.1.

487 Brannigan, L., D. Marshall, A. Naveira Garabato, and A. Nurser, 2015: The seasonal cycle of
 488 submesoscale flows. *Ocean Modelling*, **92**, 69–84, doi:10.1016/j.ocemod.2015.05.002.

489 Buckingham, C., and Coauthors, 2016: Seasonality of submesoscale flows in the ocean surface
 490 boundary layer. *Geophysical Research Letters*, **43(5)**, 2118–2126, doi:10.1002/2016GL068009.

491 Callies, J., R. Ferrari, J. Klymak, and J. Gula, 2015: Seasonality in submesoscale turbulence.
 492 *Nature Communications*, **6:6862**, doi:10.1038/ncomms7862.

493 Canuto, V., and M. Dubovikov, 2010: Mixed layer sub-mesoscale parameterization – part 1:
 494 Derivation and assessment. *Ocean Science*, **6 (3)**, 679–693, doi:10.5194/os-6-679-2010.

495 Capet, X., J. McWilliams, M. Molemaker, and A. Shchepetkin, 2008a: Mesoscale to submesoscale
 496 transition in the California Current system. Part I: Flow structure, eddy flux, and observational
 497 tests. *Journal of Physical Oceanography*, **38(1)**, 29–43, doi:10.1175/2007JPO3671.1.

498 Capet, X., J. McWilliams, M. Molemaker, and A. Shchepetkin, 2008b: Mesoscale to subme-
 499 soscale transition in the California Current system. Part II: Frontal processes. *Journal of Physi-
 500 cal Oceanography*, **38(1)**, 44–64, doi:10.1175/2007JPO3672.1.

501 Capet, X., J. McWilliams, M. Molemaker, and A. Shchepetkin, 2008c: Mesoscale to submesoscale
 502 transition in the California Current system. Part III: Energy balance and flux. *Journal of Physical
 503 Oceanography*, **38(10)**, 2256–2269, doi:10.1175/2008JPO3810.1.

504 Cerovečki, I., L. Talley, M. Mazloff, and G. Maze, 2013: Subantarctic Mode Water formation, de-
 505 struction, and export in the eddy-permitting Southern Ocean state estimate. *Journal of Physical*
 506 *Oceanography*, **43**(7), 1485–1511, doi:10.1175/JPO-D-12-0121.1.

507 Chassignet, E., and Z. Garraffo, 2001: Viscosity parameterization and the Gulf Stream separation.
 508 *Proceedings of the 'Aha Huliko'a Hawaiian Winter Workshop*, University of Hawaii at Manoa,
 509 Honolulu, Hawaii, 37–41.

510 D'Asaro, E., C. Lee, L. Rainville, R. Harcourt, and L. Thomas, 2011: Enhanced turbulence and
 511 energy dissipation at ocean fronts. *Science*, **332**(6027), 318–322, doi:10.1126/science.1201515.

512 de Boyer Montégut, C., G. Madec, A. Fischer, A. Lazar, and D. Iudicone, 2004: Mixed layer depth
 513 over the global ocean: An examination of profile data and a profile-based climatology. *Journal*
 514 *of Geophysical Research: Oceans*, **109**, C12 003, doi:10.1029/2004JC002378.

515 Dong, S., J. Sprintall, S. Gille, and L. Talley, 2008: Southern Ocean mixed-layer depth from argo
 516 float profiles. *Journal of Geophysical Research: Oceans*, **113** (C6), doi:10.1029/2006JC004051.

517 Ferrari, R., and C. Wunsch, 2009: Ocean circulation kinetic energy: Reservoirs, sources, and
 518 sinks. *Annual Review of Fluid Mechanics*, **41**, 253–282, doi:10.1146/annurev.fluid.40.111406.
 519 102139.

520 Flierl, G. R., and D. McGillicuddy, 2002: Mesoscale and submesoscale physical-biological inter-
 521 actions. *The Sea, Vol. 12: Biological-Physical Interactions in the Sea*, ed. by A. R. Robinson, J.
 522 J. McCarthy and B. J. Rothschild, Hoboken, NJ, 113–185.

523 Fox-Kemper, B., R. Ferrari, and R. Hallberg, 2008: Parameterization of mixed layer eddies. Part
 524 I: Theory and diagnosis. *Journal of Physical Oceanography*, **38**(6), 1145–1165, doi:10.1175/
 525 2007JPO3792.1.

526 Fox-Kemper, B., and D. Menemenlis, 2008: Can large eddy simulation techniques improve
 527 mesoscale rich ocean models? *Geophysical Monograph Series*, **177**, 319–337, doi:10.1029/
 528 177GM19.

529 Frenger, I., M. Münnich, N. Gruber, and R. Knutti, 2015: Southern Ocean eddy phenomenology.
 530 *Journal of Geophysical Research: Oceans*, **120(11)**, 7413–7449, doi:10.1002/2015JC011047.

531 Frölicher, T., J. Sarmiento, D. Paynter, J. Dunne, J. Krasting, and M. Winton, 2015: Dominance
 532 of the Southern Ocean in anthropogenic carbon and heat uptake in CMIP5 models. *Journal of*
 533 *Climate*, **28 (2)**, 862–886.

534 Gargett, A., and J. Marra, 2002: Effects of upper ocean physical processes—turbulence, advection,
 535 and air-sea interaction—on oceanic primary production. *The Sea, Vol. 12: Biological-Physical*
 536 *Interactions in the Sea, ed. by A. R. Robinson, J. J. McCarthy and B. J. Rothschild*, Hoboken,
 537 NJ, 19–49.

538 Holte, J., and L. Talley, 2009: A new algorithm for finding mixed layer depths with applications
 539 to Argo data and Subantarctic Mode Water formation. *Journal of Atmospheric and Oceanic*
 540 *Technology*, **26 (9)**, 1920–1939, doi:10.1175/2009JTECHO543.1.

541 Jiao, Y., and W. Dewar, 2015: The energetics of centrifugal instability. *Journal of Physical*
 542 *Oceanography*, **45 (6)**, 1554–1573.

543 Khatiwala, S., F. Primeau, and T. Hall, 2009: Reconstruction of the history of anthropogenic CO₂
 544 concentrations in the ocean. *Nature*, **462(7271)**, 346–349, doi:10.1038/nature08526.

545 Klein, P., and G. Lapeyre, 2009: The oceanic vertical pump induced by mesoscale and subme-
 546 soscale turbulence. *Annual Review of Marine Science*, **1**, 351–375, doi:10.1146/annurev.marine.
 547 010908.163704.

548 Klocker, A., and R. Abernathey, 2014: Global patterns of mesoscale eddy properties and diffusiv-
 549 ities. *Journal of Physical Oceanography*, **44**(3), 1030–1046, doi:10.1175/JPO-D-13-0159.1.

550 Lapeyre, G., and P. Klein, 2006: Dynamics of the upper oceanic layers in terms of surface quasi-
 551 geostrophy theory. *Journal of Physical Oceanography*, **36**(2), 165–176, doi:10.1175/JPO2840.
 552 1.

553 Large, W., J. McWilliams, and S. Doney, 1994: Oceanic vertical mixing: A review and a model
 554 with a nonlocal boundary layer parameterization. *Reviews of Geophysics*, **32** (4), 363–403, doi:
 555 10.1029/94RG01872.

556 Leith, C., 1996: Stochastic models of chaotic systems. *Physica D: Nonlinear Phenomena*, **98**(2),
 557 481–491, doi:10.1016/0167-2789(96)00107-8.

558 Lévy, M., D. Iovino, L. Resplandy, P. Klein, G. Madec, A. Tréguier, S. Masson, and K. Takahashi,
 559 2012: Large-scale impacts of submesoscale dynamics on phytoplankton: Local and remote
 560 effects. *Ocean Modelling*, **43**, 77–93, doi:10.1016/j.ocemod.2011.12.003.

561 Lévy, M., P. Klein, and A. Tréguier, 2001: Impact of sub-mesoscale physics on production and
 562 subduction of phytoplankton in an oligotrophic regime. *Journal of Marine Research*, **59**(4),
 563 535–565, doi:10.1357/002224001762842181.

564 Lévy, M., and A. Martin, 2013: The influence of mesoscale and submesoscale heterogeneity
 565 on ocean biogeochemical reactions. *Global Biogeochemical Cycles*, **27**(4), 1139–1150, doi:
 566 10.1002/2012GB004518.

567 Mahadevan, A., and D. Archer, 2000: Modeling the impact of fronts and mesoscale circulation on
 568 the nutrient supply and biogeochemistry of the upper ocean. *Journal of Geophysical Research:*
 569 *Oceans*, **105**, 1209–1225, doi:10.1029/1999JC900216.

570 Mahadevan, A., and A. Tandon, 2006: An analysis of mechanisms for submesoscale vertical
 571 motion at ocean fronts. *Ocean Modelling*, **14(3)**, 241–256, doi:10.1016/j.ocemod.2006.05.006.

572 Marshall, J., A. Adcroft, C. Hill, L. Perelman, and C. Heisey, 1997a: A finite-volume, incompress-
 573 ible Navier-Stokes model for studies of the ocean on parallel computers. *Journal of Geophysical*
 574 *Research: Oceans*, **102(C3)**, 5753–5766, doi:10.1029/96JC02775.

575 Marshall, J., C. Hill, L. Perelman, and A. Adcroft, 1997b: Hydrostatic, quasi-hydrostatic, and
 576 nonhydrostatic ocean modeling. *Journal of Geophysical Research: Oceans*, **102 (C3)**, 5733–
 577 5752, doi:10.1029/96JC02776.

578 McWilliams, J., J. Molemaker, and I. Yavneh, 2001: From stirring to mixing of momentum: Cas-
 579 cades from balanced flows to dissipation in the oceanic interior. *Proceedings of the 'Aha Hu-*
 580 *liko'a Hawaiian Winter Workshop*, University of Hawaii at Manoa, Honolulu, Hawaii, 59–66.

581 Mensa, J., Z. Garraffo, A. Griffa, T. Özgökmen, A. Haza, and M. Veneziani, 2013: Seasonality
 582 of the submesoscale dynamics in the Gulf Stream region. *Ocean Dynamics*, **63(8)**, 923–941,
 583 doi:10.1007/s10236-013-0633-1.

584 Molemaker, M., J. McWilliams, and I. Yavneh, 2005: Baroclinic instability and loss of balance.
 585 *Journal of Physical Oceanography*, **35(9)**, 1505–1517, doi:10.1175/JPO2770.1.

586 Naveira Garabato, A., R. Ferrari, and K. Polzin, 2011: Eddy stirring in the Southern Ocean.
 587 *Journal of Geophysical Research: Oceans*, **116(C09019)**, doi:10.1029/2010JC006818.

588 Naveira Garabato, A., K. Polzin, B. King, K. Heywood, and M. Visbeck, 2004: Widespread
 589 intense turbulent mixing in the Southern Ocean. *Science*, **303(5655)**, 210–213, doi:10.1126/
 590 science.1090929.

591 Nikurashin, M., and R. Ferrari, 2010a: Radiation and dissipation of internal waves generated by
592 geostrophic motions impinging on small-scale topography: Application to the Southern Ocean.
593 *Journal of Physical Oceanography*, **40(9)**, 2025–2042, doi:10.1175/2010JPO4315.1.

594 Nikurashin, M., and R. Ferrari, 2010b: Radiation and dissipation of internal waves generated
595 by geostrophic motions impinging on small-scale topography: Theory. *Journal of Physical*
596 *Oceanography*, **40(5)**, 1055–1074, doi:10.1175/2009JPO4199.1.

597 Nikurashin, M., and R. Ferrari, 2011: Global energy conversion rate from geostrophic flows into
598 internal lee waves in the deep ocean. *Geophysical Research Letters*, **38(8)**, L08 610, doi:10.
599 1029/2011GL046576.

600 Omand, M., E. D’Asaro, C. Lee, M. Perry, N. Briggs, I. Cetinić, and A. Mahadevan, 2015:
601 Eddy-driven subduction exports particulate organic carbon from the spring bloom. *Science*,
602 **348(6231)**, 222–225, doi:10.1126/science.1260062.

603 Pollard, R., and L. Regier, 1990: Large variations in potential vorticity at small spatial scales in
604 the upper ocean. *Nature*, **348(6298)**, 227–229, doi:10.1038/348227a0.

605 Rocha, C., T. Chereskin, S. Gille, and D. Menemenlis, 2016: Mesoscale to submesoscale
606 wavenumber spectra in Drake Passage. *Journal of Physical Oceanography*, **46(2)**, 601–620,
607 doi:10.1175/JPO-D-15-0087.1.

608 Rosso, I., A. Hogg, A. Kiss, and B. Gayen, 2015: Topographic influence on submesoscale
609 dynamics in the Southern Ocean. *Geophysical Research Letters*, **42(4)**, 1139–1147, doi:
610 10.1002/2014GL062720.

611 Rosso, I., A. Hogg, R. Matear, and P. Strutton, 2016: Quantifying the influence of sub-mesoscale
612 dynamics on the supply of iron to Southern Ocean phytoplankton blooms. *Deep Sea Research*
613 *Part I: Oceanographic Research Papers*, **115**, 199–209, doi:10.1016/j.dsr.2016.06.009.

614 Rosso, I., A. Hogg, P. Strutton, A. Kiss, R. Matear, A. Klocker, and E. van Sebille, 2014: Vertical
615 transport in the ocean due to sub-mesoscale structures: Impacts in the Kerguelen region. *Ocean*
616 *Modelling*, **80**, 10–23, doi:10.1016/j.ocemod.2014.05.001.

617 Rudnick, D., 1996: Intensive surveys of the Azores Front: 2. Inferring the geostrophic and vertical
618 velocity fields. *Journal of Geophysical Research: Oceans*, **101(C7)**, 16 291–16 303, doi:10.
619 1029/96JC01144.

620 Sallée, J.-B., R. Matear, S. Rintoul, and A. Lenton, 2012: Localized subduction of anthropogenic
621 carbon dioxide in the Southern Hemisphere oceans. *Nature Geoscience*, **5 (8)**, 579–584, doi:
622 10.1038/ngeo1523.

623 Sallée, J.-B., K. Speer, and R. Morrow, 2008: Response of the Antarctic Circumpolar Current to
624 atmospheric variability. *Journal of Climate*, **21(12)**, 3020–3039, doi:10.1175/2007JCLI1702.1.

625 Sallée, J.-B., K. Speer, S. Rintoul, and S. Wijffels, 2010: Southern Ocean thermocline ventilation.
626 *Journal of Physical Oceanography*, **40(3)**, 509–529, doi:10.1175/2009JPO4291.1.

627 Sasaki, H., P. Klein, B. Qiu, and Y. Sasai, 2014: Impact of oceanic-scale interactions on the
628 seasonal modulation of ocean dynamics by the atmosphere. *Nature Communications*, **5**, 5636,
629 doi:10.1038/ncomms6636.

630 Shakespeare, C., 2015: On the generation of waves during frontogenesis. *Doctoral dissertation*,
631 University of Cambridge, Cambridge, UK.

Shakespeare, C., and A. Hogg, 2017: Spontaneous surface generation and interior amplification of internal waves in a regional-scale ocean model. *Journal of Physical Oceanography*, **47**(4), 811–826, doi:10.1175/JPO-D-16-0188.1.

Shakespeare, C., and J. Taylor, 2014: The spontaneous generation of inertia-gravity waves generated during frontogenesis forced by large strain: Theory. *Journal of Fluid Mechanics*, **757**, 817–853, doi:10.1017/jfm.2014.514.

Shakespeare, C., and J. Taylor, 2015: The spontaneous generation of inertia-gravity waves generated during frontogenesis forced by large strain: Numerical simulations. *Journal of Fluid Mechanics*, **772**, 508–534, doi:10.1017/jfm.2015.197.

Shakespeare, C., and J. Taylor, 2016: Spontaneous wave generation at strongly strained density fronts. *Journal of Physical Oceanography*, **46**(7), 2063–2081, doi:10.1175/JPO-D-15-0043.1.

Sokolov, S., and S. Rintoul, 2009: Circumpolar structure and distribution of the Antarctic Circumpolar Current fronts: 1. Mean circumpolar paths. *Journal of Geophysical Research: Oceans*, **114** (C11), doi:10.1029/2008JC005108.

Spall, S., and K. Richards, 2000: A numerical model of mesoscale frontal instabilities and plankton dynamics – I. Model formulation and initial experiments. *Deep Sea Research Part I: Oceanographic Research Papers*, **47**(7), 1261–1301, doi:10.1016/S0967-0637(99)00081-3.

St. Laurent, L., A. Naveira Garabato, J. Ledwell, A. Thurnherr, J. Toole, and A. Watson, 2012: Turbulence and diapycnal mixing in Drake Passage. *Journal of Physical Oceanography*, **42**(12), 2143–2152, doi:10.1175/JPO-D-12-027.1.

Stone, P., 1966: On non-geostrophic baroclinic stability. *Journal of the Atmospheric Sciences*, **23**(4), 390–400, doi:10.1175/1520-0469(1966)023<0390:ONGBS>2.0.CO;2.

654 Taylor, J. R., and R. Ferrari, 2009: On the equilibration of a symmetrically unstable front
655 via a secondary shear instability. *Journal of Fluid Mechanics*, **622**(1), 103–113, doi:10.1017/
656 S002211200800527.

657 Taylor, J. R., and R. Ferrari, 2010: Buoyancy and wind-driven convection at mixed layer density
658 fronts. *Journal of Physical Oceanography*, **40**(6), 1222–1242, doi:10.1175/2010JPO4365.1.

659 Thomas, L., 2005: Destruction of potential vorticity by winds. *Journal of Physical Oceanography*,
660 **35**(12), 2457–2466, doi:10.1175/JPO2830.1.

661 Thomas, L., and R. Ferrari, 2008: Friction, frontogenesis, and the stratification of the sur-
662 face mixed layer. *Journal of Physical Oceanography*, **38**(11), 2501–2518, doi:10.1175/
663 2008JPO3797.1.

664 Thomas, L. N., A. Tandon, and A. Mahadevan, 2008: Submesoscale processes and dynamics.
665 *Ocean modeling in an eddying regime*, 17–38, doi:10.1029/177GM04.

666 Thomas, L. N., and J. R. Taylor, 2010: Reduction of the usable wind-work on the general cir-
667 culation by forced symmetric instability. *Geophysical Research Letters*, **37**(18), L18 606, doi:
668 10.1029/2010GL044680.

669 Thompson, A., A. Lazar, C. Buckingham, A. Naveira Garabato, G. Damerell, and K. Hey-
670 wood, 2016: Open-ocean submesoscale motions: A full seasonal cycle of mixed layer in-
671 stabilities from gliders. *Journal of Physical Oceanography*, **46**(4), 1285–1307, doi:10.1175/
672 JPO-D-15-0170.1.

673 Watson, A., J. Ledwell, M. Messias, B. King, N. Mackay, M. Meredith, B. Mills, and A. Garabato,
674 2013: Rapid cross-density ocean mixing at mid-depths in the Drake Passage measured by tracer
675 release. *Nature*, **501**(7467), 408–411, doi:10.1038/nature12432.

- 676 Wunsch, C., and R. Ferrari, 2004: Vertical mixing, energy, and the general circulation of the
677 oceans. *Annual Review of Fluid Mechanics*, **36**, 281–314, doi:10.1146/annurev.fluid.36.050802.
678 122121.
- 679 Young, W., 1994: The subinertial mixed layer approximation. *Journal of Physical Oceanography*,
680 **24 (8)**, 1812–1826, doi:10.1175/1520-0485(1994)024<1812:TSMLA>2.0.CO;2.

681
682
683
684
685
686

LIST OF TABLES

Table 1.	Correlation coefficients between $\langle \overline{w_{rms}} \rangle$, $\langle \overline{\Delta H_{ML}} \rangle$, and each of $\langle \overline{Ro_b} \rangle$, $\langle \overline{\frac{1}{2} \mathbf{u}_m^2} \rangle$, $\langle \overline{S_m} \rangle$, and $\langle \overline{ \nabla_h b } \rangle$ from the $1/192^\circ$ simulation. Regions are indicated by font style - boldface font indicates values measured over the whole domain, standard font indicates values measured only in R1, and italic font indicates values measured only in R2.	33
-----------------	---	----

	$\langle \overline{Ro_b} \rangle$	$\langle \frac{1}{2} \overline{\mathbf{u}_m^2} \rangle$	$\langle \overline{S_m} \rangle$	$\langle \overline{ \nabla_h b } \rangle$
	0.05	0.80	0.73	-0.18
$\langle \overline{w_{rms}} \rangle$	-0.32	0.64	0.57	-0.52
	<i>-0.21</i>	<i>0.81</i>	<i>0.72</i>	<i>-0.58</i>
	0.64	-0.12	0.20	0.66
$\langle \overline{\Delta H_{ML}} \rangle$	0.70	-0.26	-0.08	0.73
	<i>0.77</i>	<i>-0.36</i>	<i>-0.35</i>	<i>0.59</i>

687 TABLE 1. Correlation coefficients between $\langle \overline{w_{rms}} \rangle$, $\langle \overline{\Delta H_{ML}} \rangle$, and each of $\langle \overline{Ro_b} \rangle$, $\langle \frac{1}{2} \overline{\mathbf{u}_m^2} \rangle$, $\langle \overline{S_m} \rangle$, and $\langle \overline{|\nabla_h b|} \rangle$
 688 from the $1/192^\circ$ simulation. Regions are indicated by font style - boldface font indicates values measured over
 689 the whole domain, standard font indicates values measured only in R1, and italic font indicates values measured
 690 only in R2.

LIST OF FIGURES

- Fig. 1.** Domains of the nested models. The models are nested offline, running at $1/12^\circ$, $1/24^\circ$, $1/48^\circ$, $1/96^\circ$, and $1/192^\circ$ resolution. The domain size is reduced for the three highest resolution simulations. Each model features a 2° sponge region (hatched lines) where the density and velocity fields are relaxed to the open boundary conditions. The meandering black lines are the time-mean -0.7 m and -0.25 m contours of sea surface height from the $1/12^\circ$ simulation, indicating the position of the Polar Front (southern line) and Subantarctic Front (northern line) according to the altimetry-based definitions of Sallée et al. (2008). 36
- Fig. 2.** a) Time-averaged mixed layer depth, $\overline{H_{ML}}$ from the $1/192^\circ$ model. b) Change in mixed layer depth between the $1/12^\circ$ and $1/192^\circ$ models, $\Delta\overline{H_{ML}}$. The mixed layer depth decreases in most places at higher resolution, reflected by positive values of $\Delta\overline{H_{ML}}$. The amplitude of the change varies with location. c) Base-10 logarithm of the time-averaged RMS vertical velocity, $\overline{w_{rms}}$, from the $1/192^\circ$ model, which is also vertically averaged over the top 100 m of the ocean. The RMS velocity increases at higher resolution via an increasingly vigorous submesoscale field. The white line indicates the 400-m isobath. The magenta boxes indicate regions which will be analysed in Figure 5. 37
- Fig. 3.** a) Snapshot of surface ζ from 30 June 2015, in the $1/192^\circ$ model. b) Base-10 logarithm of the time-averaged Rossby number, $\overline{Ro_b}$, as defined in Section b. Regions where the depth is shallower than 400 m have been grayed out. 38
- Fig. 4.** Base-10 logarithm of the surface a) mesoscale kinetic energy, $\frac{1}{2}\overline{|\mathbf{u}_M|^2}$, b) submesoscale kinetic energy, $\frac{1}{2}\overline{|\mathbf{u}_S|^2}$, and c) mesoscale strain rate, $\overline{S_M}$ 39
- Fig. 5.** Full-depth, spatially-averaged vertical profiles of the RMS vertical velocity from (a) R1 and (b) R2. Blue lines indicate the mesoscale component, $\overline{(w_{rms})_M}$, and red lines indicate the submesoscale component, $\overline{(w_{rms})_S}$. A histogram of the ocean depth in each region is presented by the gray bars. 40
- Fig. 6.** Scatter plots showing the trend of $\langle\overline{w_{rms}}\rangle$ (left column) and $\langle\Delta\overline{H_{ML}}\rangle$ (right column) with $\langle\overline{Ro_b}\rangle$ (top row), mesoscale KE (middle row), and mesoscale strain (bottom row). Blue, red, and gray dots indicate locations in R1, R2, and throughout the rest of the domain, respectively. Correlation coefficients appear in the bottom right corner of each panel. The black vertical lines are error bars, indicating one standard deviation above and below the mean. 41
- Fig. 7.** Maps of base-10 logarithm $|\overline{\nabla_h b}|$ from the a) $1/12^\circ$ and b) $1/192^\circ$ simulations, and scatter plots showing the trend of c) $\langle\overline{w_{rms}}\rangle$ and d) $\langle\Delta\overline{H_{ML}}\rangle$ against $\langle|\overline{\nabla_h b}|\rangle$ from the $1/192^\circ$ simulation. Blue, red, and gray dots indicate locations in R1, R2, and throughout the rest of the domain, respectively. The black vertical lines are error bars, indicating one standard deviation above and below the mean. 42
- Fig. 8.** a) Map of W , the vertical velocity scaling (2) for waves generated at fronts undergoing large strain, diagnosed using the filtered mesoscale output variables from the $1/192^\circ$ simulations, compared with b) the model vertical velocity, $\overline{w_{rms}}$. c) Scatter plot showing the trend of $\langle\overline{w_{rms}}\rangle$ against $\langle W \rangle$. The black dashed line is the 1:1 line indicating perfect agreement between W and $\overline{w_{rms}}$ 43

732 **Fig. 9.** Trend of $\langle \overline{w_{rms}} \rangle$ as a function of horizontal resolution, where the spatial averaging is taken
 733 vertically over the top 400 m and horizontally over a) R1, b) R2, and c) the whole domain.
 734 Black vertical bars indicate the range of $\overline{w_{rms}}$ values within each region. The numerical
 735 values are given in the tables on the right. Also shown in the tables are the values of s , the
 736 fractional increase in $\langle \overline{w_{rms}} \rangle$ due to doubling the resolution. 44

737 FIG. 1. Domains of the nested models. The models are nested offline, running at $1/12^\circ$, $1/24^\circ$, $1/48^\circ$, $1/96^\circ$,
 738 and $1/192^\circ$ resolution. The domain size is reduced for the three highest resolution simulations. Each model
 739 features a 2° sponge region (hatched lines) where the density and velocity fields are relaxed to the open boundary
 740 conditions. The meandering black lines are the time-mean -0.7 m and -0.25 m contours of sea surface height
 741 from the $1/12^\circ$ simulation, indicating the position of the Polar Front (southern line) and Subantarctic Front
 742 (northern line) according to the altimetry-based definitions of Sallée et al. (2008).

FIG. 2. a) Time-averaged mixed layer depth, $\overline{H_{ML}}$ from the $1/192^\circ$ model. b) Change in mixed layer depth between the $1/12^\circ$ and $1/192^\circ$ models, $\Delta \overline{H_{ML}}$. The mixed layer depth decreases in most places at higher resolution, reflected by positive values of $\Delta \overline{H_{ML}}$. The amplitude of the change varies with location. c) Base-10 logarithm of the time-averaged RMS vertical velocity, $\overline{w_{rms}}$, from the $1/192^\circ$ model, which is also vertically averaged over the top 100 m of the ocean. The RMS velocity increases at higher resolution via an increasingly vigorous submesoscale field. The white line indicates the 400-m isobath. The magenta boxes indicate regions which will be analysed in Figure 5.

FIG. 3. a) Snapshot of surface ζ from 30 June 2015, in the $1/192^\circ$ model. b) Base-10 logarithm of the time-averaged Rossby number, $\overline{Ro_b}$, as defined in Section b. Regions where the depth is shallower than 400 m have been grayed out.

FIG. 4. Base-10 logarithm of the surface a) mesoscale kinetic energy, $\frac{1}{2} \overline{|\mathbf{u}_M|^2}$, b) submesoscale kinetic energy, $\frac{1}{2} \overline{|\mathbf{u}_S|^2}$, and c) mesoscale strain rate, $\overline{S_M}$.

FIG. 5. Full-depth, spatially-averaged vertical profiles of the RMS vertical velocity from (a) R1 and (b) R2. Blue lines indicate the mesoscale component, $\overline{(w_{rms})}_M$, and red lines indicate the submesoscale component, $\overline{(w_{rms})}_S$. A histogram of the ocean depth in each region is presented by the gray bars.

FIG. 6. Scatter plots showing the trend of $\langle \overline{w_{rms}} \rangle$ (left column) and $\langle \overline{\Delta H_{ML}} \rangle$ (right column) with $\langle \overline{Ro_b} \rangle$ (top row), mesoscale KE (middle row), and mesoscale strain (bottom row). Blue, red, and gray dots indicate locations in R1, R2, and throughout the rest of the domain, respectively. Correlation coefficients appear in the bottom right corner of each panel. The black vertical lines are error bars, indicating one standard deviation above and below the mean.

FIG. 7. Maps of base-10 logarithm $|\nabla_h b|$ from the a) $1/12^\circ$ and b) $1/192^\circ$ simulations, and scatter plots showing the trend of c) $\langle w_{rms} \rangle$ and d) $\langle \Delta H_{ML} \rangle$ against $\langle |\nabla_h b| \rangle$ from the $1/192^\circ$ simulation. Blue, red, and gray dots indicate locations in R1, R2, and throughout the rest of the domain, respectively. The black vertical lines are error bars, indicating one standard deviation above and below the mean.

FIG. 8. a) Map of W , the vertical velocity scaling (2) for waves generated at fronts undergoing large strain, diagnosed using the filtered mesoscale output variables from the $1/192^\circ$ simulations, compared with b) the model vertical velocity, $\overline{w_{rms}}$. c) Scatter plot showing the trend of $\langle \overline{w_{rms}} \rangle$ against $\langle W \rangle$. The black dashed line is the 1:1 line indicating perfect agreement between W and $\overline{w_{rms}}$.

FIG. 9. Trend of $\langle \overline{w_{rms}} \rangle$ as a function of horizontal resolution, where the spatial averaging is taken vertically over the top 400 m and horizontally over a) R1, b) R2, and c) the whole domain. Black vertical bars indicate the range of $\overline{w_{rms}}$ values within each region. The numerical values are given in the tables on the right. Also shown in the tables are the values of s , the fractional increase in $\langle \overline{w_{rms}} \rangle$ due to doubling the resolution.