

2018-02

Plankton as prevailing conditions: A surveillance role for plankton indicators within the Marine Strategy Framework Directive

McQuatters-Gollop, A

<http://hdl.handle.net/10026.1/10670>

10.1016/j.marpol.2017.12.021

Marine Policy

Elsevier

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Plankton as prevailing conditions: a surveillance role for plankton indicators within the Marine Strategy Framework Directive

Authors and affiliations: Jacob Bedford^a, David Johns^b, Simon Greenstreet^c and Abigail McQuatters-Gollop^a

^aCentre for Marine Conservation and Policy Research, Plymouth University, Drake Circus, Plymouth, UK, PL4 8AA

^bSir Alister Hardy Foundation for Ocean Science, The Laboratory, Citadel Hill, Plymouth, UK, PL1 2PB

^cMarine Scotland – Science, Marine Laboratory, 375 Victoria Road, Aberdeen, UK, AB11 9DB

Corresponding author: Jacob Bedford, jacob.bedford@plymouth.ac.uk

Keywords: biodiversity; ecosystem approach; climate change; assessment

Abstract

The Marine Strategy Framework Directive (MSFD) uses an indicator-based approach for ecosystem assessment; indicators of the state of ecosystem components ('state indicators') are used to determine whether, or not, these ecosystem components are at 'Good Environmental Status' relative to prevailing oceanographic conditions. Here, it is illustrated that metrics of change in plankton communities frequently provide indications of changing prevailing oceanographic conditions. Plankton indicators can therefore provide useful diagnostic information when interpreting results and determining assessment outcomes for analyses of state indicators across the food web. They can also perform a strategic role in assessing these state indicators by influencing target setting and management measures. In addition to their primary role of assessing the state of pelagic habitats against direct anthropogenic pressures, plankton community indicators can therefore also fulfil an important 'surveillance' role for other state indicators used to formally assess biodiversity status under the MSFD.

1 Introduction

An ecosystem-based approach is increasingly adopted for the management of marine ecosystems [1, 2]. Whilst previous management strategies focused on key species and habitats, they neglected the interactions and linkages between ecosystem components, as well as between ecological and social systems [3, 4]. Ecosystem-based management on the other hand, considers humans as part of the ecosystem, and aims to manage the impact of multiple anthropogenic activities to achieve a healthy ecosystem state with a sustained flow of ecosystem services to humans [4, 5]. The EU Marine Strategy Framework Directive (MSFD) takes an ecosystem approach to the management of European seas, supported by Integrated Ecosystem Assessments, where indicators are required to synthesize scientific information and formally assess progress towards the overall ecosystem objective of 'Good Environmental Status' (GES) [6, 7]. Out of the 11 qualitative descriptors that comprise the MSFD [8], the descriptors, 'Biodiversity', 'Food webs' and 'Sea Floor Integrity', describe ecosystem state. [9]

As a directive concerning direct, manageable anthropogenic pressures on the marine environment, the development of MSFD biodiversity state indicators for formal assessment initially focused on indicators with clear pressure-state relationships and associations with defined thresholds and targets. An example is a fish stock size controlled by levels of fishing pressure [10, 11]. These state indicators can follow an 'Activity'-'Pressure'-'State'-'Response' (APSR) framework of marine management, where a human activity applies a defined pressure on the system. This pressure causes a change in the state of the indicator, which can trigger a management response [12]. However, Shephard, Greenstreet, Piet, Rindorf and Dickey-Collas [12] argue that a separate class of indicators called 'surveillance indicators', where the links to defined anthropogenic pressures are not well understood and where target setting is difficult, can also contribute to ecosystem assessments under the MSFD. Surveillance indicators do not have a direct influence on the formal assessment of Good Environmental Status, but their 'surveillance' can provide information on wider ecosystem impacts of anthropogenic pressures as well as changing environmental conditions. Therefore, surveillance indicators can also result in triggering management action when pre-defined bounds are passed.

Indicators that describe the structure and functioning of plankton communities have been developed to formally assess the state of 'pelagic habitats' within the MSFD. These include indicators of bulk properties such as primary production as well as indicators of change in plankton functional groups [13]. Plankton indicator change may be

48 driven by a multitude of direct anthropogenic pressures, most notably eutrophication resulting from anthropogenic
49 nutrient pollution [14]. The assessment of these MSFD plankton indicators, therefore, can directly contribute to the
50 design of the programme of management measures needed to ensure marine ecosystems are in Good
51 Environmental Status under the MSFD, should a change in the plankton indicators be detected during assessment,
52 and linked to direct anthropogenic pressure.

53 Plankton dynamics, however, are largely driven by climate [15], particularly at the regional scale which is the focus of
54 the MSFD. Consequently, both climate variability and anthropogenic climate change can cause widespread changes
55 in the plankton [16] which are likely to manifest through changes in plankton indicators. The MSFD [8] refers to
56 these drivers of change as 'prevailing conditions' and mandates that "*the quality and occurrence of habitats and the*
57 *distribution and abundance of species are in line with prevailing physiographic, geographic and climatic conditions*".
58 Changes in the plankton driven by climate change and environmental variability, therefore, would be considered in
59 line with Good Environmental Status, with no management impetus through the MSFD.

60 Because plankton are sensitive to changes in climatic and physical oceanographic conditions however, and have
61 been shown to amplify weak climatic signals [17], they can be useful indicators for large scale changes in prevailing
62 conditions. For example, indicators of variability in volume of Atlantic inflow into the North Sea, a key forcing
63 variable for the North Sea ecosystem, can be derived from zooplankton communities [18]. Furthermore, due to the
64 key role of phytoplankton as primary producers in the marine food web, and the key role of zooplankton as prey for
65 higher trophic levels such as fish, climate-induced changes in plankton themselves may be considered as prevailing
66 conditions for other biodiversity components [19]. In this way, in addition to their use in directly assessing for Good
67 Environmental Status, plankton indicators can also be considered surveillance indicators, reflecting change in
68 prevailing conditions that can aid in the interpretation of formal biodiversity indicator assessments. Plankton
69 indicators can therefore have an additional 'surveillance role' even when the plankton indicator changes are not
70 linked to direct anthropogenic pressures.


71 The surveillance role of plankton indicators is not limited to the formally assessed MSFD plankton indicators
72 however, and can extend to the wider climate change trends identified from time-series datasets that aren't
73 formally assessed within the MSFD. For example a trend for the replacement of *Calanus finmarchicus* by its

74 congeneric warmer-water species *Calanus helgolandicus* was identified in the North Atlantic and is an indicator of
75 climate change [20]. Similarly, changes in the phenology of phytoplankton bloom dynamics, linked to the efficiency
76 of energy transfer from phytoplankton to higher trophic levels, have been identified and attributed to climate
77 change [21]. These trends are not formally assessed within the MSFD, but are derived from the same time-series
78 datasets as the assessed MSFD plankton indicators, providing useful supplementary information with no additional
79 monitoring effort.

80 Here, the surveillance indicator framework presented by Shephard, Greenstreet, Piet, Rindorf and Dickey-Collas [12]
81 is used to demonstrate the utility of plankton indicators in the surveillance role of informing on changing prevailing
82 conditions. This framework illustrates how surveillance indicators can add contextual information to formal state
83 indicator assessments within the MSFD, aiding in assessment interpretation. Specifically, here the contextual
84 information gained from the surveillance of plankton indicators is classified as 'diagnostic', which helps diagnose the
85 drivers of changes within the ecosystem, and 'strategic' which aids in setting targets and management measures for
86 Good Environmental Status.

87 **1.1 The surveillance indicator framework**

88 The surveillance indicator framework described by Shephard et al. (2015) provides a conceptual tool for integrating
89 changes in prevailing conditions into the formal biodiversity indicator assessment process. Due to their lack of clear
90 pressure-state relationships, surveillance indicators cannot follow directly an Activity-Pressure-State-Response
91 framework. Therefore, Shephard et al. modified the traditional APSR framework to include surveillance indicators
92 (Figure 1). A key feature of their surveillance indicator framework is that there are no GES targets for surveillance
93 indicators. Instead, when a surveillance indicator moves outside of a defined bound, new research is triggered as the
94 potential implication of this indicator change may not be clear. This research focuses on addressing whether the
95 change in surveillance indicators means that the targets and management measures for associated assessed
96 indicators need to be re-evaluated. Precautionary management may be implemented as a result of surveillance
97 indicator change, in respect to the management responses to changes in associated formally assessed indicators.



98 Figure 1. The 'surveillance indicator' framework used here. Diagram adapted from Shephard, Greenstreet, Piet,
 99 Rindorf and Dickey-Collas [12]. Assessed indicator (blue) change is detected. If indicator moves to being not in GES
 100 (NGES), a management measure is considered, based on the pressure-state relationship of the assessed indicator
 101 with a direct pressure. Surveillance indicators (red, bottom) are monitored simultaneously to the assessed indicator,
 102 and surveillance indicator change is detected when the surveillance indicator moves out of predefined bounds (not
 103 within bounds: NWB). This surveillance indicator change triggers research targeted at the pressure-state
 104 relationships and GES targets of associated assessed indicators (blue, top).

105 When applying plankton to this surveillance indicator framework, time-series data can be used for setting
 106 surveillance bounds [12, 22], for example based on past ranges of indicator values, or using past variability to
 107 categorize different magnitudes of change. This is because long term time-series aid in contextualising any indicator
 108 changes identified, in terms of the indicated changes in prevailing conditions. An example is the use of time-series
 109 data in the detection of regime shifts, such as the 1980s climate-driven regime shift detected in Continuous Plankton
 110 Recorder survey data that caused widespread changes in both phytoplankton and zooplankton
 111 communities, coinciding with changes across the wider food web [23-25]. Time series data can also aid in identifying
 112 whether observed changes are the continuation of longer term trends by identifying any existing trajectories of
 113 indicator change [26].

114 Often, however, the strength of coupling between hydro-climatic variation, plankton, and other food web
115 components may not be clear and instead obscured by natural variability. Thus, covariation between a plankton
116 indicator and assessed indicators at higher trophic levels would not be sufficient to trigger precautionary
117 management alone within the framework. Furthermore, the use of correlations to derive links between
118 environmental variation and higher trophic levels has been criticised [27]. Instead, within the framework, any
119 covariation identified would highlight questions that could be considered when interpreting the results of formal
120 state indicator assessments, often requiring further research and analysis. Examples of how information on
121 prevailing conditions gained through plankton surveillance provides evidence for the interpretation of formal
122 biodiversity indicator assessments are given below.

123 **2 Diagnostic role in identifying drivers of change in formally assessed biodiversity indicators**

124 A key challenge in assessing any biodiversity state indicator within the Marine Strategy Framework Directive is in the
125 attribution of observed indicator changes to either direct anthropogenic pressure or prevailing conditions [28], thus
126 'diagnosing' the cause of indicator change (Figure 2) [29]. Within pelagic habitats, it is established that an
127 understanding of climate-driven plankton trends is essential for disentangling any effect of direct pressures from
128 wider climatic influences [30]. For example, an indicator for phytoplankton community structure using functional
129 groups is formally assessed at the OSPAR level [31]. This indicator may reveal changes in phytoplankton community
130 structure as a result of human pressures, such as, for example, the effects of anthropogenic nutrient loading altering
131 the proportions of dinoflagellates and diatoms within phytoplankton communities [14]. Phytoplankton community
132 structure, however, is also influenced by climate. For example, the CPR survey reveals multi-decadal range changes
133 in multiple phytoplankton taxa in response to climate change. These responses to climate are not uniform across
134 taxa, with some taxa tracking northward movements of thermoclines closer than others, causing restructuring of
135 phytoplankton communities [32]. Understanding the climate contribution to changes in plankton communities,
136 therefore, helps diagnose the drivers of change in the assessed MSFD plankton indicators (Fig 3A).

137 As well as performing this diagnostic role in the interpretation of formally assessed pelagic habitat indicators
138 however, plankton surveillance information can also be useful for interpreting changes in assessed indicators within
139 other habitats and trophic levels. Similarly to plankton, MSFD indicators from these other components may be

driven by both direct anthropogenic pressures as well as changes in prevailing conditions, requiring a degree of attribution of the different drivers when interpreting indicator change. Plankton indicator surveillance could inform on changes in prevailing conditions affecting these assessed indicators, and therefore help diagnose when changes are not driven by direct anthropogenic pressures alone. For example, under the MSFD, benthic habitat condition is assessed at the OSPAR level for the 'Biodiversity' and 'Seafloor integrity' descriptors [33]. Multi-metric indices are used to compare the condition of benthic habitat communities over intensity gradients of different anthropogenic pressures, resulting from a range of human activities including bottom-trawling and sediment extraction allowing for the determination of the degree to which the pressures causes change in benthic condition [33].

Benthic communities, however, are also impacted by large scale climate variability, and regime shifts detected in plankton communities have coincided with changes in the benthos [34]. Changes in the abundance of the larval stages of different benthic invertebrate groups (meroplankton) in relation to climate have also been detected from plankton time-series surveys [35]. Furthermore, particularly in coastal regions, there is often tight benthic-pelagic coupling as phytoplankton production is the main source of organic supply to benthic faunal communities [36]. Phytoplankton bloom dynamics may therefore control benthic community structure by influencing food availability and levels of environmental hypoxia [37]. Clare, Spencer, Robinson and Frid [38] showed that abrupt shifts in the temporal trends of large and long-lived taxa within a benthic community time-series were attributed to increased detrital input from pelagic primary production. Increasing frequency of Harmful Algal Bloom events as a result of climate change [39, 40] may also influence benthic communities through selectively impacting both larval and post-larval survival of benthic invertebrates [41]. As the MSFD benthic condition assessment is based on quantifying pressure state relationships, changes in benthic state indicators influenced by changes in prevailing conditions may result in the influence of direct pressures being misinterpreted [42]. The surveillance of plankton indicators including bulk primary productivity and HAB dynamics (Fig 3B), can therefore aid in the interpretation of the assessment of benthic habitat condition.




Figure 2. Schematic diagram of the diagnostic role for plankton surveillance information. Change in the state of a formally assessed biodiversity state indicator can be influenced by both direct anthropogenic pressures and prevailing conditions. Plankton surveillance can aid in understanding the relative influence of prevailing conditions.




Figure 3. Examples of the diagnostic role of plankton surveillance information in MSFD implementation using the surveillance indicator framework from Shephard, Greenstreet, Piet, Rindorf and Dickey-Collas [12]. A) The role of plankton surveillance information in diagnosing drivers of change in pelagic habitat MSFD indicators. Here, range

166 *shift indicators (bottom, red) trigger research targeted at the pressure-state relationship between phytoplankton GES*
 167 *indicator and eutrophication pressure (top, blue)- 'Is change in plankton GES indicator driven by climate induced*
 168 *range shifts?' B) The potential role of plankton surveillance information in diagnosing the drivers of change in*
 169 *assessed indicators within other habitats and ecosystem components. Here, surveillance of phytoplankton indicators*
 170 *(red), trigger research targeted at the benthic pressure-state relationship, and therefore assessment of GES, between*
 171 *benthic community composition and anthropogenic benthic disturbance (blue)- 'Is change in benthic condition*
 172 *indicator influenced by climate?'*

173 **3 Strategic role in influencing targets and management measures for formally assessed biodiversity** 174 **indicators**

175 In addition to diagnosing the drivers of change in MSFD biodiversity indicators during formal assessments, plankton
 176 surveillance information could contribute to the determination of GES targets (Figure 4). For example, an indicator
 177 for recovery in the population abundance of sensitive fish species has been developed for formal assessment at the
 178 OSPAR level [43]. However, the influence of changing prevailing oceanographic conditions on population growth is
 179 required to determine the scope for population recovery [43]. Changes in plankton indicators can track trends in
 180 physical oceanographic conditions that may affect recovery, and changes in plankton community composition and
 181 phenology may affect fish recruitment independently of the size of the spawning stock biomass [44]. For example,
 182 directly after the 1980s plankton regime shift North Sea cod populations fell to historically low levels and showed
 183 weak signs of recovery [45]. Furthermore, a regime shift that occurred in the North Sea in the early 2000s was
 184 suggested as the leading candidate mechanism to explain the low herring recruitment observed between 2002 and
 185 2007, with plankton shifts providing more explanatory power than the effects of physical variables alone [46].
 186 Although the linking of fish recruitment dynamics to environmental variability is challenging [47], surveillance of
 187 plankton indicators provides information on any influence of plankton on fish recovery potential [48].

188 The method for assessing GES in respect to fish population recovery is outlined by [49]. First, targets for a given
 189 indicator are set at the individual species level, based on the indicator metric falling in the upper 25 percentile of all
 190 values in the species' reference period. These species-level indicator assessments are then aggregated to the

191 community level by comparing the number of different species achieving their target for the given indicator.
192 Therefore, changes in prevailing conditions that affect the recovery potential of stocks, despite a reduction in
193 anthropogenic pressure, may mean the GES targets may no longer be realistic. Instead, the permitted range in which
194 individual species metrics can fall may need to be increased, or the number of species required to be in GES at the
195 community level may need to be reduced [50]. In this way, plankton indicator surveillance can contribute to the
196 setting of realistic targets for the assessment of fish state indicators [51] (Figure 5A).

197 As well as affecting the feasibility of reaching a specified state target, changes in prevailing conditions detected
198 through plankton surveillance may alter the sensitivity of an ecosystem component to a specified anthropogenic
199 pressure, thus affecting the amount of pressure that will cause an assessed biodiversity indicator to move away from
200 Good Environmental Status. (Figure 4) For example, indicators of seabird population size and breeding success are
201 formally assessed at the OSPAR level within the MSFD [52, 53] and are useful indicators of the food web
202 repercussions of direct pressures targeted at the lower levels of the food web, such as fishing pressure on forage fish
203 prey [54, 55]. For effective ecosystem-based management, management of forage fish exploitation must account for
204 the need to sustain top predators and as forage fish biomass and productivity is highly variable, the setting of
205 acceptable fishing levels must remain adaptive [56, 57]. With a reduction in the recruitment success of key forage
206 fish species such as sandeel predicted under climate change [58], reducing fishing pressure on forage fish through
207 precautionary management measures may be needed to maintain Good Environmental Status of seabirds under
208 future climate conditions [59].

209 Forage fish abundance and growth has been linked to phytoplankton production [60] and zooplankton community
210 composition through changes in the distribution of copepods indicating both changes in physical oceanographic
211 conditions and influencing recruitment and growth [61 {Clausen, 2017 #333}]. There can also be direct trophic links
212 between zooplankton and seabirds, especially in the non-breeding season [62, 63]. In these ways, climate-driven
213 plankton shifts may place an additional 'unmanageable' pressure on seabirds, influencing the outcome of seabird
214 state indicator assessments, and could therefore indicate relevant prevailing conditions when setting management
215 measures (Fig. 5B). Within MSFD assessment cycles, management of direct pressures could be altered to take into
216 account trends in climatic (non-manageable) drivers [64], informed by plankton surveillance. In this way, although
217 the drivers of climate induced changes cannot be addressed by the MSFD, adaptive management of direct pressures

218 could increase the likelihood of an indicator remaining in Good Environmental Status in relation to manageable
219 pressures, as well as help increase the resilience of the ecosystem component to climate change [65-67].




Figure 4. Schematic diagram of the 'strategic' role for plankton surveillance information. Targets, and associated management measures for a formally assessed state indicator can be adapted to changing prevailing conditions. Plankton surveillance information can inform appropriate target setting and management measures.




Figure 5. Examples of the strategic role of plankton surveillance information in MSFD implementation using the surveillance indicator framework from Shephard, Greenstreet, Piet, Rindorf and Dickey-Collas [12]. A) The potential role of plankton surveillance information in setting targets for other components and descriptors. Here, plankton indicator surveillance (red) triggers research around the target representing GES for the recovery of sensitive fish communities (blue)- 'Is the current GES target still achievable under the new climate conditions?'. This research could lead to the adjustment of GES state targets. B) The potential role of plankton surveillance information in influencing the programme of measures. Here, plankton indicator changes linked to prevailing conditions (bottom, red) trigger research targeted at the pressure-state relationship between forage fish fisheries and seabird breeding success (top, blue)- 'Is the current threshold level of fishing still sustainable, considering the changed prey landscape?' This research could lead to more precautionary management measures being implemented.

4 Summary and conclusions

231 In this paper, we have illustrated a surveillance role of plankton indicators within the Marine Strategy Framework
232 Directive in addition to their primary role in formally assessing pelagic habitats for influences of direct anthropogenic
233 pressures. Plankton indicators are useful early-warning indicators of physical hydro-climatic changes and can
234 therefore inform on changes in the underlying prevailing conditions in which MSFD biodiversity indicators are
235 formally assessed. Furthermore, changes in plankton can be important prevailing conditions to consider themselves.
236 The importance of including biotic ecosystem drivers, such as changes in plankton, within marine monitoring
237 programmes has been acknowledged by the Framework for Ocean Observing (FOO) with the development of
238 'ecosystem Essential Ocean Variables (eEOVs)', which are defined biological or ecological quantities derived from
239 field observations [68]. The surveillance indicator framework presented by Shephard, Greenstreet, Piet, Rindorf and
240 Dickey-Collas [12], is a useful tool in translating this established monitoring need into the MSFD implementation
241 process.

242 This surveillance of plankton indicators provides two, newly-defined, types of contextual information for the
243 assessment of biodiversity within the MSFD. 'Diagnostic' plankton surveillance information can help disentangle the
244 influence of direct anthropogenic pressure from the influence of prevailing conditions, both within pelagic habitats,
245 and within other habitats and ecosystem components. On the other hand, plankton surveillance information can
246 have a 'strategic' role by indicating when the climate influence on the ecosystem may mean targets and
247 management measures need to be altered. Due to the highly variable nature of coupling between changes in the
248 plankton and changes in the wider marine ecosystem, both diagnostic and strategic roles of plankton surveillance
249 information are based on the triggering of targeted research questions for consideration during assessments,
250 following the observation of a change in plankton indicators and the detection of trends, thereby making an
251 important evidence contribution to allow the implementation of the MSFD to be adaptive under climate change [69].

252 Currently, changes in plankton communities linked to climate are considered as being aligned with Good
253 Environmental Status, as the changes are linked to natural variations or exogenous pressures. Limiting the
254 application of these climate-driven indicator changes in this way however, is not using monitoring effort efficiently,
255 when plankton indicators are also useful in a wider surveillance role. Progressing this surveillance role for plankton
256 indicators requires further work on understanding ecosystem interactions between plankton and other formally
257 assessed biodiversity components, as well as the consequences of changes in climatic and oceanographic conditions

258 on both plankton indicators and the wider foodweb. This in turn requires further collaboration between scientists
259 working on these different components. Ultimately, the maintenance of long-term plankton time series therefore
260 has multiple applications for ecosystem-based management of European seas within the Marine Strategy
261 Framework Directive.

264 **Acknowledgements:**

265 Thanks go to Mark Dickey-Collas for valuable insight and advice in the development of the manuscript.

266 **Funding sources:**

267 A.M-G was supported by the UK Natural Environmental Research Council Knowledge Exchange fellowship scheme
268 (NE/L002663/1).

269 **References**

- 270
271
- 272 [1] S. Jennings, J. Rice, Towards an ecosystem approach to fisheries in Europe: a perspective on existing
273 progress and future directions, *Fish and Fisheries* 12(2) (2011) 125-137.
- 274 [2] S.E. Apitz, M. Elliott, M. Fountain, T.S. Galloway, European environmental management: moving to an
275 ecosystem approach, *Integrated environmental assessment and management* 2(1) (2006) 80-85.
- 276 [3] P.S. Levin, C. Mollmann, Marine ecosystem regime shifts: challenges and opportunities for ecosystem-
277 based management, *Philosophical Transactions of the Royal Society B: Biological Sciences* 370(1659)
278 (2014) 20130275-20130275.
- 279 [4] P. Tett, R.J. Gowen, S.J. Painting, M. Elliott, R. Forster, D.K. Mills, E. Bresnan, E. Capuzzo, T.F.
280 Fernandes, J. Foden, R.J. Geider, L.C. Gilpin, M. Huxham, A.L. McQuatters-Gollop, S.J. Malcolm, S. Saux-
281 Picart, T. Platt, M.F. Racault, S. Sathyendranath, J. van der Molen, M. Wilkinson, Framework for
282 understanding marine ecosystem health, *Marine Ecology Progress Series* 494 (2013) 1-27.

- 283 [5] S. Broszeit, N.J. Beaumont, M.C. Uyarra, A.-S. Heiskanen, M. Frost, P.J. Somerfield, A.G. Rossberg, H.
 284 Teixeira, M.C. Austen, What can indicators of good environmental status tell us about ecosystem
 285 services?: Reducing efforts and increasing cost-effectiveness by reapplying biodiversity indicator data,
 286 *Ecological Indicators* 81 (2017) 409-442.
- 287 [6] P.S. Levin, M.J. Fogarty, S.A. Murawski, D. Fluharty, Integrated ecosystem assessments: developing
 288 the scientific basis for ecosystem-based management of the ocean, *PLoS biology* 7(1) (2009) e1000014.
- 289 [7] Y.M. Walther, C. Möllmann, Bringing integrated ecosystem assessments to real life: a scientific
 290 framework for ICES, *ICES Journal of Marine Science* 71(5) (2013) 1183-1186.
- 291 [8] European Commission, Marine Strategy Framework Directive 2008/56/EC, European Commission
 292 (2008).
- 293 [9] A. Borja, M. Elliott, J.H. Andersen, A.C. Cardoso, J. Carstensen, J.G. Ferreira, A.-S. Heiskanen, J.C.
 294 Marques, J.M. Neto, H. Teixeira, Good Environmental Status of marine ecosystems: What is it and how do
 295 we know when we have attained it?, *Marine Pollution Bulletin* 76(1) (2013) 16-27.
- 296 [10] J.C. Rice, M.-J. Rochet, A framework for selecting a suite of indicators for fisheries management, *ICES*
 297 *Journal of Marine Science: Journal du Conseil* 62(3) (2005) 516-527.
- 298 [11] M.-J. Rochet, V.M. Trenkel, Which community indicators can measure the impact of fishing? A review
 299 and proposals, *Canadian Journal of Fisheries and Aquatic Sciences* 60(1) (2003) 86-99.
- 300 [12] S. Shephard, S.P.R. Greenstreet, G.J. Piet, A. Rindorf, M. Dickey-Collas, Surveillance indicators and
 301 their use in implementation of the Marine Strategy Framework Directive, *ICES Journal of Marine Science:*
 302 *Journal du Conseil* 72(8) (2015) 2269-2277.
- 303 [13] A. McQuatters-Gollop, D.G. Johns, E. Bresnan, J. Skinner, I. Rombouts, R. Stern, A. Aubert, M.
 304 Johansen, J. Bedford, A. Knights, From microscope to management: The critical value of plankton
 305 taxonomy to marine policy and biodiversity conservation, *Marine Policy* 83 (2017) 1-10.
- 306 [14] A. McQuatters-Gollop, A.J. Gilbert, L.D. Mee, J.E. Vermaat, Y. Artioli, C. Humborg, F. Wulff, How well
 307 do ecosystem indicators communicate the effects of anthropogenic eutrophication?, *Estuarine, Coastal and*
 308 *Shelf Science* 82(4) (2009) 583-596.
- 309 [15] L. Zhai, T. Platt, C. Tang, S. Sathyendranath, A. Walne, The response of phytoplankton to climate
 310 variability associated with the North Atlantic Oscillation, *Deep Sea Research Part II: Topical Studies in*
 311 *Oceanography* 93 (2013) 159-168.
- 312 [16] G.C. Hays, A.J. Richardson, C. Robinson, Climate change and marine plankton, *Trends in ecology &*
 313 *evolution* 20(6) (2005) 337-344.
- 314 [17] A.H. Taylor, J.I. Allen, P.A. Clark, Extraction of a weak climatic signal by an ecosystem, *Nature*
 315 416(6881) (2002) 629-632.
- 316 [18] P.C. Reid, M. Edwards, G. Beaugrand, M. Skogen, D. Stevens, Periodic changes in the zooplankton of
 317 the North Sea during the twentieth century linked to oceanic inflow, *Fisheries Oceanography* 12(4-5) (2003)
 318 260-269.
- 319 [19] K.F. Drinkwater, G. Beaugrand, M. Kaeriyama, S. Kim, G. Ottersen, R.I. Perry, H.-O. Pörtner, J.J.
 320 Polovina, A. Takasuka, On the processes linking climate to ecosystem changes, *Journal of Marine Systems*
 321 79(3) (2010) 374-388.
- 322 [20] P. Helaouët, G. Beaugrand, Macroecology of *Calanus finmarchicus* and *C. helgolandicus* in the North
 323 Atlantic Ocean and adjacent seas, *Marine Ecology Progress Series* 345 (2007) 147-165.
- 324 [21] M. Edwards, A.J. Richardson, Impact of climate change on marine pelagic phenology and trophic
 325 mismatch, *Nature* 430(7002) (2004) 881-884.
- 326 [22] W.N. Probst, V. Stelzenmüller, A benchmarking and assessment framework to operationalise
 327 ecological indicators based on time series analysis, *Ecological Indicators* 55 (2015) 94-106.
- 328 [23] M. Lindegren, V. Dakos, J.P. Groeger, A. Gårdmark, G. Kornilovs, S.A. Otto, C. Möllmann, Early
 329 detection of ecosystem regime shifts: a multiple method evaluation for management application, *PLoS One*
 330 7(7) (2012) e38410.
- 331 [24] J.P. Gröger, M. Missong, R.A. Rountree, Analyses of interventions and structural breaks in marine and
 332 fisheries time series: Detection of shifts using iterative methods, *Ecological indicators* 11(5) (2011) 1084-
 333 1092.
- 334 [25] P.C. Reid, R.E. Hari, G. Beaugrand, D.M. Livingstone, C. Marty, D. Straile, J. Barichivich, E.
 335 Goberville, R. Adrian, Y. Aono, R. Brown, J. Foster, P. Groisman, P. Helaouët, H.-H. Hsu, R. Kirby, J.
 336 Knight, A. Kraberg, J. Li, T.-T. Lo, R.B. Myneni, R.P. North, J.A. Pounds, T. Sparks, R. Stübi, Y. Tian, K.H.
 337 Wiltshire, D. Xiao, Z. Zhu, Global impacts of the 1980s regime shift, *Global Change Biology* (2015) n/a-n/a.
- 338 [26] E. Gorokhova, M. Lehtiniemi, L. Postel, G. Rubene, C. Amid, J. Lesutiene, L. Uusitalo, S. Strake, N.
 339 Demereckiene, Indicator properties of Baltic Zooplankton for classification of environmental status within
 340 Marine Strategy Framework Directive, *PloS one* 11(7) (2016) e0158326.

- 341 [27] M. Dickey-Collas, M.R. Payne, V.M. Trenkel, R.D. Nash, Hazard warning: model misuse ahead, *ICES*
 342 *Journal of Marine Science: Journal du Conseil* 71(8) (2014) 2300-2306.
- 343 [28] A. McQuatters-Gollop, Challenges for implementing the Marine Strategy Framework Directive in a
 344 climate of macroecological change, *Philos Trans A Math Phys Eng Sci* 370(1980) (2012) 5636-55.
- 345 [29] N.J. Hardman-Mountford, J.I. Allen, M.T. Frost, S.J. Hawkins, M.A. Kendall, N. Mieszowska, K.A.
 346 Richardson, P.J. Somerfield, Diagnostic monitoring of a changing environment: An alternative UK
 347 perspective, *Marine Pollution Bulletin* 50(12) (2005) 1463-1471.
- 348 [30] A. McQuatters-Gollop, M. Edwards, P. Helaouët, D.G. Johns, N.J.P. Owens, D.E. Raitsos, D.
 349 Schroeder, J. Skinner, R.F. Stern, The Continuous Plankton Recorder survey: How can long-term
 350 phytoplankton datasets contribute to the assessment of Good Environmental Status?, *Estuarine, Coastal*
 351 *and Shelf Science* 162 (2015) 88-97.
- 352 [31] OSPAR, Changes in Phytoplankton and Zooplankton Communities., Intermediate Assessment 2017.
 353 Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017> (2017).
- 354 [32] W.J. Chivers, A.W. Walne, G.C. Hays, Mismatch between marine plankton range movements and the
 355 velocity of climate change, *Nature Communications* 8 (2017) 14434.
- 356 [33] OSPAR, Condition of Benthic Habitat Defining Communities Intermediate Assessment 2017. Available
 357 at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017>(2017).
- 358 [34] S.N. Birchenough, H. Reiss, S. Degraer, N. Mieszowska, Á. Borja, L. Buhl - Mortensen, U.
 359 Braeckman, J. Craeymeersch, I. De Mesel, F. Kerckhof, Climate change and marine benthos: a review of
 360 existing research and future directions in the North Atlantic, *Wiley Interdisciplinary Reviews: Climate*
 361 *Change* 6(2) (2015) 203-223.
- 362 [35] R.R. Kirby, G. Beaugrand, J.A. Lindley, Climate-induced effects on the meroplankton and the benthic-
 363 pelagic ecology of the North Sea, *Limnology and Oceanography* 53(5) (2008) 1805.
- 364 [36] J.R. Griffiths, M. Kadin, F.J. Nascimento, T. Tanelander, A. Törnroos, S. Bonaglia, E. Bonsdorff, V.
 365 Brüchert, A. Gårdmark, M. Järnström, The importance of benthic–pelagic coupling for marine ecosystem
 366 functioning in a changing world, *Global Change Biology* 23(6) (2017) 2179-2196.
- 367 [37] Q. Zhang, R.M. Warwick, C.L. McNeill, C.E. Widdicombe, A. Sheehan, S. Widdicombe, An unusually
 368 large phytoplankton spring bloom drives rapid changes in benthic diversity and ecosystem function,
 369 *Progress in Oceanography* 137 (2015) 533-545.
- 370 [38] D.S. Clare, M. Spencer, L.A. Robinson, C.L. Frid, Explaining ecological shifts: the roles of temperature
 371 and primary production in the long - term dynamics of benthic faunal composition, *Oikos* (2017).
- 372 [39] M. Edwards, D. Johns, S. Leterme, E. Svendsen, A. Richardson, Regional climate change and harmful
 373 algal blooms in the northeast Atlantic, *Limnology and oceanography* 51(2) (2006) 820-829.
- 374 [40] G.M. Hallegraeff, Ocean climate change, phytoplankton community responses, and harmful algal
 375 blooms: a formidable predictive challenge¹, *Journal of phycology* 46(2) (2010) 220-235.
- 376 [41] J. Grall, L. Chauvaud, Marine eutrophication and benthos: the need for new approaches and concepts,
 377 *Global Change Biology* 8(9) (2002) 813-830.
- 378 [42] I. Kroncke, H. Reiss, Influence of macrofauna long-term natural variability on benthic indices used in
 379 ecological quality assessment, *Marine Pollution Bulletin* 60(1) (2010) 58-68.
- 380 [43] OSPAR, Recovery in the Population Abundance of Sensitive Fish Species, Intermediate Assessment
 381 2017. Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017>
 382 (2017).
- 383 [44] S. Lowerre - Barbieri, G. DeCelles, P. Pepin, I.A. Catalán, B. Muhling, B. Erisman, S.X. Cadrin, J. Alós,
 384 A. Ospina - Alvarez, M.M. Stachura, Reproductive resilience: a paradigm shift in understanding spawner -
 385 recruit systems in exploited marine fish, *Fish and Fisheries* (2016).
- 386 [45] M. Lindegren, R. Diekmann, C. Möllmann, Regime shifts, resilience and recovery of a cod stock,
 387 *Marine Ecology Progress Series* 402 (2010) 239-253.
- 388 [46] M.R. Payne, E.M. Hatfield, M. Dickey-Collas, T. Falkenhaus, A. Gallego, J. Gröger, P. Licandro, M.
 389 Llope, P. Munk, C. Röckmann, Recruitment in a changing environment: the 2000s North Sea herring
 390 recruitment failure, *ICES Journal of Marine Science: Journal du Conseil* 66(2) (2009) 272-277.
- 391 [47] P. Pepin, Reconsidering the impossible—linking environmental drivers to growth, mortality, and
 392 recruitment of fish 1, *Canadian Journal of Fisheries and Aquatic Sciences* 73(2) (2015) 205-215.
- 393 [48] T. Platt, S. Sathyendranath, C. Fuentes-Yaco, Biological oceanography and fisheries management:
 394 perspective after 10 years, *ICES Journal of Marine Science: Journal du Conseil* 64(5) (2007) 863-869.
- 395 [49] S.P. Greenstreet, A.G. Rossberg, C.J. Fox, W.J. Le Quesne, T. Blasdale, P. Boulcott, I. Mitchell, C.
 396 Millar, C.F. Moffat, Demersal fish biodiversity: species-level indicators and trends-based targets for the

- 397 Marine Strategy Framework Directive, ICES Journal of Marine Science: Journal du Conseil 69(10) (2012)
398 1789-1801.
- 399 [50] W.N. Probst, M. Kloppmann, G. Kraus, Indicator-based status assessment of commercial fish species
400 in the North Sea according to the EU Marine Strategy Framework Directive (MSFD), ICES Journal of
401 Marine Science: Journal du Conseil 70(3) (2013) 694-706.
- 402 [51] A.J. Kenny, H.R. Skjoldal, G.H. Engelhard, P.J. Kershaw, J.B. Reid, An integrated approach for
403 assessing the relative significance of human pressures and environmental forcing on the status of Large
404 Marine Ecosystems, *Progress in Oceanography* 81(1) (2009) 132-148.
- 405 [52] OSPAR, Marine Bird Abundance, Intermediate Assessment 2017. Available at:
406 <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017>
407 (2017).
- 408 [53] OSPAR, Marine Bird Breeding Success or Failure, Intermediate Assessment 2017. Available at:
409 <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017>
410 (2017).
- 411 [54] I. Rombouts, G. Beaugrand, X. Fizzala, F. Gaill, S.P.R. Greenstreet, S. Lamare, F. Le Loc'h, A.
412 McQuatters-Gollop, B. Mialet, N. Niquil, J. Percelay, F. Renaud, A.G. Rossberg, J.P. Féral, Food web
413 indicators under the Marine Strategy Framework Directive: From complexity to simplicity?, *Ecological*
414 *Indicators* 29 (2013) 246-254.
- 415 [55] J.C. Tam, J.S. Link, A.G. Rossberg, S.I. Rogers, P.S. Levin, M.-J. Rochet, A. Bundy, A. Belgrano, S.
416 Libralato, M. Tomczak, Towards ecosystem-based management: identifying operational food-web
417 indicators for marine ecosystems, *ICES Journal of Marine Science* (2017) fsw230.
- 418 [56] M. Dickey-Collas, G.H. Engelhard, A. Rindorf, K. Raab, S. Smout, G. Aarts, M. van Deurs, T. Brunel,
419 A. Hoff, R.A. Lauerburg, Ecosystem-based management objectives for the North Sea: riding the forage fish
420 rollercoaster, *ICES Journal of Marine Science: Journal du Conseil* (2013) fst075.
- 421 [57] A.S. Cook, D. Dadam, I. Mitchell, V.H. Ross-Smith, R.A. Robinson, Indicators of seabird reproductive
422 performance demonstrate the impact of commercial fisheries on seabird populations in the North Sea,
423 *Ecological indicators* 38 (2014) 1-11.
- 424 [58] S.A. Arnott, G.D. Ruxton, Sandeel recruitment in the North Sea: demographic, climatic and trophic
425 effects, *Marine Ecology Progress Series* 238 (2002) 199-210.
- 426 [59] M. Carroll, A. Butler, E. Owen, S. Ewing, T. Cole, J. Green, L. Soanes, J. Arnould, S. Newton, J. Baer,
427 Effects of sea temperature and stratification changes on seabird breeding success, *Climate Research* 66(1)
428 (2015) 75-89.
- 429 [60] K. Eliassen, J. Reinert, E. Gaard, B. Hansen, J.A. Jacobsen, P. Grønkjær, J.T. Christensen, Sandeel as
430 a link between primary production and higher trophic levels on the Faroe shelf, *Marine Ecology Progress*
431 *Series* 438 (2011) 185-194.
- 432 [61] M. Frederiksen, T. Anker - Nilssen, G. Beaugrand, S. Wanless, Climate, copepods and seabirds in the
433 boreal Northeast Atlantic - current state and future outlook, *Global Change Biology* 19(2) (2013) 364-372.
- 434 [62] M.J. Jessopp, M. Cronin, T.K. Doyle, M. Wilson, A. McQuatters-Gollop, S. Newton, R.A. Phillips,
435 Transatlantic migration by post-breeding puffins: a strategy to exploit a temporarily abundant food
436 resource?, *Marine biology* 160(10) (2013) 2755-2762.
- 437 [63] T. Reiertsen, K.E. Erikstad, T. Anker-Nilssen, R. Barrett, T. Boulinier, M. Frederiksen, J. González-
438 Solís, D. Gremillet, D. Johns, B. Moe, Prey density in non-breeding areas affects adult survival of black-
439 legged kittiwakes *Rissa tridactyla*, (2014).
- 440 [64] M. Frost, G. Bayliss-Brown, P. Buckley, M. Cox, S.R. Dye, W.G. Sanderson, B. Stoker, N. Withers
441 Harvey, A review of climate change and the implementation of marine biodiversity legislation in the United
442 Kingdom, *Aquatic Conservation: Marine and Freshwater Ecosystems* (2016) n/a-n/a.
- 443 [65] M.M. Fuentes, L. Chambers, A. Chin, P. Dann, K. Dobbs, H. Marsh, E.S. Poloczanska, K. Maison, M.
444 Turner, R.L. Pressey, Adaptive management of marine mega-fauna in a changing climate, *Mitigation and*
445 *Adaptation Strategies for Global Change* 21(2) (2016) 209-224.
- 446 [66] M.D. Morecroft, H.Q. Crick, S.J. Duffield, N.A. Macgregor, Resilience to climate change: translating
447 principles into practice, *Journal of Applied Ecology* 49(3) (2012) 547-551.
- 448 [67] P.E. Hulme, Adapting to climate change: is there scope for ecological management in the face of a
449 global threat?, *Journal of Applied ecology* 42(5) (2005) 784-794.
- 450 [68] A.J. Constable, D.P. Costa, O. Schofield, L. Newman, E.R. Urban, E.A. Fulton, J. Melbourne-Thomas,
451 T. Ballerini, P.W. Boyd, A. Brandt, Developing priority variables ("ecosystem Essential Ocean Variables"—

452 eEOVs) for observing dynamics and change in Southern Ocean ecosystems, *Journal of Marine Systems*
453 161 (2016) 26-41.
454 [69] T.P. Hughes, D.R. Bellwood, C. Folke, R.S. Steneck, J. Wilson, New paradigms for supporting the
455 resilience of marine ecosystems, *Trends in ecology & evolution* 20(7) (2005) 380-386.

456