

2018-04

Waxy bitumen stranding in southern Australia: a geochemical study of multiple oil families and their likely origins

Rowland, Steven

<http://hdl.handle.net/10026.1/11071>

10.1016/j.orggeochem.2017.12.010

Organic Geochemistry

Elsevier

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Waxy bitumen stranding in southern Australia: a geochemical study of multiple oil families and their likely origins

Dianne S. Edwards^{a,1}, David M. McKirdy^a,
Steven J. Rowland^b, David J. Heath^{b,2}, Patricia S. Gray^{a,3}

a. Organic Geochemistry in Basin Analysis Group, Centre for Tectonics, Resources and Exploration (TRaX), Department of Earth Sciences, School of Physical Sciences, University of Adelaide, SA 5005, Australia

b. Petroleum and Environmental Geochemistry Group, School of Geography, Earth and Environmental Sciences, University of Plymouth, Drake Circus, Plymouth, PL4 8AA, U.K.

1. Present address: Geoscience Australia, GPO Box 378, Canberra, ACT 2601, Australia

2. Present address: Department of Environmental Sciences, Jožef Stefan Institute, Jamova cesta 39, SI-1000 Ljubljana, Slovenia

3. Present address: Department of Environment, Water and Natural Resources, South Australia, GPO Box 1047, Adelaide, SA 5001, Australia

* Corresponding author. Email address: david.mckirdy@adelaide.edu.au

Abstract

Reports of bitumen stranding on the ocean beaches of southern Australia date back to the early days of European settlement. Previous investigations have shown that this ‘coastal bitumen’ comprises three categories of stranded petroleum: waxy bitumen, asphaltite and oil slicks. All three varieties are physically and chemically distinct from each other, and bear no geochemical resemblance to any indigenous Australian crude oil. This study focuses on the most common variety, waxy bitumen, which accounted for 90% of the strandings on six South Australian beaches repeatedly surveyed during 1991–1992.

Geochemical analysis of 96 individual specimens collected from these survey sites and other beaches in South Australia and western Victoria has shown them to be variously weathered high-wax crude oils of paraffinic to aromatic-intermediate bulk composition. Elemental, isotopic and biomarker differences allow their assignment to at least five oil families with inferred source facies that range from deep freshwater lacustrine through paludal and deltaic to euxinic marine, possibly deposited within different sedimentary basins. Family 1, 2 and 3 waxy bitumens all contain biomarkers derived from the freshwater alga *Botryococcus* sp. and tropical angiosperms (notably dipterocarps). Similar biomarker assemblages are unknown in Australian sedimentary basins but are common in Cenozoic crude oils and source rocks throughout western Indonesia. Family 4 waxy bitumens lack these biomarkers but do contain dinosterane and 24-*n*-propylcholestane, indicative of a marine source affinity, while the carbon-isotopic signatures and high pristane/phytane ratios of Family 5 waxy bitumens are consistent with their origin from coal-rich source rocks deposited in fluvial to deltaic sedimentary successions.

The majority of these waxy bitumens represent an oceanic influx of non-indigenous, Southeast Asian crude oils carried into the waters of southern Australia by the Leeuwin Current. Although they are likely to originate from natural seepage within the Indonesian Archipelago, it is unknown whether the parent oils emanate from submarine seeps or from inland seepages

which are then carried to the sea by rivers. The common practice of tanker cleaning operations in the Java and Banda Seas may augment the supply of natural bitumen to the beaches of Australia.

Key words: waxy bitumen, southern Australia, biomarker alkanes, carbon isotopic signatures, oil families, ocean transport, Indonesian provenance

1. Introduction

A recent comprehensive review by Edwards et al. (2016) outlined the history and geographic distribution of bitumen stranding on ocean beaches along the southern coastline of Australia. While these ‘coastal bitumens’ can readily be distinguished from the occasional spillages of crude and refined petroleum from passing maritime traffic (Padley et al., 1993), they are by no means uniform in appearance or composition. Historical records reveal that there are two main types of this ‘ocean wanderer’ bitumen: hard asphalt-like material (‘asphaltum’, now known as asphaltite) and softer waxy bitumen (tar balls).

Arguably the most distinctive variety, asphaltites are a single family of heavy (4–18 °API gravity) aromatic-asphaltic crudes with very high sulfur contents (3–4%). Their biomarker and isotopic signatures indicate an origin from marine shale deposited during a mid-Cretaceous oceanic anoxic event (Edwards et al., 1998; Totterdell et al., 2008). They are most likely a product of low-intensity seepage from tar mats exposed by the incision of submarine canyons into the continental slope along Australia’s southern margin (Hall et al., 2014). While asphaltite appears to have been the main variety referred to in reports from the earlier part of the twentieth century, waxy bitumen is now the most common form of Australian coastal bitumen, accounting for 90% of annual strandings in South Australia and western Victoria during the early 1990s (Padley, 1995; Edwards et al., 2016).

1.1 Historical continuity of waxy bitumen stranding

Tar balls have been stranding on the beaches of southern Australia for more than 100 years, long before the advent of ocean-going tankers and local shipping powered by fuel oil (Sprigg and Woolley, 1963). Moreover, the range of chemical compositions evident in these stranded bitumens has remained relatively constant for at least 65 years (Edwards et al., 2016). They are typically found on sandy west-facing ocean beaches close to the high-water mark. However, there are exceptions to this rule.

Among the samples analysed in the present study are several examples of highly weathered 'fossil waxy bitumen'. These were found within a pumice-rich sand layer above a prominent calcrete horizon located close to the base of a coastal sand dune at the top of the present-day beach, as well as on the inland side of this dune in Beachport Conservation Park (Edwards et al., 2016, fig. 4). Fossil waxy bitumen samples were also found in blowouts between and behind barrier dunes within Canunda National Park and as far south as the Admella Dunes (Fig. 1). These Quaternary barrier shoreline deposits, which are preserved along the Coorong Coastal Plain (Bourman et al., 2016) and extend as far south as Portland in Victoria, have been extensively dated using thermoluminescence (Huntley et al., 1993) and amino acid racemisation (Murray-Wallace et al., 2001) techniques. Since the dune is located above present sea level and within the youngest Robe Range (I) coastal sand barrier system, equivalent in age to the Younghusband Peninsula that fronts the Coorong Lagoon, the pumice-rich layer and its contained bitumen may be up to 7 ka in age (Bourman et al., 2016). Geochemically, these fossil bitumen samples appear almost identical to the main family of waxy bitumens now being washed up on southern Australian beaches, thereby negating the possibility of an anthropogenic origin.

1.2 Previous geochemical studies

The first systematic attempt to characterise the geochemistry of these waxy bitumens was that of McKirdy and Horvath (1976). Using API gravity, US Bureau of Mines (USBM) correlation indices, bulk composition, elemental and isotopic analyses, gas chromatography and trace metal contents, they distinguished two types of waxy bitumen (highly paraffinic to paraffinic, and paraffinic-naphthenic), and inferred that both were derived from early mature lacustrine Cretaceous source rocks suggesting that they may have emanated from seafloor seepage in the western Otway Basin. However, the analytical techniques employed in this initial study did not permit unambiguous determination of their source affinity.

In 1983 the South Australian Department of Mines and Energy (SADME) conducted a major survey of stranded bitumen along the South Australian coastline, involving the collection of new samples for more sophisticated analysis, including by gas chromatography-mass spectrometry (GC-MS), in order to promote petroleum exploration within the adjacent offshore Otway and Duntroon basins. Interpretation of the resulting data by McKirdy (1984a, b) led to the recognition of four genetically distinct families (Families 1, 2, 3 and 4, the first three comprising waxy bitumens and the last asphaltite) based on their sulfur content, the carbon isotopic composition of their C_{12+} saturated and aromatic hydrocarbons, and various alkane biomarker ratios, criteria later augmented by other elemental data (viz. C, H and N) and bulk hydrogen and sulfur isotopic signatures (McKirdy, 1985a).

Subsequent publications (McKirdy, 1985b; McKirdy et al., 1986, 1988) adopted the aforementioned three-fold classification for the southern Australian waxy bitumens and continued to maintain that they were likely to have originated from lacustrine source beds within the Early Cretaceous rift-fill of the Otway Basin. However, failure to identify their distinctive biomarker assemblage (viz. botryococcane, bicadinanes, oleanane and 4-methylsteranes) in any organic-rich facies of this basin (Padley, 1995) necessitated a wider research area. The fact that this association of biomarkers has so far been found only in Cenozoic lacustrine source rocks and crude oils from Sumatra and Java (Seifert and Moldowan, 1981; Williams et al., 1985;

Longley et al., 1990; Schiefelbein and Cameron 1992; Hwang et al., 2002), now suggests that the waxy bitumens represent flotsam which originates from oil seeps in the Indonesian Archipelago and is carried southward into Australian waters by long-distance ocean currents (Currie et al., 1992; McKirdy et al., 1994; Padley, 1995; McGowran et al., 1997; Edwards et al., 2016). That said, there is no agreement on the number of different oil families represented (Table 1), and the locations of their parent seeps are yet to be identified.

1.3 Present investigation

This paper presents the results of geochemical analyses undertaken by Padley (1995) on a suite of 96 archival and more recently stranded waxy bitumens recovered from ocean beaches between Cape Otway, Victoria and the southern Eyre Peninsula, South Australia (Fig. 1). Most of these samples (n = 53) were collected during 1991 and 1992. When combined with elemental, isotopic, biomarker and bulk compositional information from other sources (McKirdy, 1984a, b, 1985a; Tewkesbury, 1989; Watson, 1990), the resulting compilation provided the basis for a revised genetic classification of southern Australian waxy coastal bitumens, and a clearer understanding of their weathering characteristics, source affinity and likely origins. The latter was achieved, first by comparison of our updated classification with those of bitumens found in the Northern Territory (Summons et al., 1993) and Western Australia (Alexander et al., 1994); and then with the various oil families identified in the western Indonesian Archipelago (Robinson, 1987; ten Haven and Schiefelbein, 1995; Schiefelbein and Cameron, 1997; Hwang et al., 2002).

2. Materials and methods

2.1 Sample suite

The bitumen samples examined in this study are listed in Table A1 of the supplementary material, along with their date of collection and, where available, API gravity and sulfur content. The locations of their respective stranding sites are shown in Fig. 1. Each sub-sample taken for analysis came from the interior of the tar ball in order to minimise the effect of weathering on the resulting analytical data. To document these effects in one of the larger specimens (sample 28), three sub-samples were removed from its interior, intermediate and exterior zones and analysed separately.

2.2 Sample fractionation

A weighed aliquot of pure coastal bitumen was dissolved in a minimum amount of dichloromethane and flooded with petroleum ether (30 ml/g). The solution was then refluxed for 2 hours in the presence of activated copper turnings to remove elemental sulfur. The insoluble asphaltenes were separated from the soluble maltenes (asphaltene-free oil) by filtration through pre-extracted filter paper (Whatman Grade 42). The asphaltenes were washed with petroleum ether in Soxhlet apparatus and then recovered by dissolving in dichloromethane. Solvent was removed from the asphaltene and maltene fractions under vacuum in a Büchi rotary evaporator. Each fraction, dissolved in a small volume of solvent, was transferred by Pasteur pipette to a pre-weighed glass vial. Final removal of the solvent was achieved under a stream of nitrogen.

Maltenes were separated into hydrocarbons (saturated and aromatic) and polar compounds (resins) by standard liquid chromatography on a column of activated silica over alumina (sample: adsorbent ratio = 1:100). The saturated hydrocarbon fraction was eluted with petroleum ether; mono-, di- and tri-aromatic hydrocarbon fraction with petroleum ether/dichloromethane (91:9); heavy aromatic hydrocarbon fraction with petroleum ether/dichloromethane (40:60); and resins with methanol/dichloromethane (65:35). Thin layer

chromatography on Merck Kiesel Gel 60G (0.5 mm thick) was employed whenever the amount of maltenes available was either small (<10 mg) or their content of light aromatic hydrocarbons was relatively high. The sample (~20 mg) was spotted 2 cm from the base of the plate and developed in *n*-hexane. *n*-Hexadecane, naphthalene, anthracene and pyrene were used as reference standards. The separated bands were visualised by spraying with a methanolic solution of Rhodamine 6G and viewing under UV light. Four fractions (saturated hydrocarbons; low-molecular-weight aromatic hydrocarbons; higher molecular-weight aromatic hydrocarbons; and polar compounds) were recovered from the silica gel by desorption using the same solvent mixtures as described for column chromatography. Finally, an aliquot (~10 mg) of the saturated hydrocarbon fraction was treated three times with urea to separate the normal and iso-alkanes (paraffins; adduct) from the cyclic and multibranched alkanes (naphthenes; non-adduct). Further details of this procedure may be found in Padley (1995).

2.6 Gas chromatography

The saturated hydrocarbon fraction (isolated from each of the bitumen samples attributed to source 1 in Table A1, supplementary material) was dissolved in *n*-hexane and analysed by gas chromatography (GC) in one of two laboratories, employing the following instrumental parameters.

In the Department of Geology and Geophysics, University of Adelaide, the instrument was a Varian 3400 gas chromatograph fitted with a DB-1 capillary column (30 m x 0.25 mm i.d. x 0.25 µm film thickness) and a flame ionisation detector (FID). The temperature programme of the oven was 40 °C for 2 min, 40 to 300 °C at 4 °C/min and isothermal at 300 °C for 20 min. Two injection techniques were employed: a Varian split/splitless injector operated in the split mode (ratio 20:1) at a temperature of 300 °C, with hydrogen as the carrier gas at a head pressure

of 15 psi; and a SGE on-column injector, with nitrogen as the coolant and hydrogen as the carrier at a linear flow rate of 35 cm/sec.

At the Australian Geological Survey Organisation (now Geoscience Australia), Canberra, the instrument was a Hewlett Packard Series II 5890 gas chromatograph fitted with a FID and a HP Ultra-1 capillary column (50 m x 0.2 mm i.d. x 0.33 μ m film thickness). The carrier gas was hydrogen at a linear flow rate of 35 cm/sec. The sample was injected on-column at 60 °C and the oven temperature ramped from 60 to 300 °C at 4 °C/min before being held isothermal at 300 °C for 20 min.

All quantitative work was carried out in the former laboratory using the on-column injector and 3-methylhenicosane (AC 22; 10 μ g/ml) as the internal standard.

2.7 High-temperature capillary gas chromatography

High-temperature capillary gas chromatography (HTCGC) of the total alkane fraction of six waxy bitumens (samples 28, 50, DB1, 48, 290 and EC1) was undertaken in the then Department of Environmental Sciences at the University of Plymouth, UK. Samples were dissolved in cyclohexane and heated to 55 °C prior to injection into a Carlo Erba HRGC 5300 Mega Series gas chromatograph fitted with a SGE HT5 column (12 m x 0.32 mm x 0.1 μ m film thickness). The carrier gas was helium at 49 kPa head pressure and kept at a constant flow rate of 3 ml/min using the Carlo Erba CP-CF516 module. The sample was injected on-column with air as the coolant and the oven temperature-programmed from 50 to 410 °C at 10 °C/min before being held isothermal at 410 °C for 5 min. It should be noted that no attempt was made to quantify the recoveries of the saturated hydrocarbon fractions through the purification steps undertaken prior to their analysis by HTCGC (see Heath et al., 1995, 1997).

2.8 Gas chromatography–mass spectrometry

Selected total alkane fractions were analysed by gas chromatography–mass spectrometry (GC–MS) in several modes. Routine full scanning (SCAN) and selected-ion monitoring (SIM) were carried out at the Australian Wine Research Institute, Adelaide, using a Varian 3400 gas chromatograph fitted with a DB-1 capillary column (30 m x 0.25 mm x 0.25 µm film thickness) directly connected to the ion source of a TSQ-70 quadrupole mass spectrometer. Samples were dissolved in *n*-hexane and injected via a split/splitless injector operated in split mode (ratio 20:1) at 300 °C. The carrier gas was hydrogen at a head pressure of 15 psi. The operating conditions of the mass spectrometer were as follows: electron energy 70 eV, emission current 200 µA, source temperature 250 °C, scan dwell time 70 msec, mass range 47–500 amu.

The following mass fragmentograms were examined: *m/z* 123, diterpanes; *m/z* 177, demethylated triterpanes; *m/z* 183, acyclic isoprenoid alkanes (including botryococcane); *m/z* 191, triterpanes (including hopane, moretane, gammacerane and oleanane); *m/z* 205, methylated triterpanes; *m/z* 217 and 218, steranes; *m/z* 231, 4-methyl steranes; and *m/z* 259, diasteranes. Individual compounds were identified by their mass spectra which were compared to standards and published data, particularly that of Philp (1985).

The same instrumentation was used to analyse selected aromatic hydrocarbon fractions, except that they were injected on-column at 60 °C and the carrier gas had a linear flow rate of 35 cm/sec. The following mass fragmentograms were examined: *m/z* 128, naphthalene; *m/z* 142, methylnaphthalenes; *m/z* 156, dimethylnaphthalenes; *m/z* 178, phenanthrene and anthracene; *m/z* 192, methylphenanthrenes; *m/z* 206, dimethylphenanthrenes.

To detect and confidently identify key biomarker compounds (notably oleanane, bicadinanes and C₃₀ desmethyl steranes and 4-methyl steranes) selected total alkane fractions and branched/cyclic (urea non-adduct) were analysed by metastable reaction monitoring (MRM) GC–MS on a magnetic sector double focussing VG 7070E mass spectrometer at the Australian Geological Survey Organisation, Canberra. Details of the instrumentation, operating conditions

and parent-to-daughter transitions monitored are reported in McKirdy et al. (1994), although in this study a slightly different temperature program was used (viz. 50 to 180 °C at 8 °C/min, 180 to 300 °C at 4 °C/min, and then held isothermal at 300 °C for 20 min). The concentrations of specific biological marker compounds were measured by the addition of a deuterated sterane standard 3-CD₃-5 α (H)-cholestane (10 ng/ml).

2.9 Stable carbon isotope mass spectrometry

Stable carbon isotopic compositions of the total saturated and aromatic hydrocarbon fractions were determined for the purpose of oil-oil correlations, as well as to give an indication of the bitumen's source affinity. Stable carbon isotopic analyses were carried out using a method slightly modified from that of Sofer (1984). Portions (~1.5 mg) of the saturated and aromatic hydrocarbon fractions were oxidised over cuprous oxide (CuO, 150 mg) at 900 °C for 8 hours with silver wire (to remove sulfur and halogens) in evacuated quartz combustion tubes. The resultant carbon dioxide was separated from the co-produced H₂O and other condensable gases (e.g. SO₂) on a vacuum line by cooling with liquid nitrogen-chilled ethanol. The carbon dioxide was retrieved for introduction to the mass spectrometer by condensing the gas into collection tubes using liquid nitrogen. ¹³C/¹²C ratios were measured on a Micromass 602E isotope mass spectrometer in the Department of Geology and Geophysics, University of Adelaide. The instrument was equipped with a dual inlet system, allowing the sample to be compared directly with a reference gas. The resulting data are presented as $\delta^{13}\text{C}$ values relative to the VPDB standard. Reproducibility, as determined from ten duplicate measurements of the TO2 graphite standard, was ± 0.06 ‰.

3. Results and discussion

3.1 Physical properties and bulk chemical composition

The suite of waxy coastal bitumens examined in this study are heavy to light, solid oils with densities varying between 10 and 40 °API (specific gravity = 0.9957–0.8270); pour points (43–82 °C) that reflect their high wax contents; and sulfur contents that range from low to moderate (0.1–2.6%) (Table 1). Collectively, they comprise inspissated residues of crude oils that range in composition from paraffinic, through paraffinic-naphthenic to aromatic intermediate (Fig. 2; based on data in Table A2, supplementary material). Not surprisingly, waxy C_{23+} *n*-alkanes that commonly display an odd-over-even carbon number preference in the C_{25} – C_{31} range are prominent components of their saturated hydrocarbon fractions (Figs. 3). High-temperature gas chromatograms reveal the presence of individual *n*-alkanes up to C_{60+} (Fig. 4), implying that the parent crude oils are exceptionally rich in wax.

Most of these bitumens display evidence of pronounced weathering (as discussed in Section 3.3). Accordingly, each bitumen specimen has been assigned an empirical degree of weathering (DOW) value in the range 0–5, based upon the extent of removal of its normal and acyclic isoprenoid alkanes (Tables A2 and A3, supplementary material).

3.2 Waxy bitumen families and their source affinities

Differences in the sulfur content, carbon isotopic composition and alkane biomarker geochemistry of the waxy bitumens permit their subdivision into five oil families. Based on a larger data set, the new classification (Table 5) expands the three-fold genetic classification originally proposed by McKirdy (1984a, b) and subsequently employed by McKirdy et al. (1986, 1988, 1994) to include two additional families, one marine (Family 4) and the other paludal (Family 5). The original classification invoked a similar suite of geochemical criteria, augmented by the hydrogen and sulfur isotopic compositions of the whole bitumen. Comparison of the new

classification with those based on sample suites from the Northern Territory (Summons et al., 1993) and Western Australia (Alexander et al., 1994) reveals some overlap but also significant differences (Table 6). Families 1 and 3 are common to all three classification schemes, whereas Families 2, 4 and 5 are not represented in either of these two earlier schemes.

The carbon isotopic signatures of the saturated and aromatic hydrocarbon fractions of the waxy bitumens (Table A4, supplementary material) provide the first clue to their likely origin (Fig. 5). They reveal the algal source affinity of the vast majority of the specimens, while at the same time clearly distinguishing them from asphaltite, the other variety of southern Australian coastal bitumen (Volkman et al., 1992; McKirdy et al., 1986, 1994, 2015; Padley, 1995; Edwards et al., 1998, 2016; Hall et al., 2014). The lack of uniformity of the signatures plotted in Fig. 5 highlights the fact that, unlike the asphaltites, the waxy bitumens belong to multiple oil families.

Saturated hydrocarbon chromatograms typical of the five families are shown in Fig. 3. Of particular significance is the presence of botryococcane (peak labelled *), a C₃₄ irregular acyclic isoprenoid alkane that elutes just before *n*-nonacosane (Peters et al., 2005), in 80% of the waxy bitumens examined. MRM chromatograms illustrating the distributions of triterpanes and steranes in Family 2 and 4 bitumens are shown in Figs. 6 and 7, while those of a Family 1 bitumen may be found in McKirdy et al. (1994, fig. 8). The bicadinane and methylbicadinane distributions of Family 1 and 2 bitumens are compared in Fig. 8. A range of source-dependent biomarker signatures derived from the aforementioned chromatograms of 22 representative samples is presented in Table 3 (parameters 1-16). For an explanation of the significance of these parameters the reader is referred to Peters et al. (2005) and references cited therein.

3.2.1 Family 1

The waxy bitumens of Family 1 are paraffinic to paraffinic-naphthenic crude oils with a relatively homogeneous bulk chemical composition and uniformly low sulfur contents (<0.3%: Tables A1 and A2, supplementary material; Fig. 2). Isotopically they are the heaviest of all the samples analysed in this study (mean $\delta^{13}\text{C}_{\text{sat}} = -25.7\text{‰}$, mean $\delta^{13}\text{C}_{\text{aro}} = -24.5\text{‰}$: Table 2; Fig. 5), likely due to the high content of *Botryococcus*-derived alginite in their source rocks (Boreham et al., 1994; see below). This, in turn, is consistent with the mean isotopic composition of botryococcane ($\delta^{13}\text{C} = -12.4\text{‰}$), pristane (-25.9‰) and phytane (-26.6‰) in the seven Family 1 bitumens reported by Dowling et al. (1995).

The biomarker assemblage of the least weathered Family 1 waxy bitumen (sample 28, Fig. 3a) has the following pertinent characteristics (Tables 3 and 4):

- a bimodal *n*-alkane distribution with maxima at C_{17} and C_{25}
- an intermediate pristane/phytane ratio
- a high concentration of botryococcane
- an ethylcholestane-dominant $5\alpha(\text{H})$ -sterane distribution
- a diacholestane-dominant diasterane distribution
- 4-methyl-24-ethylcholestane in abundance
- the absence of dinosterane
- a high C_{30} hopane/ C_{29} $5\alpha(\text{H})$ -sterane ratio
- the presence of 30-norhopane
- an abundance of oleanane, and
- relatively high concentrations of bicadinane and methylbicadinane.

Its C_{15} – C_{35} *n*-alkane profile (Fig. 3a) is typical of oils derived from a mixture of algal lipids (lower molecular weight *n*-alkanes) and terrestrial plant waxes (higher molecular weight *n*-alkanes), or from the freshwater chlorophyte alga *Botryococcus* sp..

With the advent of HTCGC it became possible to resolve and quantify high-molecular-weight (HMW) hydrocarbons in the range *n*- C_{40} to *n*- C_{100} (del Rio and Philp, 1992; Heath et al.,

1995, 1997). High-temperature chromatograms of the saturated hydrocarbon fractions of five Family 1 waxy bitumens (Fig. 4a–e) reveal marked differences in their C₃₅–C₇₀ *n*-alkane profiles. Making due allowance for the effect of biodegradation on the front ends of the chromatograms (discussed in Section 3.3), the differences are manifest in the relative proportions of two additional modes in their carbon number profiles (with maxima at 41–44 and ~60, respectively). Similar modes have been reported in the wax precipitated in pipes during production of Cenozoic lacustrine oil from the Redwash Field, Uinta Basin, Utah (Carlson et al., 1993). While the origin of this bimodal C₃₅₊ *n*-alkane distribution has yet to be established, the aforementioned differences likely point to variations in the mix of lacustrine algae and higher plants contributing resistant biopolymers (algaenans) and long-chain lipids to their respective source rocks (Tegelaar et al., 1989). Another feature reflecting the freshwater lacustrine source affinity of these Family 1 bitumens is the marked predominance of normal over branched and monocyclic alkanes in the C₄₀–C₆₀ range of their respective saturated hydrocarbon distributions (Fig. 4: Carlson et al., 1993).

Botryococcane is the saturated analogue of a C₃₄ isoprenoid alkene synthesized by *Botryococcus braunii* (specifically clonal race B: Metzger et al., 1985). The presence of 4-methyl-24-ethylcholestane and the absence of dinosterane are together diagnostic of the input of freshwater dinoflagellates to their precursor organic matter (Summons et al., 1987). The pristane/phytane ratios of the least weathered Family 1 bitumens (pr/ph = <1–2.6: Table A4, supplementary material) indicate that their precursor organic matter was deposited in an anoxic to sub-oxic lacustrine setting.

The sterane distribution of the Family 1 waxy bitumens is unusual in that the regular steranes are dominated by ethylcholestane, whereas diacholestane (*not* ethyldiacholestane) is in most instances the dominant diasterane homologue (parameters 1 and 2, Table 4). This anomaly does not appear to be the result of weathering because the sterane distribution is the same for the

least and most weathered samples, and ethyldiacholestane is not known to be preferentially removed by biodegradation (Connan, 1984; Peters et al., 2005).

The co-dominance of ethylcholestane and diacholestane, therefore, is likely to be the result of the dual input of algal and land-plant detritus to the source kerogen. 4-Methyl-24-ethylcholestane is abundant relative to ethylcholestane ($C_{30} 4\alpha\text{Me-ster}/C_{29} 5\alpha\text{H-ster} = 2$: parameter 12: Table 4). The Family 1 waxy bitumens lack both 24-*n*-propylcholestane (a marine algal marker: Moldowan et al., 1990) and (in most instances) dinosterane, confirming that the predominant algal contribution to their source material was from freshwater species.

The high abundance of diasteranes relative to steranes (parameter 5, Table 4) in the Family 1 waxy bitumens reflects the high clay content of their source rock (McKirdy et al., 1983). Further evidence of an argillaceous source lithofacies comes from the dominance of the C_{30} homologue among the C_{27} – C_{35} hopanes and the presence of rearranged hopanes including C_{29} – C_{35} diahopanes and C_{27} and C_{29} $18\alpha\text{H-neohopanes}$ (parameters 7–9, Table 4; ten Haven et al., 1988; Moldowan et al., 1991).

In Family 1 waxy bitumens, the hopanes are dominated by $\alpha\beta$ -hopane and the hopane/sterane ratio is high ($C_{30} \text{ hop}/C_{29} \text{ ster} = 1.7$ – 4.5 : parameter 6, Table 4) indicating a significant contribution of bacterial lipids to the parent organic matter. Such low sterane abundances relative to hopane are typical of lacustrine-sourced oils (Seifert and Moldowan, 1981). Methylhopanes are present in relatively low concentrations.

The angiosperm-derived pentacyclic triterpanes oleanane, bicadinane and methylbicadinane are abundant in the Family 1 waxy bitumens (Table 4), indicating that they are derived from source rocks of Upper Cretaceous or younger age (McKirdy et al., 1994 and references therein). Of all the waxy bitumens, those assigned to Family 1 have the highest concentrations of oleanane ($\text{ol}/\text{hop} = 0.04$ – 0.24 : parameter 13, Table 4). Methylbicadinane and four isomers of bicadinane were identified in the Family 1 waxy bitumens (Fig. 8). Using the nomenclature of Summons et al. (1993) and Murray et al. (1994a), the C_{30} bicadinane isomers are

labelled W, T, T1 and R, respectively; and the methylated bicadinane isomers of T and T1, are annotated as MeT and MeT1, respectively. Typically, the relative abundance of bicadinane isomers in Family 1 waxy bitumens is $T > T1 > R > W$ (Table 4).

Other biomarkers that provide evidence for the contributions of higher plants to the precursor organic matter of the Family 1 waxy bitumens are diterpanes, notably 16 β H-phylocladane and the C₁₈–C₂₀ labdanes (McKirdy et al., 1994). These diterpanes are derived from resin acids produced by both gymnosperms (mainly conifers) and angiosperms (Alexander et al., 1987).

Family 1 is an end-member group of high-wax, low-sulfur, paraffinic crude oils derived from an anoxic to suboxic lacustrine facies containing Type I sapropelic kerogen. In this instance, the precursors included freshwater algae (*Botryococcus* sp. and dinoflagellates) and the biodegraded remains of tropical land-plants (specifically angiosperms of the family *Dipterocarpaceae*). High concentrations of the following biomarkers are unique to the Family 1 waxy bitumens: botryococcane, 4-methyl,24-ethylcholestane, oleanane and bicadinane. *Botryococcus* sp. is common in non-marine kerogens but its diagnostic biomarker, botryococcane, is geologically rare. The limited occurrence of this saturated hydrocarbon may reflect both the rarity of *Botryococcus braunii* Race B (the only one of several different clonal races known to synthesize botryococcane) and the poor resistance of its biological precursor to bacterial and chemical degradation during deposition under oxic conditions (Derenne et al., 1988).

3.2.2 Family 2

The Family 2 waxy bitumens, like those of Family 1, also exhibit high concentrations of botryococcane (bot/*n*-C₂₉ = 0.15–0.71: Table 3). However, their sulfur contents (0.6–2% S) are much higher than those of the Family 1 bitumens, and their carbon isotopic compositions are

both lighter and far less uniform (Fig. 5). Their bulk chemical composition varies from paraffinic-naphthenic to aromatic-intermediate (Fig. 2) and the proportion of saturated to aromatic hydrocarbons also is highly variable ($\text{sat/arom} = 2.4\text{--}8.6$). In the least weathered samples the pristane/phytane ratio is low ($\text{pr/ph} = 0.7\text{--}1.7$). This second family of waxy bitumens may be subdivided into Family 2A and Family 2B, the latter distinguished by the presence of a pronounced naphthene hump in their saturated hydrocarbon chromatograms (Fig. 3b, c). In the least weathered samples of both sub-families, the *n*-alkane profile is distinctly bimodal, although the mode centred on *n*-C₁₇ is less dominant in Family 2A. It is largely on the basis of their $\text{bot}/n\text{-C}_{29}$ values that little weathered samples CB-28 and CB-92 (examples of Family 3 in McKirdy et al., 1994) are herein reassigned to Family 2B (Tables 3 and 4).

The biomarker assemblage of this family is similar to that described for Family 1, except that in addition they contain trace quantities of the marine algal biomarkers, 24-*n*-propylcholestane and dinosterane (Tables 4 and 5). Their apparent mixed source affinity could arise in one of several ways:

- Generation of hydrocarbons from Type I/II to Type II/III kerogens with the organic matter being deposited in freshwater lacustrine or deltaic facies which were subjected to episodic marine incursions.
- Acquisition of higher plant and/or marine signatures during migration of algal-derived lacustrine oils through deltaic and marine facies.
- Reservoirs that have been charged by oils from different source rocks.

3.2.3 Family 3

The waxy bitumens of Family 3 differ from those of Family 2 in exhibiting a wider spread of bulk compositions (paraffinic to aromatic-intermediate: Fig. 2); lower concentrations of botryococcane ($\text{bot}/n\text{-C}_{29} = 0.04\text{--}0.14$); and, on average, higher sulfur contents ($S = 1.1\text{--}2.6$

467 %) and somewhat lower pristane/phytane ratios (pr/ph typically <1: Table 3). These latter
468 features indicate that their source rocks were deposited beneath anoxic, saline (sulfate-bearing)
469 bottom waters starved of iron. Their isotopic signatures overlap those of Family 2 waxy
470 bitumens but are much more tightly grouped (Fig. 5). As was the case for Family 2, these
471 bitumens may be assigned to two sub-families, Family 3A and Family 3B, the saturated
472 hydrocarbon chromatograms of the latter displaying a more pronounced naphthene hump than
473 that shown in Fig. 3d.

474 The Family 3 waxy bitumens can be further distinguished from those of Families 1 and 2
475 by the dominance of cholestane and diacholestane in their C_{27} – C_{30} $5\alpha(H)$ -sterane and diasterane
476 distributions, the presence of 24-*n*-propylcholestane and dinosterane, and the fact that bicadinane
477 and oleanane are present in only trace quantities (Tables 4 and 5). The Family 3 parent oils were
478 generated from a mixture of algal lipids and higher plant waxes, with the former providing the
479 greater input to their source kerogen. The change in overall geochemical character from Family
480 1 to 3 reflects a progressive decrease in the amount of terrestrial influence and a corresponding
481 increase in marine algal source affinity. An accompanying decline in the triterpane ratio Ts/Tm ,
482 from values ≥ 3 in Family 1 to ≈ 1 in Family 3 (Table 4), is consistent with a change in source
483 lithofacies from anoxic shale for Family 1 to predominantly carbonate for Family 3 (Peters et al.,
484 2005).

486 3.2.4 Family 4

488 Family 4 comprises seven miscellaneous samples of waxy, paraffinic to paraffinic-
489 naphthenic bitumen (Fig. 2) which have bimodal *n*-alkane profiles, with maxima at C_{18} and either
490 C_{31} or C_{34} , but notably lack botryococcane. For such a little-weathered bitumen (DOW = 0.5), the
491 saturated hydrocarbon distribution of sample EC 1 (Figs. 3d and 4f) is highly unusual in that it is
492 dominated *n*-alkanes in the range C_{30} – C_{50} . It bears a striking resemblance to that of wax

precipitated from a marine-sourced crude oil during production (Philp, 1994, fig. 2).

MRM data were acquired for four of these samples (Table 4). The presence of trace quantities of 24-*n*-propylcholestane and dinosterane, but no oleanane or bicadinanes (Fig. 7) provides evidence for their marine source affinity. In the absence of appropriate MRM data showing its lack of bicadinanes, sample CB-31-1 was originally assigned to Family 2 (McKirdy et al., 1994). The relatively low abundance of diasteranes in sample EC 1 (C_{29} dia/ster = 0.18), together with the obvious presence of 30-norhopanes (C_{29}/C_{30} hopane = 1.8) and its bulk C-isotopic composition (CV <0.47; Fig. 5), implies that its parent oil was derived from a marine carbonate source rock containing predominantly algal organic matter. Higher diasterane abundances in the other three samples are more typical of an argillaceous source rock. This implied difference in source facies in turn is consistent with the fact that sample EC 1 is appreciably more depleted in ^{13}C than the other Family 4 bitumens and hence plots as an outlier in Fig. 5. It is therefore apparent that the Family 4 waxy bitumens originated from several different source rocks of varying marine lithofacies. The highly weathered state of more than half the samples assigned to this family (Table A3, supplementary material) precluded their definitive assignment to sub-families.

The waxy bitumens of Family 4 differ from those of Family 1, 2 and 3 in lacking botryococcane, oleanane and bicadinanes (Table 5). While not necessarily precluding an origin in Southeast Asia, the absence of these biomarkers raises the possibility that the Family 4 waxy bitumens could be products of a petroleum system located within the offshore Otway Basin and/or Bight Basin, although the Bremer Basin and offshore Perth Basin could also be possibilities given the west to east current direction that transports the waxy bitumens in southern Australian waters (Edwards et al, 2016). Further afield, the basins on Australia's North West Shelf are also worthy of consideration since they lie in the path of the Leeuwin Current. The highly weathered state of more than half the samples (Table A3, supplementary material) prevented their definitive assignment to sub-families.

3.2.5 Family 5

Family 5 comprises five samples of soft bitumen colonised by goose barnacles (Edwards et al., 2016, fig. 5g). These waxy bitumens have a paraffinic bulk composition (Fig. 2) and are sulfur-poor ($S = 0.2\%$; Table A1, supplementary material). Their *n*-alkane distribution is unimodal (maximum at C_{27}) with the lower molecular weight homologues (C_{15} – C_{21}) largely intact (Fig. 3). Pristane is much more abundant than phytane ($pr/ph > 4$). The other key characteristics which distinguish the Family 5 waxy bitumens from Families 1 to 4 are that they do not contain botryococcane (Fig. 3f), they are isotopically more depleted in ^{13}C , and their CV values are > 0.47 (Table 5), highlighting their land-plant source affinity (Fig. 5). Together these features indicate their likely derivation from coal-rich fluvial to deltaic sediments. MRM analysis of their saturated hydrocarbon fractions is required to further constrain the age and location of their source lithofacies.

3.3 Weathering characteristics

The weathering of an oil at or near the sea surface involves a combination of evaporation, photo-oxidation, water washing and biodegradation and causes a marked change in its chemical composition (National Research Council, 2003). The most significant effects are evaporative loss of the light ends (C_1 – C_{12}), dissolution of monoaromatic hydrocarbons (e.g. benzene, toluene) and the selective removal of the C_{12} – C_{23} *n*-alkanes by bacteria, followed at a slower rate by the low-molecular-weight isoprenoid alkanes. Overall, there is an increase in the proportion of non-hydrocarbons (resins and asphaltenes), mainly at the expense of the saturated hydrocarbons as illustrated by sub-sampling of Family 1 bitumen 28 (Table A2, supplementary material).

The aromatic hydrocarbon fractions of eleven samples (representative of all but Family 4) were analysed by SIM GC-MS to augment the limited aromatic maturity data already

545 available on southern Australian waxy bitumens (McKirdy et al., 1994). However, inspection of
546 their respective mass chromatograms revealed that in more than half the samples phenanthrene
547 and the methylphenanthrenes, along with naphthalene and the methylnaphthalenes, had largely
548 been removed by weathering. A combination of water washing and biodegradation (Palmer,
549 1993; Peters et al., 2005) is likely to have been responsible for their removal.

550 The Family 1 waxy bitumens include some of the most highly weathered samples (i.e. *n*-
551 alkanes below C₂₃ and light isoprenoid alkanes totally removed). These paraffinic to paraffinic-
552 naphthenic bitumens typically contain high concentrations of saturated hydrocarbons (Table A3,
553 supplementary material) which would not normally be expected to remain in extremely
554 weathered oil samples. A selection of these bitumens with DOW values in the range 0–5 exhibit
555 a progressive depletion of the C₁₄–C₃₂ *n*-alkanes (Fig. 4a–e) which is typical of oils that have
556 been degraded by aerobic bacteria (Bailey et al., 1973). The pristane/phytane ratio systematically
557 decreases with increased weathering as pristane is more volatile than phytane. A marked increase
558 in the pristane/*n*-heptadecane and the phytane/*n*-octadecane ratios is observed as biodegradation
559 proceeds (Table 3). The botryococcane/*n*-nonacosane ratio in both Family 1 and Family 2
560 bitumens likewise increases with increasing biodegradation.

561 In some of the Family 1 waxy bitumens analysed by MRM (viz. A923.152A, 48 Dune
562 and 68), the sterane distributions have been impacted by weathering (DOW = 2–4: Table A2,
563 supplementary material), displaying a preferential loss of the biological isomer (20R). Two of
564 these samples have also lost their bicadinanes and methylbicadinanes (Table 4). That said, there
565 are other similarly weathered samples in which these biomarkers are intact, although this may
566 also reflect variations in the input of dipterocarp resins to the source rocks.

567 The extensive biodegradation displayed by the most severely weathered waxy bitumens
568 is most likely to have occurred during their exposure to the marine environment. However,
569 another possibility is that their parent oils may have been degraded in the reservoir. For example,
570 similar highly biodegraded oils are produced from the Duri Field in Sumatra (Seifert and

Moldowan, 1981). From the biomarker information obtained for these waxy bitumen samples it is not possible to differentiate between *in situ* reservoir alteration and biodegradation in the marine environment.

The fact that the waxy bitumens form tar balls means that they can survive in the ocean for long (albeit unquantified) periods of time. Their resistance to weathering is not unique as there are many documented cases of large pieces of bitumen surviving in the sea for more than a year (Blumer et al., 1973; Butler et al., 1976; Lee, 1980; Warnock et al., 2015).

3.4 Evidence for a Southeast Asian origin

The unusual assemblage of the age-specific biomarkers botryococcane, oleanane and bicadinanes found in the Family 1, 2 and 3 waxy bitumens (Table 5), suggests that their parent oils originated from one or more Cenozoic basins somewhere in the Indonesian Archipelago (Fig. 9).

Botryococcane has previously been identified only in Cenozoic lacustrine oils and source rocks from Central Sumatra (notably the Minas, Duri and South Bekasap fields: Seifert and Moldowan, 1981) and the Eocene Maoming Oil Shale, Maonan Basin, China (Brassell et al., 1986, 1988). High concentrations of 4-methyl-24-ethylcholestane are also diagnostic of lacustrine environments. This sterane is abundant in the Messel Shale (Rubinstein and Albrecht, 1975) and many non-marine Chinese oils (Shi et al., 1982; Wolff et al., 1986a, b; Philp and Fan, 1987; Fan and Baisheng, 1988). However, the association of high concentrations of botryococcane and 4-methylsteranes has been rarely documented, a notable exception being the aforementioned Maoming Oil Shale. Oleanane first appeared during the Late Cretaceous (late Campanian) but high concentrations of this pentacyclic triterpane have only been observed in Cenozoic sediments and oils, corresponding to the time when angiosperms became the dominant flora. Bicadinane has been identified in Cenozoic oils throughout Indonesia, southern China,

Papua New Guinea, the Philippines and New Zealand (van Aarssen et al., 1990; Murray et al., 1993, 1994a, b). However, high concentrations of bicadinane appear to be confined to Cenozoic sediments and associated oils from Southeast Asia.

Bicadinanes are derived principally from tropical angiosperms of the family *Dipterocarpaceae*, trees that first appeared in the Oligocene and now dominate the lowland rain forests of Southeast Asia, but do not grow in Australia (Paijmans, 1976; Ashton, 1982; Ghazoul, 2016). Fossil evidence suggests that they originated in Africa, and subsequently migrated via India to Asia. However, no dipterocarp micro- or megafossils have been identified in Cenozoic sedimentary successions within Australia (Hill, 1994). The trace levels of bicadinanes (\pm oleanane) found in a Jurassic crude oil and source rock from the Eromanga Basin (Armanios et al., 1995), Jurassic oils and condensates from the Perth Basin (Summons et al., 1995) and Late Cretaceous-sourced oils in the Gippsland Basin (George et al., 1998; Volk et al., 2011) are attributable to a more general angiosperm flora (van Aarssen et al., 1994; Murray et al., 1994b). Therefore, it seems reasonable to conclude that these Family 1–3 waxy bitumens do not originate from Australian sedimentary basins.

High concentrations of bicadinane are in fact typical of Cenozoic oils and source rocks from Indonesia (Cox et al., 1986; Teerman et al., 1987; van Aarssen et al., 1992; Sosrowidjojo et al., 1994) and Papua New Guinea (Murray et al., 1993; Waples and Wulff, 1996). They reflect the predominance of dipterocarps in the Cenozoic flora of this region. Given that the biomarker signatures of the complete range of waxy bitumens indicate their derivation from lacustrine through marginal marine to paludal (coal-bearing) organic facies (Table 5), it is informative to note similar signatures and source affinities in oils from Indonesia (Table 7).

Waxy oils of lacustrine source affinity are confined to basins located in western Indonesia (Fig. 9). Arguably the best documented are those in the onshore Central and South Sumatra basins and the offshore Sunda and East Java basins (Pramono et al., 1990; Schiefelbein and Cameron, 1997; Hwang et al., 2002). Their respective source rocks, viz. the Brown Shale

Formation, Pematang Group (Williams et al., 1985) and the Banuwati Shale (Robinson, 1987), are of Eocene to Oligocene age. The former unit generated the oil reserves of the giant Minas (4 billion barrels) and Duri (3 billion barrels). Lacustrine oils likewise occur further north in the offshore East Natuna, West Natuna and Anambas basins (Noble et al., 2009).

Seifert and Moldowan (1981) were the first to report that lacustrine oils produced from the Central Sumatra Basin have divergent geochemical signatures. The biomarker assemblage of the heavily biodegraded Duri-1 crude oil matches that of the Minas-1 oil, both containing abundant botryococcane. In other oils, such as the South Bekasap-1 crude, the concentration of botryococcane is much lower; and in the Pematang-1 oil this freshwater algal marker is entirely absent, a more saline lagoonal setting for its source rock being indicated by the presence of gammacerane (Table 7). With access to larger datasets, subsequent workers have divided the Indonesian lacustrine oil family into subgroups: Type A, B and C (ten Haven and Schiefelbein, 1995) and Groups I–V (Hwang et al., 2002). Across Sumatra this compositional heterogeneity has been attributed to differences in water chemistry in different sub-basins (Katz and Mertani, 1989) or, alternatively, variation of the Brown Shale source rock facies within a single deep lake system (Schiefelbein and Cameron, 1997).

The biomarker assemblages of the southern Australian waxy bitumens (particularly Families 1–3) show many similarities to some of the lacustrine West Indonesian oils (Table 7), but an exact match is not evident. For example, the deep lacustrine Minas-1 oil from the Central Sumatra Basin is a high wax, low sulfur, paraffinic crude oil that closely resembles the Family 1 bitumens. As previously reported (Seifert and Moldowan, 1981; Currie et al., 1992; Summons et al., 1993; McKirdy et al. 1994; Peters et al., 2005), its biomarker signatures include unusually high concentrations of botryococcane, C₃₀ 4-methylsteranes, bicadinane and methylbicadinane; a ethylcholestane-dominant sterane distribution; the presence of oleanane; and an absence of C₃₀ 24-*n*-propylcholestane and dinosterane, all features of the Family 1 bitumens (Table 5). Moreover, its C-isotopic signatures ($\delta^{13}\text{C}_{\text{sat}} = -26.2\text{‰}$, $\delta^{13}\text{C}_{\text{aro}} = -24.7\text{‰}$, CV = -0.17: Peters et

al., 2005) also plot close to those of the Family 1 bitumens (Fig. 5). However, its bicadinane distribution differs significantly from that of a typical Family 1 bitumen (Fig. 10), as does its HMW C₄₀–C₆₀ alkane distribution and those of several other fresh/brackish water lacustrine oils from Central Sumatra (Carlson et al., 1993), preventing their exact correlation.

Other Indonesian waxy crude oils originated from coal-rich fluvial deltaic sediments of Oligocene-Miocene age and are assigned to the terrigenous family (Table 7). Source rocks in this category include the coals of the Sihapas Group, Central Sumatra Basin and the coals and shales of the Talang Akar Formation in the South Sumatra Basin, the Sunda Basin and the Ardjuna Sub-basin (Robinson, 1987; Pramono et al., 1990; Noble et al., 1991). The Australian waxy bitumens of Family 5 (Table 5) are insufficiently characterised to ascertain the degree to which they resemble the terrigenous oils of western Indonesia, but warrant further research.

With the exception of several examples from the North Sumatra and East Natuna basins, most Indonesian crude oils of marine origin are located in the eastern half of the archipelago (ten Haven and Schiefelbein, 1995; Noble et al., 2009). Here their source rocks may be of either Cenozoic or Mesozoic age and include both carbonate and shale lithofacies, whereas in western Indonesia oil-prone marine source rocks are both less common and all of Cenozoic age (Price et al., 1987; Robinson, 1987; Peters et al., 1999). It is in Eastern Indonesia that most of the known offshore oil seeps occur (Fig. 9). Those located in Seram and the Timor Sea are of marine source affinity and thought to have originated in Mesozoic carbonate and shale lithofacies (Price et al., 1987; Noble et al., 2009). In this respect, they appear similar to the Family 4 waxy bitumens which notably lack oleanane and bicadinanes (Table 5).

An alternative origin for the marine waxy bitumens of Family 4 is from the Australian North West Shelf and Perth basins where marine oils are known (Edwards and Zumberge, 2005 and references therein). In contrast to Southeast Asia, the passive margin of Australia has few documented seepage sites. Where detected, seepage appears to be predominantly gas rather than oil (Logan et al., 2010). However, in the Browse Basin on the Yampi Shelf (in the vicinity of the

Cornea field) natural oil and gas seeps have been sampled (Rollet et al., 2006; Burns et al., 2010), and a microseep of oil was also found at one site in the southern Bonaparte Basin (Stalvies et al., 2017; Fig. 9). These seeps were correlated with crude oils (Cornea and Gwydion) from the Yampi Shelf and determined to be sourced by the Early Cretaceous Echuca Shoals Formation (Logan et al., 2008; Stalvies et al., 2017). However, the C-isotopic compositions of these oils and source rocks are lighter than those displayed by the Family 4 waxy bitumens (Fig. 5), ruling out the Browse Basin as their potential source. Likewise, the Early Triassic-sourced marine oils from the Perth Basin have significantly lighter isotopic compositions than the Family 4 waxy bitumens and contain a distinctive C_{33} *n*-alkylcyclohexane (Summons et al., 1995; Thomas and Barber, 2004) not found in the bitumens, thereby precluding their correlation. The majority of Jurassic marine oils in the Northern Carnarvon and Bonaparte basins are non-waxy crudes sourced by organic-rich marine shales within the Dingo Claystone and Vulcan Formation (Edwards and Zumberge, 2005). However, since these basins also host some waxy Jurassic oils that are isotopically similar to the Family 4 waxy bitumens, further work is required to evaluate their possible common origin.

In summary, the southern Australian coastal waxy bitumens cannot yet be precisely correlated with any Southeast Asian oils. Nevertheless, the Family 1–3 waxy bitumens (comprising ~80% of the samples examined in this study) display sufficient similarities with oils from Indonesia to be most likely derived from this region. An Australian origin for the Family 4 and 5 waxy bitumens remains a possibility, notwithstanding the fact that they do not match any of the oils so far discovered in the Otway Basin, nor any of the minor oil shows (e.g. Greenly-1, Duntroon Sub-basin; Jerboa-1, Eyre Sub-basin) encountered during drilling in the Bight Basin (Ruble et al., 1991; Padley, 1995; Edwards et al., 1999). A potential origin for the Family 5 bitumens in the Gippsland Basin was considered but dismissed because, although Latrobe Group-sourced oils have similar pristane/phytane ratios ($pr/ph = 3–11$), they are significantly more enriched in ^{13}C ($\delta^{13}C_{sat} = -25.2$ to -28.0‰ ; $\delta^{13}C_{arom} = -24.6$ to -26.9‰ ; Summons et al.,

2002). Further studies of potential source rocks and generated hydrocarbons within the basins along Australia's southern margin and North West Shelf are required for definitive oil-source and oil-oil correlations.

3.5 Possible modes of origin

Accepting that the majority of the waxy coastal bitumens are the remains of weathered Southeast Asian crude oils, their occurrence in southern Australia has several possible explanations: jettisoned oil cargoes, tanker washings, leakage from oil pipelines, or natural ocean flotsam emanating from oil seeps.

Australia began importing crude oil during the mid-1950s and continues to import large quantities from Southeast Asia. The composition of many of these imported crudes is documented in a national oil-on-the-sea database (NOSID: Sandison and Edwards, 2001). Oil slicks from tanker washings and petroleum accidentally spilled from local shipping intermittently strand along the Victorian and South Australian coastline (Padley et al., 1993). The physical and chemical properties of these freshly spilled oils readily distinguish them from weathered waxy bitumen (Edwards et al., 2016). The latter is therefore thought not to be a by-product of local maritime activity in southern Australian waters. However, the Straits of Malacca and the Sunda Strait are major shipping routes where tanker cleaning operations are undertaken prior to reloading at Singapore (Fig. 9). Regular oil spills in the western Java Sea since 2004 have been linked to both the high density of maritime traffic and leakage from submarine pipelines carrying oil from offshore production platforms (Putri et al., 2016).

Active onshore oil seeps are common throughout the Indonesian Archipelago (Fig. 9) and Papua-New Guinea (Livsey et al., 1992; Waples and Wulff, 1996), but few overlie known oil fields and none of the giant fields are associated with seeps (Macgregor, 1993, 1995). In the case of the giant Minas Field, the low ratio of gas to oil has reduced the capacity for vertical

migration, making it difficult for the oil to escape to the surface (Thompson et al., 1991). Unless adjacent to major rivers, the inland locations of these surface seeps make them less likely sources for the Australian waxy bitumens.

Submarine seeps such as those recently identified by multibeam echo sounder bathymetry and backscatter imaging in combination with deep-sea piston coring (Fig. 9: Noble et al., 2009; Orange et al., 2009) are potential sources for the Australian coastal waxy bitumens. As previously mentioned, the seeps located offshore from the islands of Timor and Seram are potential points of origin for the marine Family 4 waxy bitumens. It is also worth noting that the asphalt deposit on Buton (Fig. 9) originates from a marine crude oil (Satyana et al., 2013). While no seeps have yet been identified in the offshore Sunda Basin (Macgregor, 1995), this Cenozoic lacustrine depocentre is the site of major lacustrine oil production and remains a possible source for the Family 1 bitumens. Likewise, the co-occurrence of Cenozoic lacustrine and marine oils in the offshore East Natuna Basin (Noble et al., 2009) makes it a potential source for the mixed lacustrine/marine bitumens of Families 2 and 3 (Table 5). Natural seafloor seepage of crude oil into the ocean throughout the Indonesian Archipelago is now well established and would readily give rise to the seaborne waxy bitumens (tar balls) that ultimately strand in southern Australia. However, the fact that such seeps are generally episodic and ephemeral (Kvenvolden and Cooper, 2003) adds to the difficulty of linking a given bitumen stranding to its parent seep.

The possibility that Southeast Asian oils could be transported as flotsam into Australian waters via the Leeuwin Current (McGowran et al., 1997) was invoked by Currie et al. (1992) as an explanation for the geochemical similarity of the Western Australian coastal bitumens and certain Indonesian crude oils. For details of the complex system of ocean currents that flow through the Indonesian Archipelago (Fig. 9) the reader is referred to Gordon (2005), Gordon et al. (2012) and Susanto et al. (2016), while the manner in which they feed into the headwaters of the Leeuwin Current and ultimately into the coastal current and South Australian Current that traverse Australia's southern continental margin (Middleton and Bye, 2007) is summarised in

Edwards et al. (2016). A new study, aimed at more closely linking the stranding pattern of coastal bitumen in South Australia to offshore oceanographic conditions during the last three years (2014–2016), is well advanced (Corrick et al., 2016).

4. Conclusions

Geochemical analysis of 96 waxy bitumens recovered from the coastline of southern Australia has shown them to be inspissated high-wax crude oils of paraffinic to aromatic-intermediate bulk composition. Subtle elemental, isotopic and biomarker differences evident in these bitumens allow their assignment to at least five different oil families. Their inferred source facies range from deep freshwater lacustrine through paludal to estuarine and deltaic marine, possibly deposited within different sedimentary basins.

The Family 1, 2 and 3 waxy bitumens all contain biomarkers derived from the freshwater alga *Botryococcus* sp. and tropical angiosperms (notably dipterocarps). Similar biomarker assemblages are unknown in Australian crude oils and source rocks but are common in the Cenozoic basins of Indonesia. The Family 4 bitumens lack these biomarkers but do contain dinosterane and 24-*n*-propylcholestane, indicative of a marine source affinity, while the C-isotopic signatures and high pristane/phytane ratios of Family 5 waxy bitumens are consistent with an origin from coal-rich source rocks. Hence, it is believed that the majority of the waxy bitumens represent an oceanic influx of non-indigenous, Indonesian crude oil into the waters of southern Australia. Further studies on the Family 4 and 5 waxy bitumens are required, since their origin from seepage in offshore Australian basins has not been ruled out.

The body of geochemical and oceanographic evidence assembled herein points to the origin of the Family 1–3 waxy bitumens from long-term natural seepage within the Indonesian Archipelago. However, it is unknown whether their parent oils emanate from submarine seeps or, less likely, from inland seepages which are then carried to the sea by rivers. The common

practice of tanker cleaning operations in the Java and Banda Seas may augment the supply of coastal bitumen to the beaches of Australia. Further work is required to determine whether such an anthropogenic input makes a significant contribution to the natural background level of seaborne oil within the region.

Acknowledgements

This study formed part of the doctoral research of Dianne Edwards (née Padley) while in receipt of a University of Adelaide Postgraduate Award. The HTGC analyses were undertaken by David Heath at the University of Plymouth as part of his PhD project, funded by the then UK Science and Engineering Council (SERC) and British Gas plc. Patricia Gray (née Tewkesbury) generated some of the C-isotope data during her B.Sc. Honours project at the University of Adelaide. The motivation for writing this paper came from David McKirdy's involvement in the Great Australian Bight Research Program, a collaboration between BP, CSIRO, the South Australian Research and Development Institute (SARDI), the University of Adelaide, and Flinders University that aims to provide a whole-of-system understanding of the environmental, economic and social values of the region. DMMcK acknowledges financial support from Sagasco Resources Limited (now Lattice Energy), the Australian Research Council, the University of Adelaide Research Grants Scheme and CSIRO. We thank Alex Corrick, Tony Hall, Andy Ross, Se Gong, Colin Murray-Wallace, David Christophel, Robert Hill, Robert Langford and Riko Hashimoto for their helpful comments on early drafts of the manuscript, and Junhong Chen and Tom Bernecker for their reviews. Silvio Mezzomo and Alex Corrick are thanked for preparing the figures for publication. Finally, we are indebted to John Volkman, Paul Greenwood and two anonymous reviewers whose perceptive critiques guided its final revision. DSE publishes with the permission of the CEO, Geoscience Australia. The contributions of DSE, DMMcK and PSG to this paper comprise TRaX Record xxx.

References

- van Aarssen, B. G. K., Cox, H. C., Hoogendoorn, P., de Leeuw, J. W., 1990. A cadinene biopolymer in fossil and extant dammar resins as a source for cadinanes and bicadinanes in crude oils from southeast Asia. *Geochimica et Cosmochimica Acta* 54, 3021–3031.
- van Aarssen, B.G.K., Hessels, J.K.C., Abbink, O.A., de Leeuw, J.W., 1992. The occurrence of polycyclic sesqui-, tri-, and oligoterpenoids derived from a resinous polymeric cadinene in crude oils from southeast Asia. *Geochimica et Cosmochimica Acta* 56, 1231–1246.
- van Aarssen, B.G.K., de Leeuw, J.W., Collinson, M., Boon, J., Goth, K., 1994. Occurrence of polycadinene in fossil and recent resins. *Geochimica et Cosmochimica Acta* 58, 223–229.
- Alam, M., Pearson, M.J., 1990. Bicadinanes in oils from the Surma Basin, Bangladesh. *Organic Geochemistry* 15, 461–464.
- Alexander, R., Noble, R.A., Kagi, R.I., 1987. Fossil resin biomarkers and their application in oil to source-rock correlation, Gippsland Basin, Australia. *Australian Petroleum Exploration Association Journal* 27(1), 63–71.
- Alexander, R., Currie, T.J., Kagi, R.I., 1994. The origin of coastal bitumens from Western Australia. *Australian Petroleum Exploration Association Journal* 34(1), 787–798.
- Armanios, C., Alexander, R., Sosrowidjojo, I.B., Kagi, R.I., 1995. Identification of bicadinanes in Jurassic organic matter from the Eromanga Basin, Australia. *Organic Geochemistry* 23, 837–843.
- Ashton, P.S., 1982. *Dipterocarpaceae*. *Flora Malesiana* 9(2), 237–552.
- Bailey, N.J.L., Krouse, H.R., Evans, C.R., Rogers, M.A., 1973. Alteration of crude oil by waters and bacteria – evidence from geochemical and isotopic studies. *American Association of Petroleum Geologists Bulletin* 57, 1276–1290.

- 830 Blumer, M., Ehrhardt, M., Jones, J.H., 1973. The environmental fate of stranded crude oil. Deep-
831 Sea Research 20, 239–259.
- 832 Boreham, C.J., Summons, R.E., Roksandic, Z., Dowling, L.M., Hutton, A.C., 1994. Chemical,
833 molecular and isotopic differentiation of organic facies in the Tertiary lacustrine Duaringa
834 Oil Shale deposit, Queensland, Australia. Organic Geochemistry 21, 685–712.
- 835 Bourman, R.P., Murray-Wallace, C.V., Harvey, N., 2016. Coastal Landscapes of South
836 Australia. University of Adelaide Press, Adelaide, pp. 145–176. DOI:
837 <http://dx.doi.org/10.20851/coast-sa>
- 838 Brassell, S.C., Eglinton, G., Fu, J., 1986. Biological marker compounds as indicators of the
839 depositional history of the Maoming Oil Shale. In: Leythaeuser, D., Rullkötter, J. (Eds.),
840 Advances in Organic Geochemistry 1985, Pergamon, Oxford, pp. 927–941.
- 841 Brassell, S.C., Sheng, G., Fu, J., Eglinton, G., 1988. Biological markers in lacustrine Chinese oil
842 shales. In: Kelts, K., Talbot, M., Fleet, A.J. (Eds.), Lacustrine Petroleum Source Rocks.
843 Geological Society Special Publication No. 40, Blackwell, Oxford, pp. 229–308.
- 844 Burns, K.A., Brinkman, D.L., Brunskill, G.J., Logan, G.A., Volk, H., Wasmund, K., Zagorskis,
845 I., 2010. Fluxes and fate of petroleum hydrocarbons in the Timor Sea ecosystem. Marine
846 Chemistry 118, 140–155.
- 847 Butler, J.N., Morris, B.F., Sleeter, T.D., 1976. The fate of petroleum in the open ocean. In:
848 Sources, Effects and Sinks of Hydrocarbons in the Aquatic Environment. American
849 Institute of Biological Sciences, Washington D. C., pp. 287–297.
- 850 Carlson, R.M.K., Teerman, S.C., Moldowan, J.M., Jacobson, S.R., Chan, E.I., Dorrough,
851 K.S., Seetoo, W.C., Mertani, B., 1993. High temperature gas chromatography of high-wax
852 oils. In: Proceedings of the Indonesian Petroleum Association, pp. 483–507.
- 853 Connan, J., 1984. Biodegradation in reservoirs. In Brooks, J. and Welte, D.H. (Eds.), Advances
854 in Petroleum Geochemistry, Vol. 1. Academic Press, London, pp. 299–335.

- Corrick, A.J. Hall, P.A., McKirdy, D.M., Gong, S., Trefry, C., Dyt, C., Angelini, Z., Ross, A.S., Kempton, R., Armand, S., White, C., 2016. A revised oil family classification scheme and geochemical weathering proxies for South Australian coastal bitumen strandings. 19th Australian Organic Geochemistry Conference, Conference Handbook and Abstracts, pp. 24–25.
- Cox, H.C., de Leeuw, J.W., Schenck, P.A., van Koningsveld, H., Jansen, J.C., van de Graaf, B., van Geerestein, V.J., Kanters, J.A., Kruk, C., Jans, A.W.H., 1986. Bicadinane, a C₃₀ pentacyclic isoprenoid hydrocarbon found in crude oil. *Nature*, 319, 316–318.
- Currie, T.J., Alexander, R., Kagi, R.I., 1992. Coastal bitumens from Western Australia – long distance transport by ocean currents. *Organic Geochemistry* 18, 595–601.
- Derenne, S., Largeau, C., Casadevall, E., Connan, J., 1988. Mechanism of formation and chemical structure of coorongite – II. Structure and origin of the labile fraction. Fate of botryococcenes during early diagenesis. *Organic Geochemistry* 13, 965–971.
- Dowling, L.M., Boreham, C.J., Hope, J.M., Murray, A.P., Summons, R.E., 1995. Carbon isotopic composition of hydrocarbons in ocean-transported bitumens from the coastline of australia. *Organic Geochemistry* 23, 729–737.
- Edwards, D.S., Zumberge, J.E., 2005. The oils of western Australia II. Regional petroleum geochemistry and correlation of crude oils and condensates from western Australia and Papua New Guinea. Geoscience Australia and GeoMark Research Ltd, Canberra and Houston, unpublished. http://www.ga.gov.au/metadata_gateway/metadata/record/gcat_37512
- Edwards, D.S., McKirdy, D.M., Summons, R.E., 1998. Enigmatic asphaltites from the southern Australian margin: molecular and isotopic composition. *PESA Journal* 26, 106–129.
- Edwards, D.S., Struckmeyer, H.I.M., Bradshaw, M.T., Skinner, J.E., 1999. Geochemical characteristics of Australia’s southern margin petroleum systems. *Australian Petroleum Production and Exploration Association Journal* 39(1), 297–321.

- Edwards, D.S., Vinall, D.R., Corrick, A.J., McKirdy, D.M., 2016. Natural bitumen stranding on the ocean beaches of southern Australia: a historical and geospatial review. *Transactions of the Royal Society of South Australia* 140(2), 152–185.
- Fan Pu, Baisheng, Z., 1988. Biomarker characteristics of nonmarine oils in China. *Organic Geochemistry* 13, 627–632.
- George, S.C., Eadington, P.J., Lisk, M., Quezada, R.A., 1998. Geochemical comparison of oil trapped in fluid inclusions and reservoir oil in Blackback oilfield, Gippsland Basin, Australia. *PESA Journal* No. 26, 64–81.
- Ghazoul, J., 2016. *Dipterocarp Biology, Ecology and Conservation*. Oxford University Press, Oxford, 320 p.
- Gordon, A.L., 2005. Oceanography of the Indonesian seas and their throughflow. *Oceanography* 13, 14–27.
- Gordon, A.L., Huber, B.A., Metzger, E.J., Susanto, R.D., Hurlburt, H.E, Adi, T.R., 2012. South China Sea throughflow impact on the Indonesian throughflow. *Geophysical Research Letters* 33(11), <http://dx.doi.org/10.1029/2012GL052021>.
- Grantham, P.J., Posthuma, J., Baak, A., 1983. Triterpanes in a number of Far Eastern crude oils. In: Bjørøy, M. et al. (Eds.), *Advances in Organic Geochemistry 1981*, Wiley, Chichester, pp. 675–683.
- Hall, P.A., McKirdy, D.M., Grice, K., Edwards, D.S., 2014. Australasian asphaltite strandings: their origin reviewed in light of the effects of weathering and biodegradation on their biomarker and isotopic profiles. *Marine and Petroleum Geology* 57, 572–593.
- ten Haven, H.L., Schiefelbein, C., 1995. The petroleum systems of Indonesia. In: *Proceedings of the Indonesian Petroleum Association 24th Annual Convention*, pp. 443–459.
- ten Haven, H.L., de Leeuw, J.W., Sinninghe Damsté, J.S., Schenck, P.A., Palmer, S.E., Zumberge, J.E., 1988. Application of biological markers in the recognition of palaeohypersaline environments. In: Fleet, A.J., Kelts, K, Talbot, M.R. (Eds.), *Lacustrine*

- 907 Petroleum Source Rocks. Geological Society of London Special Publication 40, Blackwell,
908 Oxford, pp. 123-130.
- 909 Heath, D., Moffatt, B., Lowry, R., Rowland, S., 1995. Quantification of the C₃₀₊ fraction of
910 North Sea gas condensates by high temperature capillary gas chromatography. Analytical
911 Proceedings 32, 485–487.
- 912 Heath, D.J., Lewis, C.A., Rowland, S.J., 1997. The use of high temperature gas chromatography
913 to study the biodegradation of high molecular weight hydrocarbons. Organic Geochemistry
914 26, 459–473.
- 915 Hill, R.S. (ed.), 1994. History of the Australian Vegetation: Cretaceous to Recent. Cambridge
916 University Press, Cambridge, 433 p.
- 917 Holba, A.G., Tegelaar, E., Ellis, L., Singletary, M.S., Albrecht, P., 2000. Tetracyclic
918 polyprenoids: Indicators of freshwater (lacustrine) algal input. Geology 28, 251–254.
- 919 Huntley, D.J., Hutton, J.T., Prescott, J.R., 1993. The stranded beach-dune sequence of south-east
920 South Australia: a test of thermoluminescence dating, 0-800 ka. Quaternary Science
921 Reviews 12, 1–20.
- 922 Hwang, R.J., Heidrick, T., Mertani, B., Qivayanti, M.L., 2002. Correlation and migration studies
923 of North Central Sumatra oils. Organic Geochemistry 33, 1361–1379.
- 924 Katz, B.Z., Mertani, B., 1989. Central Sumatra – a geochemical paradox. In: Proceedings of the
925 Indonesian Petroleum Association 18th Annual Convention, pp. 403–425.
- 926 Kvenvolden, K.A., Cooper, C.K., 2003. Natural seepage of crude oil into the marine
927 environment. Geo-Marine Letters 23, 140–146.
- 928 Lee, R.F., 1980. Processes affecting the fate of oil in the sea. In: Geyer, R.A. (Ed.), Marine
929 Environmental Pollution. 1. Hydrocarbons. Elsevier Science, Amsterdam, pp. 337–351.
- 930 Livsey, A.R., Duxbury, N., Richards, F., 1992. The geochemistry of Tertiary and Pre-Tertiary
931 source rocks and associated oils in Eastern Indonesia. In: Proceedings of the Indonesian
932 Petroleum Association 21st Annual Convention, pp. 499–520.

- 933 Logan, G.A., Jones, A.T., Ryan, G., Wettle, M., Thankappan, M., Grosjean, E., Rollet, N.,
934 Kennard, J.M., 2008. Review of Australian offshore natural hydrocarbon seepage studies.
935 Geoscience Australia Record, 2008/17.
- 936 Logan, G.A., Jones, A.T., Kennard, J.M., Ryan, G.J., Rollet, N., 2010. Australian offshore
937 natural hydrocarbon seepage studies, a review and re-evaluation. *Marine and Petroleum*
938 *Geology* 27, 26–45.
- 939 Longley, I.M., Barraclough, R., Bridden, M.A., Brown, S., 1990. Pematang lacustrine petroleum
940 source rocks from the Malacca Strait PSC, Central Sumatra, Indonesia. In: *Proceedings of*
941 *the 19th Annual Convention of the Indonesian Petroleum Association*, pp. 279–297.
- 942 Macgregor, D.S., 1993. Relationships between seepage, tectonics and subsurface petroleum
943 reserves. *Marine and Petroleum Geology* 10, 606–619.
- 944 Macgregor, D. S., 1995. The exploration significance of surface oil seepage: an Indonesian
945 perspective: In: *Proceedings of the Indonesian Petroleum Association 24th Annual*
946 *Convention*, pp. 97–109.
- 947 McGowran, B., Li, Q., Cann, J., Padley, D., McKirdy, D.M., Shafik, S., 1997. Biogeographic
948 impact of the Leeuwin Current in southern Australia since the late middle Eocene.
949 *Palaeogeography, Palaeoclimatology, Palaeoecology* 136, 19–40.
- 950 McKirdy, D.M., 1984a. Coastal bitumens and potential source rocks in the western Otway Basin,
951 South Australia and Victoria. AMDEL Report F 5840/84 for Australian Aquitaine
952 Petroleum Pty Ltd. and Ultramar Australia Inc., South Australian Department of Mines and
953 Energy Open File Envelope No. 5876, pp. 3–365.
- 954 McKirdy, D.M., 1984b. Coastal bitumens and potential source rocks, Duntroon Basin, South
955 Australia. AMDEL Report F 5769/84 for Getty Oil Development Company Ltd., South
956 Australian Department of Mines and Energy Open File Envelope No. 5876, pp. 366–402.
- 957 McKirdy, D.M., 1985a. Otway Basin coastal bitumens: elemental and stable isotopic
958 compositions, and biological marker geochemistry. AMDEL Report F 6176/85 for South

- 959 Australian Department of Mines and Energy (unpubl.). S. Aust. Dept. of Mines and Energy
- 960 Open File Envelope No. 5876, pp. 403-440.
- 961 McKirdy, D.M., 1985b. Coorongite, coastal bitumen, and their origins from the alga
- 962 *Botryococcus* in the western Otway Basin. In Otway 85 – Earth Resources of the Otway
- 963 Basin. Geological Society of Australia, Adelaide, pp. 34–49.
- 964 McKirdy, D.M., Horvath, Z., 1976. Geochemistry and significance of coastal bitumen from
- 965 southern and northern Australia. Australian Petroleum Exploration Association Journal
- 966 16(1), 123–136.
- 967 McKirdy, D.M., Aldridge, A.K., Ypma, P.J.M., 1983. A geochemical comparison of some crude
- 968 oils from Pre-Ordovician carbonate rocks. In: Bjorøy, M., Albrecht, P., Cornford, C., et al.
- 969 (Eds.), Advances in Organic Geochemistry 1981. John Wiley, Chichester, pp. 99–107.
- 970 McKirdy, D.M., Cox, R.E., Volkman, J.K., Howell, V.J., 1986. Botryococcane in a new class of
- 971 Australian non-marine crude oils. Nature 320, 57–59.
- 972 McKirdy, D.M., Cox, R.E., Morton, J.G.G., 1988. Biological marker, isotopic and geological
- 973 studies of lacustrine crude oils in the western Otway Basin, South Australia. In: Fleet, A.J.,
- 974 Kelts, K., Talbot, M.R. (Eds.), Lacustrine Petroleum Source Rocks. Geological Society of
- 975 London Special Publication 40, Blackwell, Oxford, p. 327
- 976 McKirdy, D.M., Summons, R.E., Padley, D., Serafini, K.M., Boreham, C.J., Struckmeyer,
- 977 H.I.M., 1994. Molecular fossils in coastal bitumens from southern Australia: signatures of
- 978 precursor biota and source rock environments. Organic Geochemistry 21, 265–286.
- 979 McKirdy, D.M., Corrick, A.J., Ross, A.S., Gong, S., Trefry, C., Dyt, C., Z. Angelini, Z., Jobin,
- 980 O., Hall, P.A., Edwards, D.S., 2015. Understanding the geographical distribution of
- 981 historical and modern asphaltite strandings along the South Australian coastline. AAPG
- 982 International Conference and Exhibition, Melbourne, Abstract.

- 983 Metzger, P., Casadevall, E., Pouet, M. J., Pouet, Y., 1985. Structures of some botryococcenes:
984 branched hydrocarbons from the B-race of the green alga *Botryococcus braunii*.
985 *Phytochemistry* 24, 2995–3002.
- 986 Middleton, J.F., Bye, J.A.T., 2007. A review of the shelf-slope circulation along Australia's
987 southern shelves: Cape Leeuwin to Portland. *Progress in Oceanography*, 75, 1–41.
- 988 Moldowan, J.M., Fago, F.J., Lee, C.Y., Jacobson, S.R., Watt, D.S., Slougui, N.E., Jeganathan,
989 A., Young, D.C., 1990. Sedimentary 24-*n*-propylcholestanes, molecular fossils diagnostic
990 of marine algae. *Science* 247, 309–312.
- 991 Moldowan, J.M., Fago, F.J., Carlson, R.M.K., Young, D.C., van Dutne, G., Clardy, J., Schoell,
992 M., Pillinger, C.T., Watt, D.S., 1991. Rearranged hopanes in sediments and petroleum.
993 *Geochimica et Cosmochimica Acta* 55, 3333–3353.
- 994 Murray, A.P., Summons, R.E., Bradshaw, J., Pawih, B., 1993. Cainozoic oil in Papua New
995 Guinea—evidence from geochemical analysis of two newly discovered seeps. In: Carman,
996 G.J., Carman, Z. (Eds.), *Petroleum Exploration and Development in Papua New Guinea*
997 *Proceedings of the 2nd Papua New Guinea Petroleum Convention*, Port Moresby, pp. 489–
998 498.
- 999 Murray, A., Padley, D., McKirdy, D.M., Summons, R.E., Booth, W.E., 1994a. Oceanic transport
1000 of fossil dammar resin: the chemistry of coastal resinites from South Australia.
1001 *Geochimica et Cosmochimica Acta* 58, 3049–3059.
- 1002 Murray, A.P., Summons, R.E., Boreham, C.J., Dowling, L.M., 1994b. Biomarker and *n*-alkane
1003 isotope profiles for Tertiary oils: relationship to source rock depositional setting. In:
1004 Telnæs, N., van Graas, G., Øygard, K. (Eds.), *Advances in Organic Geochemistry 1993*,
1005 Elsevier Science, Amsterdam, pp. 521–542.
- 1006 Murray-Wallace, C.V., Brooke, B.P., Cann, J.H., Belperio, A.P., Bourman, R.P., 2001. Whole-
1007 rock aminostratigraphy of the Coorong Coastal Plain, South Australia: towards a 1 million

- 1008 year record of sea-level highstands. *Journal of the Geological Society*, London 158, 111–
- 1009 124.
- 1010 National Research Council (NRC), 2003. *Oil in the Sea: Inputs, Fates, and Effects*. National
- 1011 Academies Press, Washington, D.C., 280 p.
- 1012 Noble, R.A., Wu, C.H., Atkinson, C.D., 1991. Petroleum generation and migration from Talang
- 1013 Akar coal and shales offshore N.W. Java, Indonesia. *Organic Geochemistry* 17, 363–374.
- 1014 Noble, R., Orange, D., Decker, J., Teas, P., Baillie, P., 2009. Oil and gas seeps in deep marine
- 1015 sea floor cores as indicators of active petroleum systems in Indonesia. In: *Proceedings of*
- 1016 *the Indonesian Petroleum Association 33rd Annual Convention*, pp. 385–394.
- 1017 Orange, D.L., Teas, P.A., Decker, J., Baillie, P., Johnstone, T., 2009. Using SeaSeep surveys to
- 1018 identify and sample natural hydrocarbon seeps in offshore frontier basins. In: *Proceedings*
- 1019 *of the Indonesian Petroleum Association 33rd Annual Convention*, 21 p.
- 1020 Padley, D., 1995. *Petroleum geochemistry of the Otway Basin and the significance of coastal*
- 1021 *bitumen strandings on adjacent southern Australian beaches*. Ph.D. thesis, University of
- 1022 Adelaide. <http://hdl.handle.net/2440/48539>
- 1023 Padley, D., McKirdy, D.M., Murray, A., Summons, R.E., 1993. Oil strandings on the beaches of
- 1024 southern Australia: origins from natural seepage and shipping. In: Oygard, K. (Ed.),
- 1025 *Organic Geochemistry*. Poster sessions from the 16th International Meeting on Organic
- 1026 *Geochemistry*, Stavanger. Falch Hurtigtrykk, Oslo, pp. 660–663.
- 1027 Pajmans, K. (Ed.), 1976. *New Guinea vegetation*. Australian National University Press,
- 1028 Canberra, 212 p.
- 1029 Palmer, S.E., 1993. Effect of biodegradation and water washing on crude oil composition. In:
- 1030 Engel, M.H., Macko, S.A. (Eds.), *Organic Geochemistry*. Plenum Press, New York, pp.
- 1031 511–533.

- 1032 Peters, K.E., Fraser, T.H., Amris, W., Rustanto, B., Hermanto, E., 1999. Geochemistry of crude
1033 oils from eastern Indonesia. American Association of Petroleum Geologists Bulletin 83,
1034 1927–1942.
- 1035 Peters, K.E., Walters, C.C., Moldowan, J.M., 2005. The Biomarker Guide, 2nd Edition. Volume
1036 2: Biomarkers and Isotopes in Petroleum Systems and Earth History. Cambridge
1037 University Press, pp. 475–1155.
- 1038 Philp, R.P., 1985. Fossil Fuel Biomarkers, Applications and Spectra. Methods in Geochemistry
1039 and Geophysics, 23. Elsevier, New York.
- 1040 Philp, R.P., 1994. Geochemical characteristics of oils from predominantly from terrigenous
1041 source materials. In: Scott, A.C., Fleet, A.J. (Eds.), Coal and Coal-bearing Strata as Oil-
1042 prone Source Rocks? Geological Society Special Publication 77, 71–91.
- 1043 Philp, R.P., Fan, Zhao-an, 1987. A geochemical investigation of oils and source rocks from
1044 Qianjiang depression of Jiangnan Basin, a terrigenous saline basin, China. Organic
1045 Geochemistry 11, 549–562.
- 1046 Pramono, H., Wu, C.C.H., Noble, R.A., 1990. A new oil kitchen and petroleum bearing sub-
1047 basin in the offshore northwest Java Basin. In: Proceedings of the Indonesian Petroleum
1048 Association 19th Annual Convention, pp. 254–277.
- 1049 Price, P.L., O'Sullivan, T., Alexander, R., 1987. The nature and occurrence of oil in Seram,
1050 Indonesia. In: Proceedings of the Indonesian Petroleum Association 16th Annual
1051 Convention, pp., 141–173.
- 1052 Putri, M.R., Setiawan, A., Pohlmann, T., Mayer, B., Gade, M., 2016. The assessment of oil
1053 pollution in Seribu Islands based on remote sensing and numerical models. Proceedings of
1054 the Living Planet Symposium 2016, Prague, Czech Republic, 9–13 May 2016 (ESA SP-
1055 740, August 2016).
- 1056 del Rio, J.C., Philp, R.P., 1992. High-molecular-weight hydrocarbons: a new frontier in organic
1057 geochemistry. Trends in Analytical Chemistry 11, 187–193.

- 1058 Robertson, 2017. Robertson Tellus Sedimentary Basins of the World Map,
1059 [http://www.datapages.com/gis-map-publishing-program/gis-open-files/global-
framework/robertson-tellus-sedimentary-basins-of-the-world-map](http://www.datapages.com/gis-map-publishing-program/gis-open-files/global-
1060 framework/robertson-tellus-sedimentary-basins-of-the-world-map)
- 1061 Robinson, K.M., 1987. An overview of source rocks and oils in Indonesia. In: Proceedings of the
1062 Indonesian Petroleum Association 16th Annual Convention, pp. 97–122.
- 1063 Rollet, N., Logan, G.A., Kennard, J.M., O’Brien, P.E., Jones, A.T., Sexton, M., 2006.
1064 Characterisation and correlation of active hydrocarbon seepage using geophysical data
1065 sets: an example from the tropical, carbonate Yampi Shelf, Northwest Australia. *Marine
1066 and Petroleum Geology* 23, 145–164.
- 1067 Rubinstein, I., Albrecht, P., 1975. The occurrence of nuclear methylated steranes in shale.
1068 *Journal of the Chemical Society, Chemical Communications*, 957–958.
- 1069 Ruble, T.E., Logan, G.A., Blevin, J.E., Struckmeyer, H.I.M., Liu, K., Ahmed, M., Eadington,
1070 P.J., Quezada, R.A., 2001. Geochemistry and charge history of a palaeo-oil column:
1071 Jerboa-1, Eyre Sub-basin, Great Australian Bight. In: Hill, K.C., Bernecker, T. (Eds.),
1072 Eastern Australasian Basins Symposium: a refocused energy perspective for the future.
1073 *Petroleum Exploration Society of Australia, Special Publication*, pp. 521–529.
- 1074 Sandison, C., Edwards, D., 2001. National Oil-on-the-Sea Identification Database (NOSID)
1075 Users’ Guide. Australian Geological Survey Organisation, Australian Government
1076 Analytical Laboratories and Australian Maritime Safety Authority, 39 p.
- 1077 Satyana, A.H., Irawan, C., Kurniawan, W., 2013. Revisit geology and geochemistry of Buton
1078 asphalt deposits, SE Sulawesi: implications for petroleum exploration of Buton area. In:
1079 Proceedings of the Indonesian Petroleum Association 37th Annual Convention &
1080 Exhibition. IPA13-G-170.
- 1081 Scheifelbein, C., Cameron, N., 1997. Sumatra/Java oil families. In: Fraser, A.J., Matthews, S.J.,
1082 Murphy, R.W. (Eds.), *Petroleum Geology of Southeast Asia. Geological Society Special
1083 Publication* 126, Blackwell, Oxford, pp. 143-146.

- 1084 Seifert, W.K., Moldowan, J.M., 1981. Palaeoreconstruction by biological markers. *Geochimica*
1085 *et Cosmochimica Acta* 45, 783–794.
- 1086 Shi, J.-Y., Mackenzie, A. S., Alexander, R., Eglinton, G., Gower, A.P., Wolff, G A., Maxwell,
1087 J.R., 1982. A biological marker investigation of petroleums and shales from the Shengli
1088 oilfield, the People’s Republic of China. *Chemical Geology* 35, 1–31.
- 1089 Sofer, Z., 1984. Stable isotope compositions of crude oils: application to source depositional
1090 environments and petroleum alteration. *Bulletin of the American Association of Petroleum*
1091 *Geologists* 68, 31–49.
- 1092 Sosrowidjojo, I.B., Alexander, R., Kagi, R.I., 1994. The biomarker composition of some crude
1093 oils from Sumatra. *Organic Geochemistry* 21, 303–312.
- 1094 Sprigg, R.C., Woolley, J.B., 1963. Coastal bitumen in southern Australia, with special reference
1095 to observations at Geltwood Beach, south-east South Australia. *Transactions of the Royal*
1096 *Society of South Australia* 86, 67–103.
- 1097 Stalvies, C., Talukder, A., Ross, A., Grosjean, E., Carr, A., Williams, A., Gresham, M., Binning,
1098 M., Jablonski, D., 2017. Establishing hydrocarbon seepage to the Ashmore Platform,
1099 Bonaparte Basin, Australia: a natural seeps study. *Marine and Petroleum Geology* 8, 56–
1100 68.
- 1101 Subroto, E.A., Alexander, R., Pranyoto, U., Kagi, R.I., 1992. The use of 30-Norhopanes series,
1102 a novel carbonate biomarker, in source rock to crude oil correlation in the North Sumatra
1103 Basin, Indonesia. In: *Proceedings of the Indonesian Petroleum Association, 21st Annual*
1104 *Convention*, pp. 145-163.
- 1105 Summons, R.E., Volkman, J.K., Boreham, C.J., 1987. Dinosterane and other steroidal
1106 hydrocarbons of dinoflagellate origin in sediments and petroleum. *Geochimica et*
1107 *Cosmochimica Acta* 51, 3075–3082.

- 1108 Summons, R.E., Bradshaw, J., Brooks, D.M., Goody, A.K., Andrew P. Murray, A.P., Foster,
1109 C.B., 1993. Hydrocarbon composition and origins of coastal bitumens from the Northern
1110 Territory, Australia. PESA Journal No. 21, 31–42.
- 1111 Summons, R.E., Boreham, C.J., Foster, C.B., Murray, A.P., Gorter, J.D., 1995.
1112 Chemostratigraphy and the composition of oils in the Perth Basin, Western Australia.
1113 APEA Journal 35(1), 613–632.
- 1114 Summons, R.E., Zumberge, J.E., Boreham, C.J., Bradshaw, M.T., Brown, S.W., Edwards, D.S.,
1115 Hope, J.M., Johns, N., 2002. The Oils of Eastern Australia. Petroleum Geochemistry and
1116 Correlation. Geoscience Australia, Canberra and GeoMark Research, Houston.
1117 [https://ecat.ga.gov.au/geonetwork/srv/eng/search#!a05f7892-ff6d-7506-e044-](https://ecat.ga.gov.au/geonetwork/srv/eng/search#!a05f7892-ff6d-7506-e044-00144fdd4fa6)
1118 [00144fdd4fa6](https://ecat.ga.gov.au/geonetwork/srv/eng/search#!a05f7892-ff6d-7506-e044-00144fdd4fa6)
- 1119 Susanto, R.D., Wei, Z., Adi, T.R., Zheng, Q., Fang, G., Fan, B., Supangat, A., Agustiadi, T., Li,
1120 S., Trenggono, M., Setiawan, A., 2016. Oceanography surrounding Krakatau Volcano in
1121 the Sunda Strait, Indonesia. Oceanography 29(2), 264–272.
- 1122 Sutton, P.A., Rowland, S.J., 2012. High temperature gas chromatography–time-of-flight-mass
1123 spectrometry (HTGC–ToF-MS) for high-boiling compounds. Journal of Chromatography
1124 A 1243, 69–80.
- 1125 Sutton, P.A., Wilde, M.J., Martin, S.J., Cvacka, J., Rowland, S.J., 2013. Studies of long chain
1126 lipids in insects by high temperature gas chromatography and high temperature gas
1127 chromatography–mass spectrometry. Journal of Chromatography A 1297, 236–240.
- 1128 Teerman, S.C., Hwang, R.J., Williams, H.H., 1987. Liquid hydrocarbon potential of Sumatran
1129 resinite. In: Proceedings of the Indonesian Petroleum Association 16th Annual
1130 Convention, pp. 231–240.
- 1131 Tegelaar, E.W., Matthezing, R.M., Jansen, J.B.H., Horsfield, B., de Leeuw, J.W., 1989. Possible
1132 origin of *n*-alkanes in high-wax crude oils. Nature 342, 529–531.

- 1133 Tewksbury, P.S., 1989. Organic geochemistry of oil seeps in the Otway Basin. B.Sc. Honours
1134 thesis, University of Adelaide (unpublished).
- 1135 Tissot, B.P., Welte, D.H., 1984. Petroleum Formation and Occurrence. Springer-Verlag, Berlin.
- 1136 Thomas, B.M., Barber, C.J., 2004. A re-evaluation of the hydrocarbon habitat of the northern
1137 Perth Basin. The APPEA Journal, 44(1), 13-57.
- 1138 Thompson, M., Reminton, C., Purnomo, J., MacGregor, D., 1991. Detection of liquid
1139 hydrocarbon seepage in Indonesian offshore frontier basins using airborne laser
1140 fluorosensor (ALF). The results of a Pertamina/BP Joint study. In: Proceedings of the
1141 Indonesian Petroleum Association 20th Annual Convention, pp. 663–689.
- 1142 Totterdell, J.M, Struckmeyer, H.I.M., Boreham, C.J., Mitchell, C.H., Monteil, E., & Bradshaw,
1143 B.E., 2008. Mid–Late Cretaceous organic-rich rocks from the eastern Bight Basin:
1144 implications for prospectivity. In: Blevin, J.E., Bradshaw, B.E., Uruski, C. (Eds.), Eastern
1145 Australasian Basins Symposium III, Petroleum Exploration Society of Australia, Special
1146 Publication, pp. 137–158.
- 1147 Volk, H., Ahmed, M., Gong, S., Boreham, C., Tingate, P., Edwards, D., 2011. Land plant
1148 markers in Gippsland Basin oils, Australia. 25th International Meeting on Organic
1149 Geochemistry, Interlaken, Abstracts, p. 414.
- 1150 Volkman, J.K., O’Leary, T., Summons, R.E., Bendall, M.R., 1992. Biomarker composition of
1151 some asphaltic coastal bitumens from Tasmania, Australia. Organic Geochemistry 18,
1152 669–682.
- 1153 Waples, D.W., Wulff, K.J., 1996. Genetic classification and exploration significance of oils and
1154 seeps of the Papuan Basin. In: Buchanan, P.G. (Ed.), Petroleum Exploration, Development
1155 and Production in Papua New Guinea. Proceedings of the 3rd Papua New Guinea
1156 Petroleum Convention, Port Moresby, pp. 417–430.
- 1157 Warnock, A.M., Hagen, S.C., Passeri, D.L., 2015. Marine tar residues: a review. Water, Air &
1158 Soil Pollution 226: 68. DOI 10.1007/s11270-015-2298-5

- 1159 Watson, B., 1990. Coastal bitumen study. AMDEL Report 009/130 for South Australian
1160 Department of Mines and Energy (unpubl.).
- 1161 Williams, H.H., Kelley, P.A., Janks, J.S., Christensen, R.M., 1985. The Paleogene Rift Basin
1162 source rocks of Central Sumatra. In: Proceedings of the Indonesian Petroleum Association
1163 14th Annual Convention, pp. 57–90.
- 1164 Wolff, G.A., Lamb, N.A., Maxwell, J.R., 1986a. The origin and fate of 4-methyl steroid
1165 hydrocarbons. I. Diagenesis of 4-methyl steranes. *Geochimica et Cosmochimica Acta* 50,
1166 335–342.
- 1167 Wolff, G.A., Lamb, N.A., Maxwell, J.R., 1986b. The origin and fate of 4-methyl steroid
1168 hydrocarbons. II. Dehydration of stanols and occurrence of C₃₀ 4-methyl steranes. *Organic*
1169 *Geochemistry* 10, 965–976.
- 1170

Table 1. Previous geochemical studies of Australian waxy coastal bitumen

Location	No. of samples	No. of oil families	Data source
South Australia & Northern Territory	11	2	1
South Australia & Western Victoria	27	3	2
South Australia & Western Victoria	96	5	3
Northern Territory	22	2	4
Western Australia	70	8	5

1. McKirdy & Horvath (1976)
2. McKirdy et al. (1984a, b, 1985b, 1986, 1994)
3. Padley (1995)
4. Summons et al. (1993)
5. Alexander et al. (1994)

Table 2. Summary of waxy bitumen bulk properties*

Specific gravity °API	Pour point** °C	Sulfur content wt %	Oil type***
10 – 40 mean = 22 n = 42	43 – 82 mean = 61 n = 33	0.1 – 2.6 mean = 1.3 n = 45	P (n = 16) P-N (n = 36) Ar-I (n = 21)

* Based on information compiled in Appendix Table A1 (see supplementary data)

** Measured using standard method IP 15/67

*** P = paraffinic, P-N = paraffinic-naphthenic, Ar-I = aromatic-intermediate (after Tissot and Welte, 1984)

Table 3. Source-dependent biomarker parameters of selected waxy bitumens

Sample ID	Steranes						Triterpanes		
	27 : 28 : 29 ster	27 : 28 : 29 dia	27/29 ster	30/29 ster	29 dia/ster	30/29 hop/ster	27 Ts/Tm	29/30 hop	30 dia/hop
Family 1									
28	37:26:37	51:21:28	1.0	0	0.70	2.8	3.7	0.61	0.17
4066	21:25:54	nd	0.38	0	0.70	3.5	3.6	0.84	0.19
Gelt (318)	22:20:58	45:22:33	0.38	0	0.44	3.1	2.1	0.82	0.17
Devil's (317)	29:22:49	35:31:33	0.57	0	0.20	1.7	0.77	1.0	0.09
CB-38 (378)	32:19:49	11:37:52	0.65	0	0.46	4.2	2.7	0.68	0.10
59	nd	nd	0.68	0	0.41	4.5	3.1	0.51	0.09
A923.152A	x	x	x	x	x	x	3.2	1.0	0
48 Dune	x	x	x	x	x	x	3.0	0.88	0.21
148 Dune	28:21:51	39:26:34	0.56	0	0.65	2.1	3.1	0.90	0.21
68	x	x	x	x	x	x	2.7	0.49	0.18
290	30:18:52	45:21:34	0.59	0	0.46	2.9	3.5	0.78	0.11
Family 2									
CB-37	35:11:54	nd	0.64	0.22	0.60	6.7	2.3	0.52	0.11
CB-92 (606)	30:20:50	45:25:30	0.61	0.22	0.22	1.4	1.1	1.3	0
CB-28 (605)	37:26:37	42:28:30	0.98	0.23	0.32	1.1	1.4	1.2	0.03
Family 3									
CH 8	41:22:37	37:32:31	1.1	0	0.26	0.61	1.3	1.5	0
CB-35	43:25:32	47:24:29	1.3	0	0.32	1.3	1.0	1.2	nd
CB-8	43:19:38	37:23:40	1.1	0.05	0.25	0.93	1.1	1.7	0.04
Family 4									
CB31-1 (379)	33:14:53	47:22:31	0.62	nd	0.60	0.78	3.3	1.2	nd
CB31-2 (608)	37:20:43	49:16:35	0.87	0.05	0.96	0.97	3.4	1.0	nd
EC 1	nd	nd	0.83	nd	0.18	nd	0.63	1.8	0
ST 15	nd	nd	nd	nd	0.35	1.8	1.9	2.1	0
Parameter	1	2	3	4	5	6	7	8	9

Ratios calculated from peak areas in MRM mass chromatograms as follows:

1. $C_{27} : C_{28} : C_{29} 5\alpha, 14\alpha, 17\alpha$ (H) (20R) steranes
2. $C_{27} : C_{28} : C_{29} 13\beta, 17\alpha$ (H) (20R) diasteranes
3. $C_{27} 5\alpha, 14\alpha, 17\alpha$ (H) cholestane (20R) / $C_{29} 5\alpha, 14\alpha, 17\alpha$ (H) ethylcholestane (20R)
4. $C_{30} 5\alpha, 14\alpha, 17\alpha$ (H) *n*-propylcholestane (20R) / $C_{29} 5\alpha, 14\alpha, 17\alpha$ (H) ethylcholestane (20R)
5. $C_{29} 13\beta, 17\alpha$ (H) diasteranes (20S + 20R) / $C_{29} 5\alpha, 14\alpha, 17\alpha$ (H) steranes (20S + 20R)
6. $C_{30} 17\alpha, 21\beta$ (H) hopane / $C_{29} 5\alpha, 14\alpha, 17\alpha$ (H) steranes (20S + 20R)
7. $C_{27} 18\alpha$ (H)-22,29,30-trisnorhopane / $C_{27} 17\alpha$ (H)-22,29,30-trisnorhopane
8. $C_{29} 17\alpha, 21\beta$ (H) hopane / $C_{30} 17\alpha, 21\beta$ (H) hopane
9. $C_{30} 17\alpha$ (H) diahopane / $C_{30} 17\alpha, 21\beta$ (H) hopane

Table 4 continued.

Sample ID	C ₃₀ Steranes				Triterpanes		
	24- <i>n</i> -propyl	dino	4 α -Me/ 29 ster	ol/ hop	bicad T/ hop	ol/ Σ bicad	Major bicadinanes
Family 1							
28	A	A	nd	0.09	0.66	0.09	T, MeT, R, T1, W
4066	A	A	nd	0.04	nd	nd	T, MeT, T1, R, W
Gelt (318)	A	A	nd	0.08	0.37	nd	T, T1, R, W*
Devil's (317)	A	A	nd	0.24	1.5	nd	T, MeT, T1, R, W
314	A	A	nd	nd	2.9	nd	T, T1, MeT, W
CB-38 (378)	A	A	2.1	0.08	0.48	0.09	T, MeT, T1, R, W
59	A	P?	nd	0.05	nd	nd	x
A923.152A	A	A	nd	nd	2.1	0.31	T, MeT, T1, R, W
48 Dune	A	A	nd	0.06	nd	nd	x
148 Dune	A	A	nd	0.04	1.2	0.03	T, T1, R, W*
68	A	A	nd	0.09	nd	nd	T, MeT, T1, R, W
290	A	P?	nd	0.07	0.46	0.10	T, MeT, T1, R, W
Family 2							
CB-37 (604)	P	P	5.5	0.06	0.22	0.24	T, T1, MeT, R, W
CB-92 (606)	P	P	0.88	0.05	nd	nd	T1, MeT, T, R, W
CB-28 (605)	P	P	0.61	0.05	0.07	0.61	T, MeT, T1, R, W
Family 3							
CH8	nd	nd	nd	x	0.14	x	T, MeT, T1, R, W
CB-35	A	A	0.83	nd	0.08	nd	T1, T, MeT, R, W
V2-6	P	P	nd	x	x	x	x
CB-8	P	P	0.28	0.02	nd	nd	T1, T, R, W*
Family 4							
CB-31-1 (379)	P	P	0.18	0	0	0	A
CB-31-2 (608)	P	P	nd	0	0	0	A
EC 1	nd	nd	nd	0	0	0	A
ST 15	nd	nd	nd	0	0	0	A
Parameter	10	11	12	13	14	15	16

10. 24-*n*-propylcholestane
11. dinosterane (4,23,24-trimethylcholestane)
12. 4 α -methyl-24-ethylcholestane (20*R*) + 4 α -methyl-23,24-dimethylcholestane (20*R*) / 24-ethylcholestane (20*R*)
13. oleanane / C₃₀ 17 α ,21 β (H) hopane
14. bicadinane T / C₃₀ 17 α ,21 β (H) hopane
15. oleanane / bicadinanes T + T1 + R (see Fig. 8)
16. Bicadinanes listed in order of decreasing abundance. Isomers labelled as in Grantham et al. (1983), Alam and Pearson (1990), Summons et al. (1993) and McKirdy et al. (1994). * Methylbicadinane M⁺ 426 \rightarrow *m/z* 383 not recorded.

A = absent

P = present

nd = not determined because peak not integrated or appropriate MRM chromatogram not acquired

x = not determined because compound(s) removed by severe weathering

Table 4. Diagnostic characteristics and source affinity of waxy bitumen families

Family	S wt %	Pr/Ph	BI*	Bot/ <i>n</i> -C ₂₉	Steranes	24- <i>n</i> -Propyl	Dino
1 n = 35	<0.3	0.2–2.6	530–870	0.16–0.86	29 > 27 > 28	A	A
2 n = 17	0.6–2	0.7–1.7	nd	0.15–0.71	29 ≥ 27 > 28	P	P
3 n = 33	1.1–2.6	0.5–1.1	104–173	0.04–0.14	27 > 29 > 28	P or A	P or A
4 n = 7	0.9	0.5–1.4	0	0	29 ≥ 27 > 28	P	P
5 n = 5	0.2	4–11	nd	nd	nd	nd	nd
	Hop/Ster	Oleanane	Bicadinanes	CV**	Source affinity		
1	1.7–4.5	P	P	<0.47	Lacustrine shale Anoxic–suboxic Cenozoic		
2	1.1–6.7	P	P	<0.47	Lacustrine shale > marine carbonate Anoxic–suboxic Cenozoic		
3	0.6–1.3	P	P	<0.47	Marine carbonate > lacustrine shale Anoxic/euxinic Cenozoic		
4	0.8–1.8	A	A	<0.47	Marine carbonate or calcareous shale Anoxic–suboxic Mesozoic		
5	nd	nd	nd	>0.47	Non-marine (?) coal Oxic		

Pr/Ph = pristane/phytane

Bot = C₃₄ botryococcane

* Botryococcane Index = C₃₄ botryococcane x 100 / ∑C₃₇–C₄₀ head-to-head isoprenoid alkanes (Seifert and Moldowan, 1981)

** CV = –2.538¹³C_{sat} + 2.228¹³C_{aro} – 11.65 (Sofer, 1984)

Table 5. Comparison of oil family classifications for Australian waxy coastal bitumens

Present study	Previous studies		Key biomarker characteristics	Source affinity
South Australia & western Victoria	Northern Territory	Western Australia		
Family 1	Family 1	Group 1	Botryococcane $C_{29} > C_{27} > C_{28}$ sterane C_{30} 4 α -methylsterane Bicadinanes	Lacustrine shale Cenozoic
Family 2	NR	NR	As for Family 1, plus 24- <i>n</i> -propylcholestane & dinosterane	Lacustrine shale & marine carbonate Cenozoic
NR	NR	Groups 2 & 3	Bicadinanes abundant Oleanane / C_{30} hopane >0.1	Deltaic Cenozoic
Family 3	Family 2	Group 4	C_{29}/C_{30} hopane >1 C_{29} diasterane/sterane <0.4 Bicadinanes & oleanane contents very low	Marine carbonate Cenozoic
NR	NR	Groups 5, 6 & 7	As for Group 4, but with significant oleanane & variable bicadinane contents	Marine carbonate Cenozoic
NR	NR	Group 8	$C_{27} > C_{29} > C_{28}$ sterane Bicadinanes & oleanane absent	Marine siliciclastic Mesozoic
Family 4	NR	NR	$C_{29} > C_{27} > C_{28}$ sterane C_{29}/C_{30} hopane >1 Bicadinanes & oleanane absent	Marine carbonate & calcareous shale Mesozoic
Family 5	NR	NR	Pr/Ph ≥ 4 CV >0.47	Non-marine paludal (coal)
Reference:	Summons et al. (1993)	Alexander et al. (1994)		

NR = not represented

Table 6. Biomarker characteristics of West Indonesian oil families

Family	Key biomarker signatures	Basins	Source lithofacies & age	Data sources
Fresh/deep lacustrine	Botryococcane C ₃₀ Tetracyclic polyprenoid (TPP)* C ₃₀ 4-methylsteranes C ₂₆ /C ₂₅ tricyclic terpane >>1 Ts/Tm >1 Oleanane Bicadinanes Hopanes > Steranes Pristane/phytane = 1.5–3 24- <i>n</i> -Propylcholestane absent	Central Sumatra South Sumatra Sunda	Shale Cenozoic	1, 2, 3, 4, 7
Brackish lacustrine	As above, but with minor or no botryococcane & higher abundances of terrigenous triterpanes (oleanane, bicadinanes) Gammacerane present	Central Sumatra South Sumatra Ombilin East Java	Shale Cenozoic	1, 2, 3, 4, 7
Terrigenous (fluvio-deltaic)	C ₂₉ sterane dominant Bicadinanes Oleanane C ₂₆ /C ₂₅ tricyclic terpane <1 Pristane/phytane >3	North Sumatra Central Sumatra South Sumatra West Natuna Northwest Java	Coal, carbonaceous shale Cenozoic	2, 3, 4, 5
Marine	C ₂₇ sterane dominant 24- <i>n</i> -Propylcholestane present C ₂₉ /C ₃₀ hopane >1 Pristane/phytane = 1–2.5	North Sumatra East Natuna	Carbonate Cenozoic	2, 3, 4, 6

* Reported as unknown 'lacustrine classifier' by ten Haven & Schiefelbein (1995); subsequently identified by Holba et al. (2000)

Key to data sources:

1. Seifert & Moldowan (1981)
2. Robinson (1987)
3. ten Haven & Schiefelbein (1995)
4. Schiefelbein & Cameron (1997)
5. van Aarssen et al. (1992)
6. Subroto et al. (1992)
7. Hwang et al. (2002)

Figure captions

Figure 1. Map showing the location of waxy bitumen stranding sites in South Australia and western Victoria.

Figure 2. Bulk composition of coastal waxy bitumens according to classification of Tissot and Welte (1984).

Figure 3. Gas chromatograms of the saturated hydrocarbon fractions in representative waxy coastal bitumens. Key: Pr = pristane; Ph = phytane; * = C₃₄ botryococcane; IS = internal standard, 3-methylheneicosane; numbers refer to chain length of *n*-alkanes.

Figure 4. High temperature gas chromatograms of the saturated hydrocarbon fractions in representative waxy bitumens: Family 1 (a-e), showing effects of increasing weathering; Family 4 (f). Key as for Fig. 3.

Figure 5. Carbon isotopic compositions of C₁₂₊ saturated and aromatic hydrocarbon fractions in coastal waxy bitumens and asphaltites from southern Australia. The oblique line corresponds to a CV value of 0.47, where $CV = -2.53\delta^{13}C_{\text{sat}} + 2.22\delta^{13}C_{\text{aro}} - 11.65$, which according to Sofer (1984; reinterpreted by McKirdy et al., 1986) separates oils of land-plant origin from those of algal origin.

Figure 6. MRM chromatograms showing the distribution of A) triterpanes ($M^+ \rightarrow 191$) and B) steranes ($M^+ \rightarrow 217$ and 231) in a Family 2 waxy bitumen (sample CB-28).

Figure 7. MRM chromatograms showing the distribution of A) triterpanes ($M^+ \rightarrow 191$) and B) steranes ($M^+ \rightarrow 217$ and 231) in a Family 4 waxy bitumen (sample CB-31-2).

Figure 8. MRM chromatograms showing the contrasting bicadinane and methylbicadinane signatures of representative Family 1 and Family 2 waxy bitumens.

Figure 9. Map of selected sedimentary basins within the Indonesian Archipelago showing the location of oil seeps (after Livsey et al., 1992; Macgregor, 1995; Noble et al., 2009) and pathways of the Indonesian and South China Sea throughflows (after Gordon, 2005; Gordon et al., 2012; Susanto et al., 2016). Sedimentary basin outlines are from Robertson (2017).

Figure 10. Partial MRM 412 → 369 chromatograms illustrating the C₃₀ bicadinane distributions in a Family 1 waxy bitumen and crude oils from Indonesia and Papua New Guinea.

SUPPLEMENTARY MATERIAL

Table A1. Waxy bitumen inventory (locations listed from west to east), bulk properties and classification

Sample ID	Location	Date sample collected	Specific gravity °API	Pour point °C	Sulfur content wt %	Data source
<i>Family 1</i>						
376	Anxious Bay, Eyre Pen., SA	10.10.1992	nd	nd	nd	1
CB-71	Sheringa Beach, Eyre Pen., SA	6.9.1983	nd	nd	0.2	4,8
Coles Point	Coles Point, Eyre Pen., SA	10.12.1990	nd	nd	nd	1
CB-67	Avoid Bay, Eyre Pen., SA	6.9.1983	nd	nd	0.1	4,8
A923.152A	Thistle Island, offshore Eyre Pen., SA	19.4.1984	nd	nd	nd	1
CB-51	Gym Beach, Yorke Pen., SA	6.9.1983	nd	73	0.2	4,8
296	Ravine des Casoars, K.I., SA	20.9.1991	nd	nd	nd	1
79	West Bay, K.I., SA	15.10.1990	nd	nd	nd	1
CB-84	Sandy River Beach, K.I., SA	14.12.1983	35	68	0.2	4,8
290	Hanson Bay, K.I., SA	19.9.1991	27	nd	0.1	1
VB 3	Vivonne Bay, K.I., SA	14.9.1989	nd	nd	nd	1
287	Bales Bay, K.I., SA	19.9.1991	nd	nd	nd	1
Pen 8	Pennington Bay, K.I., SA	15.9.1989	nd	nd	nd	1
CB-77	Pennington Bay, K.I., SA	12.12.1983	38	68	0.1	4,8
68	Pennington Bay, K.I., SA	14.10.1990	32	nd	0.1	1
67	Wilson River, K.I., SA	14.10.1990	22	nd	nd	1
192	False Cape, K.I., SA	22.2.1991	nd	nd	nd	1
248	Windmill Bay, K.I., SA	30.5.1991	nd	nd	nd	1
A923.158	Encounter Bay, Port Elliot, SA	26.5.1990	nd	nd	nd	1
59	The Granites, Long Beach, SA	1.10.1990	nd	nd	0.2	1
CB-38 (378)	5 km N of The Granites, Long Beach, Kingston, SA	30.8.1983	25	68	0.3	1,3,5,8
43	Beachport C.P., SA	30.9.1990	nd	nd	nd	1
48 Dune	Beachport C.P., SA	30.9.1990	11	nd	nd	1
50	Beachport C.P., SA	30.9.1990	36	nd	0.2	1
148 Dune	Beachport C.P., SA	28.11.1990	nd	nd	nd	1
358	Beachport C.P., SA	26.9.1990	nd	nd	nd	1
Gelt (318)	Geltwood Beach, Canunda N.P., SA	1960	nd	nd	nd	1
4066	Geltwood Beach, Canunda N.P., SA	1961	nd	nd	0.1	1
Devil's (317)	Devils Gap, Canunda N.P., SA	1961	nd	nd	<0.3	1
17	Canunda N.P., SA	28.9.1990	nd	nd	nd	1
28	Canunda N.P., SA	29.9.1990	40	nd	0.3	1
DB 1	Descartes Bay, Vic	17.4.1990	17	nd	nd	1
314	Descartes Bay, Vic	24.9.1991	nd	nd	nd	1
314HW	Descartes Bay, Vic	24.9.1991	nd	nd	nd	1
CB-1	Bridgewater Bay, Vic	26.8.1983	27	71	0.3	3,5,8
<i>Family 2A</i>						
A923.157	Corny Point, Yorke Pen., SA	3.199	nd	nd	nd	1,7
CB-36	Wright Bay, SA	30.8.1983	nd	nd	1	6
CB-37	The Granites, Long Beach, SA	30.8.1983	nd	nd	0.6	1,6,8
CB-34	Eve's Cove, Cape Lannes, SA	29.8.1983	nd	nd	0.8	6

CB-30	Beachport C.P., SA	29.8.1983	nd	nd	1.9	6
CB-16	Canunda N.P., SA	28.8.1983	nd	nd	1.5	6
CB-21	Canunda N.P., SA	28.8.1983	nd	nd	1.8	6
CB-11	Douglas Point, SA	27.8.1983	nd	nd	1.7	3,6
CB-93	Cape Bridgewater, Vic	11.1991	nd	nd	1.3	6
Family 2B						
CB-52	Daly Head, Yorke Pen., SA	6.9.1983	nd	74	1.4	4
CB-39	The Granites, Long Beach, SA	30.8.1983	nd	nd	1.3	6
CB-28 (605)	Beachport C.P., SA	29.8.1983	17	72	1.3	1,3,6,8
CB-29	Beachport C.P., SA	29.8.1983	nd	nd	1.3	6
CB-92 (606)	Geltwood Beach, Canunda N.P., SA	8.1961	nd	nd	2	1,3,6,8
CB-23	Canunda N.P., SA	28.8.1983	23	61	1.2	3,6
CB-25	Canunda N.P., SA	28.8.1983	nd	nd	0.8	6
CB-14	Admella Dunes, SA	27.8.1983	20	57	1.3	3,6
Family 3A						
CB-73	Petrel Cove, Fleurieu Pen., SA	12.12.1983	nd	nd	2.6	4,8
Pen 6	Pennington Bay, K.I., SA	15.9.1989	nd	nd	nd	1
Bitumen B	Unspecified, K.I., SA	3.1990	nd	nd	nd	1,7
CB-35 (607)	Guichen Bay, Robe, SA	30.8.1983	17	59	2.3	1,3,6,8
A923.99	Canunda N.P., SA	8.1983	nd	nd	nd	1
CB-17	Canunda N.P., SA	28.8.1983	nd	nd	1.1	6
12	Canunda N.P., SA	27.9.1990	17	nd	nd	1
200	Canunda N.P., SA	26.2.1991	nd	nd	nd	1
33	Beachport C.P., SA	30.9.1990	13	nd	nd	1
349	Canunda N.P., SA	26.9.1990	nd	nd	nd	1
CB-12	Douglas Point, SA	27.8.1983	16	57	2	3,6,8
3	Nene Valley C.P., SA	27.9.1990	nd	nd	nd	1
91	Picaninnie Ponds, SA	25.11.1990	nd	nd	nd	1
CB-9	Cape Northumberland, SA	26.8.1983	21	59	1.9	3,6
CB-10	Cape Northumberland, SA	26.8.1983	24	57	1.8	3,5,8
GB 22	Port Campbell N.P., SA	16.4.1990	nd	nd	nd	1
93	Nelson, Discovery Bay, Vic	25.11.1990	nd	nd	1.3	1,6
CB-7	Nobles Rocks, Discovery Bay, Vic	26.8.1983	17	58	1.9	3,6,8
CB-2	Bridgewater Bay, Vic	26.8.1983	nd	nd	nd	6
A923.198	Lake Yambuk, Vic	5.1.1991	nd	nd	nd	1
CH 1	Childers Cove, Vic	14.9.1990	nd	nd	nd	1
CH 3	Childers Cove, Vic	14.4.1990	nd	nd	nd	1
CH 8	Childers Cove, Vic	14.4.1990	20	nd	nd	1
V2-6	Castle Cove, Vic	16.4.1991	nd	nd	nd	1
ST 16	Station Beach, Cape Otway, Vic	14.4.1990	nd	nd	nd	1
CB-376	Lion Headland, Vic	13.10.1985	13	57	2.1	5,8
Family 3B						
CB-20	Canunda N.P., SA	28.8.1983	21	59	2	3,6
CB-18	Geltwood Beach, Canunda N.P., SA	28.8.1983	20	58	2.6	3,8
CB-15	Admella Dunes, SA	27.8.1983	19	55	1.8	3,6
CB-13	Nene Valley C.P., SA	27.8.1983	18	59	1.4	3,5,8
CB-8 (380)	Nelson, Vic	26.8.1983	16	58	2.4	3,5,8

CB-5	Swan Lake, Discovery Bay, Vic	26.8.1983	19	57	1.7	3,6
CB-3	Descartes Bay, Vic	26.8.1983	26	51	2.1	3,5,8
CB-4	Descartes Bay, Vic	26.8.1983	15	53	1.6	3,5,8
Family 4						
EC 1	Eve's Cove, Cape Lannes, SA	1979	10	nd	nd	1
CB-31-1 (379)	Beachport C.P., SA	29.8.1983	21	66	0.9	1,3,8
CB-31-2 (608)	Beachport C.P., SA	29.8.1983	nd	nd	nd	1,6
257	Beachport C.P., SA	26.9.1991	34	nd	nd	1
323	Nene Valley C.P., SA	25.9.1991	nd	nd	nd	1
312	Bridgewater Bay, Vic	24.9.1991	nd	nd	nd	1
ST 15	Station Beach, Cape Otway, Vic	14.4.1990	nd	nd	nd	1
Family 5						
295	Ravine des Casoars, K.I., SA	20.9.1991	nd	nd	0.2	1
375	D'Estrees Bay, K.I., SA	16.2.1992	nd	nd	nd	1
363	Little Dip C.P., SA	26.9.1991	nd	nd	nd	1
357	Beachport C.P., SA	26.9.1991	nd	nd	0.2	1
313	Descartes Bay, Vic	24.9.1991	nd	nd	nd	1
Unassigned						
MH6	Sleaford Bay, Eyre Pen., SA	16.8.1961	21	66	<0.3	2
CB-64	Sleaford Bay, Eyre Pen., SA	17.9. 1983	nd	62	0.4	4
CB-48	Sturt Bay, Yorke Pen., SA	6.9.1983	nd	nd	1.3	4
CB-89	Ravine des Casoars, K.I., SA	15.12.1983	nd	67	1.2	4
CB-82	West Bay, K.I., SA	14.12.1983	nd	67	0.5	4
CB-87	Rocky River, K.I., SA	14.12.1983	nd	78	0.2	4
MH1	Geltwood Beach, Canunda N.P., SA	6.1961	28	43	<0.3	2
MH2	Geltwood Beach, Canunda N.P., SA	8.1961	20	nd	<0.3	2
MH4	Devils Gap, Canunda N.P., SA	16.8.1961	19	nd	<0.3	2
MH5	Cape Northumberland, SA	23.8.1963	28	58	<0.3	2
MH3	Discovery Bay, Vic	8.10.1961	27	82	<0.3	2
MH7	Bridgewater Bay, Vic	9.10.1961	28	71	<0.3	2

C.P. = Conservation Park

N.P. = National Park

K.I. = Kangaroo Island

Pen. = Peninsula

nd = not determined

Key to data sources:

1. This study 2. McKirdy and Horvath (1976)

5. McKirdy (1985) 6. Tewksbury (1989)

3. McKirdy (1984a)

7. Watson (1990)

4. McKirdy (1984b)

8. McKirdy et al. (1994)

Table A2. Bulk composition, oil type and degree of weathering of waxy bitumens

Sample ID	Par %	Naph %	Arom %	Res %	Asph %	Oil type	Degree of weathering*
<i>Family 1</i>							
28 interior	52.8	28.0	11.1	4.2	3.9	P	0
28 intermediate	— 68.7 —		15.2	7.2	8.9	nd	0
28 exterior	— 64.6 —		17.2	9.1	9.1	nd	0
4066	56.7	23.3	10.0	4.8	5.4	P	0
192	39.3	29.5	7.8	4.7	18.8	P-N	0
Gelt (318)	— 72.6 —		9.2	— 18.2 —		nd	0
Devils (317)	— 20.7 —		24.4	— 54.9 —		nd	0.5
314	39.3	28.2	14.3	9.2	9.0	P-N	1
314HW	— 61.1 —		10.5	13.8	14.6	nd	1
CB-77	35.2	21.9	3.0	34.7	5.2	P-N	1
Coles Point	36.2	26.5	8.8	5.8	22.6	P-N	1
CB-84	29.5	28.4	0.5	36.2	5.4	P-N	1
VB3	— 61.4 —		8.6	7.4	22.6	nd	1
CB-67	25.1	21.3	6.0	42.7	4.9	Ar-I	1
CB-1	38.0	15.2	8.3	34.8	3.7	P-N	2
CB-71	30.8	26.3	4.1	34.8	4.0	P-N	2
CB-38 (378)	45.6	17.7	8.6	23.9	4.2	P	2
17	— 68.0 —		6.8	2.5	22.7	nd	2
CB-51	22.3	24.3	2.3	43.6	7.5	Ar-I	2
50	41.3	21.8	10.5	5.1	21.3	P	2
59	32.8	28.4	14.7	8.6	15.5	P-N	2
79	48.3	22.3	12.0	4.9	12.6	P	2
376	— 59.0 —		9.6	5.9	25.5	nd	2
A923.152A	53.9	18.1	10.1	8.6	9.4	P	3
DB 1	44.4	20.5	15.8	5.5	13.8	P	3
Pen 8	33.7	20.6	16.4	8.1	21.2	P-N	3
48 Dune	25.9	21.9	4.8	2.1	45.3	Ar-I	3
287	27.6	26.3	21.7	8.8	15.6	P-N	4
296	— 51.2 —		21.1	10.0	17.8	nd	4
358	30.3	30.3	18.6	7.9	13.1	P-N	4
148 Dune	— 65.8 —		28.5	11.6	1.2	nd	4
67	— 50.7 —		18.4	12.1	18.9	nd	4
A923.158	— 59.3 —		20.0	9.0	11.8	nd	4
248	34.8	20.4	16.2	11.1	17.6	P-N	4
68	28.2	27.9	21.4	8.0	14.6	P-N	4
43	— 50.6 —		14.9	5.0	29.5	nd	4
290	— 47.0 —		22.5	11.0	19.5	nd	5
<i>Family 2A</i>							
CB-93	26.3	21.4	20.3	21.7	10.2	Ar-I	0.5
CB-30	14.2	32.8	12.6	29.5	10.9	Ar-I	0.5
CB-37	22.1	36.1	6.8	26.3	8.7	P-N	1+
CB-16	20.6	27.6	10.7	24.4	16.7	Ar-I	2
CB-36	36.6	14.2	9.0	35.5	4.7	P-N	4
CB-34	13.5	34.6	9.0	33.7	9.2	Ar-I	4
A923.157	23.5	20.5	8.0	43.5	4.6	Ar-I	5
CB-11	18.9	23.6	18.2	26.3	13.1	Ar-I	5
CB-21	19.7	21.1	9.1	33.2	16.9	Ar-I	5
<i>Family 2B</i>							
CB-92 (606)	27.9	21.7	12.2	25.1	13.1	Ar-I	0.5
CB-28 (605)	25.9	7.7	11.1	29.7	25.7	Ar-I	0.5

CB-23	14.8	36.0	14.2	22.2	5.7	P-N	2
CB-14	19.6	42.9	15.9	18.1	3.5	P-N	2
CB-29	29.4	38.6	8.0	16.7	7.3	P-N	2
CB-25	26.7	31.9	9.7	24.1	7.6	P-N	2
CB-39	14.5	31.3	10.0	31.1	13.1	Ar-I	2
CB-52	24.0	20.2	6.0	40.2	9.6	Ar-I	2+

Family 3A

CB-10	52.0	14.0	12.9	18.2	2.9	P	0
A923.99	— 64.2 —		17.2	10.2	8.4	nd	0.5
CB-17	32.7	30.1	14.2	16.5	6.5	P-N	0.5
12	46.2	20.4	17.5	7.6	8.2	P	0.5
CH 8	42.2	26.8	19.6	5.1	6.3	P	0.5
CB-376	22.0	44.3	18.3	8.1	7.3	N	0.5
CH 1	30.3	30.3	22.7	9.8	6.9	P-N	0.5
Bitumen B	17.6	22.5	6.5	29.7	23.7	Ar-I	0.5
A923.198	30.0	26.8	18.5	18.5	6.3	P-N	0.5
200	— 49.9 —		22.1	22.0	12.6	nd	0.5
93	30.0	31.5	21.5	21.4	14.6	nd	0.5
33	34.6	29.0	14.6	15.5	6.2	P-N	0.5
3	37.4	22.8	17.9	13.2	8.7	P-N	0.5
CB-35 (607)	— 53.9 —		14.8	25.8	5.5	nd	2
CB-12	19.6	42.9	15.9	18.1	3.5	P-N	2
CB-7	13.2	29.4	14.4	36.1	6.9	Ar-I	2
CB-9	20.2	32.7	15.6	24.5	7.0	P-N	2
CH-3	17.8	15.4	40.2	16.5	10.1	Ar-I	2
Pen 6	28.0	37.5	18.4	11.7	4.8	P-N	2
349	— 52.5 —		24.6	9.8	13.0	nd	2
CB-2	16.0	27.8	21.6	23.7	10.9	Ar-I	2
V2-6	38.0	24.1	18.5	12.4	7.0	P-N	2
GB 22	29.3	27.4	22.2	14.6	6.6	P-N	2
ST 16	— 52.1 —		20.3	20.3	7.4	nd	4
91	25.8	28.8	23.3	13.8	8.3	P-N	5

Family 3B

CB-4	42.5	12.4	13.8	24.2	7.1	P	0.5
CB-18	44.8	16.3	10.0	23.0	5.9	P	0.5
CB-8 (380)	28.9	23.1	14.5	28.0	5.5	P-N	0.5
CB-20	34.6	26.0	9.5	24.2	5.7	P-N	2
CB-5	24.7	33.9	17.0	19.0	5.4	P-N	2
CB-3	23.8	20.8	14.4	33.9	7.1	Ar-I	2
CB-15	17.1	28.9	15.7	34.6	4.1	Ar-I	2
CB-13	28.9	4.2	13.3	47.4	6.2	Ar-I	2

Family 4

CB-31-1 (379)	62.7	11.1	7.2	14.9	4.1	P	0
CB-31-2 (608)	36.4	23.1	7.7	29.2	3.6	P-N	0
Eves Cove	36.8	14.3	12.9	12.8	23.1	P-N	0.5
312	34.8	19.6	11.8	11.7	22.1	P-N	3
323	37.4	30.7	12.5	7.0	12.5	P-N	3
257	45.0	18.1	11.5	7.0	18.4	P	3
ST15	— 24.5 —		6.5	4.2	64.9	nd	4

Family 5

357	— 72.7 —		16.0	7.4	3.9	nd	0.5
313	— 68.8 —		17.3	11.1	2.8	nd	0.5
375	45.2	26.1	18.5	5.6	4.8	P	0.5
363	— 74.7 —		12.9	9.6	2.8	nd	0.5
295	57.4	17.2	16.7	4.9	3.6	P	0.5

* The degree of weathering has been determined from the GC-FID trace of the saturated hydrocarbon fraction. Numerical values are defined in Table A.2.

Par = paraffins (*n*- + *iso*-alkanes)

Naph = naphthenes (multibranched + cycloalkanes)

Arom = aromatic hydrocarbons

Res = resins

Asph = asphaltenes

P = paraffinic

P-N = paraffinic-naphthenic

Ar-I = aromatic-intermediate

nd = not determined

Table A3. Extent to which weathering has altered the composition of a waxy bitumen based on the appearance of its saturated hydrocarbon chromatogram

Degree of weathering	Compounds removed	Comments
<i>Unaffected</i>		
0	Evaporative loss of <i>n</i> -alkanes up to C ₁₂	No evidence of biodegradation Light ends lost during sample preparation
<i>Slight</i>		
0.5	Depletion of <i>n</i> -alkanes up to C ₁₅	
1	Removal of <i>n</i> -alkanes up to C ₁₅	Pristane < <i>n</i> -C ₁₇ Phytane < <i>n</i> -C ₁₈
2	Depletion of <i>n</i> -alkanes up to C ₂₁	Pristane > <i>n</i> -C ₁₇ Phytane > <i>n</i> -C ₁₈
3	Removal of <i>n</i> -alkanes up to C ₂₁ , pristane & phytane	Appearance of UCM
<i>Moderate</i>		
4	Removal of <i>n</i> -alkanes up to C ₂₃	
5	Removal of <i>n</i> -alkanes up to C ₂₆	

Table A4. Alkane parameters and bulk isotopic composition of waxy bitumen families

Sample ID*	<i>n</i> -Alkane profile range maxima		Pr/Ph	Pr/ <i>n</i> -C ₁₇	Ph/ <i>n</i> -C ₁₈	Bot/ <i>n</i> -C ₂₉	BI	δ ¹³ C _{sat} ‰	δ ¹³ C _{aro} ‰
Family 1									
28 interior	10 – 40	17, 25	2.4	0.38	0.16	0.24	nd	–25.9	–24.8
28 intermediate	10 – 39	15, 27	2.9	0.35	0.13	0.26	nd	nd	nd
28 exterior	10 – 39	15, 27	2.9	0.35	0.13	0.29	nd	nd	nd
4066	12 – 44	17, 27	2.2	0.29	0.14	0.28	nd	–25.9	–24.5
192	12 – 40	20, 29	2.6	0.38	0.13	0.31	nd	nd	nd
Gelt (318)	8 – 40	17, 28	2.3	0.33	0.15	0.23	nd	nd	nd
Devils (317)	8 – 40	19	1.3	0.24	0.13	0.18	nd	nd	nd
314	15 – 40	27	1.9	0.50	0.21	0.28	nd	nd	nd
CB-77	16 – 37	23	0.56	0.90	0.18	0.28	530	–25.6	nd
Coles Point	16 – 42	27	1.4	0.81	0.40	0.39	nd	nd	nd
CB-84	15 – 38	27	1.2	1.8	0.75	0.30	530	–25.4	nd
VB3	16 – 42	27	1.6	1.3	0.57	0.33	nd	nd	nd
CB-67	15 – 39	29	0.71	0.40	0.22	0.67	870	–25.4	–24.6
CB-1	14 – 40	27	2.2	2.6	1.0	0.27	740	–25.1	–24.3
CB-71	15 – 40	27	0.22	0.93	2.4	0.35	600	–25.5	–24.5
CB-38 (378)	16 – 40	27	2.0	7.3	3.8	0.45	560	–25.2	–24.2
17	18 – 43	27	1.8	x	x	0.41	nd	nd	nd
CB-51	17 – 39	29	0.74	3.9	5.9	0.50	580	–25.1	–24.4
50	17 – 44	29	1.4	4.1	x	0.31	nd	–25.6	–24.7
59	22 – 44	27	1.1	x	x	0.36	nd	–25.5	–24.8
79	16 – 42	27	0.70	x	1.4	0.16	nd	nd	nd
376	18 – 42	27	x	x	x	0.40	nd	nd	nd
A923.152A	19 – 44	28	x	x	x	0.14	nd	nd	nd
DB1	21 – 50	35	x	x	x	0.56	nd	–26.4	–25.5
Pen 8	22 – 44	29	x	x	x	0.79	nd	nd	nd
48 Dune	22 – 44	29	x	x	x	0.36	nd	–26.0	–24.6
287	23 – 44	29	x	x	x	0.62	nd	nd	nd
296	22 – 44	20	x	x	x	0.70	nd	nd	nd
358	23 – 42	29	x	x	x	0.24	nd	nd	nd
148 Dune	24 – 44	29	x	x	x	0.65	nd	nd	nd
67	23 – 44	29	x	x	x	0.18	nd	–25.9	–24.8
A923.158	24 – 44	29	x	x	x	0.57	nd	nd	nd
248	25 – 44	29	x	x	x	0.86	nd	–25.5	–24.4
68	25 – 44	31	x	x	x	0.24	nd	–26.2	–24.5
43	25 – 43	29	x	x	x	0.65	nd	nd	nd
290	25 – 44	31	x	x	x	0.68	nd	–26.0	–24.7
Family 2A									
CB-93	9 – 40	17, 29	1.1	0.47	0.49	0.25	nd	–26.9	–26.4
CB-30	11 – 40	16, 29	1.2	0.58	0.52	0.50	nd	–26.1	–26.2
CB-37	9 – 40	29	1.7	0.99	0.78	0.63	nd	–26.0	–25.2
CB-16	9 – 40	11, 28	1.0	0.97	2.0	0.53	nd	–26.0	–25.4
CB-36	23 – 40	29	x	x	x	0.48	nd	–26.0	–25.1
CB-34	24 – 40	29	x	x	x	0.56	nd	–25.9	–24.3
A923.157	24 – 36	28	x	x	x	0.19	nd	nd	nd
CB-11	25 – 40	32	x	x	x	0.15	nd	–27.2	–26.5
CB-21	24 – 40	29	x	x	x	0.71	nd	–26.0	–25.0
Family 2B									
CB-92 (606)	13 – 39	29	0.70	0.71	0.73	0.19	nd	–26.9	–27.6
CB-28 (605)	14 – 40	17, 27	1.2	0.93	0.82	0.24	nd	–27.2	–27.1
CB-23	14 – 40	20, 30	1.0	0.84	1.1	nd	nd	–27.2	–26.8

CB-14	15 – 40	29	0.98	2.8	2.8	0.38	nd	–26.0	–24.9
CB-29	17 – 40	29	0.70	2.4	0.97	0.50	nd	–26.0	–25.3
CB-25	15 – 40	27	0.90	1.3	1.1	0.42	nd	–25.6	–24.9
CB-39	9 – 40	29	0.88	1.4	2.0	0.42	nd	–25.9	–26.2
CB-52	14 – 41	27	0.95	6.2	2.7	0.28	460	–25.7	–25.5
<i>Family 3A</i>									
CB-10	12 – 40	18, 30	1.0	0.57	0.54	0.08	113	–26.4	–26.0
A923.99	13 – 42	18, 30	1.1	0.32	0.30	0.05	nd	nd	nd
CB-73	14 – 40	18, 29	1.5	0.30	0.19	0.22	1200	–26.6	–25.7
CB-17	12 – 40	19, 29	0.97	0.46	0.52	0.09	nd	–26.9	–26.4
12	13 – 44	32	0.95	0.56	0.58	0.06	nd	–26.5	–26.5
CH 8	13 – 44	34	0.77	0.41	0.41	0.07	nd	–26.7	–26.4
CB-376	13 – 44	19, 31	0.81	0.90	0.79	0.05	129	–26.7	–26.4
CH1	13 – 44	32	1.0	0.78	0.82	0.09	nd	nd	nd
Bitumen B	15 – 36	29	0.56	0.25	0.71	0.12	nd	nd	nd
A923.198	14 – 44	32	0.98	0.96	1.0	0.06	nd	nd	nd
200	14 – 42	30	0.82	0.87	1.0	0.06	nd	nd	nd
93	13 – 42	31	0.88	0.94	1.0	0.06	nd	nd	nd
33	14 – 42	32	0.96	1.4	1.2	0.05	nd	nd	nd
3	16 – 43	32	0.85	0.87	0.83	0.04	nd	nd	nd
CB-35 (607)	14 – 41	18, 31	0.99	1.6	1.6	0.07	nd	–27.1	–25.4
CB-12	14 – 40	31	0.96	2.4	2.2	0.06	nd	–26.9	–26.6
CB-7	13 – 40	30	0.77	4.9	5.7	0.08	nd	–27.9	–26.5
CB-9	16 – 40	31	0.88	5.5	4.6	0.07	nd	–27.0	–25.2
CH-3	14 – 42	32	0.93	1.6	2.8	0.06	nd	nd	nd
Pen 6	17 – 42	31	0.82	2.9	4.0	0.09	nd	nd	nd
349	16 – 40	30	0.61	3.2	4.1	0.11	nd	nd	nd
CB-2	23 – 40	32	0.73	x	x	0.14	nd	–27.2	–26.5
V2-6	18 – 44	33	0.55	1.0	x	0.11	nd	–26.9	–26.3
GB 22	22 – 44	32	x	x	x	0.06	nd	nd	nd
ST 16	24 – 44	32	0.49	x	3.8	0.09	nd	–26.7	–26.3
91	24 – 44	32	x	x	x	0.06	nd	nd	nd
<i>Family 3B</i>									
CB-4	14 – 40	18, 31	1.1	0.79	0.67	0.04	137	–26.4	–26.0
CB-18	13 – 40	18, 30	1.0	0.99	0.95	0.06	104	–26.6	–26.1
CB-8 (380)	14 – 40	18, 31	0.90	0.93	0.93	0.05	137	–26.1	–26.0
CB-20	14 – 40	19, 30	0.88	1.1	1.2	0.07	nd	nd	nd
CB-5	14 – 40	18, 30	0.88	1.3	1.2	0.06	nd	–26.8	–26.3
CB-3	14 – 40	19, 31	1.0	1.6	1.6	0.06	173	–26.5	–25.9
CB-15	14 – 40	18, 31	0.88	2.5	2.4	0.10	nd	nd	nd
CB-13	17 – 40	31	0.86	7.2	7.7	0.06	119	–26.2	–25.9
<i>Family 4</i>									
CB-31-1 (379)	14 – 40	19, 31	0.49	0.13	0.19	[0.02]	[10]	–26.1	–25.3
CB-31-2 (608)	14 – 40	19, 31	0.90	0.13	0.19	0	0	nd	nd
EC 1	13 – 44	18, 34	0.77	0.77	0.41	0	0	–27.4	–26.3
312	20 – 40	34	1.4	x	x	0	0	nd	nd
323	22 – 40	27	x	x	x	0	0	–25.7	–25.0
257	22 – 44	35	x	x	x	0	0	–26.3	–25.6
ST 15	25 – 50	39	x	x	x	0	0	–26.2	–26.0
<i>Family 5</i>									
357	11 – 40	17, 27	4.0	0.62	0.15	nd	nd	–28.5	–26.9
313	11 – 40	15, 27	5.1	0.53	0.12	nd	nd	–28.4	–26.8
375	10 – 40	15, 27	11.1	0.51	0.07	nd	nd	nd	nd
363	11 – 40	16, 27	4.0	0.48	0.14	nd	nd	nd	nd
295	11 – 40	15, 27	4.4	0.59	0.14	nd	nd	–28.5	–26.9

*Samples in each family are listed in order of increasing degree of weathering

x = compounds removed by weathering

na = not applicable

nd = not determined

[] = unreliable

Pr = pristane

Ph = phytane

Bot = C₃₄ botryococcane

BI = botryococcane index (Seifert and Moldowan, 1981)