

2018-09-05

Functional Imagery Training versus Motivational Interviewing for Weight Loss: A randomised controlled trial of brief individual interventions for overweight and obesity

Solbrig, L

<http://hdl.handle.net/10026.1/11340>

10.1038/s41366-018-0122-1

International Journal of Obesity

Nature Publishing Group

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

**Functional Imagery Training versus Motivational Interviewing for Weight Loss:
A randomised controlled trial of brief individual interventions for overweight and
obesity**

Linda Solbrig^{1,2}, Ben Whalley¹, David J. Kavanagh^{3,4}, Jon May¹, Tracey Parkin⁵, Ray Jones⁶,
*Jackie Andrade¹

¹School of Psychology, Cognition Institute, University of Plymouth

²NIHR CLAHRC South-West Peninsula

³Institute for Health & Biomedical Innovation, Centre for Children’s Health Research

⁴School of Psychology & Counselling, Queensland University of Technology

⁵School of Health Professions (Dietetics), University of Plymouth

⁶School of Nursing & Midwifery, University of Plymouth

Word count: 4000

Keywords: Cognition; Counseling; Diet, reducing; Motivation; Motivational Interviewing; Imagery; Weight loss

*Address for correspondence: Professor Jackie Andrade, School of Psychology, Cognition Institute, University of Plymouth, Drake Circus, Plymouth, PL4 8AA, UK

Phone: +44 1752 584807

Email: jackie.andrade@plymouth.ac.uk

28 **Funding statement:** This research was funded by the National Institute for Health
29 Research (NIHR) Collaboration for Leadership in Applied Health Research and Care
30 South West Peninsula. The views expressed are those of the author(s) and not necessarily
31 those of the NHS, the NIHR or the Department of Health.

32

33 **Acknowledgements:** The authors would like to thank the research assistants who
34 have tirelessly supported this project: Despina Djama, Lloyd Taylor, Kirsten Woodman,
35 and Marina Khalil.

36

37

38

39 **Functional Imagery Training versus Motivational Interviewing for weight loss:**

40 **A randomised controlled trial**

41

42 **Objective:** Functional Imagery Training (FIT) is a new brief motivational intervention based
43 on the Elaborated Intrusion theory of desire. FIT trains the habitual use of personalised,
44 affective, goal-directed mental imagery to plan behaviours, anticipate obstacles, and mentally
45 try out solutions from previous successes. It is delivered in the client-centred style of
46 Motivational Interviewing (MI). We tested the impact of FIT on weight loss, compared with
47 time- and contact-matched MI.

48 **Design:** We recruited 141 adults with BMI (kg/m²) ≥ 25, via a community newspaper, to a
49 single-centre randomised controlled trial. Participants were allocated to one of two active
50 interventions: FIT or MI. Primary data collection and analyses were conducted by researchers
51 blind to interventions. All participants received two sessions of their allocated intervention;
52 the first face-to-face (1 hour), the second by phone (maximum 45 min). Booster calls of up to
53 15 mins were provided every 2 weeks for 3 months, then once-monthly until 6 months.
54 Maximum contact time was 4 hours of individual consultation. Participants were assessed at
55 Baseline, at the end of the intervention phase (6 months), and again 12 months post-baseline.

56 **Main outcome measures:** Weight (kg) and waist circumference (WC, cm) reductions at 6
57 and 12 months.

58 **Results:** FIT participants (N=62) lost 4.11 kg and 7.02cm of WC, compared to .74kg and
59 2.72cm in the MI group (N=58) at 6 months (weight mean difference (WMD)=3.37kg,
60 $p<.001$, 95%CI[-5.2, -2.1], waist-circumference mean difference (WCMD)=4.3cm, $p<.001$,
61 95%CI[-6.3,-2.6]). Between-group differences were maintained and increased at month 12:
62 FIT participants lost 6.44kg (W) and 9.1cm (WC) compared to the MI who lost .67kg and

63 2.46cm (WMD=5.77kg, $p<.001$, 95% CI[-7.5, -4.4], WCMD=6.64cm, $p<.001$, 95%[-7.5, -
64 4.4]).

65 **Conclusion:** FIT is a theoretically-informed motivational intervention which offers
66 substantial benefits for weight loss and maintenance of weight reduction, compared with MI
67 alone, despite including no lifestyle education or advice.

68 **Trial registration:** ISRCTN registry: ISRCTN17292316 DOI
69 10.1186/ISRCTN17292316,18/07/2016.

70 **Introduction**

71 Obesity is a leading cause of premature morbidity and mortality worldwide (1,2), but
72 relatively small changes in weight (5% reduction) produce substantial health benefits (3–5).
73 In individuals with overweight and obesity, a sustained reduction of only 2-5 kg reduces
74 cardiovascular risk factors and can prevent progression to type 2 diabetes mellitus (6–8).
75 Reductions in waist circumference bring their own health benefits, as excess abdominal fat
76 increases diabetes risk fivefold in men with waist >102 cm and threefold in women with
77 waist >88cm (9).

78 Most weight loss programs focus on lifestyle education and peer support. These
79 interventions produce some reduction in weight (10,11), but reductions are often reversed
80 within 1-2 years: typically 40-50% of the original loss (12,13). Few studies document weight
81 loss past the end of treatment (14). Maintenance of weight loss has been reported primarily
82 after high-intensity lifestyle interventions that provided comprehensive weight loss
83 counselling for up to 8 years (15), or in trials that delivered at least 6-18 months of extended
84 care, with therapist contact, after the successful completion of a weight loss program (16).

85 Current weight loss programmes place little emphasis on maintaining motivation (17),
86 despite its central role in long-term weight control (18–20). Participants struggle to stay
87 motivated when trying to maintain weight loss (21–23). Motivational interventions may
88 therefore be as effective as skills based and educational approaches. Motivational
89 Interviewing (MI) (24) is the best-established motivational intervention, but trials that include
90 MI to support weight loss or physical activity typically achieve only modest effects
91 (15,25,26). In consequence, MI's clinical significance is limited (25,27,28). There is a need
92 for more effective motivational support.

93 Two features of MI may produce sub-optimal impact. Participants are not taught how
94 to apply MI strategies by themselves, reducing its effectiveness in real-life decision situations

95 (29,30). The latest cognitive models of desire (29,31–33) and emotion (34–36) highlight the
96 critical role of episodic multi-sensory mental imagery in motivation (37–41), but MI remains
97 a heavily verbal style of counselling.

98 Functional Imagery Training (FIT) (42) is a motivational intervention based on
99 Elaborated Intrusion theory, which recognises the importance of vivid mental imagery in
100 desires (32). FIT uses imagery to increase desire and self-efficacy for change. It is delivered
101 in the empathic, accepting and respectful spirit of MI (24), trusting participants to be experts
102 on themselves, and assisting them to identify their own goals and related behaviours rather
103 than trying to convince them to adopt a preset regimen. The FIT protocol prescribes similar
104 steps to MI (24): elicitation of the person’s incentives for change, exploring discrepancies
105 between core values and current behaviour, boosting self-efficacy, and, when people are
106 committed to change, developing specific action plans for implementing this commitment. At
107 each step, FIT also invites participants to develop personalised multisensory imagery to
108 maximize the veridicality and emotional impact of each aspect. In essence, FIT is imagery-
109 based MI, with a strong additional focus on training self-motivation through the use of
110 imagery. When participants have developed plans for change, they transition into training to
111 become their own FIT therapist, supporting their autonomy and ability to flexibly respond to
112 self-management challenges in the natural environment using imagery. They are shown how
113 to develop a cognitive habit of practising emotive goal-related imagery in response to cues
114 from a routine behaviour, and encouraged to generate this imagery whenever motivation
115 needs to be strengthened. Because imagery is more emotive than verbal thought (34–36), this
116 vivid, affectively charged functional imagery sustains desire for change until each new
117 behaviour becomes habitual. Functional imagery also interferes with cravings when
118 temptations occur, by competing for working memory with craving imagery (43). Detailed
119 episodic imagery that is firmly grounded in experience allows participants to anticipate

120 problematic situations, and plan and rehearse effective responses to them, increasing self-
121 efficacy through ‘symbolic practice’ (44). Initial small-scale tests of FIT have shown benefits
122 for reducing snacking (42).

123 In this study we recruited members of the public with overweight and obesity to
124 compare the effects of MI and FIT on weight loss over 6 months of low-intensity treatment,
125 and an additional 6 months of follow-up. We predicted that FIT would produce greater initial
126 weight loss, because imagery amplifies the effective components of MI, and better
127 maintenance at 12 months because FIT teaches the cognitive skills individuals need to stay
128 motivated.

129 Method

130 *Ethics and trial registration*

131 Study approval was granted by the Faculty of Health and Human Sciences research
132 ethics committee, University of Plymouth, on March 23rd, 2015. The International Standard
133 Randomised Controlled Trials registration was <http://www.isrctn.com/ISRCTN17292316>,
134 18/07/2016.

135 *Participants and recruitment*

136 We advertised once for potential participants in the *Plymouth Shopper*, a free local
137 community newspaper, reaching around 64,000 homes. The advertisement sought adult
138 participants with a BMI ≥ 25 kg/m², to test motivational interventions for losing weight and
139 becoming more active. Exclusion criteria were current pregnancy, diagnosed eating disorder,
140 and failure to complete baseline assessments.

141 *Sample size:* A recommended sample size of 191 was estimated using G*Power
142 3.1.9.2, assuming power of .8 to detect a small effect size of .2 for between-group difference.

143 *Design and overview*

144 The trial was a single-centre, two-armed, single-blind, randomised controlled parallel
145 design with matched therapist contact time, comparing FIT with MI (1:1 ratio of
146 participants). The trial had 6 months of active intervention, followed by a 6-month follow-up
147 with no therapist contact. Analyses are reported by intention-to-treat.

148 *Randomisation and masking*

149 Participants completed demographic details and assessments before randomisation.
150 Participants were randomized to MI or FIT by the lead researcher using
151 <https://www.randomizer.org/> (random pairs option). In the two post-treatment assessment
152 sessions, research assistants (RAs) who were blind to the intervention group, collected and
153 recorded primary outcomes. Participants were informed that new RAs would take their
154 measurements, and were asked not to give away whether they were in the FIT or MI group.
155 The lead researcher was present in the room and could verify and record if un-blinding
156 occurred. Analysis of primary outcomes and quality of life was performed by an analyst who
157 was blind to intervention and not otherwise involved in the trial.

158 *Assessment measures*

159 Primary outcomes were body weight, BMI and waistline. Weight (kg) was measured
160 in street clothes, with shoes removed, using Omron BF511 Family Body Composition
161 Monitor. Height was measured to the nearest centimetre, allowing calculation of BMI
162 (kg/m²). For waistline measurement, participants removed coats and sweatshirts, but no other
163 clothing. Waistline was measured to the nearest centimetre, at the height of the umbilicus,
164 using a tailoring tape measure.

165 A secondary outcome was the global quality of life, measured using the 1-item
166 Global Quality of Life Scale (GQOL; 43). Participants in both groups were asked if they
167 would recommend the treatment they had been allocated to a family member, friend or
168 colleague.

169 We collected data on participants' experiences and process variables, including
170 frequency of motivational cognitions, self-efficacy for diet and physical activity, self-reported
171 diet and physical activity. These results will be reported separately.

172 *Interventions*

173 Both interventions were delivered individually by the lead author. Face-to face
174 sessions were all conducted in the same counselling room on the University of Plymouth
175 campus. Session 1 immediately followed the collection of baseline assessments and
176 randomisation, and lasted 1 hour. Session 2 (approximately 35 min) was delivered by
177 telephone a week later. Participants then received fortnightly 5-15 minute-long 'booster'
178 phone calls until 3 months, followed by monthly calls to 6 months post-baseline.

179 We developed scripts to guide delivery of MI and FIT and ensure treatment fidelity
180 and consistency (available from the authors), but the order of segments was flexible and
181 guided by the participant's needs and responses, in keeping with the spirit of MI. Active
182 listening (open questions, affirmation, simple and complex reflections, summarising) was
183 used in both treatments. The initial session of MI and FIT had the same general structure,
184 incorporating a negotiated agenda, discussion of the treatment allocation, assessment
185 feedback, existing or potential goals, incentives for adoption of those goals, and past
186 successes with weight loss efforts. The therapist checked degree of goal commitment. Once
187 participants were committed to behaviour change, they developed a plan for action over the
188 following few days, including strategies to address potential barriers to its implementation.

189 Session 2 reviewed and developed the themes from Session 1, in the light of
190 experiences since the initial session. Booster calls provided opportunities to review
191 progress, reaffirm successful aspects of performance and incentives for behaviour
192 changes, and set additional sub-goals. Exclusively in FIT, all sessions and booster
193 calls included mental imagery exercises.

194 If participants had requested advice on diet or physical activity they would
195 have been referred to the UK's National Health Service's (NHS) publicly available
196 and accessible NHS Choices website which provides general information and advice
197 on health, for example staying fit, losing weight, or beating stress
198 (<https://www.nhs.uk/pages/home.aspx>). We did not have to refer anyone to this
199 option however. Qualitative participant experience data will be published in a
200 separate paper, delivering insight on how participants felt about not being provided
201 with pre-set regimen, diets, or information and advice.

202 *FIT*

203 After discussion of assessment feedback in Session 1, the therapist explained the
204 rationale for using imagery and gave an experience of affectively-charged imagery. After
205 discussing incentives for potential behaviour change, participants imagined these outcomes
206 occurring, as specific future events that were created as vividly as possible. Similar images
207 were elicited about past successes and about detailed plans for the coming days, including
208 successful achievement of each step and success in reaching their ultimate goal.
209 Participants nominated a routine behaviour that could prompt their imagery practice. They
210 carried out this behaviour in the counselling room, while imagining their action plan and
211 goal. They were also encouraged to practise imagery before engaging in their chosen
212 behaviour, and offered the simple 'Goal in Mind' app
213 (<https://itunes.apple.com/au/app/goal-in-mind/id1289557359?mt=8;>
214 <https://play.google.com/store/apps/details?id=com.goalinmind&hl=en>) to download. They
215 could use the app to upload motivational photos, tick off when they had remembered to do
216 imagery practise (the app did not provide reminders), input a goal they would like to achieve
217 and access a five minute audio that guided them through imagining how they would work on

218 their goal today and how good it would feel to achieve it. The audio was emailed as an MP3
219 file if participants did not want to use the app, but wanted the audio to practise.

220 Session 2 reviewed progress, including participants' efforts at practising imagery.
221 Imagery was used to help solve any problems with progress towards their goal and to
222 motivate new sub-goals.

223 Booster calls developed imagery about recent successes, problem solutions, or new
224 goals behaviours. If required, additional imagery exercises included: 'Cravings Buster'
225 (deliberate switching of attention from craving imagery to goal imagery) and 'Plateau'
226 (reflecting on benefits experienced so far and exploring additional ways of working on goals).

227 *MI*

228 The therapist did not explicitly evoke imagery, and avoided language that
229 was likely to trigger it. Some additional questions were added to the manual, to
230 ensure that the MI sessions had similar time and intensity as the FIT sessions. A few
231 examples are: "When you think about that list of things, how does it make you
232 feel?", "Would you mind summarising the things that are likely to get better if you
233 change your behaviour?" and "Is there anyone who could help you follow it
234 through?" MI participants were offered a goal sheet with the action plan they
235 developed with the therapist in the first session; FIT did not have this. They took the
236 goal sheet home and were encouraged to review their statements, goals and
237 strategies, especially when they felt they needed extra motivation. Two examples of
238 this additional exercise from the script are: "Would you like to write that down so
239 you have a summary to take away?" and "If you need a bit of a boost to your
240 motivation over the next few days, you could try reading that over to remind
241 yourself about what you said." The sections were as follows: What I am going to
242 do...Why I want to do it...How I'll do it...I know I can do it because.... Participants

243 were encouraged to add to the sheet as their goals or reasons for change evolved.

244 During the booster calls, they were asked if they had added any new goals or ticked

245 off achievements on their sheet.

246 *Intervention fidelity*

247 The therapist was trained in FIT by two of the creators of FIT and undertook a 3-day

248 MI course. She attended weekly clinical supervision meetings with the senior author, to

249 review individual sessions. A random 20% sample of initial FIT and MI sessions was rated

250 on the Motivational Interviewing Treatment Integrity (MITI) 3.1.1 (46) by an RA not

251 involved in the project. FIT sessions were rated on a 15-item checklist based on the manual.

252 Additionally, two RAs listened to the session recordings independently and categorised them

253 according to the intervention they thought the participants had received.

254 *Procedure*

255 Participants gave informed consent a week before their initial session. After

256 completing demographic details and all assessments, they were randomly allocated to MI or

257 FIT. The treatment sessions followed, as described above. After the first booster call, all

258 participants were asked to complete reassessments of the expected process variables (results

259 reported separately). At the end of treatment (6m), they attended a 15-minute post-treatment

260 assessment session in the counselling room. Quality of Life was assessed via emailed

261 questionnaire one week before this session. RAs blind to intervention measured waist and

262 weight, and participants completed process measures online. They were told that the therapist

263 would be available if they were experiencing distress (none took up this offer). Participants

264 were reminded that they were entering the unsupported maintenance period and that the

265 therapist would be in touch 2-3 weeks before the final weigh-in, to arrange the appointment.
266 They received £15 for their time and travel.

267 At 12 months, participants returned for the final weigh-in. This session did not
268 include self-report instruments. They received £5 for completing this assessment.

269 *Data analysis*

270 Weight and waist circumference were analysed separately. To estimate differences
271 between MI and FIT, outcome measurements were regressed onto baseline score (kg or cm),
272 a time indicator (6/12 months), and a group indicator (MI/FIT), using linear mixed-effects
273 models (47). These models also included baseline BMI and its interaction with time and
274 group; baseline BMI was included because it captures additional information about the
275 severity of participants' condition when entering the study. These models are analogous to
276 repeated measures ANCOVA, but allowed us to make efficient use of all available data
277 without imputation of missing values. Of primary interest were the between-groups contrasts
278 for weight and waist circumference at 6 and 12 months. Tests of parameter values and other
279 contrasts are reported with Satterthwaite approximation for degrees of freedom. In a
280 secondary analysis, GQOL scores at 6 months were regressed on baseline GQOL scores,
281 baseline BMI, group, and the interaction of baseline scores with group. Alternative model
282 parameterisations, in which treatment effects were estimated as a linear slope from baseline
283 to 6 or 12 months, produced equivalent inferences.

284 To support probability statements about the average effect of FIT vs. MI, and likely
285 prognoses of future participants selecting FIT or MI, we re-ran our mixed models using a
286 Bayesian estimation procedure with pessimistic but weakly informative priors (48); full
287 details are available in our data supplement, but for regression coefficients these priors were
288 Gaussian, zero-centred, and with a scale adjusted to $2.5 \times SD(y) / SD$ (Gabry, J., & Goodrich,
289 B. (2016). *rstanarm: Bayesian applied regression modeling via stan. R package version, 2(1).*)

290 Based on these models, we provide summaries of the posterior density for the average
291 treatment effect, and for the predicted prognoses of new individuals selecting FIT or MI. All
292 models appeared to converge satisfactorily based on visual inspection of MCMC traces and
293 parameter R-hat statistics (49). All data and R code for the analyses presented here are
294 available in an online supplement.

295 *Economic costing:*

296 We used Public Health England's weight management economic assessment tool
297 No.2 (50), to estimate the increase in quality-adjusted life-years (QALY) associated with the
298 additional weight lost by FIT participants at 12-months compared with MI.

299 **Results:**

300 A total of 141 participants was recruited in the time available (March -May 2016). One
301 hundred and 20 were randomised, 58 to MI and 62 to FIT. One hundred and 14 were included
302 in the analysis of the 6-month follow-up, and 112 completed both 6 and 12 month follow-ups
303 (Consort Diagram, figure 1; Table A in the supplementary materials). No statistically
304 significant differences were found between groups at baseline (Table 1). 21 out of 58 FIT
305 participants reported having used the app or audio when asked at 6 months. 25 out of 55 MI
306 participants had continued to use their goal-sheet past the first MI session when asked at 6
307 months. [insert table 1]

308 *MI and FIT fidelity checks*

309 MI skills were rated on the MITI's (44) 5-point scale: 1: Never, 2: Rarely, 3:
310 Sometimes, 4: Often, 5: Always. For MI, ratings for Evocation, Collaboration, Autonomy,
311 Support, Direction, and Empathy ranged from 3.9 to 4.9 (median=4.5). For FIT, they ranged
312 from 3.8-4.7 (median =4.5). For FIT, 15 session elements were rated as 0 absent, or 1 present.
313 Totals ranged from 13-15 (median = 15). Independent raters correctly assigned 100% of
314 audio recordings to intervention.

315

[insert figure 1]

316 To visualise changes in weight and waist, we plotted unadjusted means and 95%
317 confidence intervals for each group. Figure 2 indicates that participants treated with MI
318 experienced little to no reduction in weight or waist from baseline to either 6 or 12-month
319 follow-ups. Those treated with FIT experienced large reductions in weight and waist
320 circumference. Relative to both the MI group and baseline, participants treated with FIT
321 continued to lose weight after treatment ended.

322 [insert figure 2]

323 Our primary statistical models estimated the differences between-groups, conditional
324 on baseline BMI, at month 6 and 12; results are presented in Table 2. We found substantial
325 and statistically significant differences between the MI and FIT groups at both follow-ups.

326 To make probability statements about the size of benefit obtained by participants
327 undergoing FIT, we re-estimated our mixed models using a Bayesian procedure. Table 2
328 shows treatment effects, and associated 95% credible intervals from these models.

329

330 [insert table 2]

For weight and waist circumference, there was overwhelming evidence that FIT was beneficial. Similarly, for GQOL the difference between groups at month 6 was statistically significant. In the FIT group, 58/59 participants would recommend the intervention to others; one might recommend. In MI, 53/55 would recommend, 2 might.

To help clinicians and others evaluate the likely benefit of FIT in clinical practice, we computed the posterior-probability that the benefit of FIT for a new participant would exceed a range of values between 0 and 15kg lost, and between 0-15cm of waist reduction (Figure 3): [insert figure 3]

When considering the risks or benefits of interventions, clinicians, participants and researchers benefit from probability information presented as 'natural frequencies' or in 'pictographs' (51). Consequently, we used the same model-based simulations to calculate the range of likely prognoses from the participants' perspective (Figure 4). After treatment, only 22% of MI participants were predicted to lose 5% or more of their initial weight, compared with the NICE target (52) that 30% do; after 12 months this figure was 23%. In contrast, 54% of new FIT participants were predicted to lose at least 5% of their initial weight after treatment, and 75% are predicted to lose this much by 12-months. Stated differently, nine of every 10 participants would have benefitted more from FIT than from MI (probability $FIT > MI$ for kg lost at 12m = 0.94; for cm lost it is 0.85); for half of these participants, the expected additional benefit is substantial (> 5kg difference in projected outcomes, see data supplement).

[insert figure 4]

Economic assessment:

We based our inputs to the PHE assessment tool (50) on the conservative assumption that MI was equivalent to no-treatment, and that FIT participants would begin to gain weight

immediately after the 12-month follow-up. We modelled costs on the basis that 58 new patients would be treated with FIT and that they would reduce their BMI by an average of 2.148 kg/m² over 1 year (based on our primary outcome models). We very conservatively assumed that each hour of individual treatment would cost £250 to deliver, and entered a worst-case per-participant cost for FIT at £1000.

Even based on these highly conservative assumptions, the PHE model suggests that FIT would be cost-effective from the healthcare perspective within three years, judged by NICE's conventional willingness-to-pay threshold of between £20,000 and £30,000 per QALY. Cost-per-QALY after three years was £22,036, falling to £12,363 after five years, and £7,229 after 10 years. Including costs of social care and the prospect of increased employment, cost per QALY was only £12,968 after three years, £3,739 after 5 years, and was cost-saving from a 10-year perspective.

Discussion

In this first randomised controlled trial of FIT, we have shown that two FIT interviews and nine brief booster phone calls, amounting to under 4 hours of therapist contact over six months, resulted in substantially greater and clinically meaningful weight loss and waistline reductions at six months, compared with MI. Participants in the FIT arm, but not MI, continued to lose weight and waist circumference in the unsupported 6-month maintenance phase. Participants in both treatment groups reported improved quality of life at 6 months, but FIT participants reported greater improvements. Seventy-one percent of the FIT group lost more than 5% of their initial weight, easily exceeding the NICE weight management target that at least 30% of service-users should do so (52). MI did not meet the NICE target, with only 23% losing 5%.

Importantly, we found that FIT was acceptable to participants and that they would recommend FIT to a family member, friend or colleague. Because the delivery of FIT closely matches the protocol of an existing intervention (MI), scaling delivery to larger numbers of participants should be straightforward, with existing MI practitioners requiring only minimal additional training.

To put these findings into a broader context, FIT performed favourably compared to a longer, more intensive intervention in a recent UK trial: Ahern and colleagues (53) tested weight loss in participants with overweight and obesity referred by GPs to Weight Watchers. Participants randomised to the Weight Watchers programme for 12 months had lost an average of 6.8kg at 12 months, only 0.4 kg more than participants in our RCT, who received less than 4 hours of FIT spread over 6 months. Participants on the standard 12-week Weight Watchers programme lost less than FIT participants in this study: only 4.8kg on average (53).

The approach tested in this trial, of providing solely motivational support, differs from the strategy recommended by Public Health England (54,55) of combining behaviour change techniques with lifestyle education and advice. NICE recommends including MI and imagery in behaviour change strategies. FIT combines both in a coherent, structured intervention that trains users to become their own therapist; the present results support this approach. It remains to be determined if combining FIT with diet and physical activity education would generate superior outcomes.

Sustained reductions of around 5% of body weight can effect significant health improvements, such as decreased blood lipids, precursors of Type 2 diabetes and improved blood pressure (3,4). Weight loss of between 5% and 10% is associated with significant improvements in cardiovascular disease risk factors (5). The reduction in waist circumference, from an average of 106 cm to 97 cm by 12 months in FIT, brings its own health risk reductions: excess abdominal fat indicated by a high waist circumference in men

(>102 cm), presents a fivefold increase in the risk of developing diabetes (9). For women, across a range of baseline BMIs between 25-50 kg/m², waist reduction of 5-10cm is associated with a reduction in cholesterol and systolic blood pressure (56). Results from the Public Health England economic assessment tool suggest that these health benefits would make delivery of FIT cost effective, although detailed cost-effectiveness evaluation must form part of additional large-scale evaluations of FIT.

The fact that FIT outperformed the best established motivational intervention by such a margin is encouraging. We demonstrated a mean weight difference of 2kg for FIT, compared with the active MI intervention group at 6 months. This benefit of FIT over MI was larger than the benefit of MI over minimal control interventions in Armstrong et al's., (24) meta-analysis of MI for weight loss: mean difference = 1.47 kg 95% CI = -2.05 to -0.88 at around 6 months. However, we note that weight in the MI group stabilised after six months, and successfully preventing weight gain is an important health focus (57,58). That FIT demonstrates such an improvement over an existing treatment, and produces continued weight loss after the end of the intervention period, highlights the benefit of developing and adapting existing interventions based on recent developments in cognitive science.

Limitations and future directions

We focused purely on motivation, providing no lifestyle advice or information. We did not assess participants' knowledge of nutrition or physical activity. First, this was in keeping with the autonomous spirit of MI which FIT retains. The ethos of both interventions is to motivate participants to seek the information or support they need to achieve their goals. Second, we focused on process outcomes in this trial, rather than content outcomes. Assessing previous knowledge at baseline could have led participants to believe they should discuss their knowledge, or to expect advice or prescribed diet and exercise plans. Although we recruited from the general public, we acknowledge that we may have attracted a well-

informed sample (e.g., the kind of person who would sign up for a study conducted at a university). Perhaps this sample was more knowledgeable than a random sample of people with overweight, but this certainly did not make them more successful; they still struggled with overweight or obesity when they signed up for the study. There is little evidence that knowledge alone motivates behaviour. For example, a recent, large cross-sectional study on self-management of glycemic control found no correlation between knowledge about type 2 diabetes and actual self-management behaviours (59). Even with healthcare professionals, education did not lead to behaviour change. McCluskey and Lovarini, (2005) tested an educational intervention to improve evidence-based practise amongst allied health professionals. Improved knowledge was maintained at 8 month follow-up but behaviour change was very limited, with nearly two thirds of health professionals still not reading any research literature. It is conceivable however that education is beneficial in populations with very limited knowledge of nutrition and exercise. Future studies should therefore assess whether FIT works best as an alternative to established weight-loss programmes based on lifestyle education and advice, or as an adjunct to them.

As far as possible, we matched MI and FIT for intervention intensity. There were the same number of sessions, scheduled for the same duration, in both conditions. Although the dialogue for FIT incorporated the same essential components as for MI, we added some extra elements to the MI manual to add depth to the interview and equate the time taken. In FIT, all participants were offered the Goal in Mind app to guide their imagery practice. In MI, all participants were given goal sheets to review at home. If anything, MI participants had more time to talk to the therapist because they did not do imagery exercises.

FIT achieved larger effect sizes than expected. Because of practical constraints on recruitment, our sample was smaller than that recommended by our power calculations to detect modest to moderate effect sizes, therefore it is plausible that our trial over-estimates

the true effect of FIT, i.e. a 'Type M' (magnitude) Error (61). Nonetheless, the posterior probability for an effect of FIT < 1kg lost was extremely low, and the evidence that FIT was preferable to MI was substantial.

This was the first efficacy trial of FIT for weight loss, and our results must be replicated in a larger multi-centre trial to be confident that FIT can be delivered effectively at scale by other therapists. However, we have achieved proof of concept, and compared the intervention with a well-established intervention for weight loss. Although the FIT group maintained their weight loss at our 12- month follow-up, further trials should follow participants for several years to ensure that FIT does indeed help participants maintain the weight loss achieved during treatment, and to properly assess the health economic benefit of these reductions.

Conclusion:

Less than 4 hours of Functional Imagery Training, a novel intervention that combines motivational interviewing with mental imagery training, led to substantially greater weight loss over 6 months than MI alone, despite the fact that specific education on lifestyle and activity was absent from this intervention. The benefits of FIT persisted to 12 months; participants continued to make substantial reductions in weight even after therapy ended.

References:

1. Eastwood P. Statistics on Obesity , Physical Activity and Diet : Heal Soc Care Inf Centre, Lifestyles Stat. 2013;2:15–35.
2. Health and Social Care Information Centre. Statistics on Obesity, Physical Activity and Diet: England 2015. Heal Soc Care Inf Cent [Internet]. 2015;(March):103.
Available from: <http://www.hscic.gov.uk/catalogue/PUB16988/obes-phys-acti-diet-eng-2015.pdf>
3. Jakicic JM, Clark K, Coleman E, Donnelly JE, Foreyt J, Melanson E, et al. Appropriate strategies for intervention weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc.* 2001;33:2145–56.
4. Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exerc.* 2009;41(2):459–71.
5. Wing R, Lang W, Wadden T, Safford M, Knowler W, Bertoni A, et al. Benefits of Modest Weight Loss in Improving Cardiovascular Risk Factors in Overweight and Obese Individuals With Type 2 Diabetes. *Diabetes Care* [Internet]. 2011;34(7):1481–6.
Available from:
<http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=ovftm&NEWS=N&AN=00003458-201107000-00007>
6. Espeland M. Reduction in Weight and Cardiovascular Disease Risk Factors in Individuals With Type 2 Diabetes: One-Year Results of the Look AHEAD Trial. *Diabetes Care.* 2007;
7. Tuomilehto J, Lindström J, Eriksson JG, Valle TT, Hämäläinen H, Ilanne-Parikka P, et al. Prevention of Type 2 Diabetes Mellitus by Changes in Lifestyle among Subjects with Impaired Glucose Tolerance. *N Engl J Med* [Internet]. Massachusetts Medical

- Society ; 2001 May 3 [cited 2016 Nov 18];344(18):1343–50. Available from:
<http://www.nejm.org/doi/abs/10.1056/NEJM200105033441801>
8. Williams G, Hamm MP, Shulhan J, Vandermeer B, Hartling L, Avenell A, et al. Reduction in Weight and Cardiovascular Disease Risk Factors in Individuals With Type 2 Diabetes: One-Year Results of the Look AHEAD Trial Received. *Natl Inst Heal Clin Excell* [Internet]. 2012;8(6):1585–93. Available from:
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3996838&tool=pmcentrez&rendertype=abstract>
 9. Gatineau Mary, Hancock Caroline, Holman Naomi, Outhwaite Helen, Oldridge Lorraine CA and EL. Adult obesity and type 2 diabetes *About Public Health England*. 2014;1–39.
 10. Hartmann-Boyce J, Johns DJ, Jebb SA, Aveyard P, Ogden J, Onakpoya I, et al. Effect of behavioural techniques and delivery mode on effectiveness of weight management: Systematic review, meta-analysis and meta-regression. *Obes Rev*. 2014;15(7):598–609.
 11. Truby H, Baic S, DeLooy A, Fox KR, Livingstone MBE, Logan CM, et al. Randomised controlled trial of four commercial weight loss programmes in the UK: initial findings from the BBC “diet trials”. *BMJ* [Internet]. 2006;332(7553):1309–14. Available from:
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1473108&tool=pmcentrez&rendertype=abstract>
 12. Wadden TA, Butryn ML, Byrne KJ. Efficacy of Lifestyle Modification for Long-Term Weight Control. *Obes Res* [Internet]. 2004;12(S12):151S–162S. Available from:
<http://doi.wiley.com/10.1038/oby.2004.282>
 13. Bessesen DH. Systematic Review: An Evaluation of Major Commercial Weight Loss

- Programs in the United States. *Yearb Endocrinol.* 2006;2006:153–7.
14. Kozica S, Lombard C, Teede H, Ilic D, Murphy K, Harrison C. Initiating and continuing behaviour change within a weight gain prevention trial a qualitative investigation. *PLoS One* [Internet]. 2015;10(4):1–14. Available from: <http://dx.doi.org/10.1371/journal.pone.0119773>
 15. Wadden TA. Eight-year weight losses with an intensive lifestyle intervention: The look AHEAD study. *Obesity.* 2014;22(1):5–13.
 16. Ross Middleton KM, Patidar SM, Perri MG. The impact of extended care on the long-term maintenance of weight loss: A systematic review and meta-analysis. *Obes Rev.* 2012;13(6):509–17.
 17. West DS, Gorin AA, Subak LL, Foster G, Bragg C, Hecht J, et al. A motivation-focused weight loss maintenance program is an effective alternative to a skill-based approach. *Int J Obes (Lond)* [Internet]. Nature Publishing Group; 2011;35(2):259–69. Available from: <http://www.scopus.com/inward/record.url?eid=2-s2.0-79951683411&partnerID=tZOtx3y1>
 18. Elfhag K, Rössner S, Rossner S, Wu T, Gao X, Chen M, et al. Who succeeds in maintaining weight loss? A conceptual review of factors associated with weight loss maintenance and weight regain. *Obes Rev* [Internet]. 2005;6(1):67–85. Available from: <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=ovfte&NEWS=N&AN=00005407-200103070-00025>
 19. Silva MN, Markland D, Carra??a E V., Vieira PN, Coutinho SR, Minderico CS, et al. Exercise autonomous motivation predicts 3-yr weight loss in women. *Med Sci Sports Exerc.* 2011;43(4):728–37.
 20. Teixeira PJ, Going SB, Sardinha LB, Lohman TG. A review of psychological pre-

- treatment predictors of weight control. *Obes Rev.* 2005;6(7):43–65.
21. Sabinsky MS, Toft U, Raben A, Holm L. Overweight men's motivations and perceived barriers towards weight loss. *Eur J Clin Nutr.* 2007;61(4):526–31.
 22. Sharifi N, Mahdavi R, Ebrahimi-Mameghani M. Perceived Barriers to Weight loss Programs for Overweight or Obese Women. *Heal Promot Perspect [Internet].* 2013;3(1):11–22. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3963684&tool=pmcentrez&rendertype=abstract>
 23. Solbrig L, Jones R, Kavanagh D, May J, Parkin T, Andrade J. People trying to lose weight dislike calorie counting apps and want motivational support to help them achieve their goals. *Internet Interv [Internet].* The Authors; 2017;7:23–31. Available from: <http://dx.doi.org/10.1016/j.invent.2016.12.003>
 24. Miller W, Rollnick S. *Motivational Interviewing: Helping People Change.* Guilfordpress; 2012.
 25. Armstrong MJ, Mottershead T a., Ronksley PE, Sigal RJ, Campbell TS, Hemmelgarn BR. Motivational interviewing to improve weight loss in overweight and/or obese patients: A systematic review and meta-analysis of randomized controlled trials. *Obes Rev.* 2011;12(4):709–23.
 26. Hardcastle SJ, Taylor AH, Bailey MP, Harley R a, Hagger MS. Effectiveness of a motivational interviewing intervention on weight loss, physical activity and cardiovascular disease risk factors: a randomised controlled trial with a 12-month post-intervention follow-up. *Int J Behav Nutr Phys Act [Internet].* 2013;10:40. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23537492>
 27. Martins RK, McNeil DW. Review of Motivational Interviewing in promoting health behaviors. *Clin Psychol Rev [Internet].* Elsevier B.V.; 2009;29(4):283–93. Available

- from: <http://dx.doi.org/10.1016/j.cpr.2009.02.001>
28. VanWormer JJ, Boucher JL. Motivational Interviewing and Diet Modification: A Review of the Evidence. *Diabetes Educ.* 2004;30(3):404–19.
 29. Kavanagh DJ, Andrade J, May J, Connor JP. Motivational interventions may have greater sustained impact if they trained imagery-based self-management. *Addiction.* 2014;109(7):1062–3.
 30. Olander EK, Fletcher H, Williams S, Atkinson L, Turner A, French DP. What are the most effective techniques in changing obese individuals' physical activity self-efficacy and behaviour: A systematic review and meta-analysis. *Int J Behav Nutr Phys Act* [Internet]. 2013;10(29):29. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3639155&tool=pmcentrez&rendertype=abstract>
 31. Andrade J, May J, Kavanagh D. Sensory Imagery in Craving: From Cognitive Psychology to New Treatments for Addiction. *J Exp Psychopathol.* 2012;3(2):127–45.
 32. Kavanagh DJ, Andrade J, May J. Imaginary relish and exquisite torture: the elaborated intrusion theory of desire. *Psychol Rev.* 2005;112(2):446–67.
 33. Kavanagh, David J. ; Andrade, Jackie ; May, Jon ; Connor JP. Do brief alcohol motivational interventions work like we think they do? *Addiction.* 2014;109(7):1062–3.
 34. Blackwell SE, Rius-Ottenheim N, Schulte-van Maaren YWM, Carlier IVE, Middelkoop VD, Zitman FG, et al. Optimism and mental imagery: A possible cognitive marker to promote well-being? *Psychiatry Res* [Internet]. Elsevier; 2013;206(1):56–61. Available from: <http://dx.doi.org/10.1016/j.psychres.2012.09.047>
 35. Holmes E, Mathews A. Mental Imagery and Emotion: A Special Relationship? *Emotion.* 2005;5(4):489–97.

36. Renner F, Ji JL, Pictet A, Holmes EA, Blackwell SE. Effects of Engaging in Repeated Mental Imagery of Future Positive Events on Behavioural Activation in Individuals with Major Depressive Disorder. *Cognit Ther Res*. Springer US; 2017;41(3):369–80.
37. Parham SC, Kavanagh DJ, Gericke CA, King N, May J, Andrade J. Assessment of Motivational Cognitions in Diabetes Self-Care: the Motivation Thought Frequency Scales for Glucose Testing, Physical Activity and Healthy Eating. *Int J Behav Med* [Internet]. Springer US; 2016 Nov 7 [cited 2017 Feb 4];1–10. Available from: <http://link.springer.com/10.1007/s12529-016-9607-2>
38. Knäuper B, Roseman M, Johnson PJ, Krantz LH. Using mental imagery to enhance the effectiveness of implementation intentions. *Curr Psychol*. 2009;28(3):181–6.
39. Knäuper B, McCollam A, Rosen-Brown A, Lacaille J, Kelso E, Roseman M. Fruitful plans: adding targeted mental imagery to implementation intentions increases fruit consumption. *Psychol Heal*. 2011;26(5):601–17.
40. Michie S, Ashford S, Sniehotta F, Dombrowski S, Bishop a, French D. A refined taxonomy of behaviour change techniques to help people change their physical activity and healthy eating behaviours: the CALO-RE taxonomy. 2011; Available from: <http://discovery.ucl.ac.uk/1311528/>
41. Neill JO, Oluyomi T, Epstein LH. Eating Behaviors Episodic future thinking reduces eating in a food court. *Eat Behav* [Internet]. Elsevier Ltd; 2016;20:9–13. Available from: <http://dx.doi.org/10.1016/j.eatbeh.2015.10.002>
42. Andrade J, Khalil M, Dickson J, May J, Kavanagh DJ. Functional Imagery Training to reduce snacking: Testing a novel motivational intervention based on Elaborated Intrusion theory. *Appetite* [Internet]. Elsevier Ltd; 2016;100:256–62. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0195666316300502>
43. May J, Andrade J, Panabokke N, Kavanagh D. Visuospatial tasks suppress craving for

- cigarettes. *Behav Res Ther* [Internet]. Elsevier Ltd; 2010;48(6):476–85. Available from: <http://dx.doi.org/10.1016/j.brat.2010.02.001>
44. Bandura A. Self-efficacy mechanism in human agency. *Amer Psych*. 1982;37(2):122–47.
 45. Hyland ME, Sodergren SC. Development of a New Type of Global Quality of Life Scale, and Comparison of Performance and Preference for 12 Global Scales [Internet]. *Quality of Life Research*. Springer; [cited 2017 Nov 6]. p. 469–80. Available from: <http://www.jstor.org/stable/pdf/4034400.pdf>
 46. Moyers TB, Martin T, Manuel JK, Miller WR, Ernst D. Revised Global Scales : (MITI 3 . 1 . 1). 2010;1(January):1–29.
 47. Bates D, Mächler M, Bolker BM, Walker SC. Fitting linear mixed-effects models using lme4. *J Stat Softw* [Internet]. 2015;67(1):1–48. Available from: <http://cran.r-project.org/package=lme4><http://www.jstatsoft.org/index.php/jss/article/view/v067i01/v67i01.pdf>
 48. Gabry J, Goodrich B. rstanarm: Bayesian applied regression modeling via stan [Computer software manual] [Internet]. R package version. 2016. p. 0–3. Available from: <http://cran.r-project.org/package=rstanarm>
 49. Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DD. *Bayesian Data Analysis*. CRC Press. 2014.
 50. Copley V. User guide : weight management economic assessment tool *Economic assessment of adult weight management interventions About Public Health England*. Public Heal Engl. 2016;
 51. Tait AR, Voepel-Lewis T, Zikmund-Fisher BJ, Fagerlin A. The effect of format on parents' understanding of the risks and benefits of clinical research: A comparison between text, tables, and graphics. *J Health Commun*. 2010;15(5):487–501.

52. NICE. Obesity: identification, assessment and management. Natl Institue Heal Care Excell [Internet]. 2017;(November 2014). Available from:
<https://www.nice.org.uk/guidance/cg189/ifp/chapter/obesity-and-being-overweight>
53. Ahern AL, Wheeler GM, Aveyard P, Boyland EJ, Halford JCG, Mander AP, et al. Extended and standard duration weight-loss programme referrals for adults in primary care (WRAP): a randomised controlled trial. *Lancet* [Internet]. The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license; 2017;389(10085):2214–25. Available from: [http://dx.doi.org/10.1016/S0140-6736\(17\)30647-5](http://dx.doi.org/10.1016/S0140-6736(17)30647-5)
54. Thompson L, Aveyard P, Jebb S, Blackshaw J, Coulton V. Let’s Talk About Weight: A step-by-step guide to brief interventions with adults for health and care professionals. 2017; Available from:
www.gov.uk/phe%0Awww.facebook.com/PublicHealthEngland
55. NICE. Weight management : lifestyle services for o ov verweight or obese children and y oung people. Natl Inst Heal Clin Excell [Internet]. 2017;(October). Available from: nice.org.uk/guidance/ph53
56. Han TS, Richmond P, Avenell A, Lean MEJ. Waist circumference reduction and cardiovascular benefits during weight loss in women. *Int J Obes* [Internet]. 1997;21(2):127–34. Available from:
<http://www.nature.com/doi/10.1038/sj.ijo.0800377>
57. Lloyd-Jones DM, Liu K, Colangelo LA, Yan LL, Klein L, Loria CM, et al. Consistently stable or decreased body mass index in young adulthood and longitudinal changes in metabolic syndrome components: The coronary artery risk development in young adults study. *Circulation*. 2007;115(8):1004–11.
58. Strong K a, Parks SL, Anderson E, Winett R, Davy BM. NIH Public Access. *J Am*

- Diet Assoc. 2008;108(10):1708–15.
59. CHEN Q, WANG H, WANG Y, WANG Z, ZHAO D, CAI Y. Exploring Self-Management on Glycemic Control Using a Modified Information–Motivation–Behavioral Skills Model in Type 2 Diabetes Mellitus Patients in Shanghai, China: a Cross-Sectional Study IMB model of self-management on HbA. *J Diabetes*. 2018;86(21).
 60. McCluskey A, Lovarini M. Providing education on evidence-based practice improved knowledge but did not change behaviour: a before and after study. *BMC Med Educ* [Internet]. BioMed Central; 2005 Dec 19 [cited 2018 Feb 27];5(1):40. Available from: <http://bmcmededuc.biomedcentral.com/articles/10.1186/1472-6920-5-40>
 61. Gelman A, Carlin J. Beyond Power Calculations. *Perspect Psychol Sci* [Internet]. 2014;9(6):641–51. Available from: <http://journals.sagepub.com/doi/10.1177/1745691614551642>

Table 1. Baseline demographics, split by intervention:

Group/intervention	MI (Range, Median, Mean)		FIT (Range, Median, Mean)	
N	55		59	
Gender (N)	Female	40	Female	43
	Male	15	Male	16
Age (N years)	Range	19-70	Range	20-72
	Median	43	Median	45
	Mean	42	Mean	45
BMI (kg/m ²)	Range:	24.51-53.33	Range	25.98-47.97
	Median	31.34	Median	31.85
	Mean	32.54	Mean	33.21
Weight (kg)	Range	131-59	Range	62.30-140.5
	Median	87.40	Median	89.40
	Mean	89.66	Mean	91.46
Waistline (cm)	Range	80-148	Range	79-144
	Median	105	Median	103
	Mean	106.01	Mean	106.78
Employment status (N)	Employed full-time	12	Employed full-time	20
	Employed part-time	18	Employed part-time	14
	Retired	5	Retired	6
	In education	4	In education	7
	Unemployed	3	Unemployed	5
	Other	11	Other	4
	Self-employed	1	Self-employed	2
Highest level of education (N)	GCSE	15	GCSE	14
	NVQ/Diploma	4	NVQ/Diploma	5
	Trade	8	Trade	6
	A or O-Levels	14	A or O-Levels	15
	Access course	2	Access course	4
	Foundation degree	2	Foundation degree	1
	Degree		Degree	
	Postgraduate	4	Postgraduate	3
	Degree	1	Degree	4
	No info given	4	No info given	5 ¹

¹ There were no statistically significant differences between groups at baseline

Table 2. Between group contrasts (with Satterthwaite corrected degrees of freedom for Kg and Cm) and posterior mean differences (and 95% credible intervals) for the effect of FIT vs. MI at month 6 and 12. MCMC = Markov Chain Monte Carlo estimates from Bayesian model fits.

Outcome	Follow up	FIT (mean)	FIT (sd)	MI (mean)	MI (sd)	df	t	p	Treatment effect (MCMC)	lower	upper
Cm	Baseline	106.07	13.75	105.50	12.51						
Cm	Month 6	99.05	12.61	102.78	13.37	205.4	-4.727	<.001	-4.444	6.328	2.560
Cm	Month 12	96.97	12.59	103.04	12.45	206.1	-7.012	<.001	-6.697	8.602	4.801
Kg	Baseline	90.48	15.90	89.13	14.76						
Kg	Month 6	86.37	15.07	88.39	15.72	161.5	-4.877	<.001	-3.670	5.203	2.139
Kg	Month 12	84.04	15.96	88.46	15.34	163.4	-7.707	<.001	-5.929	7.482	4.418
GQOL	Baseline	62.13	10.59	61.71	14.51						
GQOL	Month 6	75.81	11.66	72.53	10.42	109.0	2.107	.037	2.831	0.091	5.565

^{2, 3}

² Cm=waist circumference in cm, kg=mean weight in kg, GQOL=mean score of Global Quality of Life assessment

³ Footnote: MCMC: Markov Chain Monte Carlo estimates from Bayesian model fits.

Figure 1. CONSORT Flow Diagram




Figure 2. Mean waist circumference and weight with 95% confidence interval, by group.




Figure 2: Unadjusted weight and waist circumference by group: mean and 95% confidence interval.

Figure 3. Prognosis for new participants randomised to FIT vs. MI, expressed as the probability the benefit will equal or exceed the value on the x-axis.




Figure 4. 12 month prognosis for 100 new participants undergoing MI or FIT.

