The Plymouth Student Scientist - Volume 01 - 2008

The Plymouth Student Scientist - Volume 1, No. 2 - 2008

2008

The effect of starvation on the size of mussels selected by the Green Shore Crab, Carcinus maenas

Morris, Brecht

Morris, B. (2008) 'The effect of starvation on the size of mussels selected by the Green Shore Crab, Carcinus maenas', The Plymouth Student Scientist, 1(2), pp. 4-18. http://hdl.handle.net/10026.1/13155

The Plymouth Student Scientist University of Plymouth

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.





The effect of starvation on the size of mussels selected by the Green Shore Crab, *Carcinus maenas*

Brecht Morris

2008

Project Advisor: <u>Mick Uttley</u>, Faculty of Science, University of Plymouth, Drake Circus, Plymouth, PL4 8AA

Abstract

The foraging behaviour of the green shore crabs, *Carcinus maenas*, of 24-26mm carapace width was observed over a 6 day starvation period. Edible mussels, *Mytilus edulis*, for each millimetre interval from 5-25mm were used to determine how prey size selection was effected by starvation. The most profitable mussel size for the studied crab size is approximately 9mm. At low starvation levels mussel size selection was relatively specific with the majority of selected mussels ranging in size from 7-12mm. Increased starvation caused greater variation in size selection with the majority of selected mussels ranging in size from 7-12mm. Increased starvation random foraging behaviour and subsequent size selection was exhibited as mean mussel size values were close to the predicted random selection value of 12.5mm. Mussels were selected after progressively fewer encounters as starvation increased and consequently the maximum size of a mussel deemed profitable increased with elevated starvation. The findings are in accordance with the optimal foraging theory, since increased starvation is associated with decreasing encounter rates and low encounter rates cause broadening of a forager's diet due to the selection of successively less optimal prey.

1. Introduction

There is considerable experimental evidence that some organisms can make decisions with respect to their foraging activity, in order to achieve optimal nutritional gain from available foods. MacArthur and Pianka (1966) were the first to put forward the optimal foraging theory, which states that an organism shows preference towards a prey item which provides a high profitability. The profitability of a food item is determined by E/h (where E= calorific value of the prey item, h= handling time). Therefore an organism which displays optimal foraging behaviour would forage for a highly profitable food source which is of high calorific value and requires little handling time. It is suggested that upon encountering a new prey item, an optimal foraging organism will assess whether the new prey is more than or equally as profitable in terms of calorific value and handling time as the average profitability of an item which is already included in its diet (i.e. E new/h $new \ge E_{average}/h_{average}$). Furthermore it is suggested by the MacArthur and Pianka (1966) that optimal foraging is influenced by prey abundance. When food is abundant a foraging organism's diet is specific as the consumer can afford to reject inferior prey since there is a high probability of encountering a more profitable prey item in the time it would take to capture and handle the previous one. Conversely when prey is scarce diets are broadened as there is

a low probability of encountering an optimal prey type.

Optimal foraging is thought to have evolved from an organism's innate necessity to maximise fitness and subsequent reproductive success (Stephens and Krebs 1986). The utilisation of a food source which provides all dietary requirements for minimal exertion would allow an organism to expend less energy on foraging activities and more energy on The reproduction. fundamental principle for maximising fitness is the survival of reproductively viable offspring (Darwin 1859). Therefore if a greater rate of energy gain allows for greater reproductive success then foraging beneficial optimal is in maximising fitness.

Optimal foraging can be related to food choice i.e. foods that convey the maximum net benefit (Scheel 1992, Jones 1990, Fryxell and Lundberg 1994). For example Mascaro and Seed (2001) found that Carcinus maenas, showed a preference towards species which bivalve gave the greatest profitability. However optimal foraging can also be related to the size of the selected optimal prey type. Several studies have investigated the effect of prey size on foraging behaviour. Werner and Hall (1974) studied prey size selection in bluegill sunfish and found that when daphnia were abundant only smaller individuals were selected, as prey abundance decreased larger individuals were selected in addition to the smaller organisms. Similarly Juanes and Conover (1994) found that when given a choice, blue fish consumed primarily small silverside fish due to the low times and hiah handling attack success rates. When small prev items became less abundant it was found that larger sub-optimal sized silverside fish were incorporated into their diet in addition to the more optimal small sized prey. An apparent conformity in the aforementioned studies is clear. When a prey species is abundant, smaller more profitable individuals are selected and larger individuals are rejected. As prey abundance individuals decreases larger are included in the diet, even though they may be less profitable. This concurs with a prediction from optimal foraging theory which states that a predator should reject less profitable prey items from its diet at higher densities of profitable prey (MacArthur and Pianka 1966, Bence and Murdoch 1986).

Molluscivore crabs have been at the forefront of studies testing the optimal foraging theory in relation to prey size selection (Elner and Hughes 1978, Hughes 1979, Hughes and Elner 1979, Hughes and Seed 1981, Blundon and Kennedy 1982, Arnold 1984). It has been shown that, in marine systems, the pattern of preference for smallsized molluscan prey is wide spread among decapod crustacean predators (Micheli 1995, Juanes 1992). Elner and Hughes (1978) observed that under unlimited prey conditions crabs chose mussels of а small to

intermediate size. close to the predicted optimum. This occurred after a brief period of manipulation which suggested that the crabs could assess the profitability of a prey item before selection occurred (concurrent with the foraging model suggested bv MacArthur and Pianka 1966). As the optimum sized mussels are depleted. crabs selected prey both above and below the optimum size. A possible explanation for the reluctance of less optimal size selection is that large mussels require the expenditure of excessive amounts of energy in order to gain access to the food source and very small mussels are often mishandled by the chelae. thus increasing handling times (Rheinallt 1986). Furthermore Hughes and Seed (1981) found that the blue crab, Callinectes sapidus, preferred smaller mussel sizes since it simultaneously minimised handling time and maximised net gain of energy intake. This proves that time minimisation plays an important role in prey size selection. Lawton and Hughes (1985) observed similar behaviour in juvenile Cancer pagurus feeding on Littorina littorea. Smaller prey were incorporated into the diet first as they succumbed more easily to an attack, this reduced the energy requirement armour for penetrating the and consequently reduced handling time.

Prey size selection is affected by prey abundance, but can also be affected by satiation levels. A study by Hart and Gill (1992) found that only small *A. aquaticus* were consumed when satiation was reached in the threespined stickleback. Thus prey choice becomes more selective as the necessitv to feed decreases. Conversely as hunger levels increase prev size selection becomes less specific and larger A. aquaticus are consumed, despite the fact that this size of prey might not result in optimal nutritional gains. The trade offs hunger between and prey size selection is well documented in literature for many organisms (Marti 1979. Molles and Hoque and Pietruszka 1986, Bence and Murdoch 1986). In contrast, while Carcinus maenas has been shown to select prey in a way which maximises the energy intake in accordance with the optimal foraging theory (Elner and Hughes 1978), hunger dependent prey size selection in this species has yet to be studied.

This presents study paper а investigating the effect of starvation on size selection in the green shore crab, Carcinus maenas, feeding on the edible mussel, Mytilus edulis. The green shore crab has been chosen for this study as it has a widespread distribution in estuarine and coastal waters and because is also known to forage extensively on commercially exploited bivalves (Ropes 1968). Consequently a greater understanding of the factors which effect prey selection may help control predation in commercial prey species. Subsequent laboratory experiments were designed ascertain whether prey size to

selection is affected by starvation level.

2. Methods

Carcinus maenas, of 24-26mm carapace width were collected from The Hoe, Plymouth (Nat. Grid Ref. SX 477 539). Only male crabs were selected, identified by а narrow abdomen. Mvtilus edulis ranging from 5-25mm in length were collected from mussel beds in Whitsand bav. Cornwall (Nat. Grid Ref. SW 362 275). Each crab was placed in a separate 15x20x15cm plastic holding aquarium, filled with aerated seawater and kept at 11°C. Each crab was fed a single lance fish, everyday for five days to standardise starvation. Any remnants of lance fish left over from the previous dav were removed before a fresh lance fish was added. The mussels were measured to the nearest mm using callipers and the length was marked on each shell using a graphite Cunningham and Hughes pencil. (1984) determined that C. maenas would only consume mussels up to half their size, however their experiment did not use crabs which had been starved for long periods of time, therefore mussels ranging from 5-25mm were used in the experiments.

All experiments were conducted using the same procedure. An experimental tank (30x20x15cm) was set up with aerated sea water and kept at 11°C. Observations were recorded over 6 consecutive days, with starvation increasing with each day. At the start of each observation a Perspex partition was placed to the right of the aquarium to restrain the crab. Twenty one mussels were added haphazardly on the other side, one for each mm interval ranging from 5-25mm. The partition was removed and the crabs' foraging behaviour was observed. In most cases the crab would handle various mussels before one was selected. To determine whether a mussel had been selected. observations were only recorded once an attempt was made to open the mussel using the mouthparts and mandibles to chip the posterior edges of the mussels' valves (Ameyaw-Akumifi and Hughes 1987). Any foraging behaviour observed up until this point was not considered as prey selection. Two series of experiments were performed; firstly to investigate hunger dependent diet selection, secondly to look at hunger dependent diet rejection.

2.1 Hunger dependent prey selection

Twelve male crabs were used, each with a carapace width between 24-26mm. As prev selection occurred, the crab was restrained by the partition and all the mussels were removed. including the one which was selected. The selected mussel was replaced with a new mussel of an equal size to avoid any bias in subsequent experiments, caused by a weakened shell. The observations were repeated three times for all twelve crabs for 6 starvation days.

2.2 Hunger dependent prey rejection

In the previous prey selection experiment it was observed that at low starvation, it was rarely the first mussel encountered which was subsequently selected. Frequently mussels were encountered and handled without any attempt to open them. Four crabs were used in this experiment due to time and resource limitation. Rather than recording the selected mussel size, in investigation this the foraging behaviour of each crab was observed and the size of anv mussel handled. encountered and before selection occurred, was recorded. This study was continued for 6 days of starvation.

Subsequent data analysis was performed using MINITAB version 15; graphs were created using the same software.

3. Results

3.1 Mussel size selection

Since triplicate observations were carried out per crab, a one way ANOVA for each starvation day was carried out to determine whether there was any significant difference in the three observations (Bartletts tests were applied in each case and no significant difference in variance was found therefore a one way ANOVA is a suitable statistical test).

Six out of seven starvation days showed no significant difference for the triplicate observations (Day 0 P=0.922, Day 1 P=0.403, Day 2 P=0.291, Day 3 P=0.404, Day 5 P=0.819, Day 6 P=0.494). Starvation day 4 did show a significant difference (P=0.036), however after reviewing the triplicate results for day 4 no clear trend was apparent. Furthermore since the majority of starvation days showed a vast insignificance in the triplicate

data this significance can be treated as an anomaly and is due to a type 1 error. Therefore all data can be pooled for subsequent statistical analysis.



Fig. 1- Seven graphs which show the frequency of selected mussel size with increased starvation time.

Crabs exhibited clear differences in mussel size selection with increased time. starvation There are two observed effects; firstly a change in mean mussel size and secondly a change in the variation of selected mussel size (see Fig 1). Relatively small mussels were selected at a low starvation time, with most crabs showing preference towards mussels <13mm during these periods. As starvation increases larger mussels are selected, although very large mussels (i.e. >20 mm) are always rejected. Mean median and modal sizes suggest that the size of mussels selected increases with starvation extreme time, however starvation appears to decrease the average selected mussel size (see fig 2).

A test for equal variance proved negative (Bartlett's Test P < 0.001) therefore a Kruskal-Wallis test was used as an alternative to a one way analysis of variance. There is a highly significant difference in the median size of mussels selected with starvation time (Kruskal-Wallis, H=62.63, DF=6, P <0.001). The median size of selected mussels increases with starvation until day 4 median where the mussel size decreases back to the day 1 starvation value by day 6 (i.e. 11 mm).

The mean mussel size increases with starvation time until day 4 where the mean size of mussels selected begins to decrease (see fig 2). For figure 2 a quadratic model (F=8.91, DF=2

P=0.034, R-sq = 81.7) proved far more accurate in calculating a regression line than a linear model (R-sq = 56.8). A regression analysis (F= 42.59, DF= 1,241, P <0.001) showed there is a highly significant difference in the mean size of mussels selected. Mean mussel size increases with starvation until day 4 where there is an apparent reduction in size selection.

Although there appears to be a parabolic relationship in figure 2, it is likely that the curve plateaus after 4 days. At low starvation levels, mussel selection is determined bv crab preference. As starvation increases mussel selection becomes more random. as preferences towards optimal prey sizes are negated by hunger mechanisms. If a crab is exhibiting random selection then the average selected mussel size would be a similar numerical value as the average size of mussels available for selection, ranging from 5-25mm (average=15mm). However mussels over 20mm were never selected, therefore the real range of mussel sizes is 5-20mm with an average of 12.5mm. At day 4 the mean size of mussels selected (12.53mm) is very close with the expected random selection size value (see fig 2), showing that this level of starvation can be associated with a random foraging behaviour. Therefore increased starvation (> 4 days) would cause similar randomised foraging behaviour and the apparent decrease in mean mussel size can be attributed to the random probability of encountering small prey sizes.

At low starvation times size selection is relatively specific with a small range in selected mussel size. As starvation increases a greater proportion of larger mussels are selected and selection becomes less specific and more random therefore increasing the range of selected mussel sizes (see fig 3).



Fig. 2- The mean size of mussels selected with starvation time. Quadratic regression line (Equation-Average Mussel size (mm) = 9.797 + 1.195 Ts - 0.1387 Ts²) (where Ts= starvation time) (R-sq = 81.7%). Confidence intervals (95%) included. Dotted line showing the mean size of mussel selected at day 4 starvation determined by the regression line (12.53mm).



Fig. 3- The standard deviation of the mussel size selected with starvation time. Curve fitted using a quadratic model (Equation- standard deviation = 1.075 + 0.0843 Ts + 0.04629 Ts²) which shows an exponential increase in standard deviation with starvation time. Confidence intervals included (95%)

ISSN 1754-2383 [Online] ©University of Plymouth The standard deviation of selected mussel size increases with starvation time. For figure 3 a quadratic model (F= 40.91, DF= 2 P< 0.001 R-Sq = 98.2) proved more accurate for a fitted line plot compared with a linear model (R-Sq = 92.8). The exponential increase in standard deviation (see fig 3) shows an increase in the variation of the size of mussels selected with starvation time.

3.2 Mussel size rejected

There is a strong positive correlation between the size of rejected mussels and starvation time (see figure 4). Regression analysis (F = 38.31, DF = 1, 91, P < 0.001) shows that as starvation time increases there is a highly significant difference in mussel size rejection. At low starvation levels relatively small mussels are rejected. As starvation levels increase the mean size of rejected mussels increases since smaller mussels are subsequently selected.

Since the size of rejected mussels increases with elevated starvation it could be expected that the number of mussels which were rejected by a crab would decrease with increased starvation. This is due to progressively larger mussels being selected at elevated starvation levels which would have been rejected at lower starvation levels. As the maximum size of a mussel which could be selected increases with starvation. fewer mussels which are deemed too large for consumption will be encountered, thus decreasing the number of rejections upon encounter.



Fig. 4 – The size of mussels rejected before one is selected with increasing starvation time. Liner regression line (Equation- Size of mussels rejected = 17.55 + 0.7603 Ts) showing a positive correlation. Confidence intervals included (95%).



Fig. 5- The mean number of mussels rejected with increased starvation time. Linear regression line (Equation- No. of mussels rejected = 1.777 - 0.2232 Ts) showing a negative correlation. Confidence intervals included (95%)

correlation between The negative number of mussels rejected and starvation time (see fig. 5) agrees with the expected observations. Regression analysis (F = 14.71, DF = 1, 82, P < 0.001) shows that there is a highly significant difference between number of mussels rejected and starvation time. At low starvation levels a greater number of mussels are rejected than at high starvation levels. This confirms selection that becomes less discriminate as starvation increases

4. Discussion

Carcinus maenas foraging behaviour becomes less specific with increased starvation. At low starvation levels prey size selection is specific, with relatively small mussels being selected, since these gave the most profitable gains for the handling time required (Elner and Hughes 1978). As starvation increased selection became more random, with a proportion of larger, less profitable sizes being selected even though optimal sized mussels were still available. This behaviour has been observed in literature for many other organisms. In a study of Screech Owls, Marti and Hougue (1979) found that when starved for 36 hours a greater proportion of large mice (30-49g) were selected, although a range of sizes were selected. In comparison, satiated owls displayed specific size selection with a high proportion of small mice (10-19g) being consumed. Molles and Pietruszka (1986)observed similar foraging behaviour in

the stonefly, *Hesperoperla pacifica*. Fasted stoneflies were found to consume a wide range of prey sizes, whereas satiated stoneflies concentrated attacks on intermediate prey.

The differences in foraging behaviour displayed by a satiated and a fasted organism can be explained by the probability of encountering optimal prey sizes. A satiated organism would have experienced foraging success, suggesting that there is an abundance of prey items in its foraging patch. Therefore there would be a high probability of subsequent encounters with optimal prey sizes, allowing the satiated organism to be selective in its foraging activities, rejecting prey types which were deemed less profitable. As hunger levels increase, an organism would have experienced a period of low/no prey abundance and therefore it can be assumed that there would be a low probability of encountering an optimal prey type. It must therefore approach foraging activity in a less specific manner, consuming organisms which would be deemed unprofitable at high satiation (MacArthur and Pianka 1966).

At low starvation *Carcinus maenas* were observed to consume relatively small mussels (7-12mm). Larger mussels (>14mm) and very small mussels (\leq 6mm) were rarely selected. If foraging behaviour in *C. maenas* was random at low starvation the mean selected mussel size would be similar to the average size of mussels available for selection. 12.5mm (>20mm mussels were excluded as they were never selected). However the mean selected mussel size for low starvation levels were considerably lower. This suggests that C. maenas assesses the profitability of a previtem selection occurs before and is concurrent with previous predictions of optimal foraging behaviour (MacArthur and Pianka 1966. Elner and Hughes 1978). As starvation increases foraging becomes increasingly random and by day 4 size selection is in accordance with predicted random prey size selection (i.e. 12.5mm). It can be assumed that if increasingly random foraging behaviour is associated with elevated starvation levels that the most profitable mussel size is selected at day 0 where satiation is at a maximum. Therefore the optimal mussel size for C. maenas 24-26mm width of carapace is approximately 9mm.

The increase in mean selected mussel size up until day 4 is associated with increasingly random foraging behaviour and is due to the lack of available prey species with elevated starvation. In the experiment increasing starvation times inevitably progressively fewer caused prev encounters by the crab and caused the selection of prey items larger than the predicted optimum. This is explained by the negative correlation between the number of mussels rejected and starvation time. As starvation increases a crab selects a mussel after fewer encounters and at high starvation levels selection generally occurred upon the first encounter regardless of size (with the exception of mussels >20mm). Random foraging elevated starvation occurs with the assessment of because the profitability of a mussel size is limited and the first mussel encountered is selected regardless of optimal size preference.

The towards preference smaller mussel sizes at low starvation can be explained by the optimal foraging theory in that a smaller mussel provides optimal nutritional gains per unit time (Schoener 1971). However Juanes (1992) suggests that size selection may be primarily due to chelae preservation. In this study it is suggested that crabs select smaller mussel sizes, since they require less force to be broken open, thus there is less risk of chelae damage. Chelae are an important asset for a crab as they are used in mating displays, defence of territory and defence from predators (Abello et al. 2004). Therefore it is suggested that the preservation of chelae is of high importance to the crab. Another explanation for the prey size selection is the consequences of a low handling time associated with small mussel sizes. Hughes and Seed (1981) suggest that predation stresses experienced by molluscivore crabs in the wild cause them to forage for smaller shelled molluscs, since handling times are greatly reduced. This allows crabs to minimise time spent foraging exposed in an

environment, thus reducing the risk of predation.

While the most probable explanation for an average increase in mussel size selection with lower starvation levels (1-4 days) is that foraging behaviour becomes increasingly random, another explanation could be that mussel size selection is dependent on gut satiation. Turesson et al. (2002) determined that prey size increase was disproportional to gut satiation. The limited size of the gut, might bias selection towards small prey because satiation allows only the consumption of small but not large Consequently starvation prev. as increases more space becomes available in the gut to accommodate larger prey items.

The smallest mussels (5mm) were selected throughout never the experiments regardless of starvation time. The low handling time, due to weak shell defences, and nutritional gains would theoretically make this size class suitable for selection when starvation causes a deviation from the optimal size. One explanation for this lack of selection is the limitation of chelae dexterity (Rheinallt 1986). Large size differences between mussel and crab chelae, causes increased difficulties in prey manipulation and therefore increased handling time. An increased handling time makes the nutritional gains from such a small prey negligible, therefore very small mussels are not selected.

Similarly the largest mussels (>20mm) were never selected throughout the

experiment regardless of starvation time. Furthermore the rejection of the largest mussels rarely occurred as handling and subsequent size assessment of large prey did not take place. This suggests that crabs use some visual cues when foraging for mussels and tactile foraging is not exclusive, this conclusion is contrary to that reached by Hughes and Seed (1995). Although crabs have the ability to access large mussels using an uneconomical boring technique (see Elner 1978), the energy requirements and long handling times for such make nutritional techniques gains negligible in comparison to the increased risk of predation.

The increase in the size of mussels rejected with starvation time is an expected observation in accordance with the optimal foraging theory. One of the main predictions of the model is that forager should accept а successively less profitably prey only when encounter and subsequent consumption rates with highly profitable prey fall below a critical level (MacArthur and Pianka 1966). When a crab experienced starvation in the experiment it encountered no prev items for a prolonged period of time. Furthermore encounter and consumption rates became increasingly scarce as starvation days increased. Therefore larger less optimal mussels progressively are selected as starvation increases.

In conclusion starvation has a profound effect on mussel size

selection in the shore crab, Carcinus maenas. At low starvation specific foraging behaviours can be observed which result in a size selection close to the predicted optimum. As starvation increases prev encounters are greatly reduced and selection becomes less specific as resulting in the selection of less profitable mussel sizes. Mussel sizes at both extremes are never selected since increased handling times make these sizes unprofitable. Rejection of mussels is less frequent as starvation time increases as the maximum size of a mussel deemed to be profitable increases with hunger level.

Further investigations should focus on the effect of starvation on foraging Since this study focused times. primarily on prey size selection, future should research be aimed at determining whether foraging times reduced when starvation are is experienced.

Acknowledgements

With thanks to Dr. M Uttley for his advice and assistance.

LITERATURE CITED

- Abello, P., C.G. Warman, D.G. Reid and E. Naylor. Chelae loss in the shore crab *Carcinus maenas* and its effect on mating success. *Marine Biology*. Vol 121. pg 247-252
- Ameyaw-Akumfi, C. and R.N. Hughes. 1987. Behaviour of *Carcinus Maenas* feeding on large *Mytilus edulis*. How do they assess an optimal diet?. *Marine Ecology Progress Series*. Vol 38. Pg 213-216.
- Arnold, W S. 1984. The effects of prey size, predator size, and sediment composition on the rate of predation of

the blue crab, *Callinectes sapidus Rathbun,* on the hard clam, *Mercenaria mercenaria. Journal of Experimental Marine Biology and Ecology.* Vol 80. Pg 207-219

- Bence, J.R. and W.W. Murdoch. 1986. Prey selection by the mosquitofish: Relation to Optimal Diet Theory. *Ecology*. Vol 67. No 2. Pg 324-336
- Blundon, J.A. and V.S. Kennedy. 1982.
 Mechanical and behavioural aspects of blue crab, *Callinectes sapidus*, predation on Chesapeake Bay bivalves. *Journal of Experimental Marine Biology and Ecology*. Vol 65. Pg 47-65
- Cunningham, P.N., and R.N. Hughes. 1984. Learning of predatory skills by shorecrab *Carcinus maenas* feeding on mussels and dog whelks. *Marine Ecology Progress Series*. Vol 16. Pg 21-26
- Darwin, C. 1859. Origin of the Species.
- Elner, R.W. 1978. The mechanics of predation by the shore crab, *Carcinus maenas*, on the edible mussel, *Mytilus edulis*. *Oecologia*. Vol 36. Pg 333-344
- Elner, R.W and R.N. Hughes. 1978. Energy maximization in the diet of the shore crab, *Carcinus maenas. Journal of Animal Ecology*. Vol 47. Pg 103-116
- Fryxell, J.M. and P. Lundberg. 1994. Diet choice and predator-prey dynamics. *Evolutionary ecology*. Vol 8. No 4. Pg 407-421
- Hughes, R.N. 1979. Optimal diets under the energy maximization premise: the effects of recognition time and learning. *American Naturalist*. Vol 113. Pg 209-221
- Hughes, R. N. and R.W. Elner. 1979. Tactics of a predator, *Carcinus maenas*, and morphological responses of the prey, *Nucella lapillus*. *Journal of Animal Ecology*. Vol 48. Pg 65-78
- Hughes, R.N. and R. Seed.1981. Size selection of Mussels by the Blue Crab *Callinectes sapidus*: Energy Maximizer or Time Minimizer?. *Marine Ecology Progress Series*. Vol 6. Pg 83-89
- Hughes, R.N., and R. Seed. 1995. Behavioural mechanisms of prey selection in crabs.

Journal of Experimental Marine Biology and Ecology. Vol 193. Pg 225-238

- Juanes, F. 1992. Why do decapod crustaceans prefer small-sized molluscan prey?. *Marine Ecology Progress Series*. Vol 87. Pg 239-249
- Juanes, F. and D. O. Conover. 1994. Piscivory and prey size selection in young-of-theyear bluefish: predator preference or size-dependent capture success?. *Marine Ecology Progress Series*. Vol 114. Pg 59-69
- Jones, G. 1990. Prey selection by the Greater Horseshoe Bat: Optimal foraging by Echolocation. *The Journal of Animal Ecology*. Vol 59. No 2. Pg 587-602
- Lawton, P and R.N. Hughes. 1985. Foraging behaviour of the crab *Cancer pagurus* feeding on the gastropods *Nucella lapillus* and *Littorina littorea:* comparisons with optimal foraging theory. *Marine Ecology Progress Series*. Vol 27. Pg 143-154
- MacArthur, R.H, and E.R. Pianka. 1966. On Optimal Use of a Patchy Environment. *The American Naturalist*. Vol 100. No 916. Pg 603-609
- Marti, C.D. and J.G. Hogue. 1979. Selection of prey by size in Screech Owls. *The Auk*. Vol 96. No 2. Pg 319-327
- Mascaro, M. and R. Seed. 2001. Choice of prey size and species in *Carcinus maenas* feeding on four bivalves of contrasting shell morphology. *Hydrobiologia*. Vol 449. Pg 159-170
- Micheli, F. 1995. Behavioural plasticity in prey–size selectivity of the blue crab *Callinectes sapidus* feeding on bivalve prey. *Journal of Animal Ecology*. Vol 64. Pg 63-74
- Molles Jr. M.C. and R.D. Pietruszka. 1986. Prey selection by a stonefly: influence of hunger and prey size. *Oecologia*. Vol 72. No 3. Pg 473-478
- Rheinallt, T. 1986. Size selection by the crab Liocarcinus puber feeding on mussels Mytilus edulis and on shore crabs Carcinus maenas: the importance of mechanical factors. Marine Ecology Progress Series. Vol 29. Pg 45-53

Ropes, J.W. The feeding habits of the green shore crab, *Carcinus maenas. Fisheries Bulletin.* Vol 67. Pg 183-203

Rovero. F., R.N. Hughes and G. Chelazzi. 2000. When time is of the essence: choosing a currency for prey handling costs. Journal of Animal Ecology. Vol 69. pg 683-689

Schoener, T.W. 1971. Theory of feeding strategies. *Annual review of Ecology and Sytematics*. Vol 2. Pg 369-404

Scheel, D. 1992. Profitability, encounter rates, and prey choice of African lions.

Behavioural Ecology. Vol 4. No 1. Pg 90-97

Stephens, D.W., and Krebs J. R. 1986. Foraging Theory. Priceton University Press. Princeton New Jersy

Turesson, H., A. Persson, C Bronmark. 2002. Prey size selection in piscivorous pikeperch (*Stizostedion lucioperca*) includes active prey choice. *Ecology of Freshwater Fish*. Vol 11. Pg 223-233

Werner, E.E. and D.J. Hall. Optimal foraging and the size selection of prey by the Bluegill Sunfish. *Ecology*. Vol 55. No 5. Pg 1042-1052