01 University of Plymouth Research Outputs

University of Plymouth Research Outputs

2019-02-05

A variant in the MICA gene is associated with liver fibrosis progression in chronic hepatitis C through TGF-1 dependent mechanisms.

Sharkawy, RE

http://hdl.handle.net/10026.1/13285

10.1038/s41598-018-35736-2 Scientific Reports Nature Research (part of Springer Nature)

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

A variant in the MICA gene is associated with liver fibrosis progression in chronic hepatitis C through TGF-β1 dependent mechanisms

Rasha El Sharkawy^{1*}, Ali Bayoumi^{1*}, Mayada Metwally¹, Alessandra Mangia², Thomas Berg³, Manuel Romero-Gomez⁴, Maria Lorena Abate⁵,William L. Irving⁶, David Sheridan⁷, Gregory J. Dore⁸, Ulrich Spengler⁹, Pietro Lampertico¹⁰, Elisabetta Bugianesi⁵, Martin Weltman¹¹, Lindsay Mollison¹², Wendy Cheng¹³, Stephen Riordan¹⁴, Rosanna Santoro², Rocío Gallego-Durán⁴, Janett Fischer³, Jacob Nattermann⁹, RobertaD'Ambrosio¹⁰, Duncan McLeod¹⁵, Elizabeth Powell¹⁶, Olivier latchoumanin¹, Khaled Thabet^{1,17}, Mustafa A.M. Najim^{1,18}, Mark W. Douglas^{1,19}, Christopher Liddle¹, LiangQiao¹, Jacob George¹ and Mohammed Eslam¹, for the International Liver Disease Genetics Consortium (ILDGC).

* These authors contributed equally

¹Storr Liver Centre, The Westmead Millennium Institute for Medical Researchand Westmead Hospital, University of Sydney, and Westmead Hospital NSW, Australia

²Division of Hepatology, Ospedale Casa Sollievo della Sofferenza, IRCCS, San Giovanni Rotondo, Italy

³Section of Hepatology, Clinic for Gastroenterology and Rheumatology, University Clinic Leipzig, Leipzig, Germany

⁴Unit for The Clinical Management of Digestive Diseases and CIBERehd, Hospital Universitario Virgen del Rocío, University of Seville, Sevilla, Spain.

⁵Division of Gastroenterology and Hepatology, Department of Medical Science, University of Turin, Turin, Italy

⁶NIHR Biomedical Research Unit in Gastroenterology and the Liver, University of Nottingham, Nottingham, United Kingdom

⁷Institute of Translational and Stratified Medicine, Plymouth University, United Kingdom

⁸Kirby Institute, The University of New South Wales, Sydney, NSW, Australia.

⁹Department of Internal Medicine I, University of Bonn, Bonn, Germany

¹⁰ Università degli Studi di Milano Centro A.M. e A. Migliavacca, First Division of Gastroenterology, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Department of Pathophysiology and Transplantation Milan Italy

¹¹Department of Gastroenterology and Hepatology, Nepean Hospital, Sydney, NSW, Australia

¹²Department of Gastroenterology and Hepatology, Fremantle Hospital, Fremantle, WA, Australia

¹³ Department of Gastroenterology & Hepatology, Royal Perth Hospital, WA, Australia.

¹⁴Gastrointestinal and Liver Unit, Prince of Wales Hospital and University of New South Wales, Sydney, NSW, Australia

¹⁵Department of Anatomical Pathology, Institute of Clinical Pathology and Medical Research (ICPMR), Westmead Hospital, Sydney, Australia

¹⁶The University of Queensland, School of Medicine, Princess Alexandra Hospital, Woolloongabba, QLD, Australia

¹⁷Biochemistry Department, Faculty of Pharmacy, Minia University, Minia, Egypt

¹⁸Department of Medical Laboratories Technology, Faculty of Applied Medical Sciences, Taibah University, Medina, Saudi Arabia

¹⁹Centre for Infectious Diseases and Microbiology, Marie Bashir Institute for Infectious Diseases and Biosecurity, University of Sydney at Westmead Hospital, Westmead NSW, Australia

Corresponding Author

Jacob George

Department of Medicine

Westmead Hospital

Westmead, NSW 2145

Ph: 61-2-98457705; Fx 61-2-96357582

Email: jacob.george@sydney.edu.au

Running title: MICA and liver fibrosis

Key words: Chronic hepatitis C, fibrosis, DEPDC-5, MICA, TGFβ1

Conflicts of interest: All authors disclose no conflicts or competing interests.

Acknowledgements: We would like to thank all the patients for their participation in this study. ME, CL, MD and JG are supported by the Robert W. Storr Bequest to the Sydney Medical Foundation, University of Sydney; a National Health and Medical Research Council of Australia (NHMRC) Program Grant (APP1053206, APP1149976) and Project grants (APP1107178 and APP1108422). GD is supported by an NHMRC Fellowship (1028432). US and JN are supported by DFG SFB-TR57 and the Hector-Foundation (M63). RE and KT are supported by a scholarship from the Egyptian government. AB is supported by an Australian Government Research Training Program (RTP) scholarship.

Authors' contribution: R.E., A.B., M.M., M.E. and J.G. conceived the research; patient enrolment, clinical phenotype data collation and sample acquisition/DNA preparation was performed by R.E., A.B., M.M, M.E, J.G., M.R.-G., A.M., W.L.I, T.B., G.J.D., D.S., M.L.A., E.B., M.W., L.M., W.C., S.R., R.S., J.F., U.S., J.N., P.L., R.D., R.G-D., M.W.D, E.B., C.L., O.L and L.Q; genotyping was performed by K.T, M.M; histological analysis of tissues was conducted by D.M.; statistical analysis and interpretation of results was performed by R.E., A.B., M.M., M.E. and J.G.; RT–PCR: R.E.; ELISA: A.B.; Tissue culture: R.E, JFH-1 data: M.AM.N., the manuscript was written and revised by R.E., A.B., M.M, M.E. and J.G. All authors critically reviewed the manuscript and approved the final submitted manuscript.

Data availability: All data are provided within the main text and supplementary file.

Abstract (N=185 words)

Hepatocarcinogenesis is tightly linked to liver fibrosis. Recently, two GWAS variants, MICA rs2596542 and DEPDC5 rs1012068 were identified as being associated with the development of HCV-induced hepatocellular carcinoma (HCC) in Japanese patients. The role of these variants on hepatic inflammation and fibrosis that are closely associated with HCC development is not known, nor are the biological mechanisms underlying their impact on the liver. Here, we demonstrate in 1689 patients with chronic hepatitis C (CHC) (1,501 with CHC and 188 with HCV-related HCC), that the MICA (T) allele, despite not being associated with HCC susceptibility, is associated with increased fibrosis stage (OR: 1.47, 95% CI: 1.05-2.06, p=0.02) and fibrosis progression rate (hazards ratio: 1.41, 95% CI: 1.04-1.90, p=0.02). The DEPDC5 variant was not associated with any of these phenotypes. MICA expression was downregulated in advanced fibrosis stages. Further, (T) allele carriage was associated with lower MICA expression in liver and serum. Transforming growth factor- β 1 (TGF- β 1) expression suppresses MICA expression in hepatic stellate cells. Our findings suggest a novel mechanism linking susceptibility to advanced fibrosis and subsequently indirectly to HCC, to the level of MICA expression through TGF- β 1-dependent mechanisms.

Introduction

Chronic hepatitis C virus (HCV) infection is a leading cause of cirrhosis, hepatocellular carcinoma (HCC) and liver transplantation, with an estimated 339,000 people dying annually from complications ¹. Advanced liver fibrosis or cirrhosis represents the major risk factor for developing liver-related complications and mortality, but currently there are no approved anti-fibrotic therapies ².

The rate of liver fibrosis progression varies greatly according to disease etiology and also between individuals; the latter is at least partially attributable to genetic factors. In chronic HCV infection, host genetics play a pivotal role in shaping the immune response, virus-host interactions and ultimately the predilection to and progress of liver fibrosis ^{3,4}. This risk is likely polygenic and dependent on multiple genetic and epigenetic factors since variations in single loci are usually of modest effect size and explain only a small fraction of the phenotype ⁵. This has led in the recent past to a shift towards the discovery of novel variants with limited effects which ultimately could guide the development of polygenic scores with high predictive value.

Two genome wide association studies (GWAS) have investigated the risk of HCV-related HCC in Japanese patients. The first identified a locus in the 5' flanking region of the MHC class I-related chain A (*MICA*) on 6p21.33 (rs2596542) to be strongly associated with HCC and the progression from CHC to HCC ⁶. MICA is a ligand for natural-killer group 2 member D (NKG2D), a highly conserved C-type lectin-like membrane glycoprotein and one of the major activating receptors on NK cells ⁷. The second GWAS identified a susceptibility locus near *DEPDC5* (rs1012068) ⁸. The function of DEPDC5 is not well understood, but has been linked with hereditary forms of epilepsy ⁹, bladder cancer ¹⁰ and malignant glioblastomas ¹¹.

Complicating the interpretation of these findings however is that HCC development in chronic HCV infection is tightly linked to hepatic fibrosis with 90% of cases arising in cirrhotic livers ¹². Hence, risk variants that predispose to fibrosis could be associated with HCC formation without direct causality and disentangling the two is problematic. In this regard, little is known about the potential effect of variants in *MICA* and *DEPDC5* on liver fibrosis since both GWAS were done in patients in whom liver biopsy was not available. Importantly, functional data on the role of these variants with regard to both fibrosis pathways or HCC development are limited. The available literature is also restricted to Japanese populations with chronic HCV infection and HCC, while a single report in Caucasians ¹³ suggests that *DEPDC5* but not *MICA* is associated with fibrosis progression.

Here we sought to dissect the role of *MICA* rs2596542 and *DEPDC5* rs1012068 to liver fibrosis and to HCV-related HCC. To do this, these variants were assessed in 1,501 patients with CHC of Caucasian ancestry in whom liver biopsy was available and were compared to 188 patients with CHC-related HCC. We undertook functional studies to explore the mechanisms that might underlie the genetic association with fibrosis.

Results

Patient characteristics

The clinical, demographic and biochemical characteristic of the patients in the cohort with chronic HCV infection (n=1501) are presented in **supplementary table 1**. The genotype distribution of *MICA* rs2596542 and *DEPDC5* rs1012068 was in Hardy-Weinberg equilibrium and is presented in **supplementary table 2**. The minor allele frequency (MAF) for the two variants was similar to that observed in a healthy European population from the 1000 genome project (<u>http://browser.1000genomes.org</u>), which has some difference from the Japanese population.

Association of *MICA* rs2596542 and *DEPDC5* rs1012068 with viral and clinical characteristics

To explore if baseline clinical variables differ between chronic HCV patients according to *MICA* rs2596542 or *DEPDC5* rs1012068 genotype, we examined the association of the genotypes with baseline clinical variables; the results are presented in **supplementary tables 3 and 4, respectively**. There was no evidence of association between either rs2596542 or rs1012068 genotype with any clinical variable (i.e. age, BMI, baseline levels of ALT, AST, GGT, ALP, platelets, leukocytes, HCV-RNA quantification, gender frequency or HCV genotype distribution).

MICA rs2596542, but not DEPDC5 rs1012068 is associated with fibrosis severity

We next assessed the association between *MICA* rs2596542 and *DEPDC5* rs1012068 and liver damage (hepatic inflammation and fibrosis). The distribution of rs2596542 and rs1012068 genotypes according to histological features is depicted in **Figure 1**.

First, we examined the association of the two genetic variants with fibrosis. In multivariate analyses adopting an additive model adjusted for age, gender, BMI, HCV genotype and alcohol intake, *MICA* rs2596542 was significantly associated with stage of fibrosis (adjusted estimate, 0.072, SE, 0.051; p=0.01) (**Supplementary Table 5**). In further analysis when the cohort was segregated into those with mild (F0–1) versus advanced fibrosis (F2–4), carriage of each copy of the *MICA* rs2596542 (T) allele was associated consistently with a significant increased risk of advanced fibrosis, independent of age, gender, BMI, HCV genotype and alcohol intake (OR: 1.47, 95% CI: 1.05-2.06, p=0.02) (**Supplementary Table 6**). Similarly, rs2596542 was associated with severe fibrosis (F3-F4) (OR: 1.43, 95% CI: 1.03-1.99, p=0.03). On the contrary, we did not observe any association between *DEPDC5* rs1012068 and stage of fibrosis (**Supplementary Table 5**); the distribution of rs1012068 genotypes was also not significantly different according to the presence or absence of significant or severe fibrosis (**Supplementary Table 6**).

We next examined the association of the two genetic variants with hepatic inflammation defined by liver histopathology scored according to METAVIR. Neither *MICA* rs2596542 nor *DEPDC5* rs1012068 demonstrated any association with inflammation grade (**Supplementary Table 5**). No association was also observed when the cohort was dichotomized into absent/mild (A0-A1) versus moderate/severe (A2-A3) inflammation (**Supplementary Table 6**). Consistently, no difference was noted in serum liver enzymes (ALT or AST; as indices of liver injury) according to rs2596542 or rs1012068 (**supplementary tables 3 and 4, respectively**).

Association between MICA rs2596542 and fibrosis progression

To validate these observations, we undertook analysis in the 815 patients from the entire cohort with chronic HCV infection and a known duration of infection. This allowed assessment

of the relationship to fibrosis progression without the inherent biases of cross sectional analyses. The baseline characteristics of the cohort were similar among subjects included and not included in the fibrosis progression sub-analysis (**Supplementary Table 7**). As the fibrosis progression rate (FPR) may not be constant over time ¹⁴, we used Cox-proportional hazards analysis. In this analysis, rs2596542 was associated with an increased hazard of progression to significant fibrosis (\geq F2) in a multivariate model that included age, gender, BMI and HCV genotype (hazards ratio: 1.41, 95% CI: 1.04-1.90, p=0.02 per each T allele). In contrast, *DEPDC5* rs1012068 was not associated with FPR (hazards ratio: 1.06, 95% CI: 0.76-1.48, p=0.7) (**Supplementary Table 8**). In sum, these data suggest that *MICA* rs2596542 but not *DEPDC5* rs1012068 is associated with fibrosis severity and fibrosis progression.

Chronic HCV infection increases hepatic MICA expression

In the subsequent studies reported below, we sought to define the functional basis for the association of variants in *MICA* to liver fibrosis. The expression of MICA is low in healthy tissues but can be induced by cellular stressors such as viral infection ¹⁵; the effect of HCV infection on hepatic MICA expression is not known. We therefore compared MICA expression in liver from patients with HCV and from control, non-infected subjects. RT-PCR demonstrated increased MICA mRNA expression in patients with chronic HCV infection (**Figure 2a**). We then assessed MICA expression in the JFH1/Huh7 *in vitro* model of replicating HCV. Consistently, Huh7 cells infected with the JFH1 strain of HCV demonstrated a 3-fold upregulation of MICA mRNA compared to control uninfected cells (**Figure 2b**), p<0.05).

The biological effect of MICA in HCV-infected liver likely reflects a complex interaction between the different cell types within an inflamed liver that ultimately leads to fibrosis. Hence, we examined the expression of MICA in four primary human hepatic cell types, namely hepatocytes, Kupffer cells, hepatic sinusoidal endothelial cells and stellate cells by real time PCR. The highest expression of MICA was in stellate cells followed by Kupffer cells, while sinusoidal cells and hepatocytes had low expression (**Supplementary Figure 1**).

Membranous and soluble MICA levels associate with MICA genotype and fibrosis severity

Whether the *MICA* risk allele is associated with altered membrane bound (mMICA) expression in liver tissue from patients with chronic HCV infection is not known as all available current data are based on the soluble form (sMICA) present in serum ^{6,16}. As shown in **Figure 2c**, we observed a significant association between rs2596542 genotype and MICA mRNA expression; the (T) allele associated with lower hepatic MICA mRNA in the 94 patients with available liver tissue (P = 0.01, p=0.02 after adjustment for age and sex).

We then measured soluble MICA (sMICA) by ELISA in serum samples from a subcohort of 214 chronic HCV infection patients; their characteristics are summarized in **Supplementary Table 9** and matched the overall cohort. Similar to mMICA, compared to patients harboring the *MICA* rs2596542 CC genotype, sMICA levels were significantly reduced in those with the risk T allele (**Figure 2d**). We determined the correlation of mMICA to sMICA in 58 patients in whom paired liver and serum data was available. In this analysis, levels of sMICA associated with the expression of mMICA (p = 0.0001) (**Figure 2e**).

In total, MICA expression is upregulated in HCV infection in an rs2596542 genotype dependent manner with individuals carrying rs2596542 (T) expressing lower levels of mMICA and sMICA. This suggests that differences in expression levels of mMICA according to genotype are not a consequence of differing shedding rates of MICA.

MICA expression and liver fibrosis

The overall contribution of the MICA/NKG2D pathway to fibrosis is unclear. Interestingly and consistent with the genetic data, hepatic MICA expression was significantly lower in subjects with significant hepatic fibrosis (F2–F4) compared to those with no or mild hepatic fibrosis (F0/1) (n=94) (**Figure 2f**). MICA expression did not correlate with hepatic inflammation, liver enzymes, HCV-RNA levels and genotype, age or gender. Similarly, sMICA levels were decreased in patients with significant fibrosis compared to those without, though this was not significant (**supplementary Figure 2**).

MICA expression is regulated by TGF-B1 in hepatic stellate cells

We next thought to explore how differential expression of hepatic MICA could be linked to fibrosis progression. Hepatic stellate cells (HSC) are responsible for the deposition of extracellular matrix that is evident histologically as fibrosis ¹⁷ and as we have shown, also have the highest expression of MICA. Killing of activated HSCs by NK cells to ameliorate liver fibrosis is in part by a MICA/NKG2D-dependent manner ¹⁸. Thus, we investigated the regulation of MICA expression on HSCs in a fibrotic context.

Surface expression of MICA has been reported to be down-regulated by TGF- β 1¹⁹, an established growth factor mediating liver injury and fibrosis¹². Hence, we characterized TGF- β 1 mediated regulation of MICA expression on HSCs. We first assessed expression of TGF- β 1 mRNA after HCV infection *in vitro* using qRT-PCR. This revealed a significant increase in TGF- β 1 in HCV-infected JFH-1/Huh7 cells compared to non-infected cells (**Figure 3a**). Consistently, we observed increased TGF- β 1 transcription in liver from chronic HCV infected liver samples compared to healthy controls (**Figure 3b**). Within the chronic HCV infection group, the expression of TGF- β 1 mRNA was significantly up-regulated with advancing of

fibrosis stage (**Figure 3b**) and this was consistently observed in the sub-cohort with available serum described above (n=214) (**Figure 3c**).

Given these results, we examined for the effect of TGF- β 1 on MICA expression on a human HSC-derived cell line (LX2). We observed a marked repression of MICA mRNA transcripts in TGF- β 1 stimulated LX2 cells (**Figure 3d**), as well as reduced surface expression of MICA, by flow cytometry (**Figure 3e**). For confirmation that this effect was mediated through the TGF β R, we used a specific pharmacological inhibitor of TGF β R-I/II (using LY2109761) and this as expected, reversed the inhibitory effect of exogenous TGF β 1 on MICA expression (**Figure 3d and e**). To characterize the role of endogenous HSC-derived TGF- β 1 in the regulation of MICA expression, we used LX2 cells treated with LY2109761 and demonstrated a significant induction of MICA mRNA transcripts and surface expression levels (**Figure 3d and e**).

Lastly, previous reports have observed a positive effect of TGF- β 1 treatment on the expression of some proteases including MMP-9. Since MICA surface expression can be modulated at a post-translational level by proteolytic cleavage ²⁰, we asked whether shedding of sMICA was enhanced in TGF- β 1 stimulated cells, thus contributing to the attenuation of surface expression on these cells. To this end, we measured sMICA in cell culture supernatant and showed that the level of sMICA released by TGF- β 1 stimulated cells was not different compared with control cells suggesting that cleavage of MICA is not mediated by TGF- β 1-dependent proteases (**Figure 3f**).

MICA polymorphism is not associated with HCC

As *MICA* was originally discovered as a risk locus for HCC in Japanese patients, we investigated the role of *MICA* in hepatocarcinogenesis. We compared *MICA* genotype

distribution in patients with chronic HCV-HCC (n=188) to non-HCC from the entire chronic HCV infection cohort described above (n=1,501; **Supplementary Table 10**). We observed no difference in *MICA* rs2596542 distribution between subjects with HCC and those without. Because HCC mainly occurs in the context of cirrhosis, we undertook similar analysis restricting the comparison to those with HCC versus those with chronic HCV infection and cirrhosis (n=210). Again, no difference in *MICA* rs2596542 distribution was observed between subjects with HCC and those with cirrhosis (p=0.77). Results were not changed after adjustment for age and sex. These findings imply that *MICA* rs2596542 polymorphism is unlikely to be directly associated with the occurrence of HCC.

Discussion

Here, we report that a genetic variant in *MICA* originally identified by GWAS as a risk variant for susceptibility to HCV-associated HCC in Japanese patients ⁶ associates with fibrosis severity and progression, but not with HCC. Functionally our data suggest that the association with HCC is plausibly mediated by indirect effects through modulation of fibrosis risk. Following the data for effects on fibrosis progression, we show *MICA* genotype-dependent reductions in hepatic MICA gene expression and soluble MICA protein levels. Tying these findings to fibrosis, TGF- β 1 was increased with advancement of fibrosis and there was a corresponding reduction in MICA expression. *In vitro*, TGF- β 1 treatment resulted in MICA transcriptional and surface protein repression on HSC cells. These effects we propose leads to an increased likelihood of fibrosis in those with the risk genotype. Our findings therefore suggest a molecular explanation for the association of rs2596542 with liver fibrosis (**Figure 4**).

Consistent with the present findings in the setting of chronic HCV infection, a recent report suggests that *MICA* polymorphisms (MICA*012:01/02, MICA*017 and MICA*027) are associated with liver fibrosis in schistosomiasis ²¹. It is noteworthy that in the original GWAS report of *MICA* in HCV-related HCC, Kumar et al., did not use HCV-related cirrhosis without HCC as their control cohort, masking any potential confounding by fibrosis ⁶. This and the failure to find an association with *MICA* in a similar Japanese study published 2 months later ⁸, as well as the functional data we provide, suggest that the reported association with HCC actually relates to liver fibrosis risk.

MICA is a stress inducible protein present at very low levels in healthy tissues, but can be induced by cellular stresses such as viral infection ¹⁵. Consistently, we demonstrated increases in MICA expression during the early stages of HCV infection, while it decreases with increasing

fibrosis in an rs2596542 genotype dependent manner. Furthermore, the (T) risk allele correlated with lower mMICA in liver. Consistent with other groups 6,16 , we demonstrated decreased expression levels of sMICA in the serum of patients with the (T) rs2596542 allele.

The mechanisms that link *MICA* genotypes with fibrosis risk have not been previously elucidated. MICA is a ligand for the activating NK cell receptor NKG2D and this interaction shapes the anti-fibrotic capacity of NK-cells via elimination of activated HSCs ^{18,22} We verified that HCV infection is associated with an increase in TGF- β 1 expression in liver ²³ and the latter induces marked repression of MICA expression in HSC. This suggests that *MICA* genotype might be associated with alterations in TGF- β 1 (through as yet undefined mechanisms) and may underlie the accelerated fibrosis we observed.

We did not observe any association between *DEPDC5* rs1012068, the other variant identified by GWAS as a risk variant for HCC ⁸, with either fibrosis or HCC. A large Japanese study of 2,266 HCV patients who had eradicated HCV and with long term follow up to 20 years, was also unable to find any association between *DEPDC5* SNPs at rs1012068 and the cumulative risk of HCC ²⁴. Some studies have reported an association of *DEPDC5* with fibrosis but with opposite effects of the risk allele ^{13,25,26}. A recent smaller study conducted in Europeans suggested an association between *DEPDC5* rs1012068 but not *MICA* rs2596542 and risk of cirrhosis in chronic HCV infection ¹³. The reasons for the discrepancy between our study and this work are unclear, but may relate to differences in baseline characteristics and that the association analyses for fibrosis were performed differently, as those authors compared F0-F1 vs. F4. Interestingly, in that study, *MICA* rs2596542 showed a tendency for significance with cirrhosis (p=0.07) in the discovery cohort (n=477) but was not investigated in the validation cohort. The function of DEPDC5 in liver needs to be defined.

Our study has some limitations, including the fact that the sample size of the PCR and ELISA sub-cohorts was relatively modest. In addition, whether *MICA* rs2596542 is the functional polymorphism or if there are other variants in linkage disequilibrium and the detailed functional mechanisms for the effects are still not fully understood.

In conclusion, we have demonstrated that *MICA* rs2596542 is associated with liver fibrosis progression, likely mediated and amplified in part through TGF β -1 dependent mechanisms.

Figures legends:

Figure 1: Association of *MICA* rs2596542 and *DEPDC5* rs1012068 with inflammation, **fibrosis stage.** Association of *MICA* rs2596542 with fibrosis stage (a) and inflammation (b) and *DEPDC5* rs1012068 with fibrosis stage (c) and inflammation (d) in the CHC cohort (n=1501). P-values are univariate and provided for the additive model of inheritance. *MICA* rs2596542 (T) allele is the risk allele, the same risk allele in the GWAS by Kumar et al (6). *DEPDC5* rs1012068 (G) allele is the risk allele, the same risk allele in the GWAS by Miki et al (8).

Figure 2: Hepatic and serum MICA expression. Relative hepatic MICA mRNA expression in 94 patients with chronic hepatitis C virus infection and 28 controls (a); relative MICA mRNA expression in Huh7 cells infected with the JFH1 strain hepatitis C virus as compared to mock infected control cells (b); correlation between *MICA* rs2596542 genotype and hepatic MICA mRNA (c) and soluble MICA levels (n =214) (d). The x axis shows the genotypes at rs2596542 and the y axis shows MICA expression level relative to GAPDH by quantitative real-time PCR or the concentration of soluble MICA in pg/ml by ELISA, respectively. The level of hepatic MICA mRNA correlates positively with serum soluble MICA levels in a sub-cohort of CHC with available RNA and serum (n=58) (e); hepatic MICA mRNA according to hepatic fibrosis (f). The x axis shows hepatic fibrosis dichotomized as absent/mild (METAVIR stage F0–F1) or moderate/severe (METAVIR stage F2–F4), and the y axis shows hepatic MICA expression. The number of independent samples tested in each group is shown below the figure and P value was calculated using the two-tailed Student's t-test or ANOVA test with Tukey test for correction for multiple comparisons.

Figure 3: Transforming growth factor beta influences MICA expression. Relative TGF- β 1 mRNA expression in Huh7 cells infected with the JFH1 strain of the hepatitis C virus compared to mock infected control cells (a); relative hepatic TGF- β 1 mRNA expression in 94 patients with chronic hepatitis C virus infection and 28 controls according to fibrosis stage (b); serum concentration of TGF- β 1 in pg/ml levels according to hepatic fibrosis (n=214) (c); MICA mRNA expression and protein levels at the hepatic stellate cell surface modulation by TGF- β 1. MICA mRNA expression was assessed in control, TGF- β 1 (2ng/ml) with or without LY2109761 (a TGF β R-I/II inhibitor) (100nM, was added 90 minutes before TGF- β 1), or LY2109761 (100nM)

treated cells for 24 hours by real-time PCR (d), or 48 hours for flow cytometry (e); the level of soluble MICA was assessed in the supernatant of control or TGF- β 1 (2ng/ml) treated LX2 cells for 24 and 48 hours by ELISA (f). Results are expressed, as mean ± sem (n = 3) and P value was calculated using the two-tailed Student's t-test or ANOVA test with Tukey test for correction for multiple comparisons. *P < 0.05,** P<0.001, *** P<0.001.

Figure 4: Proposed model for the effect of *MICA* **rs2596542 in modulating liver fibrosis risk.** Patients carrying the *MICA* risk genotype have low hepatic and serum MICA expression. TGF-β1 further suppresses MICA expression on hepatic stellate cells, leading to increased liver fibrosis.

Methods

Patient cohort

The study comprised 1689 consecutive patients (1,501 with CHC and 188 with HCVrelated HCC) from the International Liver Disease Genetics Consortium (ILDGC) database. Details of the cohort and inclusion criteria have been reported ²⁷⁻²⁹. Briefly, all subjects who had a liver biopsy with scoring for fibrosis stage and disease activity before anti-viral treatment, and genomic DNA were included. For those with CHC, those with HCV RNA serum positivity of Self-reported Caucasian descent were included. Patients were excluded if they had evidence of co-infection with other hepatitis viruses, other liver diseases by standard tests or current or previous hepatic decompensation. Diagnosis of HCC was based on the EASL–EORTC Clinical Practice Guidelines ³⁰.

The study protocol was conformed to the ethical guidelines of the 1975 Declaration of Helsinki. Ethics approval was obtained from the Human Research Ethics Committees of the Western Sydney Local Health District and the University of Sydney. All other sites had ethics approval from their respective ethics committees and study was performed according to the recommendations of the centers involved. Written informed consent for genetic testing was obtained from all participants.

Clinical and laboratory assessment

The following data were collected at time of liver biopsy from all patients: gender, age, ethnicity, recruitment center, alcohol intake (g/day), body mass index (BMI) and routine laboratory tests. BMI was calculated as weight divided by the square of the height (kg/m^2).

Methods to estimate the duration of infection

Fibrosis progression was examined in 815 chronic HCV infection patients with a reliable estimated duration of infection, as previously reported ^{31,32}. Briefly, for subjects with a history of injecting drug use, the time of infection was estimated using the reported "first year of injection". For patients with a history of blood transfusion, the onset of infection was assumed to be the year of transfusion. For patients with a history of occupational exposure, the onset of infection was assumed to be the first year of needle stick exposure. The duration of infection was calculated by subtracting the estimated age at infection from age at biopsy.

Genotyping

Genotyping for *MICA* rs2596542 and *DEPDC5* rs1012068 was contracted to the Australian Genome Research Facility (AGRF; QLD, Australia). 1051 samples were genotyped using the Sequenom MassARRAY system and iPLEX Gold chemistry while 638 samples were genotyped using the TaqMan SNP genotyping allelic discrimination method (Applied Biosystems, Foster City, CA, USA). All genotyping was blinded to clinical variables.

Liver Histopathology

Liver histopathology was scored according to METAVIR³³. Fibrosis was staged from F0 (no fibrosis) to F4 (cirrhosis). Necroinflammation (A) was graded as A0 (absent), A1 (mild), A2 (moderate) or A3 (severe). The inter-observer agreement between pathologists was studied previously and was good ($\kappa = 77.5$) for METAVIR staging using κ statistics ³⁴.

Other Methods

Methods for cell culture, real-time PCR, ELISA and flow Cytometry are described in Materials and Methods of the Supporting Information

Statistical Analysis

Statistical analyses were performed using the statistical software package SPSS for Windows, version 21 (SPSS, Chicago, IL). All tests were two-tailed and p values <0.05 were considered significant. Details of statistical analyses are described in the Materials and Methods of the Supporting Information.

References

- 1. WHO. Hepatitis C. Fact sheet. Updated April 2017.
 - http://www.who.int/mediacentre/factssheets/fs164/en/ (accessed 30 July 2017).
- 2. Iredale, J.P. Models of liver fibrosis: exploring the dynamic nature of inflammation and repair in a solid organ. *J Clin Invest* **117**, 539-48 (2007).
- 3. Romero-Gomez, M., Eslam, M., Ruiz, A. & Maraver, M. Genes and hepatitis C: susceptibility, fibrosis progression and response to treatment. *Liver Int* **31**, 443-60 (2011).
- 4. Eslam, M. & George, J. Genome-Wide Association Studies and Hepatitis C: Harvesting the Benefits of the Genomic Revolution. *Semin Liver Dis* **35**, 402-20 (2015).
- 5. Gibson, G. Rare and common variants: twenty arguments. *Nat Rev Genet* **13**, 135-45 (2011).
- 6. Kumar, V. *et al.* Genome-wide association study identifies a susceptibility locus for HCV-induced hepatocellular carcinoma. *Nat Genet* **43**, 455-8 (2011).
- 7. Raulet, D.H. Roles of the NKG2D immunoreceptor and its ligands. *Nat Rev Immunol* **3**, 781-90 (2003).
- 8. Miki, D. *et al.* Variation in the DEPDC5 locus is associated with progression to hepatocellular carcinoma in chronic hepatitis C virus carriers. *Nat Genet* **43**, 797-800 (2011).
- 9. Ishida, S. *et al.* Mutations of DEPDC5 cause autosomal dominant focal epilepsies. *Nat Genet* **45**, 552-5 (2013).
- 10. Harada, Y. *et al.* Cell-permeable peptide DEPDC1-ZNF224 interferes with transcriptional repression and oncogenicity in bladder cancer cells. *Cancer Res* **70**, 5829-39 (2010).
- 11. Seng, T.J. *et al.* Complex chromosome 22 rearrangements in astrocytic tumors identified using microsatellite and chromosome 22 tile path array analysis. *Genes Chromosomes & Cancer* **43**, 181-193 (2005).
- 12. Zhang, D.Y. & Friedman, S.L. Fibrosis-dependent mechanisms of hepatocarcinogenesis. *Hepatology* **56**, 769-75 (2012).
- 13. Burza, M.A. *et al.* DEPDC5 variants increase fibrosis progression in Europeans with chronic hepatitis C virus infection. *Hepatology* **63**, 418-27 (2016).
- 14. Poynard, T. *et al.* Rates and risk factors of liver fibrosis progression in patients with chronic hepatitis c. *J Hepatol* **34**, 730-9 (2001).
- 15. Raulet, D.H., Gasser, S., Gowen, B.G., Deng, W. & Jung, H. Regulation of ligands for the NKG2D activating receptor. *Annu Rev Immunol* **31**, 413-41 (2013).
- 16. Huang, C.F. *et al.* Genetics Variants and Serum Levels of MHC Class I Chain-related A in Predicting Hepatocellular Carcinoma Development in Chronic Hepatitis C Patients Post Antiviral Treatment. *EBioMedicine* **15**, 81-89 (2017).
- 17. Tsuchida, T. & Friedman, S.L. Mechanisms of hepatic stellate cell activation. *Nat Rev Gastroenterol Hepatol* **14**, 397-411 (2017).
- 18. Krizhanovsky, V. *et al.* Senescence of activated stellate cells limits liver fibrosis. *Cell* **134**, 657-67 (2008).
- 19. Friese, M.A. *et al.* RNA interference targeting transforming growth factor-beta enhances NKG2D-mediated antiglioma immune response, inhibits glioma cell migration and invasiveness, and abrogates tumorigenicity in vivo. *Cancer Res* **64**, 7596-603 (2004).
- 20. Groh, V., Wu, J., Yee, C. & Spies, T. Tumour-derived soluble MIC ligands impair expression of NKG2D and T-cell activation. *Nature* **419**, 734-8 (2002).
- 21. Gong, Z. *et al.* Association of MICA gene polymorphisms with liver fibrosis in schistosomiasis patients in the Dongting Lake region. *Brazilian Journal of Medical and Biological Research* **45**, 222-229 (2012).

- 22. Radaeva, S. *et al.* Natural killer cells ameliorate liver fibrosis by killing activated stellate cells in NKG2D-dependent and tumor necrosis factor-related apoptosis-inducing ligand-dependent manners. *Gastroenterology* **130**, 435-52 (2006).
- 23. Taniguchi, H. *et al.* Hepatitis C virus core protein upregulates transforming growth factor-beta 1 transcription. *J Med Virol* **72**, 52-9 (2004).
- Nagaoki, Y. *et al.* Hepatocellular carcinoma development of hepatitis C virus patients with eradication to interferon therapy: a large scale, long-term study of 2266 patients. *Hepatology* 60, 1162a-1163a (2014).
- 25. Motomura, T. *et al.* Neither MICA Nor DEPDC5 Genetic Polymorphisms Correlate with Hepatocellular Carcinoma Recurrence following Hepatectomy. *HPB Surg* **2012**, 185496 (2012).
- 26. Al-Anazi, M.R. *et al.* Variations in DEPDC5 gene and its association with chronic hepatitis C virus infection in Saudi Arabia. *BMC Infect Dis* **14**, 632 (2014).
- 27. Eslam, M. *et al.* IFN-lambda 3, not IFN-lambda 4, likely mediates IFNL3-IFNL4 haplotypedependent hepatic inflammation and fibrosis. *Nature Genetics* **49**, 795-+ (2017).
- 28. Thabet, K. *et al.* MBOAT7 rs641738 increases risk of liver inflammation and transition to fibrosis in chronic hepatitis C. *Nat Commun* **7**, 12757 (2016).
- 29. Eslam, M. *et al.* Diverse impacts of the rs58542926 E167K variant in TM6SF2 on viral and metabolic liver disease phenotypes. *Hepatology* **64**, 34-46 (2016).
- 30. European Association For The Study Of The, L., European Organisation For, R. & Treatment Of, C. EASL-EORTC clinical practice guidelines: management of hepatocellular carcinoma. *J Hepatol* **56**, 908-43 (2012).
- 31. Eslam, M. *et al.* Interferon-lambda rs12979860 genotype and liver fibrosis in viral and non-viral chronic liver disease. *Nat Commun* **6**, 6422 (2015).
- 32. Eslam, M. *et al.* FibroGENE: A gene-based model for staging liver fibrosis. *J Hepatol* **64**, 390-398 (2016).
- 33. Bedossa, P. & Poynard, T. An algorithm for the grading of activity in chronic hepatitis C. The METAVIR Cooperative Study Group. *Hepatology* **24**, 289-93 (1996).
- 34. Eslam, M. *et al.* IFNL3 polymorphisms predict response to therapy in chronic hepatitis C genotype 2/3 infection. *J Hepatol* **61**, 235-41 (2014).

Collaborators: The International Liver Disease Genetics Consortium (ILDGC)

Rose white¹, Angela Rojas⁴, Margaret Bassendine²⁰, Chiara Rosso⁵, Lavinia Mezzabotta ⁷, Reynold Leung¹, Barbara Malik³, Gail Matthews⁸, Jason Grebely⁸, Vincenzo Fragomeli¹²,

Julie R. Jonsson⁸.

²⁰ Institute of Translational Medicine, Newcastle University, UK













