

2019-03-17

# Searching for the "Active Ingredients" in Physical Rehabilitation Programs Across Europe, Necessary to Improve Mobility in People With Multiple Sclerosis: A Multicenter Study

Kalron, A

<http://hdl.handle.net/10026.1/13574>

---

10.1177/1545968319834893

Neurorehabilitation and Neural Repair

SAGE Publications

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

**Article published in *Neurorehabil Neural Repair* 1545968319834893 17 Mar 2019**

**Searching for the “active ingredients” in physical rehabilitation programs across Europe, necessary to improve mobility in people with multiple sclerosis: a multicenter study.**

**Alon Kalron PT,PhD<sup>1,2</sup>, Peter Feys PT,PhD<sup>3</sup>, Ulrik Dalgas PhD<sup>4</sup>, Tori Smedal PT,PhD<sup>5,6</sup>, Jennifer Freeman PT,PhD<sup>7</sup>, Anders Romberg PT,PhD<sup>8</sup>, Helen Conyers PT<sup>9</sup>, Iratxe Elorriaga PT<sup>10</sup>, Benoit Gebara PT<sup>11</sup>, Johanna Merilainen PT<sup>12</sup>, Adnan Heric-Mansrud PT<sup>13</sup>, Ellen Jensen PT<sup>14,15</sup>, Kari Jones PT<sup>6</sup>, Kathy Knuts PT<sup>16</sup>, Benoit Maertens de Noordhout MD<sup>17</sup>, Andrej Martic PT<sup>18</sup>, Britt Normann PT,PhD<sup>19</sup>, Bert O Eijnde PT,PhD<sup>20</sup>, Kamila Rasova PhD<sup>21</sup>, Carme Santoyo Medina MSc.PT<sup>22</sup>, Ilse Baert PT,PhD<sup>3</sup>**

<sup>1</sup>Department of Physical Therapy, Sackler Faculty of Medicine and Sagol School of Neurosciences, Tel-Aviv University, Tel-Aviv, Israel

<sup>2</sup>Multiple Sclerosis Center, Sheba Medical Center, Tel-Hashomer, Israel

<sup>3</sup>REVAL Rehabilitation Research Center, BIOMED Biomedical Research Institute, Faculty of Rehabilitation Sciences, Hasselt University, Diepenbeek, Belgium

<sup>4</sup>Section of Sport Science, Dep. Public Health, Aarhus University, Aarhus, Denmark

<sup>5</sup>Norwegian Multiple Sclerosis Competence Centre, Department of Neurology, Haukeland University Hospital, Bergen, Norway

<sup>6</sup>Department of Physiotherapy, Haukeland University Hospital, Bergen, Norway.

<sup>7</sup>Faculty of Health and Human Sciences, Plymouth University, Devon, UK

<sup>8</sup>Masku Neurological Rehabilitation Center, Masku, Finland

<sup>9</sup>Poole Hospital, NHS Foundation Trust, Dorset, UK

<sup>10</sup>Eugenia Epalza Rehabilitation Center, Bilbao, Spain

<sup>11</sup>National Multiple Sclerosis Center, Melsbroek, Belgium

<sup>12</sup>Open Rehabilitation Center Aksoni, Helsinki, Finland

<sup>13</sup>Multiple Sclerosis Center, Hakadal AS, Norway

<sup>14</sup>Multiple Sclerosis Hospital, Haslev, Denmark

<sup>15</sup>Multiple Sclerosis Hospital, Ry, Denmark

<sup>16</sup>Rehabilitation and Multiple Sclerosis Center, Overpelt, Belgium

<sup>17</sup>Centre Neurologique et de Readaptation Fonctionnelle, Fraiture-en-Condroz, Belgium

<sup>18</sup>University Medical Center, Ljubljana, Slovenia

<sup>19</sup>University of Tromsø the Arctic University of Norway/Nordland Hospital Trust, Bodø, Norway

<sup>20</sup>REVAL Rehabilitation Research Center, BIOMED Biomedical Research Institute, Faculty of Medicine & Life Sciences, Hasselt University, Diepenbeek, Belgium

<sup>21</sup>Department of Rehabilitation, Third Faculty of Medicine, Charles University, Prague, Czech Republic

<sup>22</sup>MS Center of Catalonia (Cemcat), Vall Hebron University Hospital, Barcelona, Spain

**Corresponding author:** Alon Kalron, PT, PhD, Department of Physical Therapy, Sackler

Faculty of Medicine, Tel-Aviv University, Tel-Aviv, Israel, Tel: 972-9-9512726;

Mobile phone: 972-52-2436839; E-mail address: [alkalron@gmail.com](mailto:alkalron@gmail.com)

Word count of the text: 2966

Number of figures: 2

Number of tables: 5

## **Abstract**

*Background.* Physical rehabilitation programs can lead to improvements in mobility in people with multiple sclerosis (PwMS). *Objective:* Identify which rehabilitation program elements are employed in real life and how they might impact mobility improvement in PwMS.

*Methods.* Participants were divided into improved and non-improved mobility groups based on changes observed in the Multiple Sclerosis Walking Scale-12 following multimodal physical rehabilitation programs. Analyses were performed at group and subgroup (mild and moderate-severe disability) levels. Rehabilitation program elements included: setting; number of weeks; number of sessions; total duration, therapy format (individual, group, autonomous), therapy goals and therapeutic approaches. *Results.* The study comprised 279 PwMS from 17 European centers. PwMS in the improved group received more sessions of individual therapy in both subgroups. In the mildly disabled group, 60.9% of the improved received resistance training, whereas, 68.5% of the non-improved, received self-stretching. In the moderately-severely disabled group, 31.4% of the improved, received aerobic training, while 50.4% of the non-improved, received passive mobilization/stretching. *Conclusions.* We believe that our findings are an important step in opening the black-box of physical rehabilitation, imparting guidance and assisting future research in defining characteristics of effective physical rehabilitation.

## **Keywords**

multiple sclerosis, physical rehabilitation, mobility, walking, treatment elements

## ***Introduction***

Although numerous trials have established the benefits of immunomodulatory drugs in decelerating the inflammatory-related progression of multiple sclerosis (MS),<sup>1</sup> the disease remains incurable with ambulation difficulties worsening with disease progression. Consequently, physical rehabilitation remains a key factor in improving (or maintaining) mobility in people with MS (PwMS).

Rehabilitation programs comprise various therapeutic approaches (e.g. resistance training, balance training, etc.), intensity (e.g. how hard the body is taxed), volume (e.g. frequency, number and duration of sessions) and format (e.g. individual or group based; home or center-based). It has been reported that the content and approach of physical therapy for PwMS differs throughout certain European countries.<sup>2</sup> Nevertheless, very few studies have hitherto investigated which elements are currently being employed during rehabilitation sessions and which are of sufficient importance in improving mobility in PwMS. In terms of therapeutic approaches, several systematic reviews have examined the effect of physical rehabilitation programs on ambulatory outcomes in PwMS.<sup>3-7</sup> Although there is strong evidence showing that exercise therapy improves mobility related activities,<sup>3,4</sup> others have demonstrated only a limited beneficial impact.<sup>5,8</sup>

Recently, Khan & Amatya published a systematic review of systematic reviews examining rehabilitation in PwMS<sup>9</sup> reporting strong evidence for physical rehabilitation programs and moderate evidence for a range of rehabilitative treatments and approaches available for PwMS. Nevertheless, there is still a lack of evidence for most modalities and limited comparative knowledge of their efficacy across the disability spectrum due to the use of different outcome measures and limited standardized reporting of the elements in rehabilitation programs (e.g. combination of modalities and delivery modus). As such, it is difficult to provide sufficiently detailed guidelines to clinicians and patients. Additional

information is needed in order to identify core elements in physical rehabilitation programs, aimed at improving mobility in PwMS.

In our previous publication of 290 PwMS from 17 European centers, we examined the responsiveness and clinically meaningful improvement of five walking measures.<sup>10</sup> The participating centers collaborated with the European Rehabilitation within the Multiple Sclerosis (RIMS) network. Our main finding was that long walking tests and the self-reported MS walking scale-12 (MSWS-12) could detect clinically meaningful improvement after physical rehabilitation.<sup>10</sup> The aim of the present study, (a secondary analysis), was to assess real-life physical rehabilitation programs across Europe and explore the elements (format, volume, therapy goals and approaches) which could positively affect mobility in PwMS. Our hypothesis was that active treatment approaches (e.g. muscle strengthening, aerobic training) would greatly impact mobility compared to passive approaches (e.g. stretching).

## ***Methods***

### ***Participants***

A convenience sample of 290 PwMS were recruited from 17 centers within the RIMS network. A full description of participating centers was presented in our previous publication.<sup>10</sup> This study was approved by the Ethical Committee of the Hasselt University, Belgium and local ethics committees from each participating center. Inclusion criteria comprised a definite diagnosis of MS<sup>11</sup> and the Expanded Disability Status Scale (EDSS)<sup>12</sup> score of  $\geq 2$  and  $\leq 6.5$  as determined by neurologists. All patients participated in a physical rehabilitation program for 3 to 12 weeks. Inclusion criteria included a minimum of 10 sessions and a maximum duration of 3 months. Subjects were excluded if afflicted with other medical conditions interfering with walking. All subjects provided written informed consent.

### ***Rehabilitation Program Format and Volume***

The physical rehabilitation programs varied from center to center and included guided or supervised sessions by physiotherapists and/or sport/fitness instructors. We acknowledge that there are differences in the educational level of professionals providing physical therapy in MS across Europe.<sup>13</sup> Each participating site supplied information regarding their program, ie, setting; number of weeks; number of sessions; total duration and therapy format (individual, group or autonomous). Settings were neuro-rehabilitation centers with or without an overnight stay, hospitals with or without an overnight stay, private physical therapy practices, community centers, fitness centers, research facilities and patient residences. These were dichotomized as in- and outpatient rehabilitation settings. A maximum of two settings were indicated when settings were combined. Individual therapy was defined as a 1:1 ratio between therapist and patient. The duration of each session lasted 40-50 minutes. Number of sessions and total duration were recorded for each therapy format.

### ***Therapy Goals and Therapeutic Approaches***

Classification of therapy goals and therapeutic approaches were based on Rasova et al's study.<sup>14,15</sup> Therapists were required to describe for each patient one primary goal of the physical rehabilitation program: "improving balance", "improving walking capacity", "maintenance of balance and walking" and "others, unrelated to balance and walking". In order to identify the main therapeutic approaches employed during the intervention program, the therapist selected a maximum of 5 items (out of 21) from a list of approaches (e.g resistance training with equipment, balance training, gait training, passive mobilisation, aerobic training, etc).

### ***Walking Measures***

Clinical walking tests used at entry and discharge of the rehabilitation program included:

- 1) Two- and six-minute walk test (2mWT, 6mWT). The participants were instructed to complete the test 'at their fastest speed' and cover as much distance as possible by

walking up and down a 30 meter hallway, using their own walking aid. The 2mWT and 6mWT have been validated and used extensively in PwMS.<sup>16</sup>

2) The MSWS-12 is a valid patient-reported questionnaire rating walking ability in PwMS.<sup>17,18</sup> Each item is scored on a 1 to 5 scale; the higher the score, the more perceived walking difficulties. A total score is generated and converted to a 0 to 100 scale with negative change scores indicating improvement.

3) Timed 25-Foot Walk (T25FW) was performed at a normal and fastest walking speed. Participants were instructed to walk at their own comfortable pace for the normal trial and as quickly as possible for the fast trial. The T25FW has been validated as one of the three components of the Multiple Sclerosis Functional Composite.<sup>19</sup>

### ***Statistical Analysis***

The sample group was divided into two groups: improved and non-improved walkers. Allocation was determined according to the MSWS-12's minimally important change (MIC) scores as presented in our previous publication.<sup>10</sup> The MSWS-12 was selected due to its acceptance in clinical trials investigating rehabilitation and pharmacological interventions in the MS population. Additionally, its validity has been confirmed in community-residing and hospital outpatient samples of PwMS<sup>17,18</sup>.

The MIC cut-off point for the total group was -11.35; in the mildly disabled subgroup -10.7 and in the moderately-severely disabled -11.85. PwMS with change scores equal or above the MIC were assigned to the improved group and those scoring below the MIC were assigned to the non-improved group. Furthermore, the participants were categorized as either "mildly" (EDSS  $\leq 4$ ) or "moderately-severely" disabled (EDSS  $>4$ ), according to their disability level. Classification of disability subgroups corresponded with our previous publications on walking measures in PwMS.<sup>10,20</sup> All data followed a normal distribution according to the Kolmogorov-Smirnov test. Box plots determined outliers for each outcome.



Descriptive statistics were used for demographic, clinical characteristics, rehabilitation elements and mobility measures. Differences between the improved and non-improved walkers were examined by the chi-square test for MS type, gender, setting, therapy goals and by analysis of variance tests (ANOVA) for age, disease duration, EDSS, mobility measures, volume of therapy and therapeutic approaches.

Binary logistic regression analyses, with a forward method, examined the relationship between the improved/non-improved status (dependent variable) and the rehabilitation elements (setting, therapy goal, volume of therapy and therapeutic approaches) (independent variables). The regression analysis was performed separately according to disability subgroups. For all models, the assumptions underlying regression were tested by inspecting the distribution of the error term. All analyses were carried out using the SPSS software (version 25.0 for Windows; SPSS Inc., Chicago, Illinois, USA). Reported P values were two-tailed. The level of significance was set at  $p < 0.05$ .

## ***Results***

Out of the total 290 PwMS, 11 participants were missing data, therefore, only a total of 279 PwMS were included in the final analysis: 131 were assigned to the improved; 148 to the non-improved group. No significant differences between groups were observed for age, gender distribution, type of MS, disease duration and mean EDSS score. Descriptive characteristics of the sample according to group allocation and disability level are provided in Table 1.

Place Table 1 here

In terms of disability subgroups, 46 patients (47.9%) in the mildly disabled and 70 (38.3%) in the moderately-severely disabled, improved in mobility. No differences were detected between the improved and non-improved subgroups as to demographic and clinical characteristics on the moderately-severely disability level. A similar observation was recorded in the mildly disabled group, with the exception of the mean EDSS score which was higher in

those who improved compared to those who did not, 3.3 (S.D.=0.8) vs. 2.8 (S.D.=1.0),  $p=0.013$ . Descriptive and clinical characteristics according to group allocation and disability level are provided in Table 1.

Table 2 and figure 1 present the mobility outcome measures according to group allocation and disability level. For the total sample, significant differences were found in favor of the improved compared to the non-improved group for the 2mWT and the 6mWT. No differences between groups were found for the T25FW. According to the 2- and 6mWT, PwMS improved their walking distance by approximately twice the length of the non-improved PwMS. Although, individuals in the non-improved group showed improvements in the long distance walking tests, it was below the MIC cutoff score.

Place Table 2 and Figure 1 here

No differences were found between the improved and non-improved groups in the T25FW, 2 and 6MWT mobility tests for people with mild MS. In contrast, significant differences in favor of the improved group were found for the long distance walking tests in moderately-severe PwMS. As for the MSWS-12 questionnaire, participants in the improved group (both disability levels) presented with a ~26% improvement at discharge. Conversely, people with mild or moderately-severe MS, categorized as non-improved, reported more mobility difficulties (according to the MSWS-12) following the physical rehabilitation program.

The majority (72%) of the MS sample received rehabilitation in an outpatient setting. No significant differences in the amount of therapy was found between the mildly and moderately-severely disabled subgroups. PwMS in the improved group received a greater amount of individual therapy compared to the non-improved in both disability subgroups. People with mild MS who had improved, received approximately twice the number of individual therapy sessions and double the duration of therapy hours compared to those who

had not improved. Participants in the improved group, classified as moderately-severely disabled, received approximately 25% more individual therapy compared to those in the non-improved group. No differences were observed between the improved and non-improved groups in program settings and amount of autonomous therapy, regardless of disability level. Table 3 presents the rehabilitation program elements according to group allocation and disability level.

Place Table 3 here

Approximately 42% of the therapists, selected "improvement in walking capacity" as the primary goal for participants who had improved, irrespective of disability level. According to the chi-square tests, there were significant differences in therapy goals between the improved and non-improved groups in the mildly disabled subgroup;  $\chi^2(3)=18.537, p<0.001$  and in the moderately-severely disabled subgroup;  $\chi^2(3)=16.538, p=0.001$ . Figure 2 illustrates therapy goal distribution according to group allocation and disability level.

Place Figure 2 here

The therapist was instructed to select a maximum of 5 items (out of 21) from a list of therapeutic approaches. Significant differences were found in the mildly disabled group for two items; 60.9% of the participants who had improved, had received resistance training compared with 26.0% who had not improved; whereas, 68.5% of those who had not improved, had received self-stretching compared to 39.1% who had improved. As for the moderately-severely disabled group, 31.4% who had improved had received aerobic training compared to 14.2% who had not improved). In patients who had not improved, 50.4% received passive mobilization/stretching compared to 31.4% who had improved), 38.9% received resistance training compared to 21.4% who had improved. Table 4 presents the therapy approaches according to group allocation and disability level.

Place Table 4 here

Table 5 presents the results of the binary regression analysis in the mildly and moderately-severely disabled subgroups. A significant relationship was found between group allocation with resistance training and self-stretching therapeutic approaches in the mildly disabled subgroup. These variables explained 21.8% of the variance related to improved/non-improved group allocation. As for the moderately-severely disabled subgroup, passive mobilization/stretching and individual therapy volume explained 14.3% of the variance related to improved/non-improved status.

Place Table 5 here

### ***Discussion***

We report herein on a secondary analyses of a RIMS multicenter study primarily aimed at investigating the responsiveness of outcome measures in rehabilitation. Our aim was to assess physical active rehabilitation programs and explore the elements of these programs which positively affect mobility in PwMS. Upon conclusion of the rehabilitation program, 47% of PwMS participants demonstrated a clinical meaningful improvement in the MSWS-12. A novel feature of the current study was that while most of the previous studies investigating physical rehabilitation in PwMS had observed improvements in mobility according to the standard significance level, we defined mobility improvements according to meaningful clinical values.<sup>10</sup>

To the best of our knowledge, this is the first multicenter study encompassing a wide range of physical rehabilitation programs applied in the MS population, using identical outcome measures and standardized reporting terminology. We combined the data obtained from 17 MS rehabilitation centers in 9 countries. Each center had constructed their program according to the needs of their local MS population, rehabilitation facilities and the regulations/guidelines of its health care providers. Mobility assessments were performed in a standardized manner at all centers according to a detailed instruction booklet provided by the project

steering group. Although the diverse rehabilitation programs and the possibility that therapists employed multiple approaches per patient may affect the data and can be considered a limitation, we feel that it might strengthen the ecological validity of our findings.

Furthermore, in the current study we did not want to change the existing physical rehabilitation programs. The format was to observe and record the current status of physical rehabilitation across Europe in PwMS<sup>14,15</sup>, without formulating up a well-designed (controlling co-founders), scientifically strong robust study. We believe that our findings are an important step in opening the black-box of physical rehabilitation, imparting guidance and assisting future research in defining characteristics of effective physical rehabilitation.

There is a wide-ranging consensus as to the amount of physical rehabilitation needed for PwMS: more practice is probably better, but “how much more” remains an unanswered question. We found that the volume of individual therapy was significantly higher in improved versus non-improved patients, consistent in both disability groups, suggesting that the volume of rehabilitation is a key factor for a successful rehabilitation in PwMS. This statement is in accordance with a Cochrane review examining physical rehabilitation approaches in patients recovering function and mobility following stroke.<sup>21</sup> Nevertheless, our findings partially contrast with Snook & Motl’s systematic review who reported on the effect of exercise on mobility performance in PwMS. They found no differences in effect size according to the number of sessions per week and minutes per session.<sup>4</sup> Unfortunately, the authors did not calculate the total amount of exercise therapy and consequently, a direct comparison with our findings is not possible.

Findings related to the therapy goals were somewhat confusing. The goal "Other" was the main choice in the mildly disabled participants who did not improve, however, this was also the main goal of choice in the moderate-severe disabled participants. According to the study protocol, the therapists were asked to record only the primary goal, although the training

sessions may have included additional secondary goals (reflecting real life practice), such as improving walking capacity and/or balance. For instance, in cases where improving aerobic capacity was the main goal, the therapist chose the option "Other". However, this choice does not necessarily mean that secondary objectives such as improving walking and/or balance, were not included in the session. We therefore, advise to consider this finding with caution.

Currently, there is no consensus as to whether group or individual therapy (or a combination of both) sessions are preferred when seeking to improve mobility functions in PwMS.<sup>22,23</sup> Although, group-based physical rehabilitation sessions may be more efficient by potentially allowing more visits per patient than individual physical therapy sessions, there are still several potential disadvantages to this approach. One disadvantage is the lack of flexibility in tailoring interventions according to the varied functional levels of individual MS patients. Moreover, it is more difficult to match the patient's cognitive and psychological function in group therapy compared with individual therapy treatment goals. On the other hand, advantages of group therapy include social interaction and peer support between participants. Our findings suggest that individual therapy should be preferred over group or autonomous therapy in order to improve mobility in the MS population. Nevertheless, future research is warranted with a direct comparison of an identical rehabilitation program provided in individual vs. group therapy settings.

In the mildly disabled group, a significant higher proportion of patients received resistance training in the improved group versus the non-improved. The opposite result was seen in the moderately-severely disabled group (Table 4). A positive contribution of resistance training on mobility has been well-documented in the MS population.<sup>24-26</sup> The mechanisms involved may include increases in the neural drive,<sup>27</sup> a larger muscle fiber size<sup>28</sup> and enhanced neuromuscular adaptations.<sup>29</sup> For patients with a moderate-severe disability, aerobic training, rather than resistance training, was associated with clinical meaningful mobility

improvements. This finding supports the results of previous studies investigating the effect of aerobic exercise in PwMS with severe mobility deficits.<sup>30,31</sup>

Self-stretching and passive mobilization/stretching were more commonly used as a treatment approach in PwMS who did not improve. Passive movements, joint mobilization and stretching techniques generally produced only modest beneficial effects or even detrimental effects on performance in athletes<sup>32</sup> Our results also indicate that these therapy modalities appear to be less beneficial compared to more physically demanding modalities aiming to improve mobility in PwMS. The fact that stretching did not emerge as most effective, may be related to the patient sample composition, with limited patients demonstrating a need for stretching.

Group classification was based on the MSWS-12, the most widely used patient-reported measure of perceived limitation in walking due to MS. This decision was reached due to evidence from multiple studies supporting its robust measurement performance<sup>10,17-18,20,33-35</sup>. Furthermore, the MSWS-12 includes running, stair climbing, balance, concentration and effort needed to walk, therefore, measuring broader aspects of mobility. Moreover, Pilutti et al. have shown that the MSWS-12 scores correlate with the T25FW, 6mWT and gait kinematics in PwMS.<sup>32</sup> Some argue that this scale might not be an ideal choice for measuring walking impairment in mildly disabled PwMS or those who walk without a device.<sup>34</sup> Nevertheless, Langeskov-Christensen et al recently reported that the MSWS-12 captures impairments more gradually than the 2MWT and 6MWT in people with mild MS, thus, suggesting that the MSWS-12 is acceptable when assessing walking in PwMS with a low disability status.<sup>35</sup>

Our study has some limitations. Firstly, mobility was evaluated only by clinical walking tests and a patient-reported questionnaire. Utilizing instrumented gait devices that provide definite gait characteristics, might have expanded our knowledge. However, only a few of the

centers participating in this study possessed these tools.<sup>36</sup> Secondly, we did not explore the impact of different combinations of treatment modalities on an individual level. An alternative could have been to apply a standardized recording form of rehabilitation taxonomy which requires time recordings of different components per session.<sup>37</sup> More detailed data would have strengthened our findings, however, recording such detailed information after each session was not feasible for the majority of involved therapists. Finally, the aim was not to improve mobility in all patients. As such, applied intervention modalities may have also suited other therapy goals.

### ***Conclusion***

This report presents data of the core elements of physical rehabilitation programs associated with improving mobility in PwMS from 17 MS centers across Europe. Suggested elements to improve mobility include the volume of individual therapy sessions, resistance and aerobic therapeutic approaches and placing less emphasis on passive therapeutic techniques. Future studies should investigate these treatments.



### ***Acknowledgements***

Coordination of this study was partially funded via an unrestricted educational grant from Novartis Pharma AG to the European RIMS network, acknowledged for facilitating inter-European consultation and testing. The staff of each participating site and participating subjects are acknowledged for their cooperation.

### ***Conflict of Interest Statement***

The authors declare that there is no conflicts of interest.

### ***Funding***

This study was partially funded via an unrestricted educational grant from Novartis Pharma AG to RIMS.

## References

1. Martin R, Sospedra M, Rosito M, et al. Current multiple sclerosis treatments have improved our understanding of MS autoimmune pathogenesis. *Eur J Immunol.* 2016;46: 2078-2090.
2. Rasova K, Freeman J, Martinkova P, et al. The organisation of physiotherapy for people with multiple sclerosis across Europe: a multicentre questionnaire survey. *BMC Health Serv Res.* 2016; 16: 552.
3. Pearson M, Dieberg G, Smart N. Exercise as a therapy for improvement of walking ability in adults with multiple sclerosis: a meta-analysis. *Arch Phys Med Rehabil.* 2015; 96: 1339-1348.
4. Snook EM and Motl RW. Effect of exercise training on walking mobility in multiple sclerosis: a meta-analysis. *Neurorehabil Neural Repair.* 2009; 23: 108-116.
5. Latimer-Cheung AE, Pilutti LA, Hicks AL, et al. Effects of exercise training on fitness, mobility, fatigue, and health-related quality of life among adults with multiple sclerosis: a systematic review to inform guideline development. *Arch Phys Med Rehabil.* 2013; 94: 1800-1828.
6. Kjolhede T, Vissing K, Dalgas U. Multiple sclerosis and progressive resistance training: a systematic review. *Mult Scler.* 2012;18:1215-1228.
7. Cruickshank TM, Reyes AR, Ziman MR. A systematic review and meta-analysis of strength training in individuals with multiple sclerosis or Parkinson disease. *Medicine.* 2015; 94: e411.
8. Motl RW, Sandroff BM, Kwakkel G, et al. Exercise in patients with multiple sclerosis. *Lancet Neurology.* 2017;16:848-856.
9. Khan F and Amatya B. Rehabilitation in multiple sclerosis: A systematic review of systematic reviews. *Arch Phys Med Rehab.* 2017;98:353-367.

10. Baert I, Freeman J, Smedal T, et al. Responsiveness and clinically meaningful improvement, according to disability level, of five walking measures after rehabilitation in multiple sclerosis: a European multicenter study. *Neurorehabil Neural Repair*. 2014; 28:621-631.
11. Polman CH, Reingold SC, Banwell B, et al. Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald criteria. *Ann Neurol*. 2011;69:292-302.
12. Kurtzke JF. Rating neurologic impairment in multiple sclerosis: an Expanded Disability Status Scale (EDSS). *Neurology*. 1983;33:1444-1452.
13. Rasova K, Matinkova P, Pavlinkova M, et al. Physical therapy provision in multiple sclerosis across Europe: a regional lottery? *Eur J Phys Rehabil Med*. 2015;51(6):80-52.
14. Rasova K, Martinkova P, Cattaneo D, et al. Physical therapy in multiple sclerosis differs across Europe: information regarding an ongoing study. *J Int Med Res*. 2014;42:1185.
15. Martinokova P, Freeman J, Drabinova A, et al. Physiotherapeutic interventions in multiple sclerosis across Europe: Regions and other factors that matter. *Mult Scler Relat Disord* 2018;22:59-67.
16. Gijbels D, Alders G, Van Hoof E, et al. Predicting habitual walking performance in multiple sclerosis: relevance of capacity and self-report measures. *Mult Scler*. 2010;16: 618-626.
17. Hobart JC, Riazi A, Lamping DL, et al. Measuring the impact of MS on walking ability: the 12-Item MS Walking scale (MSWS-12). *Neurology*. 2003;60:31-36.
18. McGhuigan C, Hutchinson M. Confirming the validity and responsiveness of the Multiple Sclerosis Walking Scale-12 (MSWS012). *Neurology*. 2004;62:2103-2105.
19. Fisher J, Rudick G, Cutter S, et al. The Multiple Sclerosis Functional Composite Measure (MSFC): an integrated approach to MS clinical outcome assessment. *Mult Scler*. 1999;5:244-250.

20. Ilse Baert, Smedal T, Kalron A, et al. Responsiveness and meaningful improvement of mobility measures following MS rehabilitation. *Neurology*. 2018;91:e1880-e1892.
21. Pollock A, Baer G, Campbell P, et al. Physical rehabilitation approaches for the recovery of function and mobility following stroke. *Cochrane Database Syst Rev*. 2014;22:CD001920.
22. Edwards T, Pilutti LA. The effects of exercise training in adults with multiple sclerosis with severe mobility disability A systematic review and future directions. *Mult Scler. Relat Disord* 2017;16:31-39.
23. Motl RW, Learmonth YC, Pilutti L, et al. Top 10 research questions related to physical activity and multiple sclerosis. *Res Q Exerc Sport*. 2015;86:117-129.
24. Dodd KJ, Taylor NF, Denisenko S, et al. A qualitative analysis of a progressive resistance exercise programme for people with multiple sclerosis. *Disabil Rehabil*. 2006;28:1127-1134.
25. Patrocinio de Oliveira CE, Moreira OC, Carrión-Yagual ZM, et al. Effects of classic progressive resistance training Versus eccentric-enhanced resistance training in people with multiple sclerosis. *Arch Phys Med Rehabil*. 2017 [Epub].
26. Dalgas U, Stenager E, Jakobsen J, et al. Resistance training improves muscle strength and functional capacity in multiple sclerosis. *Neurology*. 2009;73:1478-1484.
27. Dalgas U, Stenager E, Lund C, et al. Neural drive increases following resistance training in patients with multiple sclerosis. *J Neurol*. 2013;260:1822-1832.
28. Dalgas U, Stenager E, Jakobsen J, et al. Muscle fiber size increases following resistance training in multiple sclerosis. *Mult Scler*. 2010;16:1367-1376.
29. Kjølhede T, Vissing K, de Place L, et al. Neuromuscular adaptations to long-term progressive resistance training translates to improved functional capacity for people with multiple sclerosis and is maintained at follow-up. *Mult Scler*. 2015;21:599-611.

30. Jackson K, Edginton-Bigelow K, Cooper C, et al. A group kickboxing program for balance, mobility, and quality of life in individuals with multiple sclerosis: a pilot study. *J Neurol Phys Ther.* 2012;36:131-137.
31. Skjerbæk AG, Næsby M, Lützen K, et al. Endurance training is feasible in severely disabled patients with progressive multiple sclerosis. *Mult Scler.* 2014;20:627-630.
32. Peck E, Chomko G, Gaz DV, et al. The effects of stretching on performance. *Curr Sports Med Rep.* 2014;13:179-185.
33. Pilutti LA, Dlugonski D, Sandroff BM, et al. Further validation of Multiple Sclerosis Walking Scale-12 scores based on spatiotemporal gait parameters. *Arch Phys Med Rehab.* 2013;94:575-578.
34. Learmonth YC, Dlugonski DD, Pilutti L, Sandroff BM, Motl RW. The reliability, precision and clinically meaningful change of walking assessments in multiple sclerosis. *Mult Scler J.* 2013;19:1784-1791.
35. Langeskov-Christensen D, Feys P, Baert I, Riemenschneider M, Stenager E, dalgas U. Performed and perceived walking ability in relation to the Expanded disability Status Scale in persons with multiple sclerosis. *J Neurol Sci.* 2017;382:131-136.
36. Leone C, Kalron A, Smedal T, Normann B, Wens I, O Eijnde, Feys P. Effects of rehabilitation on gait pattern at usual and fast speed depend on walking impairment level in multiple sclerosis. *Int J MS Care.* 2018;20(5):199-209.
37. Dejong G, Horn SD, Gassaway JA, Slavin MD, Dijkers MP. Toward a taxonomy of rehabilitation interventions: Using an inductive approach to examine the "black box" of rehabilitation. *Arch Phys Med Rehabil.* 2004;85:678-686.

## Figures

Figure 1. Mobility outcome measures according to group allocation of the total group (a), the mildly disabled group (b) and the moderately-severely disabled group (c).

Figure 2. Therapy goal distribution according to group allocation in the mildly disabled group (a) and the moderately-severely disabled group (b).

Figure 1a




Figure 1b




Figure 1c




Figure 2a





Figure 2b





**Table 1.** Descriptive Characteristics According to Study Groups and Disability Level.

Variable	Mean (SD), Frequencies, Proportions		P-Value
	Improved	Non-improved	
<i>Total group</i>			
Number, (%)	131 (47.0%)	148 (53.0%)	---
Age (y)	49.4 (11.6)	49.9 (10.2)	0.719
Gender (F/M)	86/45	95/53	0.800
Type of MS (RR/PP/SP)	77/34/20	70/55/23	0.169
Disease duration (y)	11.5 (8.3)	12.1 (8.1)	0.546
EDSS (median/range)	5.0 (4.0-6.0)	5.0 (4.0-6.0)	0.215
<i>Mild disability group (EDSS=2.0-4.0)</i>			
Number, (%)	46 (47.9%)	50 (52.1%)	---
Age (y)	47.3 (9.6)	44.8 (10.1)	0.213
Gender (F/M)	27/19	29/21	0.946
Type of MS (RR/PP/SP)	31/9/6	36/10/4	0.153
Disease duration (y)	9.5 (7.5)	7.5 (6.2)	0.524
EDSS (median/range)	3.5 (2.0-4.0)	3.0 (2.5-4.0)	0.013*
<i>Moderate-severe disability group (EDSS=4.5-6.5)</i>			
number	70 (38.3%)	113 (61.7%)	---
Age (y)	51.0 (12.2)	51.9 (9.9)	0.583
Gender (F/M)	48/22	77/36	0.952
Type of MS (RR/PP/SP)	38/20/12	42/50/21	0.410
Disease duration (y)	13.0 (8.3)	14.0 (8.2)	0.101

EDSS (median/range)	6.0 (5.0-6.0)	6.0 (5.0-6.5)	0.243
---------------------	---------------	---------------	-------

---

RR, Relapsing-remitting; PP, Primary progressive; SP, Secondary progressive; EDSS, Expanded disability status scale.

**Table 2.** Mobility Outcome Measures According to Study Groups and Disability Level.

Mobility parameter	Delta Post-Pre (Mean, SD)		<i>P</i> -value
	Improved	Non-improved	
<i>Total group</i>			
T25FW (usual) (s)	-1.13 (1.90)	-0.96 (2.05)	0.512
T25FW (fastest) (s)	-0.91 (1.42)	-0.53 (2.03)	0.081
2MWT (m)	11.90 (16.13)	5.75 (13.61)	0.001*
6MWT (m)	36.25 (40.87)	16.53 (40.77)	<0.001*
MSWS-12 (score, 0-100)	-24.65 (14.86)	4.95 (12.36)	<0.001*
<i>Mild disability group (EDSS=2.0-4.0)</i>			
T25FW (usual) (s)	-0.65 (1.71)	-0.44 (0.83)	0.504
T25FW (fastest) (s)	-0.63 (1.61)	-0.30 (0.65)	0.186
2MWT (m)	13.08 (14.36)	7.65 (13.14)	0.059
6MWT (m)	36.16 (43.39)	28.98 (33.66)	0.368
MSWS-12 (score, 0-100)	-27.94 (16.02)	2.21 (11.83)	<0.001*
<i>Moderate-severe disability group (EDSS=4.5-6.5)</i>			
T25FW (usual) (s)	-1.31 (2.03)	-1.22 (2.23)	0.808
T25FW (fastest) (s)	-1.09 (1.40)	-0.69 (2.30)	0.205
2MWT (m)	11.88 (18.11)	5.24 (13.46)	0.006*
6MWT (m)	34.19 (41.64)	14.87 (42.83)	0.004*
MSWS-12 (score, 0-100)	-25.54 (14.01)	4.13 (12.92)	<0.001*

**Table 3.** Rehabilitation Program Format and Volume According to Study Groups and Disability Level.

Variable	Mean (S.D)		P-value
	Improved	Non-improved	
Mild disability group (EDSS=2.0-4.0)			
Settings			
Inpatient	8	9	0.939
Outpatient	38	41	
Amount of therapy			
Number of weeks	7.3 (4.0)	8.7 (3.8)	0.073
Number of sessions	35.3 (34.1)	31.3 (30.7)	0.521
Total duration (h)	22.3 (15.4)	22.7 (14.2)	0.896
Format of therapy			
Individual therapy			
Number of sessions	15.0 (15.8)	7.9 (13.9)	0.021*
Total duration (h)	10.0 (8.7)	5.0 (8.0)	0.004*
Group therapy			
Number of sessions	13.7 (20.7)	17.8 (15.7)	0.275
Total duration (h)	9.8 (12.7)	15.4 (12.8)	0.033*
Autonomous therapy			
Number of sessions	6.9 (17.7)	5.6 (15.8)	0.715
Total duration (h)	2.5 (5.2)	2.2 (6.4)	0.828

---

 Moderate-severe disability group (*EDSS*=4.5-6.5)
*Settings*

Inpatient	25	36	0.593
Outpatient	45	77	

*Amount of therapy*

Number of weeks	6.0 (3.6)	7.1 (3.6)	0.046*
Number of sessions	33.0 (22.2)	30.2 (24.0)	0.424
Total duration (h)	23.0 (15.4)	18.6 (12.6)	0.037*

*Format of therapy**Individual therapy*

Number of sessions	12.3 (7.8)	10.1 (6.8)	0.050*
Total duration (h)	10.0 (7.9)	7.5 (4.9)	0.009*

*Group therapy*

Number of sessions	10.5 (11.8)	8.2 (11.4)	0.190
Total duration (h)	8.6 (10.6)	6.6 (8.5)	0.165

*Autonomous therapy*

Number of sessions	10.2 (19.9)	11.9 (21.0)	0.606
Total duration (h)	4.5 (8.4)	4.5 (9.3)	0.954

---

**Table 4.** Intervention Approaches According to Study Groups and Disability Level.

Variable	% (n)		<i>P</i> -value
	Improved	Non-improved	
Mild disability group ( <i>EDSS</i> =2.0-4.0)	n=46	n=50	
Balance training	60.9% (28)	44.0% (22)	0.100
Gait training - functional approach	30.4% (14)	18.0% (9)	0.157
Aerobic training	41.3% (19)	42.0% (21)	0.946
Muscle strengthening -without equipment	45.7% (21)	30.0% (15)	0.116
Resistance training – with equipment	<b>60.9%</b> (28)	26.0% (13)	0.006*
Self-stretching	39.1% (18)	<b>68.5%</b> (34)	0.004*
Passive mobilization/stretching	8.7% (4)	12.7% (6)	0.601
Moderately-severe disability group ( <i>EDSS</i> =4.5-6.5)	n=70	n=113	
Balance training	77.1% (54)	69.9% (79)	0.310
Gait training - functional approach	60.0% (42)	62.8% (71)	0.755
Aerobic training	<b>31.4%</b> (22)	14.2% (16)	0.008*
Muscle strengthening -without equipment	41.4% (29)	43.4% (49)	0.878
Resistance training – with equipment	21.4% (15)	<b>38.9%</b> (44)	0.010*
Self-stretching	31.4% (22)	29.2% (33)	0.744
Passive mobilization/stretching	31.4% (22)	<b>50.4%</b> (57)	0.014*

**Table 5.** Binary Logistic Regression Analysis in the Mild and Moderate Disability

Subgroups.

		B (95% CI)	R <sup>2</sup>	P-Value
Mild disability group ( <i>EDSS</i> =2.0-4.0)				
Step 1	Resistance training	-1.461 (0.098, 0.549)	0.154	0.001
Step 2	Resistance training	-1.286 (0.113, 0.674)	0.218	0.005
	Self-stretching	-1.046 (0.144, 0.860)		0.022
Moderate-severe disability group ( <i>EDSS</i> =4.5-6.5)				
Step 1	Passive mobilization/stretching	-0.945 (0.210, 0.719)	0.067	0.003
Step 2	Passive mobilization/stretching	-1.157 (0.163, 0.605)	0.143	0.001
	Individual therapy volume	0.083 (1.033, 1.144)		0.001