

2017-12-11

# Quantifying the Erosional Response to Tectonic Forcing: Be10 Catchment Average Erosion Rates from the Taupo Rift, New Zealand.

Boulton, SJ

<http://hdl.handle.net/10026.1/13605>

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

## **Quantifying the erosional response to tectonic forcing: Be<sup>10</sup> Catchment average erosion rates from the Taupo Rift, New Zealand.**

Sarah J. Boulton<sup>1,2</sup> and Kevin P. Norton<sup>2</sup>

School of Geography, Earth and Environmental Sciences, University of Plymouth, Plymouth, U.K.

School of Geography, Environment and Earth Sciences, Victoria University of Wellington, Wellington, N.Z.

While it has long been known that denudation is greater in tectonically active areas, few studies have explicitly compared the relationship between uplift rates, catchment average erosion rates and fluvial geomorphic indices (i.e., the normalised steepness index,  $k_{sn}$ ). Here we present data from the actively extending Taupo Rift (North Island, New Zealand). At the southern end of the Taupo Rift, NE-striking faults primarily delineate the rift margins, uplifting the Kaimanawa Mountains to the east at around  $1 \text{ mmyr}^{-1}$ . Geologic mapping and topographic analysis indicates that the Kaimanawa Mountains are a tilted fault block, allowing the extrapolation of the uplift rate across the study area. River long profiles were extracted for the whole area and contain both vertical step knickpoint associated with lithologic variation and slope-break knickpoints high up in the catchments, probably related to the initiation of faulting and also base level changes in Lake Taupo. Samples for Be<sup>10</sup> catchment average denudation rate determination were collected both along the strike of the NE-SW trending faults and across the fault block. The three datasets show strong correlations and demonstrate the utility of combing different approaches to understand the geomorphology and geology of actively deforming regions.