

2019-04-12

Family-Centered Care Improves Clinical Outcomes of Very-Low-Birth-Weight Infants: A Quasi-Experimental Study

Lv, B

<http://hdl.handle.net/10026.1/13707>

10.3389/fped.2019.00138

Frontiers in Pediatrics

Frontiers Media

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.



Family-Centered Care Improves Clinical Outcomes of Very-Low-Birth-Weight Infants: A Quasi-Experimental Study

Bo Lv^{1†}, Xi-rong Gao^{1†}, Jing Sun², Tao-tao Li¹, Zhen-ye Liu¹, Li-hui Zhu^{3*} and Jos M. Latour^{3,4}

¹ Division of Neonatal Medicine, Hunan Children's Hospital, Changsha, China, ² Nursing School, Hunan University of Chinese Medicine, Changsha, China, ³ Nursing Department, Hunan Children's Hospital, Changsha, China, ⁴ Faculty of Health and Human Sciences, School of Nursing and Midwifery, University of Plymouth, Plymouth, United Kingdom

OPEN ACCESS

Edited by:

Henry J. Rozycki,
Virginia Commonwealth University,
United States

Reviewed by:

Hercília Guimarães,
Universidade do Porto, Portugal
Ömer Erdeve,
Ankara University Medical School,
Turkey

*Correspondence:

Li-hui Zhu
877845375@qq.com

[†]First co-authors

Specialty section:

This article was submitted to
Neonatology,
a section of the journal
Frontiers in Pediatrics

Received: 23 January 2019

Accepted: 25 March 2019

Published: 12 April 2019

Citation:

Lv B, Gao X, Sun J, Li T, Liu Z, Zhu L
and Latour JM (2019)
Family-Centered Care Improves
Clinical Outcomes of
Very-Low-Birth-Weight Infants: A
Quasi-Experimental Study.
Front. Pediatr. 7:138.
doi: 10.3389/fped.2019.00138

Background: Survival of very-low-birth-weight infants is improving in neonatology and family-centered-care might contribute to premature infants' clinical outcomes.

Aim: To evaluate a family-centered care intervention on clinical outcomes of very-low-birth-weight infants.

Methods: A quasi-experimental study was conducted in a Chinese NICU between June 2016 and June 2017. The intervention included parental education of basic care knowledge and skills followed by active participation in care for at least 4 h a day. A total of 319 very-low-birth-weight infants were recruited by convenience sampling; intervention group $n = 156$ and control group $n = 163$. Primary outcome measures were weight at discharge, length-of-stay, breastfeeding, nasal feeding, total parental nutrition, re-admission, hospital expenses. Secondary outcome measures were infant complications.

Results: Infants' weight at discharge was higher in the interventions group (2,654 g vs. 2,325 g, $p < 0.001$). Nutritional outcomes improved significantly: breastfeeding rate 139 vs. 91, $p < 0.001$; days of total parental nutrition 25 d vs. 32 d, $p < 0.001$; gastric feeding days 23 d vs. 35 d, $p < 0.001$. Length-of-stay and hospital expenses did not differ between both groups. Improved infants' complications were bronchopulmonary dysplasia (32 vs. 51, $p = 0.031$), retinopathy of prematurity (between groups no/mild and moderate/severe, $p = 0.003$), necrotizing enterocolitis (6 vs. 18, $p = 0.019$), and re-admission rate (21 vs. 38, $p = 0.023$). No differences were observed in intraventricular hemorrhage and nosocomial infections.

Conclusion: Very-low-birth-weight premature infants might experience improved clinical health outcomes when parents are present and caring for them. Family-centered care is as a beneficial care model for premature infants and should be recognized and implemented by NICUs where parents have currently limited access.

Keywords: neonatology, infants, very low birth weight, family-centered care, parents, patient outcomes

INTRODUCTION

It is estimated that 15 million preterm neonates are born every year worldwide (1). With the implementation of the two-child policy in China in 2015, more preterm neonates are expected to be born (2). Preterm birth complications are the most frequent causes of deaths in children under 5 years of age (3, 4). Preterm infants born with very low birth weight (VLBW) experience several disadvantages and developmental challenges (5–7). Long-term growth development is one of these challenges. A 20-year follow-up study identified that males born with VLBW were significantly shorter and lighter as their female counterpart at 20 years of age (8). Although advances in medical technology has increased survival rates in neonatology, morbidity remains high and imposes emotional and financial burdens on families, society and healthcare system.

Family-centered care (FCC) is a care model in Neonatal Intensive Care Units (NICU) that has been implemented mainly in developed countries (9–11). The Institute of Patient- and Family Centered Care defines FCC as an approach to the planning, delivery, and evaluation of healthcare based on partnerships between health professionals, patients, and families (12). In pediatrics, FCC supports the involvement of parents in the care of their infant in the NICU (13). In NICU settings, the Newborn Individualized Developmental Care and Assessment Program (NIDCAP) is another approach to involve parents in the care (14). Evidence suggests that parents feel more empowered to care for their infant and improves short-term neurodevelopment of preterm infants (15). However, more studies are needed to prove the benefit of NIDCAP on long-term neurodevelopment (16).

Recent studies have demonstrated that providing parents training about the basic care of their infant improves infants' clinical outcomes (17, 18) Allowing parents to be involved in caregiving and becoming the primary caregiver might improve weight gain and breastfeeding rates (19). Despite the growing evidence of the benefit of FCC practices in NICU, it is not yet widely implemented in NICUs across the world (20–22).

In China, FCC in Chinese NICUs are at a developmental stage. In many Chinese NICUs parents are restricted to visit their infant or to participate in caregiving (23). However, recent studies from China reveal a growing trend in FCC practices and report clinical benefits to preterm infant (24–26). To date, there are limited reports presenting evidence of FCC practices in VLBW infants. Therefore, the aim of our study was to implement and test an FCC intervention providing parental education and participation in care among parents with VLBW infants. The hypothesis was defined as: parent education and parent participation in care improve clinical outcomes of VLBW infants.

MATERIALS AND METHODS

Design

This study adopted a quasi-experimental design using convenience sampling. Due to the nature of delivering the intervention by nurses to parents and data collection, blinding

was not possible. The study was conducted between June 2016 and June 2017.

Setting

This study was conducted at the stand-alone Hunan Children's Hospital, Changsha, China. The Neonatology department includes five NICUs: Two level 3 tertiary care NICUs for term infants (45–60 beds) and preterm infants (45–60 beds) and three level 2 NICUs each 70 beds. The study was performed at the level 3 NICU for preterm infants with an annual admission rate of 600 infants. At time of the study our level 3 NICU had 60 beds operational due to high demands. The NICU is designed in three open space units with 20 incubators per unit. The NICU has one room with two incubators in case isolation for infection prevention is needed. Outside the NICU is a parent waiting room. Parents of VLBW infants are only allowed to visit their infant via video camera three times a week. In recent years, the NICU implemented the FCC caring model allowing only parents of stable preterm infants to visit the NICU and participate in the care. Based on the promising results of two studies among preterm infants (20, 21), the NICU decided to further implement FCC with parents of VLBW infants.

Participants and Recruitment

Inclusion criteria were: preterm infants with a birth weight <1,500 g; non-invasive oxygen support; parents willing to participate in the care for at least 4 h a day. Exclusion criteria were: Infants with life-threatening congenital anomalies; surgery; palliative care; expected discharge within 1 week.

Infants and parents were recruited and assigned upon admission. At the start of the study, the first infant was recruited and assigned to the intervention group, the second infant to the control group and further recruitment took place in subsequent order.

Intervention and Standard Care

The FCC intervention started by teaching parents the theoretical knowledge of basic care, infant development, hand hygiene, feeding methods, skin-to-skin contact, infection control. More specifically, basic care was instructed to involvement parents in bathing the infant, changing diapers, temperature measurement, and other basic care. Hand hygiene was instructed including the steps of handwashing and the times when to wash their hands. Breastfeeding was taught and promoted to parents. Skin-to-skin contact was taught to parents including kangaroo care and at the same time to communicate with the infant to promote infant-parent bonding. The teaching sessions lasted around 90 min and ongoing support to parents was provided throughout the admission period. After the maternal skills were assessed by the nurses, parents participated in the care for at least 4 h a day between 10.00 and 16.00 h. Nurses were trained from April to May 2016 to deliver the teaching sessions.

The standard care group followed the routine caring model and adhered to the hospital regulations. The routine standard care for parents of VLBW infants was restricted by the visiting policies; parents were not allowed to visit their infant and thus not involved in the care of their infant. Parents could come to the

hospital 3 days a week; Monday, Wednesday, and Friday. During these visits, the neonatologist would meet the parents in a special room outside the NICU. During these meetings, an update of the health status of their infant was provided and parents were able to see their infant by video connection.

Data Collection

Data were extracted from the infant's hospital records. Basic demographics of parents (mode of delivery, education and income levels) were collected. The primary outcome measures were: weight at discharge, NICU length-of-stay, breastfeeding rate, days of nasal feeding, days of total parental nutrition, re-admission within 1 month, hospital expenses. The secondary outcome measures were the infant complications: nosocomial infection rate, Bronchopulmonary dysplasia (BPD), Retinopathy of prematurity (ROP), Necrotizing enterocolitis (NEC), Intraventricular hemorrhage (IVH).

Data Analysis

Data analysis was performed with IBM Corp. Released 2012. IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY: IBM Corp. The Kolmogorov-Smirnow test was used to determine normal distribution of data. For descriptive statistics, mean, standard deviation percentages were applied. The Student *t*-test was used for continuous variable and the chi-square test for categorical variable. We defined statistical significance as $p < 0.05$. Data are presented as FCC group vs. Standard Care (SC) group.

Ethics Considerations

This study was carried out in accordance with the recommendations of improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: the TREND statement, the Trend Group (27). The protocol was approved by the Ethics Committee of Hunan Children's Hospital (HCHLL-2015-33).

Parents were assured that their decision to withdraw or refuse to participate would not impact the care and treatment of their infant. All parents in the FCC group gave written informed consent in accordance with the Declaration of Helsinki. The infants and parents in the SC group received the routine care and

parents were verbally informed and written informed consent were waived by the Ethics Committee.

RESULTS

Characteristics of Infants and Parents

Of the 524 VLBW infants admitted to the NICU during the 12-month recruitment period. 319 met the inclusion criteria and were assigned to the FCC group ($n = 156$) or standard care group ($n = 163$). Both groups represented equal gender (male: 101, 64.76% vs. 102, 62.6%). The mean birth weight was not different between both groups (1,164 g, SD 211 vs. 1,204 g, SD = 196, $p = 0.085$). Gestational age of infants was significantly less the FCC group (28.9 weeks, SD = 1.6 vs. 29.4 weeks, SD = 2.3, $p = 0.013$). Apgar scores at 1 min and antenatal corticosteroid therapy did not reveal any statistical differences between both groups.

The characteristics of the parents did not show statistical differences in the mode of delivery (spontaneous vs. sectio-caesarea) and education levels. However, parents in the FCC groups had a higher monthly income than those in the SC group (<2,000 RMB, 11 vs. 26; 2,000–5,000 RMB, 69 vs. 81; >5,000 RMB, 76 vs. 56, $p = 0.007$).

Infants' Clinical Outcomes

Discharge weight of infants in the FCC group was significantly higher than in the SC group. Similar statistical differences were observed with the NICU length-of-stay resulting in a shorter stay in the FCC group (Table 1). Three nutritional support outcome measures significantly improved in the FCC group; breastfeeding rate was higher, and the days of total parental nutrition and nasal feeding were lower in the FCC group (Table 1). The hospital expenses were lower in the FCC group, but not significant.

Infants' Complications

Table 2 presents the outcomes of the infant complication variables. Nosocomial infection and IVH rates did not present significant differences. In the FCC group were significantly less infants ($n = 32$) developing BPD compared to the SC group ($n = 51$). The FCC group had also less infants with no or mild ROP, less NEC and re-admissions compared to the SC group.

TABLE 1 | Infants' Clinical outcomes.

Outcomes	FCC group $n = 156$	SC group $n = 163$	Mean difference (95% CI)	P-value
Discharge weight (g; mean; SD)	2653.9 (672.2)	2324.9 (911.9)	329.0 (147.9–510.2)	< 0.001
Length-of-stay (days; mean; SD)	60.6 (32.1)	63.2 (25.4)	–2.3 (–8.7 to 4.0)	0.474
Breastfeeding rate (n ; %)	139 (89.1)	91 (55.8)	6.5 (3.6–11.7)	< 0.001
TPN (days; mean; SD)	24.94 (19.5)	32.23 (19.9)	–7.3 (–11.4 to –3.2)	< 0.001
Nasal feeding (days; mean; SD)	22.8 (16.4)	34.6 (27.2)	–11.8 (–16.9 to –6.7)	< 0.001
Expenses (RMB; mean; SD)	83068 (36233.2)	88595 (18433.2)	–5527.0 (–20618.6 to 9564.7)	0.472

FCC, Family-Centered Care; SC, Standard Care; TPN, total parenteral nutrition; Length-of-Stay, NICU Length-of-Stay; RMB, RenMinBi (Chinese Yuan).

TABLE 2 | Infants' complications.

Complications	FCC group <i>n</i> = 156	SC group <i>n</i> = 163	Mean difference (95% CI)	<i>p</i> -value
Nosocomial Infection (<i>n</i> , %)	24 (15.3)	19 (11.6)	1.4 (0.7–2.6)	0.413
BPD (<i>n</i> , %)	32 (20.5)	51 (31.3)	0.6 (0.3–0.9)	0.031
ROP (<i>n</i>, %)				
No-ROP to mild ROP	135 (86.5)	119 (73.0)	reference	0.003
Moderate to severe ROP	21 (13.5)	44 (27.0)	2.4 (1.3–4.2)	
NEC (<i>n</i> , %)	6 (3.8)	18 (11.0)	0.3 (0.1–0.8)	0.019
Re-admission within 1 month (<i>n</i> , %)	21 (13.4)	38 (23.3)	0.5 (0.3–0.9)	0.023
IVH (<i>n</i>, %)				
No IVH, grade 1	146 (93.6)	147 (90.2)	reference	0.310
Grade 2–3	10 (6.4)	16 (9.8)	1.6(0.7–3.6)	

FCC, Family-Centered Care; SC, Standard Care; BPD, Bronchopulmonary dysplasia; ROP, Retinopathy of prematurity; NEC, Necrotizing enterocolitis; IVH, Intraventricular hemorrhage.

DISCUSSION

In the past decades, family-centered care has been implemented and further refined in NICUs in developed countries (22, 28, 29). In the Chinese context, barriers to implement FCC in NICUs have been vocalized as not enough space around incubators and lack of trained staff (30). However, FCC is gaining momentum among Chinese NICU clinicians. Our study is one of the few Chinese studies evaluating FCC in the NICU. We were able to successfully train nurses providing education and support to parents of VLBW infants. The infants' clinical outcome measures were promising in terms of infants' weight and nutritional support while complication rates of BPD, ROP and NEC have been less in the FCC group.

Several randomized controlled trials (RCTs) have been testing FCC interventions with parent educational programs in the NICU (22). Our results show some similarities with these studies. The recently published multi-center, multi-national FICare trial documented that weight gain of premature infants between the intervention and standard care groups increased faster and remained higher in the first 21 days of the trial (19). Although we did not include daily weight gain in our study, the VLBW infants in the FCC group had significantly higher weight at discharge while length-of-stay did not differ between both groups. However, three RCTs from China testing an educational FCC intervention confirmed also improved daily weight gain (31–33).

There is growing evidence that parental presence and involvement in care improves breastfeeding practices and nutritional outcomes of premature infants. Our study demonstrated an improvement in breastfeeding practices and less total parental nutrition and gastric feeding was necessary. A similar study confirmed our findings documenting that breastfeeding rates were higher in FCC group; 80.4 vs. 66.7%, $p = 0.007$ (17). A study in Taiwan showed that feeding practices improved in the FCC group; infants in the FCC group were significantly younger when they achieved full enteral feeding (95% CI = -1.9 to -0.2 weeks, $p = 0.02$) (18). Therefore, it can be assumed that parents can positively influence the feeding

practices when providing them with sufficient educational and clinical support.

Allowing parents to be active partners in care might safeguard the preterm infants from complications. We demonstrated that VLBW infants in the FCC group had significantly less BPD, ROP, and NEC. The Stockholm Neonatal Family Centered Care Study, providing a new NICU environment for parents with individual rooms, reported that infants in the FCC unit had less moderate-to-severe BPD compared to the standard care unit; 3 vs. 11% (34). Rooming-in can be considered as a valuable contribution to FCC and it is noticed that many countries are transforming their NICUs into single-bedded rooms to allow rooming-in. Limited studies have explored the benefit of rooming-in in NICUs. A study looking at the impact of individual rooms demonstrated no significant short-term effect on parental stress and depression (35). However, another study documented a significant reduction in readmission rates (36). The clinical significance of rooming-in and single bedded designed NICUs cannot be underestimated anymore. With the results of our study and the vision to provide optimal care to infants and parents, our hospital is currently redesigning the NICU into more single-bedded FCC rooms.

The incidence of NEC in our study decreased from 18 to 6 in the FCC group. We have not tested the association of NEC and the nutritional variables. However, a meta-analysis of the effect of standardized feeding protocols and NEC, including nine studies with 4,755 infants, provide strong evidence to recommend early parental involvement in feeding practices (37). Additionally, providing parents with information might also affect their ability to care for their infant after discharge. The re-admission rate 1 month after discharge decreased significantly in our study. This is consistent in two other similar studies investigating an educational FCC intervention (25, 26).

Our study has several limitations. The sample size calculation and use of convenience sampling are limitations to address. Sample size was not calculated because we determined to run the data collection over a 12-month period using convenience sampling. The critique on this sampling strategy is that the study participants are not chosen at random which limits the generalizability of our results. Blinding of the intervention was

not possible for parents and NICU staff due to the nature of the intervention where nurses provided education to parents and were on the unit all times. We did not collect any parent outcome measure which could strengthen the possible benefit of our intervention.

In conclusion, an FCC intervention, providing education to parents and empowering them to be involved in the daily care might improve clinical outcomes of VLBW infants. Despite the quasi-experimental design of our study, FCC is a beneficial care model for premature infants and should be recognized and implemented by NICUs where parents have limited access.

ETHICS STATEMENT

This study was carried out in accordance with the recommendations of improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: the TREND statement, the Trend Group (23). The protocol was approved by the Ethics Committee of Hunan Children's Hospital (HCHLL-2015-33). Parents were assured that their decision to withdraw or refuse to participate would not impact the care and treatment of their infant. All parents in the FCC group gave written informed consent in accordance with the Declaration of Helsinki. The infants and parents in the

SC group received the routine care and parents were verbally informed and written informed consent were waived by the Ethics Committee.

AUTHOR CONTRIBUTIONS

BL, LZ, XG, TL, and JML contributed to the design of the study, data collection, data analysis, and interpretation. JS and JML drafted the first manuscript. BL, ZL, and XG provided revisions. All authors contributed to manuscript revision, read and approved the submitted version.

FUNDING

The study was partially supported by the Health and Family Planning Commission of Hunan Province (B2016031). The study was partially supported by Futang Children's Medical Center (201533).

ACKNOWLEDGMENTS

The authors thank the NICU staffs for their willingness to learn and implement the family-centered care intervention. The authors thank all parents who participated in the study.

REFERENCES

- Blencowe H, Cousens S, Chou D, Oestergaard M, Say L, Moller AB, et al. Born too soon: the global epidemiology of 15 million preterm births. *Reprod Health.* (2013) 10(Suppl 1):S2. doi: 10.1186/1742-4755-10-S1-S2
- Zeng L. Investigation and analysis on the incidence and outcome of premature babies after the release of two-child policy. *J Xiangnan Univ.* (2017) 19:53–55. doi: 10.16500/j.cnki.1673-498x.2017.04.017
- Rudan I, Chan KY, Zhang JS, Theodoratou E, Feng XL, Salomon JA, et al. Causes of deaths in children younger than 5 years in China in 2008. *Lancet.* (2010) 375:1083–9. doi: 10.1016/S0140-6736(10)60060-8
- Han W, Song J, Liu A, Huo K, Xu F, Cui S, et al. Trends in live births in the past 20 years in Zhengzhou, China. *Acta Obstet Gynecol Scand.* (2011) 90:332–7. doi: 10.1111/j.1600-0412.2010.01065.x
- Chen LC, Wu YC, Hsieh WS, Hsu CH, Leng CH, Chen WJ, et al. The effect of in-hospital developmental care on neonatal morbidity, growth and development of preterm Taiwanese infants: a randomized controlled trial. *Early Hum Dev.* (2013) 89:301–6. doi: 10.1016/j.earlhumdev.2012.10.008
- Peng Y, Huang B, Biro F, Feng L, Guo Z, Slap G. Outcome of low birthweight in China: a 16-year longitudinal study. *Acta Paediatr.* (2005) 94:843–9. doi: 10.1080/08035250510027589
- Afjeh SA, Sabzehei MK, Fallahi M, Esmaili F. Outcome of very low birth weight infants over 3 years report from an Iranian center. *Iran J Pediatr.* (2013) 23:579–87.
- Hack M, Schluchter M, Cartar L, Rahman M, Cuttler L, Borawski E. Growth of very low birth weight infants to age 20 years. *Pediatrics.* (2003) 112:e30–8. doi: 10.1542/peds.112.1.e30
- Latour JM, Haines C. Families in the ICU: do we truly consider their needs, experiences and satisfaction? *Nurs Crit Care.* (2007) 12:173–4. doi: 10.1111/j.1478-5153.2007.00234.x
- Separation and Closeness Experiences in Neonatal Environment (SCENE) research group. Parent and nurse perceptions on the quality of family-centred care in 11 European NICUs. *Aust Crit Care.* (2016) 29:201–9. doi: 10.1016/j.aucc.2016.09.003
- Dall'Oglio I, Mascolo R, Tiozzoa E, Portanovac A, Fiori M, Gawronski O, et al. The current practice of family-centred care in Italian neonatal intensive care units: a multicentre descriptive study. *Intensive Crit Care Nurs.* (2019) 50:36–43. doi: 10.1016/j.iccn.2018.07.005
- Institute for patient- and family-centered care. *Patient- and Family-Centered Care.* Available online at: www.ipfcc.org/about/pfcc.html (accessed March 1, 2019).
- Committee on hospital care and Institute for Patient- and Family-Centered Care. Patient- and family-centered care and the pediatrician's role. *Pediatrics.* (2012) 129:394–404. doi: 10.1542/peds.2011-3084
- Als H, McAnulty GB. The Newborn Individualized Developmental Care and Assessment Program (NIDCAP) with Kangaroo Mother Care (KMC): comprehensive care for preterm infants. *Curr Womens Health Rev.* (2011) 7:288–301. doi: 10.2174/157340411796355216
- Sannino P, Gianni ML, De Bon G, Fontana C, Picciolini O, Plevani L, et al. Support to mothers of premature babies using NIDCAP method: a non-randomized controlled trial. *Early Hum Dev.* (2016) 95:15–20. doi: 10.1016/j.earlhumdev.2016.01.016
- Ohlsson A, Jacobs SE. NIDCAP: a systematic review and meta-analyses of randomized controlled trials. *Pediatrics.* (2013) 131:e881–93. doi: 10.1542/peds.2012-2121
- Verma A, Maria A, Ravindra AM, Pandey M, Hans S, Verma A, et al. Family-centered care to complement care of sick newborns: a randomized controlled trial. *India Pediatr.* (2017) 54:455–9. doi: 10.1007/s13312-017-1047-9
- Yu YT, Hsieh WS, Hsu CH, Lin YJ, Linet CH, Hsieh S, et al. Family-centered care improved neonatal medical and neurobehavioral outcomes in preterm infants: randomized controlled trial. *Phys Ther.* (2017) 97:1158–68. doi: 10.1093/ptj/pzx089
- O'Brien K, Robson K, Bracht M, Cruz M, Lui K, Alvaroet R, et al. Effectiveness of Family Integrated Care in neonatal intensive care units on infant and parent outcomes: a multicentre, multinational, cluster-randomised controlled trial. *Lancet Child Adolesc Health.* (2018) 2:245–54. doi: 10.1016/S2352-4642(18)30039-7
- Latour JM, Haines C. Is family-centred care in critical care units that difficult? A view from Europe. *Nurs Crit Care.* (2005) 10:51–3. doi: 10.1111/j.1362-1017.2005.0108b.x

21. Al-Motlaq MA, Shields L. Family-centered care as a western-centric model in developing countries: luxury versus necessity. *Holist Nurs Pract.* (2017) 31:343–7. doi: 10.1097/HNP.0000000000000228
22. Ding X, Zhu L, Zhang R, Wang L, Wang TT, Latour JM. Effects of family-centered care interventions on preterm infants and parents in neonatal intensive care units: a systematic review and meta-analysis of randomised controlled trials. *Aust Crit Care.* (2019) 32:63–75. doi: 10.1016/j.aucc.2018.10.007
23. He Y, Chen JL. The current state of family-centered care in the neonatal intensive care units. *Nurs J Chin PLA.* (2014) 31:29–32. doi: 10.3969/j.issn.1008-9993.2014.07.008
24. Li HZ, Ren YM. Application effects of NICU family participation nursing mode on premature infants. *Chin J Mod Nurs.* (2016) 22:5255–7. doi: 10.3760/cma.j.issn.1674-2907.2016.36.018
25. He S, Xiong Y, Zhu LH, Lv B, Gao X, Xiong H, et al. Impact of family integrated care on infant's clinical outcomes in two children's hospitals in China: a pre-post intervention study. *Ital J Pediatr.* (2018) 44:65. doi: 10.1186/s13052-018-0506-9
26. Zhang R, Huang RW, Gao XR, Peng XM, Zhu LH, Rangasamy R, et al. Involvements of parents in the care of preterm infants: a pilot study evaluating a family-centered care intervention in a Chinese neonatal ICU. *Pediatr Crit Care Med.* (2018) 19:741–7. doi: 10.1097/PCC.0000000000001586
27. Des Jarlais DC, Lyles C, Crepaz N, Trend Group. Improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: the TREND statement. *Am J Public Health.* (2004) 94:361–6. doi: 10.2105/AJPH.94.3.361
28. Craig JW, Glick C, Phillips R, Hall SL, Smith J, Browne J. Recommendations for involving the family in developmental care of the NICU baby. *J Perinatol.* (2015) 35(Suppl 1):S5–8. doi: 10.1038/jp.2015.142
29. Lee LA, Carter M, Stevenson SB, Harrison HA. Improving family-centered care practices in the NICU. *Neonatal Netw.* (2014) 33:125–32. doi: 10.1891/0730-0832.33.3.125
30. Zhu LH, Wang L. Research progress of family-centered care in pediatrics. *Intern J Nurs.* (2012) 31:1557–9. doi: 10.3760/cma.j.issn.1673-4351.2012.09.002
31. Ying XQ, Lin XY, Yan SY, Ding LL, Zhu F. Effects of developmental care nursing model on growth and development of premature infants. *Nurs J Chin PLA.* (2012) 29(2A):15–7.
32. Chang WZ. Effects of family centered care applied in preterm baby nursing. *World Latest Med Inf.* (2013) 13:414–5.
33. Xiao AX, Huang RW, Zhang R, Shen P, Zhu LH. Application of family-centered care in the high quality nursing in neonatal ward. *Nurs J Chin PLA.* (2016) 33:70–3.
34. Ortenstrand A, Westrup B, Broström EB, Sarman I, Akerström S, Brune T, et al. The Stockholm Neonatal Family Centered Care Study: effects on length of stay and infant morbidity. *Pediatrics.* (2010) 125:e278–85. doi: 10.1542/peds.2009-1511
35. Erdev O, Arsan S, Canpolat FE, Ertem IO, Karagol BS, Atasay B, et al. Does individual room implemented family-centered care contribute to mother-infant interaction in preterm deliveries necessitating neonatal intensive care unit hospitalization? *Am J Perinatol.* (2009) 26:159–64. doi: 10.1055/s-0028-1095186
36. Erdev O, Arsan S, Yigit S, Armangil D, Atasay B, Korkmaz A. The impact of individual room on rehospitalization and health service utilization in preterms after discharge. *Acta Paediatr.* (2008) 97:1351–7. doi: 10.1111/j.1651-2227.2008.00889.x
37. Gephart SM, Hanson C, Wetzel CM, Fleiner M, Umberger E, Martin L, et al. NEC-zero recommendations from scoping review of evidence to prevent and foster timely recognition of necrotizing enterocolitis. *Matern Health Neonatol Perinatol.* (2017) 3:23. doi: 10.1186/s40748-017-0062-0

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Lv, Gao, Sun, Li, Liu, Zhu and Latour. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.