

2011-06-05

The response of foraminifera to modern seawater acidification: A real-time proxy for Paleogene hypothermal events

Hall-Spencer, Jason

<http://hdl.handle.net/10026.1/1376>

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

The response of foraminifera to modern seawater acidification: A real-time proxy for Paleogene hypothermal events

**Malcolm B. Hart¹, Bruna B. Dias², Christopher W. Smart¹,
Jason M. Hall-Spencer¹, Laura Pettit¹**

¹ Faculty of Science & Technology, University of Plymouth, Plymouth PL4 8AA, U.K.

² Universidad Federal de Santa Catarina, Florianópolis, Brazil

The seas around the island of Ischia (Italy) have a variable and, on average, lowered pH as a result of volcanic gas vents that emit carbon dioxide from the sea floor at ambient seawater temperatures. These areas of acidified seawater provide natural laboratories in which to study the long-term biological response to rising CO₂ levels. Benthic foraminifera are routinely used to interpret the effects of climate change as they have short life histories, are environmentally sensitive and have an excellent fossil record. Here, we examined changes in foraminiferal assemblages along gradients in pH at CO₂ vents on the coast of Ischia as they may provide a useful model on which to base future predictions of the consequences of ocean acidification (Dias *et al.*, 2010). We show that foraminiferal abundance, diversity and ability to calcify decreased markedly in living and dead assemblages as pH decreases, the result of CO₂ percolating through the seawater. Recent work by de Nooijer *et al.* (2009) has shown that foraminifera increase their internal pH to ~9.0 in the area of chamber calcification. If the ambient pH in the enclosing seawater is reduced to 7.8 (or below) the foraminifera with calcareous tests appear to be unable to complete the test construction process. These results are in accord with the responses recorded by coralline algae, corals, molluscs, barnacles and echinoderms at the same sites (Hall-Spencer *et al.*, 2008).

Samples from the normal (pH8.17) environments around Ischia contain a diverse fauna dominated by miliolid foraminifera (e.g., *Peneroplis planatus*, *P. pertusus*, *Quinqueloculina* spp.) while those from areas with reduced pH (7.8 to 7.6) have faunas that are progressively less diverse and composed of <100% agglutinated taxa (e.g., *Ammoglobigerina globigeriniformis*, *Miliammina fusca*, *Trochammina inflata*, *Textularia* sp. cf. *T. bocki*). The changes in the benthic foraminifera are quite dramatic for only a slight reduction in pH and confirm the possibility that events, such as the PETM, could quite easily record a widespread loss of diversity or extinction as a result of ocean acidification.

Work on Ischia is on-going and there is a comparable site being investigated in the Gulf of California. A site on the south-east coast of the island of Vulcano (Mediterranean Sea) is also being considered.

References:

- Dias, B.D., Hart, M.B., Smart, C.W. & Hall-Spencer, J.M. 2010. Modern seawater acidification: the response of foraminifera to high-CO₂ conditions in the Mediterranean Sea. *Journal of the Geological Society, London*, **167**, 843-846.
- Hall-Spencer, J.M., Rodolfo-Metalpa, R., Martin, S. et al., 2008. Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature*, **454**, 96-99, doi:10.1038/nature07051.
- De Nooijer, L.J., Toyofuku, T. & Kitazato, H. 2009. Foraminifera promote calcification by elevating their intracellular pH. *Proceedings of the National Academy of Sciences*, **106**, 15374-15378.