

2019-05-01

Holocene demographic fluctuations, climate and erosion in the Mediterranean: A meta data-analysis

Walsh, K

<http://hdl.handle.net/10026.1/13767>

10.1177/0959683619826637

The Holocene

SAGE Publications

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Holocene demographic fluctuations, climate and erosion in the Mediterranean: a meta data-analysis

Walsh, K¹., Berger, J_F²., Roberts, N³., Vanni re, B.⁴, Ghilardi, M.⁵, Brown, A.G.⁶, Woodbridge, J.³, Lespez, L.⁷, Estrany, J.⁸, Glais, A.⁹., Palmisano, A.¹⁰, Finn , M.¹¹, Verstraeten, G¹².

Affiliations

¹Department of Archaeology, University of York, York, YO1 7EP, UK and IM RA -Institute for Advanced Study, Aix-Marseille University, Marseille, France, kevin.walsh@york.ac.uk

²CNRS, UMR 5600 EVS/IRG, Universit  de Lyon, France. Jean-Francois.Berger@univ-lyon2.fr

³ School of Geography, Earth and Environmental Sciences, University of Plymouth, Drake Circus, Plymouth PL48AA, UK. C.N.Roberts@plymouth.ac.uk, jessie.woodbridge@plymouth.ac.uk

⁴ Chrono-environnement - Universit  de Franche-Comt , 16 route de Gray F-25030 Besan on cedex FRANCE, boris.vanniere@univ-fcomte.fr

⁵ CEREGE - AMU, CNRS UMR 7330, IRD, Coll ge de France, INRA, Europ le de l'Arbois BP 80 13545 Aix-en-Provence CEDEX 04 France. ghilardi@cerege.fr

⁶ Troms  University Museum, University of the Arctic, Troms , Norway and Geography & Environment, Highfields Campus, Southampton SO171BJ, UK. Tony.Brown@soton.ac.uk

⁷ LGP-CNRS UMR 8591, Department of Geography, University of Paris East-Cr teil, France. laurent.lespez@u-pec.fr

⁸ Department of Geography, University of the Balearic Islands, E-07122 Palma de Mallorca, Balearic Islands, Spain. Mediterranean Ecogeomorphological and Hydrological Connectivity Research Team (<http://medhycon.uib.cat>) and Institute of Agro-Environmental and Water Economy Research –INAGEA, University of the Balearic Islands, E-07122, Palma, Mallorca, Balearic Islands, Spain. joan.estrany@uib.cat

⁹ LETG-Caen-CNRS-UMR 6554, University of Caen-Normandy, France. arthur.glais@unicaen.fr

¹⁰ UCL Institute of Archaeology, 31-34 Gordon Square, London WC1H 0PY, UK, a.palmisano@ucl.ac.uk

¹¹ Department of Archaeology and Ancient History, Uppsala University, Box 626, 751 26 Uppsala. Email: martin.finne@antiken.uu.se, tel. 0046 18 471 6238. ORCID: 0000-0001-7433-268X

¹²Department of Earth and Environmental Sciences, Division of Geography and Tourism, KU Leuven, gert.verstraeten@kuleuven.be

Abstract: As part of the *Changing the Face of the Mediterranean Project*, we consider how human pressure and concomitant erosion has affected a range of Mediterranean landscapes has between the Neolithic and, in some cases, the post-medieval period. Part of this assessment comprises an investigation of relationships between palaeodemographic data, evidence for vegetation change and some consideration of rapid climate change events. The erosion data includes recent or hitherto unpublished work from the authors. Where possible, we consider summed probabilities of 14C dates as well as the first published synthesis of all known optically stimulated luminescence dated sequences. The results suggest that while there were some periods when erosion took place contemporaneously across a number of regions, possibly induced by climate changes, more often than not, we see a complex and heterogeneous interplay of demographic and environmental changes that result in a mixed pattern of erosional activity across the Mediterranean.

Keywords: Mediterranean, Holocene, erosion, geoarchaeology, demography, human impact

1 INTRODUCTION

Ever since humans started to have an impact on vegetation cover through agricultural and/or fire-setting activities, the rates at which landscapes erode have increased significantly (Montgomery, 2007). In many regions around the world that face or have faced anthropogenic accelerated erosion, this process has been linked with a decline in agricultural productivity to unsustainable levels, and even societal collapse (Diamond, 2005; Montgomery, 2007). As has been argued by researchers for several decades, this is all the more true for large parts of the Mediterranean, a region that has experienced significant human impact for several millennia, leading to landscape degradation through soil erosion (Lowdermilk, 1953). However, this viewpoint on the linkage between anthropogenic induced land degradation and collapse has also been challenged. First, resilient societies are thought to have adopted new strategies to cope with these environmental changes (Costanza, 2012; Grove and Rackham, 2001; Walsh, 2013). Secondly, the relative importance of the role of humans in controlling soil erosion compared to natural drivers of erosion, such as climate change, has been debated as well. Within his pioneer study on Mediterranean erosion, Vita-Finzi (1969) established a broad chronological framework with two significant phases of erosion, one dated to the Upper Palaeolithic, and the second dated to the Historic Period. Vita-Finzi stressed the importance of climate in controlling historical alluvial sedimentation. Van Andel and others have challenged this hypothesis, pointing towards the dominant role humans have exerted on sediment fluxes in antique Greece (van Andel et al., 1986; van Andel and Zangger, 1990). In more recent years, some have questioned these often-simplistic mono-causation models (Bintliff, 2002). Indeed, recent reviews of historic erosion and sedimentation for several parts of the Mediterranean have argued that there are periods when human impact can be identified from sediment records, however, the role of climate cannot be neglected either for other moments in the past, especially when regional scales are considered (Bellin et al., 2013; Dusaar et al., 2011a; Faust and Wolf, 2017). Large-scale phases of Rapid Climate Change (RCCs) suggested by Mayewski et al. (2004), i.e. periods when there are relatively significant changes in temperature and humidity, have been extensively discussed in relation to erosion in a Mediterranean context (for example - Bintliff 1992; Casana 2008; Fletcher & Zielhofer 2013; Macklin et al. 2015). Disentangling climatically caused erosion periods from erosion caused by human activity, however, is not straightforward since these two controlling factors are often coupled and increased sediment dynamics can only be explained through an interplay between both controlling factors (e.g. Fuchs, 2007).

We are fully aware that meta-analyses, such as those presented here, do not necessarily engage with the complexity of the many spatial and temporal scales over which human-environment interactions operate. This is especially true when assessing sedimentation, which reflects a wide range of processes and scales across any given catchment or drainage basin (Macklin et al., 2015; Vanmaercke et al., 2015; Verstraeten, 2015). Therefore, we aim to provide a synthesis and assessment that can be tested in the future against other data that operate at sub-regional and local scales. In order to resolve how past Mediterranean societies impacted geomorphic systems, it has been suggested that there is a need for more detailed and quantitative reconstruction of the palaeo-vegetation, demographic and geomorphic records, particularly at the regional scale (de Vente et al., 2014; Dugar et al., 2011a). Within this study, we address this complex issue of how Mediterranean societies have influenced geomorphic systems through the assessment of the relationship between demographic indices, palaeoecological data indicative of agricultural activity, and erosion data. This contribution is part of the *Changing the Face of the Mediterranean Project*, which has investigated human-environment interactions across the Mediterranean environments from the Neolithic to the Medieval and Post Medieval periods (Fig. 1). The broader project employs new models of demographic change, based on site data derived from many different published sources, along with the modelling of radiocarbon dates. We compare these data with the palaeoecological data, derived from almost all available pollen diagrams from five case study areas (within Turkey, Greece, Italy, South of France and Spain) presented within this special issue. This paper focusses on sedimentation data, comparing this, in the discussion section, with demographic and palaeoecological information. While the palynological and archaeological data lend themselves to continuous, diachronic profiling and modelling, geoarchaeological and geomorphological data rarely possess the temporal resolution or precision possessed by these other data, although, where possible, we have included existing summed probabilities of radiocarbon dates from across the region.

2 METHODS

This paper presents a combination of region-specific geoarchaeological case studies, many of which have been carried out by the authors of this paper. The study regions presented here run broadly from east to west, i.e. from Anatolia and Greece to Italy, the south of France, and ending with Spain (Fig. 2). Our fundamental aim is to assess the relationships between demographic data, the nature of human activities and their impact on the environment, as well as how these processes intersect with climate change. The geomorphic data are derived from alluvial, colluvial and lacustrine records spread across a range of landscape types, i.e. from hillslope records to river systems and coastal areas. We specifically address these various landscape settings, as it is impossible to define one isolated relationship between erosion histories and sediment stratigraphy for a given type of sediment record. Indeed, many studies have shown that colluvial, alluvial, lacustrine and coastal records all respond differently to human-induced erosion depending on the timing, duration, intensity and spatial extent of human impact (Dugar et al., 2011b; Fryirs, 2013; Houben et al., 2013; Verstraeten et al., 2017; Zolitschka et al., 2003). A critical comparison of how the various sedimentary environments responded to changes in demography, anthropogenic land cover and/or climate change may include therefore valuable information on the complexity of past human-environment interactions.

The case studies below unsurprisingly rely on ^{14}C dates; however, this paper also includes the first comprehensive meta-analysis of optically-stimulated-luminescence (hereinafter OSL) dates from across the Mediterranean.

Traditionally, rates of erosion and sedimentation, as well as the frequency of erosion/sedimentation events, have been established using radiocarbon dating of plant macrofossils or charcoal preserved in sediment archives (Brown, 1999, 1997). However, while this has been successful in temperate Europe and elsewhere, it can be far more problematic in the Mediterranean. This is primarily due to the lack of suitable organic matter within alluvial and colluvial sequences due to the semi-arid climate (in parts of the Mediterranean) and the deep penetration by roots of phreatophyte trees and shrubs. Furthermore, both charred and uncharred macrofossils may be reworked. A good example is the dating of alluvial fills in Malta, where nearly 90% of radiocarbon AMS dates were in error, with numerous time reversals and common age inversion in the late Holocene. More subtly, these problems also impose a potential bias on sampling as it is increasingly unlikely that suitable material will occur higher up the sedimentary sequence and for the last two centuries radiocarbon dates also suffer from the De Vries effect. This is also reflected in the demographic proxy records presented in this paper for Turkey (Fig. 3), Italy (Fig. 7) and Spain (Fig. 9) which lack enough dates over the last 2000 years for SPD analysis. However, suitable material can sometimes be found, and analysis of AMS dates for flood units does exist for regions such as Spain (Thorndycraft and Benito, 2006).

There are now several alternatives to ^{14}C , including OSL, which due to high insolation should theoretically be more suited to the Mediterranean than northern Europe, as well as palaeomagnetic dating and most recently cosmogenic burial dating (Granger and Muzikar, 2001). Palaeomagnetic dating has been used in Central Italy (Ellis and Brown, 1999) and the Central Medjerda basin, Tunisia (Faust et al. 2004). Cosmogenic dating has so far only been used for dating Pleistocene terraces (Rixhon et al., 2011). In order to counteract the bias in the ^{14}C record, an OSL dataset is presented in this paper. This has been derived from 17 regional studies using predominantly quartz luminescence but in some cases infra-red luminescence of feldspar (Fig. 10). All studies involved the etching of grains, used Rizø OSL counters and followed internationally accepted protocols and multiple aliquot methods. It is expected that the largest source of inter-laboratory variance is probably the estimates of dosimeter but that this should vary less than 15% (Hilger et al. 2001) Given the uncertainties of the method an uncertainty of at least 10-15% must be assumed. Also given the relatively low number of OSL dates available (139 in this study) the data are described and discussed in one dedicated section rather than distributed across the individual regional sections.

The discussion section incorporates meta-analyses of demographic proxies and pollen-count data. Demographic proxies have been derived from archaeological data (radiocarbon dates and settlement data) either harmonized from online sources and existing databases (e.g. BANADORA, RADON, University of Oxford's ORAU, EUROEVOL's project) or, manually inputted from a wide range of publications. The probabilities from each calibrated date are combined to produce a summed probability distribution (SPD). Following previous works (Weninger and Harper, 2015; Williams, 2012) showing that normalized calibrated dates emphasize narrow artificial peaks in SPDs due to abrupt peaks of the radiocarbon calibration curve, we opted to use non-normalized dates prior to summation and calibrated via IntCal13 curve (see Palmisano *et al.* 2017; Roberts *et al.* 2018). Two different proxies derived from archaeological settlement data have been used to assess demographic trends over the long run: raw site counts and

summed settlement area. To address issues related to the temporal uncertainty of occupation periods, we adopted “aoristic”, and Monte Carlo approaches (for a more detailed explanation of the methodology see Palmisano *et al.* 2017). We present a brief synthesis of key demographic trends for each region, while the detail is contained in the region-specific papers in this special issue.

We obtained pollen count data from the European modern (Davis *et al.*, 2013; Fyfe *et al.*, 2018; Palmisano *et al.*, 2017; Woodbridge *et al.*, (n.d.)) and fossil pollen databases (EPD version: Oct. 2017) (Leydet, 2007-17). Descriptions of the methodological approaches developed and applied to the pollen datasets are provided in Woodbridge *et al.* (in press) and Fyfe *et al.* (2018). The pollen count data from each site were summed into 200-year time windows and analyses were applied to the entire Mediterranean region in order to identify key vegetation types (Giesecke *et al.*, 2014). Pollen indicator groups were used to summarize key changes in the datasets. This included calculating the average arboreal pollen sum (%AP), a sum of tree crop indicators OJC (*Olea*, *Juglans*, *Castanea*) (Mercuri *et al.*, 2013a) (also combined with *Vitis*), calculation on an anthropogenic pollen index (API: *Artemisia*, *Centaurea*, cichorieae and *Plantago*, cereals, *Urtica* and *Trifolium* type) (Mercuri *et al.*, 2013b), and a sum of pastoral indicators (Asteroideae, Cichorioideae, *Cirsium*-type, *Galium*-type, Ranunculaceae and *Potentilla*-type pollen) (Mazier *et al.*, 2006).

Much of the geomorphic evidence presented below is derived from studies of alluvial/fluvial deposits and the dates assigned to phases of flooding, which might include overbank deposits and/or fluvio-torrential units (see table 1 for record types). We also refer to alluvial/fluvial units deposited in coastal zones that result in progradation or aggradation of the coastline. More than other environments, late Holocene Mediterranean coastal areas have evolved through an interplay of natural and anthropogenic factors (Anthony *et al.*, 2014; Marriner and Morhange, 2007, Arnaud-Fassetta 2002). Sometimes delayed responses of river systems in lower or coastal plains mark the delay of the stony load from the head of the watershed, which further complicates the search for causalities. Some lake-based studies are considered, in particular, the work in Anatolia where clastic units have been trapped in between the laminated sediment of Nar Lake. Finally, some examples of geoarchaeological work on colluvial units and soil erosion are presented, the most notable example of this being Fuchs’ work in the northeast Peloponnese and Notebaert & Berger in the Middle Rhone Valley (Fuchs, 2007a; Notebaert and Berger, 2014b).

Type of Record	Anatolia	Greece	Italy	France	Spain
Lake records (incl varved)	Nar Lake	Lake Kournas (Crete)	Lake Ledro		
Alluvial/fluvial		Rivers in Nth Greece/Macedonia (aggradation), Rivers on Crete	Turano basin	Mid-Rhone Valley	Slackwater deposits, overbank facies and alluvial gravels**, aggradation (Guadalete).

Colluvial		Sections (Peloponnese),		Mid-Rhone Valley	
OSL	Colluvium & alluvium*	Colluvium & alluvium	Alluvium*		Slackwater deposits

Table 1: Resumé of sedimentary record types employed in the principal analytical-curves by study region. OSL: x* - inadequate number of dates for a reliable curve. **Most of the Spanish sediments for the southeast region, which corresponds with the region studied as a part of this project, comprize overbank and alluvial deposits. A range of process types was assessed with those providing the most secure chronological controls chosen as the key datasets and curves employed in the regional syntheses.

3 EXAMPLES BY COUNTRY/REGION

3.1 ANATOLIA

3.1.1 Coastal

There are very few studies of coastal and delta progradation on the south coast of Anatolia, with most research having focussed on the rivers that enter the western Anatolian (Aegean Sea) coast. Here some major archaeological sites, such as Troy and Ephesus, are now located inland, but originally lay on, or very close to, the coastline (Brückner et al., 2005; Kayan, 1999). The Classical city of Miletus, for example, initially lay at the mouth of the Meander river, (Brückner et al., 2006, 2017), but this has pushed its mouth seawards in recent millennia. Although delta progradation can be partly explained through natural long-term geomorphological dynamics (including sea-level change impacting on accommodation space), evidence suggests that the rate of progradation has accelerated since the first millennium BC as a consequence of human-induced increases in erosion and sediment discharge (Lambrianides and Spencer, 1998). The interaction between naturally varying rates of progradation, such as the mid-Holocene slowdown in relative sea level rise (the 'Stanley effect' cf. Stanley and Warne 1997) is geographically unique due to isostatic, geomorphic (including soil) and local population history but it can have important human implications (see discussion).

3.1.2 Lake Records

Within lakes, gross lake sedimentation rate data may not accurately reflect flux rates from catchment erosion. A good example is provided by Nar Lake, a small volcanic maar in Cappadocia, central Anatolia. Most of the Nar lake sediments are annually laminated, allowing rather precise chronological resolution, but these varves are comprized primarily of endogenic carbonate, organic matter and diatoms, with only a minor detrital component. Hence overall sedimentation rate changes do not reflect catchment erosion very strongly. Interbedded between regular varves are periodic clastic layers, interpreted as turbidites for the most part. Visual recording, Itrax elemental chemistry and thin section analysis show that these clastic

layers were rare before the late Holocene (Fig. 3). On the other hand, after 2600 BP there are numerous inwash layers, notably during times when pollen data indicate land-use intensification (Roberts et al., in review).

3.1.3 Alluvial and Colluvial slope fills

There have been many geoarchaeological and geomorphological studies of Holocene alluvial and colluvial sediments in southern Anatolia (e.g. Boyer et al., 2006; Dreibrodt et al., 2014). Some of the most detailed studies have taken place linked to archaeological studies in the territory of the ancient city of Sagalassos, and which are also linked to pollen studies in the same area (e.g. Bakker et al., 2012). Dating of basin fill and fluvial sequences by 178 radiocarbon ages for geomorphological activity and their analysis using summed probability distributions shows overall low rates of sediment flux before the late Holocene (Dusar et al., 2012). This changed dramatically at 2800-2400 BP when a significant pulse of colluvial inwash entered some basins (e.g. Gravgaz) from surrounding hard limestone hillslopes. At the same time, pollen data indicate the first large-scale woodland clearance followed after ~2400 BP by the cultivation of tree crops and cereals, marking the onset of the so-called Beyşehir Occupation phase in this area (Fig. 3). The increased sedimentation rates coincided with the initial significant anthropogenic disturbance of the landscape, and sediment accumulation decreased afterwards, mainly as a result of the depletion of hillslope sediment sources (Van Loo et al., 2017). Sedimentation rates during the most prominent occupation of the area, i.e. during Roman Imperial times, were consequently much lower. By contrast, downstream river floodplain sedimentation has been substantial up until recent times, indicating not only a significant time lag in sediment delivery to the lower reaches of the river valleys but also significant reworking of previously deposited fluvial sediment (Verstraeten et al., 2017).

3.1.4 Short resumé of demographic trends

The first increase in human activity (as represented by the SPD of radiocarbon dates) took place in south-central Anatolia at ~10,300 BP (Roberts et al., 2018). This phase of Neolithic population growth ended after ~7500 BP. A more substantial demographic increase, as represented by raw counts and total areas of sites, occurs at 5000 BP during the Early Bronze Age, from this point, the randomized duration index also increases (Fig. 3). Then the Hellenistic-Roman-early Byzantine (H-R-B) phase, centred on 2000 BP is characterized by a significant increase in population, followed by a dramatic decrease just after 1500 BP.

3.2 GREECE

A synthesis of OSL and ¹⁴C dates from the Eastern Mediterranean (Jones et al., 2015) reveals interesting patterns. Most of the dated fluvial and colluvial records in this Eastern Mediterranean dataset are from Crete or southern mainland Greece, so it is appropriate to compare them with the land cover and demographics from this region (Figs. 4 and 5 and the Discussion section).

3.2.1 Coastal

The Greek mainland shoreline has been intensively investigated over the last five decades with a view to reconstructing shoreline migration and human activity during the Holocene (Runnels et al., 2005; Tourloukis and Karkanis, 2012). Records from some deltaic plains and lagoons indicate a detrital input dated from ~5500. BP, part of the early phase of delta development in the Eastern Mediterranean and Greece, however, climate controlled detrital inputs are observed earlier (Brückner et al., 2005; Psomiadis et al., 2014; Styllas and Ghilardi, 2017). The rhythms of delta progradation linked to major alluvial phases

are spatially varied due to local socio-cultural contexts and variability in regional climate, latitude, type of relief and exposure of oceanic currents.

On the Thessaloniki (or Macedonian) plain, the Neolithic site of Nea Nikomedeia, now located ~30 km inland, would have been located on the edge of an embayment at the beginning of the 7th millennium BP. During the Late Neolithic (6300 – 5900 BP) and Bronze Age (5000 – 3050 BP), there were phases of significant alluviation, a phase which broadly corresponds with regional erosion signals (Benito et al., 2015a; Jones et al., 2015). Recent research (Styllas and Ghilardi, 2017) reveals how the early development of the Aliakmon-Axios deltas also took place during the Late Neolithic 6100 – 5400 BP/4100-3400 cal. BC (Ghilardi et al., 2008a, 2008b, 2010). The second phase of important alluvial deposition is dated to 4200 – 3750 BP. A similar phase of Bronze Age (Early Helladic) erosion has also been identified on Euboea (around Chalkis to Amarynthos). This was followed by delta development during the Geometric period when urban centres flourished. Finally, the last major phase of delta development occurred from the Classical period until the Late Byzantine period.

3.2.2 Alluvial

Although just to the east of the zone for which we have demographic data, recent work in Eastern Macedonia and Thrace reveals evidence for localized anthropogenic soil erosion from the early Neolithic period (~8000 BP) at the bottom of the Dikili Tash site (Glais et al., 2016; Lespez and Glais, 2017). Soil erosion was far more significant during the Thracian period (Iron Age to Classical Greek period in southern Hellenic areas) and into the Roman period (Lespez, 2003, 2008). The impact of Thracian (~2400 BP) settlement has often been underestimated, but it is now apparent that it played an important role, with the extension and intensification of farming and grazing, as seen in the pollen data for this region (Fig. 4). Interestingly, this phase of erosion does not appear in the Eastern Mediterranean regional record (Jones et al., 2015). The second period of soil erosion and alluvial aggradation took place during the Ottoman period, linked to the intensification of farming in the plain and the valleys, along with the increase of grazing in the surrounding mountainous areas (Lespez, 2003). Therefore, we are sure that erosion in these small river basins in Northern Greece was triggered by local human activities and exacerbated by meteorological events, especially storms and their associated flash floods (Lespez, 2007; Glais et al., 2016, 2017).

Along the Angitis River, which drains the Philippos-Drama plain, alluvial aggradation accelerated early on; a significant alluvial event is recorded in the Lower Valley just after 6000 BP (Fig. 4), this may well have partly been caused by climate changes between 6500-5000 BP (Lespez et al., 2016). Some other short phases of alluvial aggradation, such as that along the Xeropotamos, might be explained by an increase in flood flows corresponding to wetter periods recorded at the global scale during the Early Byzantine (1500 – 1200 BP; Lespez, 2003; Glais et al., 2016). Studies of the main river valleys and floodplains show two historical phases of alluvial aggradation and soil erosion dated to the Thracian and Roman periods (2700 – 1800 BP) and more significantly during the Ottoman period (beginning of the 15th to the 20th ~AD).

3.2.3 Colluvial Sedimentation

Pope et al. (2003) considered the difficulty of dating alluvial fan units in the context of their work in the Sparta Basin, where several phases of deposition date to the Helladic period, with more sedimentation during the early to middle Hellenistic period and then again during the Late Roman to Byzantine (2003). These phases do correspond with other results from the area, but it is Fuchs' work in the Philious basin

(included in the meta-analysis of OSL dates below and see Figs. 5 & 10) that provides us with a rare example of well-dated series of sedimentary sequences. With OSL dates, we can assess localized processes and geosystem responses to anthropogenic and weather events. The colluvial record reveals high sedimentation from ~8700 BP, which may slightly pre-date the establishment of farming in this area. Work within small drainage basins on the island of Corfu reveals erosion dated to 8500/8400 and 8000/7900 BP. This corresponds to the arrival of Early Neolithic peoples (producing red monochrome pottery) and coincides with 8.2 ka RCC event (Berger et al. 2016). On the Peloponnese, there is a decrease in sedimentation from 6400 BP, and this relatively low level of sedimentation continues until ~4000 BP, reflecting an unusual absence of evidence for enhanced erosion during the Early Bronze Age, a period that we know witnessed significant settlement and economic activity. There was an increase in sedimentation during the Bronze Age (4000 – 3200 BP), an Early Iron Age peak (~ 3000 BP) and then significant erosion during the Roman Period (2000 – 1500 BP) (Fuchs, 2007b).

3.2.4 Crete

Coastal changes at Malia (Eastern Crete) suggest that human impact on the landscape, including burning, took place at ~ 4600 - 4250 BP. However, a more significant phase of erosion is dated to the Late Minoan IA or Hellenistic Period (Lespez et al., 2003). The most recent and detailed lake sedimentary record is derived from Lake Kournas, Northern Crete (Vanniere and Jouffroy-Bapicot, in prep'). The sedimentary evidence from this well-dated lake core reveals increased flood frequency centred on 8000 BP, 5600 BP, 3900 BP, 3200 BP, and the highest peak at ~ 2400 BP.

On southern-central Crete, widespread, coarse-grained aggradation occurred between ~4900 and 4200 BP and then 3400 to 3000 BP, processes probably driven by climate change (a phase of regional cooling between 5000 and 3000 BP) (Macklin et al., 2010; Theodorakopoulou and Bassiakos, 2017). However, we should also consider the probable impact of Bronze Age extension and intensification of arable and pastoral activity. One of the most significant phases of hydrological activity occurred between 5000 and 3070 BP, broadly corresponding with the Minoan period (Ghilardi et al., 2018). Drier conditions are reported during the 1st millennium BC in South Central (Macklin et al., 2010) and Eastern Crete (Theodorakopoulou and Bassiakos, 2017). A significant phase of flooding is observed from ~2700 – 2500 BP on the Messara Plain (South Central Crete) (Ghilardi et al., 2018).

3.2.5 Short resumé of demographic trends

In both the northern and southern study areas, there is a noticeable increase in demographic indices from 8500 BP, followed by a temporary decline starting at ~8200 BP. Population levels seem to fluctuate within an overall declining trend until ~7400 BP, meanwhile population increases in the south from ~7400 BP until ~7100 BP. In the north, this same period sees oscillations within an overall increasing trend. In both areas, there is then a significant decline. The period between 5500 and 4900 BP is characterized by relatively low demographic indicator values, with the index increasing for the Early Bronze Age (Early Helladic). There is a peak in the north centred on 4400 BP, while in the south there is a significant peak at 4200. In both areas, this period is succeeded by decline. At ~5000 and 4500 BP (Early Helladic). In the North, demographic the indices start to increase at ~4100; the first phase of the middle Helladic on the mainland, while in the south, demographic indicators increase from 3900 BP, the start of the Proto-Palatial Period on Crete. Then, in both the northern and southern regions, there are fluctuations, with population indices remaining high in both areas until ~3000 BP with a significant decline following this and lasting

until ~2700 BP (the so-called Bronze Age Dark Ages). For the southern area, the raw counts of sites curve suggests a high level of human activity during the Classical and Roman periods.

3.3 ITALY

The Italian study focusses on the central, western regions (Tuscany and Lazio), with one example coming from beyond these regions.

3.3.1 Coastal

The coastal zone just to the west of Pisa, supplied by sediments from the Rivers Serchio and Arno, developed as a lagoon environment between 5000 – 4000 BP. During the Bronze to Iron Ages, this area then emerged as a poorly drained floodplain, with enhanced sedimentation linked to an increase in human activity (Amorosi et al., 2013). At roughly the same time (Etruscan Period onwards) coastal progradation of the Apuo-Versilian plain was significant (Fig. 6), while the most noteworthy phase of erosion took place during the Little Ice Age (Bini et al., 2013).

3.3.2 Lake Records

Although Lake Ledro (southern-central Italian Alps at 650 m) is located to the northeast of our Italian study region, it provides us with a well-dated sedimentary sequence that has been the object of detailed sedimentological analyses. While there were relatively ephemeral phases of erosion between 8000 and ~4200 BP, the period between 4200 and ~3000 BP was a period of complex climatic oscillations that saw a significant increase in erosion (Magny et al., 2009; Simonneau et al., 2013). Undoubtedly, this was a consequence of the combined effects of climatic variations, but also the exposure of more extensive areas of soil after forest clearance by people. After this period, there is evidence for flooding centred on 1900 BP; then, there was an increase in flooding and erosion during the later medieval period and the Little Ice Age (Joannin et al., 2014; Magny et al., 2009).

3.3.3 Alluvial

Based on 44 dates from alluvial zones in the Sabine mountains, evidence from central western Italy reveals the onset of a sedimentation phase at ~5500 BP. After 4200 BP, erosion increased, with one phase starting at ~4000 BP and a more significant period of erosion centred on ~3000 BP (Fig. 7). Up until 2300 BP, these signals mainly come from the central parts of the river catchments. From the beginning of the Roman period, sediment from the headwaters of the Turano basin was deposited in the Ricetto and Fosso Pechiera and the Ovito catchments. After this, there are a series of peaks in sedimentation (at 1750 BP, 1300 BP, 1000 BP and 700 BP) (Borrelli et al., 2014; De Vincenzo et al., 2013). The period around 1100 BP, saw the highest levels of sedimentation, probably enhanced due to the extended exposure of land surfaces (Mensing et al., 2015a). Within this study area, the palynological work at Lago Lungo (Monti Reatini), provides a useful assessment of human engagement with this landscape from the Archaic Period (2700 BP) to the modern era (Fig. 7B). Evidence for pasture and some impact on the forest is seen during the Imperial Period. This evidence for relatively limited Roman impact fits with the story of significant variation in the levels of anthropogenic impact on Roman landscapes across the Mediterranean. The Late Roman period saw the increased impact on the forest and evidence for intensive human activity. The Medieval to Late Medieval periods (~1100 – 600 BP) constitute a phase of significant human impact on the landscape, with the Little Ice Age seeing the recovery of forest (and reduced sedimentation) (Mensing et al., 2015a).

Palaeomagnetic and OSL dating of sediments from three adjacent catchments of a similar size, geology and relief; the Fiora, Marta and Treia (Brown and Ellis, 1996), is of particular interest as we can assume that these catchments experienced similar climatic conditions during the Holocene. The alluvial sequences of all three valleys show increased deposition at the beginning and end of the Roman period (2400 – 1600 BP), which also coincides with both raised lake levels in lake Bolsena and increased population growth in S Etruria (Potter 1979; Brown and Ellis 1996). Then, there is clear evidence for erosion and extensive alluviation again during the medieval-Renaissance period (900 – 300 BP) (Op cit.). Both phases broadly coincide with the record of wetter episodes within the predominantly dry Late Holocene record for central Italy as recorded both by the Tiber and Po flood records and by high levels of Lake Bolsena (Dragoni 1998).

In the Central Apennines (60 km northeast of Rome), the analysis of sediments from cores demonstrates significant phases of flooding between 2200 BP and 1500 BP with the Fosso delle Rosce's alluvial plain extending upstream at ~ 1400 BP (Borrelli et al., 2013). The Roman phase of extensive flooding in the Central Apennines has also been identified by Giraudi (2005) and on the Tiber delta (Salomon et al., 2014). The main period of noteworthy flooding was from the 2nd century BC through to the 2nd century AD, this was followed by another phase from the 5th to 9th centuries AD (Op cit.). Also, we should note that Roman historical records reveal high flood-frequencies during the Late Republican/Early Imperial periods (Aldrete, 2007), which may well have been a consequence of significant forest clearance in some areas (Borrelli et al., 2013).

The most recent peak of flooding, 550 – 350 BP, coincides with rapid growth in the population of renaissance Rome and corresponds to both the Little Ice Age and increased agricultural pressure on its hinterland in order to feed Rome. This trend continued into the Little Ice Age. The sedimentary record reflects the flood and lake records, with the evidence suggesting a primary role for climate change, probably via a solar mechanism. The records also show that the sedimentary response was enhanced in the catchment closer to Rome (e.g. in the Marta and Treia catchments).

3.3.4 Colluvial – Geoarchaeological/Geomorphological sections

There are relatively few studies of colluvial processes in our study area. We know that erosion from the Alban Hills dated to Late Bronze Age to Iron Age resulted in deposition across the Pontine Marshes; probably the consequence of increased agriculture in the hills. To some extent, this colluviation allowed Imperial Rome to extend agricultural activity across the marshes, but this attempt at wetland management appears to have had limited success (Attema, 2017; Attema & De Haas., 2005; Sevink et al., 2013; Walsh et al., 2014). Slope erosion and enhanced geomorphological instability in the Italian Alps and the Apennines, are associated with the acceleration of landslide phenomena (Cremaschi and Nicosia, 2012) during the phase of climate variations between ~ 4000 BP and 2000 BP, coupled with a significant shift in land management and soil use.

3.3.5 Short resumé of demographic trends

The demographic indices for this study area suggest a sudden increase in population (especially in the raw count statistic) from 6500 BP. The next period of demographic increase corresponds to the Late Neolithic and Chalcolithic (centred on 5000 BP). This is followed by a reduction in all demographic indices and then a Mid - Late Bronze Age increase across all values (from ~3500 BP onwards). Population increase is also apparent during the Early Iron Age (3000 – 2700 BP). The Late Republican/Early Imperial Roman period

(2100 – 1900 BP) is unsurprisingly a time of significant human activity, specifically seen in the Archaeological site numbers statistic (Palmisano et al., 2017).

3.4 FRANCE

3.4.1 Coast

In the Camargue area, progradation took place during the Late Neolithic and into the Copper and Bronze Ages (4500 – 3000 BP). However, the fastest rates of coastal advance occurred during the Roman Period (Arnaud-Fassetta and Suc, 2015). On specific archaeological sites, such as the Greek/Roman site of La Capelière, we see evidence for Iron Age to Roman flooding events, which resulted in the temporary abandonment of this part of the Camargue (Arnaud-Fassetta and Landuré, 2003). A broadly similar sedimentary history characterizes the coast further to the east around Fréjus where the deltaic zone of the river Argens expanded rapidly during the first century BC and prograded by about 1 m/yr up until the second century AD (Devillers et al., 2007). A similar pattern is observed around Marseille, where there is evidence for Bronze Age sedimentation. However, here it is with the advent of Greek, then Roman activity, that significant sedimentation occurred (Morhange et al., 1999).

3.4.2 Alluvial

A detailed high-resolution geochemical study of Lake Bourget, based on measurements of terrigenous input, reveals four principal phases of erosion caused by climate changes and exacerbated by human activity: 4400, 2700, 2200 BP and at the end of the Wet Medieval Period around 1000 BP (3 centuries before the LIA climatic degradation) (Arnaud et al., 2012). In the southwestern Alps, detrital fluxes in lakes record the same trend with a general step around 4000/3500 BP when anthropogenic pollen increases (Brisset et al., 2017). For the central part of our Southern French region of study, we have chosen the most recent and most comprehensive dataset available. The cumulative probability distribution function (CPDF) for the Mid-Rhone Valley is founded upon a 25-yearlong study of palaeosols and sediments. The erosion/aggradation curve in Figure 8 is based on 423 radiocarbon and chronocultural dates covering 13 millennia. This provides us with an exceptional example that can be contrasted with high-quality archaeological settlement/demographic data (for a detailed explanation of the methodology and statistical analyses, see Berger 2015 and Berger and Brochier submitted).

Most of the ^{14}C dates come from the alluvial sequences of the lower valley of the Drôme River, Valdaine and Tricastin areas. The construction of the CPDF chronology is founded on the approach of extensive comparisons of sediments from fluvial systems, alluvial plains up to mountain slopes. It incorporates (1), signatures for the rapid filling of riverbeds, palaeomeanders, aggradation of levées to distal floodplains (referred to below as “hydrosedimentary” activity), (2) Periods of sandy-gravel progradation associated with high-energy tractive currents that were the product of increased connectivity between the slopes and upper basins (Fig. 8). This second category is described as “coarse sedimentation”. There are phases where we see the overlap of the erosional activity curves and the pedogenesis curve. In most cases, these reflect cumulative dates with wide standard deviations (thus the overlap). However, some cases probably reveal the complexity and variation of spatially heterogeneous processes (including variations in human impact) across a set of complex topographic contexts.

During the Early Neolithic, there was an initial phase of hydrosedimentary and coarse-sedimentary erosion centred on ~7200 BP; this segued into a phase of pedogenesis. The period between 6900 and 5600 BP saw some hydrosedimentary activity and particularly marked pedogenesis/stability, which peaked at ~5.5 BP. The Late Neolithic/Copper Age (centred on 5200 BP) included phases of enhanced hydrosedimentary activity, leading into a period from ~4900 BP characterized by relatively low levels of erosion and pedogenesis/stability, especially from 4500 to 4300 BP. This phase transitioned to another complicated period (4200 to 3600 BP) comprising both hydrosedimentary and coarse erosion. Hill-slope erosion was relatively common from the Bronze Age to the start of the Iron Age (4000 – 2700 BP) (Notebaert et al., 2014). The Early Bronze Age period witnessed the development of extended and intensified agricultural practices. Between 3500 and 2900 BP, there were two phases of hydrosedimentary activity with some coarse sedimentation, this phase also comprises evidence for pedogenesis, especially between 3000 and 2700 BP. This Late Bronze Age to Early Iron Age phase was immediately followed by one of the two most significant periods of hydrosedimentary and coarse sedimentation, lasting until ~2400 BP. This Early to Middle Iron Age period, with the development of a highly-structured economic landscape, coincided with a wetter, cooler climatic period (Göschenen I glacial advance) and high lake levels in central Europe (Holzhauser et al. 2005). Less significant erosion and pedogenesis succeeded this period. The end of the Gaulish and the high Roman Empire (2100 BP – 1500 BP) coincided with the second significant phase of hydrosedimentary and coarse-sedimentary erosion (Notebaert and Berger, 2014a). This period corresponds to the Roman Climatic Optimum. The erosion dated to the Little Ice Age and Medieval to modern periods appears less significant than that recorded for the preceding periods, but this difference is mainly due to the lack of datable material in the coarse gravels formations of the rhodanian braided-style rivers. Comparison of the sediment budget with data on settlement density indicates the influence of settlement density and associated land use on soil erosion and sediment deposition.

3.4.3 Colluvial – Geoarchaeological/Geomorphological sections

Well-dated colluvial sequences are relatively rare, except in the Mid-Rhone valley. One synthesis of geoarchaeological research from across Provence, based on units dated via ¹⁴C or associations with archaeological material, clearly indicates phases of erosion dated to the Late Neolithic/Early Bronze Age (4000 BP), then the Early Iron Age and the Late Roman to Early Medieval Period (Provansal, 1995). A sediment budget study on slope colluvial storage was undertaken across the Valdain Basin (Notebaert et al. 2013, Notebaert & Berger 2014). Here, slope erosion decreased during the Bronze Age before an increase at the Bronze-Iron Age transition, which lasted until the Roman period.

3.4.4 Corsica

On Corsica, Geoarchaeological investigations in specific coastal zones reveal significant phases of alluviation broadly dated to the Late Neolithic to Chalcolithic times (centred around the 5500 -5200 BP RCC event as suggested by Mayewski et al. (2004)) and the Bronze Age (centred around the 4200 and 3200 BP RCC events). Evidence from the Tavignano River reveals deltaic plain development dated to the Late Neolithic-Early Chalcolithic (Ghilardi et al., 2017; Guttman et al., 2013). On the Taravo River (SW Corsica), we see a significant phase of alluvial aggradation from the Middle Bronze Age (ca. 3200 cal. BP) with the Canniccia pond completely in-filled during the Late Bronze Age (3200-2800 cal. BP) (Ghilardi et al., 2017). Here, sedimentological and magnetic parameters reveal a significant phase of alluviation (~3 m in three centuries) during the Late Bronze Age, which was a period of intense cultural re-organisation in

the valleys of the island (Bagolini, 1996; Cesari and Peche-Quilichini, 2015). Research on the Sagone River (W Corsica) indicates that significant deltaic progradation occurred at the end of the Late Bronze Age; a period with relatively low levels of human activity (Ghilardi et al., 2017).

3.4.5 Short resumé of demographic trends

From 7500 BP, there is an increase in all demographic indices (raw counts, aoristic weight, and randomized duration). Demographic indices then fluctuate, followed by a definite increase and then plateau across all demographic measures from the Late Neolithic through to the Early Bronze Age (5600 to 4100 BP). The Early Bronze Age (~4200 BP) is one of the most interesting; we see an increase in demographic signals leading into this period. The demographic indices suggest lower populations for the Middle Bronze Age through to the start of the Iron Age. The Imperial Period compizes evidence for increased population levels.

3.5 SPAIN

3.5.1 Coast

In the Valencia area, sedimentation from the Turia River has resulted in coastal progradation, high magnitude flooding events between ~2800 and 2300 BP resulted in significant sand deposition, which was followed by levée formation from 2200 to ~1500 BP. The next significant processes comprized high magnitude flooding dated to ~900 BP (Carmona and Ruiz, 2011).

3.5.2 Alluvial

Thorndycraft and Benito's synthesis of fluvial records from Spain (with most examples located in the eastern regions of the country) revealed clusters of alluvial/fluvial sedimentation events dated to 9630–8785; 7980–6860; 5800–4800; 3880–3085; 2895–1820; 1300–0 BP. Since these are predominantly slackwater deposits, they reflect climate rather than land use effects. Of direct interest, is the phase between 3880-3085 BP that comprizes two peaks of slack water flood deposits and alluvial overbank facies. There then followed a period from 2895–1820 BP with the highest pre/protohistoric peaks for slackwater deposits in conjunction with alluvial channel (gravel) facies (Thorndycraft and Benito, 2006). Wolf et al's work in the Jarama river also revealed phases of erosion (Central Spain), most of which broadly correspond with those identified by Thorndycraft and Benito's, i.e., periods of sedimentation between 8000 and 7000 BP, 5000 and 3100 BP, 2200 and 1800 BP, then at around 1000 BP and finally, at 400 BP (Wolf et al., 2015). The Guadalete river (southern Spain) records phases of almost synchronous aggradation at the Holocene scale around 8000-7000, 6600-5500, 4500-3800, 2300-1900, 1000-800 and 400 BP (Faust and Wolf, 2017) (see Fig. 9).

Work on the Lorca alluvial fan (southeast Spain), produced little evidence for sedimentary activity during the Neolithic, and stability and pedogenesis seemingly characterize the Late Chalcolithic/start of the Bronze Age (soil horizons dated to 4400 to 4200 cal. BP). This is followed by a period of alluvial sedimentation during the later Bronze Age. Alluvial sedimentation is also dated to the Late Iron Age. In summary, the data analysed indicate that sedimentation over the ancient fan-surface was negligible during the Neolithic and Chalcolithic (before 4500 BP) (Silva et al., 2008). In southeast Spain, it seems likely that enhanced sedimentation was caused by increased aridity during the late Holocene (reduced vegetation cover exposing soils and sediments), the principal phase dated to ~ 1300/1200 cal. BP (Bellin

et al., 2013). In the Librilla “Rambla” (Murcia), there is evidence for higher sedimentation–erosion rates during the Copper, Bronze and Earlier Iron Age (Calmel-Avila, 2002). A slightly different sequence of processes has been identified in the lower part of Guadalentín; here the important sedimentation phases are dated to; 7500 – 5000 BP, 3400 – 1600 BP, 700 and the 400 BP (Baartman et al., 2011). Schulte’s work on historic period flooding reveals enhanced fluvial dynamics during the period 500 – 100 BP (2003), while the assessment of Late Holocene flood frequency and magnitude in this area by Benito et al. reveals a significant increase in high-magnitude floods during the nineteenth century (2010).

3.5.3 Colluvial – Geoarchaeological/Geomorphological sections

One study from the Lower Huerva Valley, Ebro Basin, NE Spain, identified slope deposits, where the lower unit is dated to 4346–4134 BP (late Chalcolithic) (Pérez-Lambán et al., 2014). While work in Central Spain (with OSL dates) reveals Early to High Medieval Central Spain erosion (Ortega et al., 2017).

3.5.4 Short resumé of demographic trends

The demographic indices for the southern Spanish study zone suggest an Early Neolithic increase in human activity, but this is less apparent than in the north. From the Late Neolithic, through the Chalcolithic and into the Early Bronze Age there was a steady increase in population. This trend came to a relatively dramatic end from 3600 BP onwards. There was a renewed increase during the Iron Age from ~ 2600 BP. Although we do not have SPD date for the Roman Period, we know that activity across the Spanish study area increased significantly during this phase, and this is reflected in the anthropogenic pollen signal.

3.6 AN OSL SPD ANALYSIS

A database of 137 OSL dates was assembled from published and unpublished sources in Spain, Italy, Hungary, Turkey, Greece and the islands of Corsica, Malta and Crete. Here, we focus on those regions that produced more than 30 dates (Table 3). While this is lower than is ideal (which is over 250 dates), the pattern is far from uniform over time. As figure 10 shows, the full dataset has distinctive peaks at 7800, 2400 and 780 BP. In the current dataset, the dates from Spain are the youngest with the youngest peak from the entire dataset at 520 BP. These data can only at this stage be regarded as preliminary due to the small dataset and therefore strong biases from the effects of individual studies. For example, the Spanish data are dominated (18 of 34) by the studies of slackwater sediments (Machado et al., 2017; Medialdea et al., 2014). These studies had different aims and therefore, different dating strategies. However, as more studies using OSL are undertaken, we will be able to judge if this east-west trend is a real artefact of site selection and apply more stringent quality control to the dataset.

Area	No of dates	Pre-Holocene	Peak 1	Peak 2	Peak 3
<i>All</i>	137	~13-15 Ka	7800	2400	780
<i>Spain</i>	34	~12-13 Ka	5750	2600	520
<i>Greece</i>	59	-	4300	3000	1300

Table 2: SPD peak dates from alluvial OSL dates in the Mediterranean (initial dataset). The “All” dates row refers to the accumulated value from Spain, Italy, Hungary, Turkey, Greece and the islands of Corsica, Malta and Crete. Spain and Greece are detailed as the two zones with more than 30 dates each. Dates in bold are the highest SPD peak in the set.

4 DISCUSSION

Here, we consider the pan-Mediterranean chronological trends via reference to the SPDs of ^{14}C and OSL dates for erosion/sedimentation, the demographic and palynological evidence along with current assessments of climate changes during the Holocene.

Overall, geomorphological and geoarchaeological data indicate a marked increase in Mediterranean sediment dynamics during the last four millennia. However, there were substantial regional variations in timing, and significant oscillations in sediment fluxes were superimposed on the long-term upward trend in sediment flux (Fig. 10). The late Holocene acceleration in the rate of soil loss in the Mediterranean has parallels in other regions such as northwest Europe, where it has been linked primarily to human-induced forest clearance for agriculture on loess substrate (Dearing and Jones, 2003; Dotterweich, 2008; Hoffmann et al., 2010). In addition to land cover change, the increase in Mediterranean erosion has also been linked to Holocene aridification of the climate. More specifically, between 8500 and 6500 BP, southern and eastern regions were overall wetter than today, and then became progressively drier, with maximum aridity typically around 3000 BP (summarized as nMDS axis 1 in figure 10). In northern and western Mediterranean regions, such as parts of Italy and Northern Iberia, a different and less coherent pattern of Holocene climatic change is registered (see Finné et al. this volume).

4.1.1 Human-landscape dynamics from the Neolithic onwards

In the eastern Mediterranean, the period between 7000 and 6000 BP saw initial aridification, followed by fluctuations between arid and humid phases. From 8000-7000, the summary curve of isotope (e.g. $\delta^{18}\text{O}$, $\delta^{13}\text{C}$, δD) data from across the Mediterranean, represented as a number of sub-regional z-score means seems to show wetter conditions in all regions except for Northern Iberia. Here, conditions were dry from 8000-7400 BP followed by wetter conditions (see Finné et al. this volume).

The Eastern Mediterranean regional sedimentary signal suggests erosion during the seventh millennia BP. However, more localized studies yield little evidence for significant erosion during the Early Neolithic; one exception being southern Greece, where the palynological evidence, the demographic and specific erosion indices intersect and suggest the emergence of an agricultural landscape, including pastoral activity, which resulted in some erosion (Fig. 4). This period, during the ninth and eighth millennia BP also coincides with the 8.2 ka event. In some areas, it is possible, that even with relatively low population levels, the initial exposure of soil after deforestation was a moment when the geosystem was at its most vulnerable, especially in hilly/mountainous zones such as those in the Peloponnese.

The evidence from the South of France also allows us to suggest a change in the dynamics of the geomorphic system during the Early Neolithic (Cardial) (centred on ~ 7300 BP) after an extended period of pedogenesis under moister conditions (Berger et al. 2016). The Anthropogenic Pollen Index for the South of France fluctuates from ~ 7400 BP; increasing significantly then falling off to a low point at ~ 6200 BP. This combined evidence for human activity intersects with coarse-grained sedimentary erosion at ~ 7200 BP, during the initial intensification of Neolithic activity (Late Cardial) (Berger et al. 2016) (Fig. 8). A similar scenario is also possible for Spain, during the phase of Cardial neolithization. From about 7500 BP we see the start of human impact on forest/woodland, with regional pastoral indicator species increasing from ~ 6500 BP. There is some evidence for erosion at a regional scale where we have the intersection of an increase in population and some, albeit limited, erosion (Fig. 9).

4.1.2 Late Neolithic to Bronze Age - "Secondary Products Revolution".

After the sixth millennium BP, agricultural activity extended into the mountainous areas, limestone hinterlands and plateaux across much of the Mediterranean (Walsh 2013, chapter 5). In many areas, this economic configuration comprised the "Secondary Products Revolution"; the intensified exploitation of milk, wool and animal traction for the ploughing of hitherto unexploited parts of the landscape (Evershed et al., 2008; Greenfield, 2010; Sherratt, 1981). There is relatively little evidence for significant erosion in the southwest Turkey case study area despite the significant rise in the Anthropogenic Pollen Index centred on 5500 BP, along with an increase in population centred on ~5000 BP (Fig. 3). This was a period, which was relatively humid across Turkey, while other parts of the eastern Mediterranean transitioned to a more arid phase.

Delta development across parts of Greece, Italy and Southern France during the Late Neolithic may represent sedimentation initiated by human activity in the hinterlands, although the demographic and palynological data do not imply significant levels of human activity, but delta development is also strongly related to the mid-Holocene slowdown in relative sea level rise (Stanley and Warne 1993, 1997). Here, we should also remember that erosion, as the removal and redeposition of sediment, can contribute to the creation of valuable land, and any assumption that erosion is by default a "problem" should be questioned (Bintliff, 2002). Whatever the balance of causes, which must have varied geographically, the result was of human significance in producing an increase in useable, and later reclaimable, land largely due to the change in channel systems from braided to anastomosing and straight patterns with levees; in the case of the Nile this had a dramatic effect on the potential productivity of the lower Nile valley (Penington et al. 2017).

For Southern Greece, the palynological data reveal an increase in agricultural/pastoral activity during the Chalcolithic and Early Bronze Age but this did not engender significant erosion. However, after ~4000 BP, the pollen and demographic indicators all record an increase in settlement activity and we do see a phase of erosion (Fig. 5). This signal is also reflected at a regional level and is clearly represented in a peak in the summed probability analysis of OSL dates (Fig. 10). There is a possibility that the extension and intensification of arable/pastoral activities into hitherto unexploited zones coincided with a period of climate change; a combination that resulted in significant erosion in some areas. In parts of the western Mediterranean, a humid phase started at ~5200 followed by aridification from ~4300 BP until ~3000 BP. An arid phase is also apparent in the east, centred on 4200 BP but was soon succeeded by relative humidity and then a new phase of aridity centred on ~3000 BP (Brayshaw et al., 2011; Roberts et al., 2011). The nMDS curves indeed suggest an increasing trend towards aridity from this point (Fig. 10) as underlined by Jallut et al. (2009) and Magny et al. (2009); a product of orbital forcing. We should consider a scenario where pasture and ploughing (the introduction of new technologies, e.g. metal ploughs) moved into vulnerable landscapes. This is, in all likelihood, the period when significant agricultural terracing emerged to facilitate hill-slope agriculture within vulnerable, topographically complex, landscapes (Krahtopoulou and Frederick, 2008). In Greece and Crete, we see some evidence for erosion at around 4300 - 4200 BP (peak 2 on the OSL curve for Greece). A "Minoan Little Ice Age" (ca. 5000 - 3000 BP) has been suggested by Moody (2000), a period when most of the investigated Cretan river valleys record several flood events, with Minoan pottery embedded within the coarse alluvial deposits. There is no doubt that across much of the Mediterranean, this period saw the complex intersection of climatic variation, or unpredictability, coincide with the emergence of complex socio-economic systems. It is possible that some

complex societies, with centralized ideological and socio-economic frameworks, stifled adaptiveness or resilience during periods of climatic stress. Therefore, it may not have been climate *per se* that caused erosion, but rather an intersection of complex socio-ecological processes (Weiberg & Finné, 2013).

In Italy, there is a short-lived spike in regional pastoral indicators during the Late Neolithic. The next period of demographic increase corresponds to the Late Neolithic and Chalcolithic (centred on 5000 BP). The period from ~3300 BP saw an increase in arboriculture (OJVC curve) and a relatively high demographic index (Fig. 7). As with other regions, the phase of erosion seen here is in all likelihood the product of the combination of extended agriculture and a climatic context that exacerbated erosion, in particular, a dry-humid-dry oscillation centred on ~4000 BP (Finné et al. this volume; Magny et al. 2009; (Zanchetta et al., 2016)). In southern France, from the northern to the southern Alps, and in the Middle Rhone Valley, this story is repeated, with erosion centred on 4200 BP (Figs 8 & 10). In Italy and France, there is also a slight downturn in demographic indices. A similar scenario existed in Spain, although here, the phase of erosion and decline in population was between 5000 and 4500 BP (Chalcolithic); the period when complex fortified sites such as Los Millares developed.

Moving towards the end of the Bronze Age, in Greece, there is a relatively short-lived, albeit significant, peak in erosion during the Late Bronze Age (~3200 BP, represented by peak 2 on the Greek curve for OSL SPD), a period with relatively high population levels and intensive agricultural activity. One of the most significant phases of hydrological activity occurred during the Minoan period (Ghilardi et al., 2018). In Italy, a phase of erosion just after 3000 BP precedes a peak in the SPD of ¹⁴C for demography. In France, while there is evidence for some Late Bronze Age erosion, it remains moderate and the archaeodemographics and palynological indicators do suggest a significant change in land use intensity, a precursor to the significant phase of erosion at ~2800 BP that was amplified by a regional climatic degradation (Arnaud et al. 2012, Berger 2015). In Spain, particularly in the south, there is a noticeable decline in activity that coincides with erosion and aridity, this evidence possibly reflecting the demise of the Argaric Culture (Castro et al., 2000, 1999; Gilman et al., 2015). From the demographic proxies, it seems that there was also a decrease in population after 3200 BP in some regions, such as Greece and Anatolia. In Italy, there is a downturn in demographic indicators (SPD of cal 14C dates) after 3000 BP (Fig. 7). One well-known example, with collapse centred on 3200 BP, is the demise of the Terramare Culture (Bernabò Brea et al., 1997). However, this population decline is less apparent in France other than a small decline at around 3000-2900 BP towards the end of the Bronze Age (Berger et al. 2007).

Reduction of activity in some areas can be partly explained by complex systems theory (Cline, 2014). Climate change may well have been one of several problems that affected these societies, but the evidence for erosion, rather than being a simple consequence of climate change, might represent a form of maladaptation, which included the exploitation of hitherto underexploited landscape types; i.e. complex, possibly rigid, socio-economic systems that failed to adapt to a range of problems, including networks of trade exchange focused on copper and bronze, new populations that undermined the palace economies of the eastern Mediterranean, possible repercussions for other copper-producing areas, including the Alps (Carozza et al., 2015; Sherratt, 1998).

4.1.3 Iron Age

There is evidence for significant erosion across much of the Mediterranean during the third millennium BP (Fig. 10). In Turkey and Greece, lake records and colluvial sedimentary sequences reflect this; the

Turkish record is in some ways the most interesting as this moment constitutes the first major phase of erosion in the records considered here. In Anatolia, pollen evidence suggests human impact on the landscape during the Late Bronze into the Iron Age. Comparison of geomorphological evidence from Nar Lake and Sagalassos with regional pollen and archaeo-demographic data (Fig. 3) show that erosion rates increased in these two areas after ~3000 BP, associated with the significant land cover change of the Beyşehir Occupation pollen phase, and the Hellenistic-Roman-early Byzantine (H-R-B) phase. Of particular significance was the pulse of colluvial inwash recorded in some basins within the region at ~2500 BP, i.e. in the first centuries following major anthropogenic land cover change.

Interestingly, erosion appears to be less significant at the height of the Roman Period. Other studies, especially that of Casana, in the southernmost area of modern Turkey, reveal a similar pattern, albeit with a slightly later start date. Significant erosion and aggradation started after 1800 BP, suggesting that initial agricultural practices were sustainable and it was a phase of enhanced storminess that initiated severe erosion (Casana, 2008). In Greece, the OSL curve has a peak at 3000 BP, and evidence from alluvial records in northern Greece suggest significant erosion a few centuries later, during the Thracian Period. On Crete, the mid-third millennium BP sediments at Lake Kournas reveal the highest number of flood events for the entire Holocene. In Greece, the Archaic period, centred on 2600 BP, demographic indices are higher for the south than for the north. However, evidence in the north for “Thracian” erosion is particularly evident. In Italy, population increase is also apparent during the Early Iron Age (3000 – 2700 BP), which coincides with an increase in pastoral indicator species as well as an increase in erosion (Fig. 4).

Significant erosion is also apparent in the south of France and Spain. As population levels and anthropogenic indicators of economic activity fluctuate across these regions, it is difficult to argue for a purely anthropogenic cause for this phase of erosion; it seems likely that climate change, shifting from arid to relatively humid in some areas, may well have driven much erosion. An important phase of flooding, which broadly corresponds with the Iberian Period (Benito et al., 2015b; Jones et al., 2015; Thorndycraft and Benito, 2006), may well represent the combination of increased precipitation and human impact on and exposure of land surfaces in central and southern Spain. There is a peak in the SPD OSL curve and a peak in flooding (~2600 BP) (Fig. 10). These trends also intersect with a transition from relatively arid to a humid climate in the south, this move to humidity taking place slightly later in the north (Finné et al. this volume). We should, however, consider the possibility of socio-economic instability, changing economic activities and practices, associated with the movement of Celtic peoples in parts of the region.

In France, the Late Bronze Age to Early Iron Age comprized two of the most significant periods of hydrosedimentary and coarse sedimentation at the Rhone basin scale and lasted until ~2600/2400 BP; the soil erosion rate significantly increased compared to previous periods (Notebaert and Berger 2014, Berger 2015). It resulted in a generalized metamorphosis of rivers towards braided forms across pre-Alpine basins. This process was very much a product of dense, coarse sedimentary loads that far exceeded the discharge capacity of rivers (Berger et al. 2008). This Early-Middle Iron Age period, with the development of a highly-structured economic landscape, coincided with the Göschenen I glacial advance and the major lake transgressive period of Bourget (Magny et al., 2017). Finally, demographic growth occurred across our Iberian case study areas. In fact, the period centred on 2600 BP compizes an ephemeral peak on the anthropogenic palynological indicator species as well as pastoral species.

4.1.4 Roman Republican and Imperial Period

The Roman Period presents us with some interesting variations across the wider Mediterranean region. First, the demographic evidence from Turkey suggests a peak of activity centred on ~1900 BP. Pastoral activity was relatively stable and remained so for much of the Roman Period, which also saw an increase in arboriculture (represented by increases in the OJCV curves (Fig. 3). Despite this, the studies presented in this paper suggest relatively low levels of erosion, although the regional summed probability analyses of floods and alluvial events for the Eastern Mediterranean show an increase in these events centred on 2000 BP (Benito et al., 2015a). In southern Greece, Fuchs' work reveals the most significant phase of erosion for the entire Holocene, while on Crete, the peak of flood events seen during the Classical Period waned by ~ 2000 BP. In the north (Macedonia, Thrace), there is some erosion, but again, this is less significant than that dated to the Thracian Iron Age. Unsurprisingly, this is a period of extensive and intensive economic activity, as revealed by the palynological and demographic data. However, moving into the Greco-Roman Period, there is evidence for a decline in population, or indeed, a reduction in site numbers in some areas, including the Peloponnese (Weiberg et al., 2016; Weiberg et al. this Volume). In Italy, the Late Republican/Early Imperial Roman period (2100 – 1900 BP) is unsurprisingly a time of substantial human activity (Palmisano et al., 2017). The evidence for erosion in some areas is limited (e.g. The Rieti Basin (Mensing et al., 2015)), although significant floods and erosion are apparent in other records, and are certainly recorded in historical documents referring to the Tiber River (Aldrete, 2007). In the Rhone Valley-South of France area, there is a clear set of coinciding signals of population increase, intensification of all forms of agricultural practice, some of which were highly-erodible vineyards, and erosion across all parts of the landscape with a definite increase of hydrosedimentary connectivity across the Rhone catchments. Across the Iberian study areas, although there is evidence for erosion dated to the Roman period, this may not have been as significant as in other parts of the Mediterranean. Perhaps one of the most noteworthy elements in the Northern Iberian story is the evidence for an extremely wet climatic phase centred on the centuries either side of 2000 BP.

We must consider the possibility that much erosion during the Roman Period was caused by the intensification of economic activity, coupled with enhanced late-summer storms that resulted in some significant erosion events. The Roman Climatic Optimum/Roman humid Period (a relatively warm and humid period centred on 2200 – 1900 BP) is considered a relatively “moist” period (Goudeau et al., 2015; McCormick et al., 2012; Reale and Dirmeyer, 2000), with higher temperatures and humidity (Chen et al., 2012). However, the increase in climatic moisture did not exist across all parts of the Mediterranean (Finné et al. this volume). It is also possible that late summer and autumn periods saw significant storm events that may well have exaggerated erosion signals in some areas, as observed in the French Alps (Giguet-Covex et al., 2012). A high flood frequency is recorded in the languedocian rivers during the Gallo-Roman Period (Berger et al. 2002). In addition, there was rapid progradation of continental river deltas such as the Rhone as well the Argens, Aude, Tiber (Arnaud-Fassetta et Landuré 2003). The causes of these changes were a combination of human forcing and regular rainfall, especially around 2000 BP (Berger et Bravard 2012). We know that Roman writers/agronomists were aware of erosion (Columella et al., 1948; Dotterweich, 2013), or at least, variation in soil quality (e.g Varro, see Loeb et al., 1934), but it is difficult to demonstrate the demise of agricultural production. Any shortfalls in agricultural yields could, of course, be mitigated by the extensive and comprehensive connected economic network which guaranteed the movement of surpluses around the empire (Horden and Purcell, 2000). Despite the evidence for Roman impact and management of the environment, claims for extreme, generalized, forest clearance and

concomitant environmental degradation is misplaced (Harris, 2013; Hughes, 1994, 2005). Based on our assessment, it is difficult to assign 2000 BP as the inception of the Anthropocene (Certini and Scalenghe, 2011), but there is no doubt that some areas, including parts of Gallia Narbonensis, did witness significant anthropogenic erosion at this time.

4.1.5 Late Roman to Medieval Period and the Little Ice Age

In the Eastern Mediterranean, the Late Roman to Early Medieval Periods witnessed relatively high levels of erosion. The final and highest peak in the OSL curve for Greece falls at 1300 BP (Fig. 10). Erosion in Turkey and Greece falls off after ~1300 BP but increases as we move into the Little Ice Age. The data from Italy reveals high levels of erosion at ~ 1400 BP, then again at ~1000 and 800 BP. Surprisingly, despite some Little Ice Age erosion; this does not appear as significant as some other areas in the Mediterranean. In the South of France, the Late Roman to Early Medieval periods also saw high levels of erosion at ~ 1400 BP, albeit, less significant than during the preceding centuries (but with regular high energy dynamics). The High Medieval Period (~1000 – 800 BP), comprising a phase of wetter climatic conditions in some eastern parts of the Mediterranean, but drier conditions in Iberia and parts of the Balkans, saw little erosion with concomitant pedogenesis/stability. Finally, the Little Ice Age is unsurprisingly characterized by a period of hydrosedimentary and erosion comprising coarse sediments, although, despite extensive and intensive agriculture, this erosion is less significant than in some previous phases. This is possibly due to the chronological and geographical interests of geomorphologists. In the south of France, combined climatic and anthropogenic forces led to erosion across entire catchments; from the mountains to the plains (Notebaert and Berger 2014a, Bravard 2010). The evidence for southern Spain a phase of erosion centred on 1000 BP, then relative stability followed by significant erosion/floods during the late and post-medieval periods, reflected in the final and most prominent peak in the OSL curve centred on 520 BP. This, as elsewhere, is undoubtedly a response to intensive agricultural activity and unstable climate.

5 CONCLUSIONS

Our regional syntheses are innovative, but only a point for departure towards more focussed sub-regional and catchment-based integrated projects. Although there are some broadly contemporary phases of erosion across the Mediterranean, such as that centred on ~4000 BP, the current state of our data does not allow or support simple causalities, i.e. the role of climate versus people. There is little doubt that this (~4000 BP) period can be characterized in some areas as a tipping point, or a significant moment for many landscape trajectories. In the case of the coastal zone, the natural slowdown in RSL rise combined with human-induced sedimentation had a significant effect in the production of productive marsh and agricultural land, which in some cases may have had significant demographic consequences (Bruckner et al. 2004; 2006; Penington et al. 2017). Then, between 3000 and 2000 BP, the fluctuations in climate combined with economic intensification across complex societies did result in significant landscape changes and increased soil degradation, and some areas crossed an irreversible threshold. What is clear from figure 10, is that the mid-Holocene aridification of the Mediterranean climate was not, on its own, enough, to increase erosion rates to the levels experienced in pre-modern times. It was only when there was a substantial and sustained increase in rural population and in woodland conversion to farmland and pasture, and the perturbation of soil structure via the extension in use of metal ploughs, that sediment flux rates increased to well above their long-term geological norm. This transition typically took place

during the 3rd millennium BP. Thus, while climate change may have been a key pacemaker for erosion events and a necessary condition for increased erosion, it was not a sufficient condition in the absence of human impact. Although many local studies indeed suggest that sedimentation phases in the Late Holocene are poorly correlated to climate signals, the role of climate cannot be neglected for several periods of increased geomorphic activity, especially when regional scales are considered (Faust et al., 2004; Bravard 2010; Bellin et al., 2013). For arid to semi-arid regions like south-eastern Spain or Tunisia, geomorphic instability is often correlated to drier periods that further degrade the vegetation cover making the landscape more vulnerable to the increasing frequency of torrential rain events, and the intensification of regressive erosion due to lower aquifers that often accompany a dry period. Such landscapes are also less resilient to human pressure, and thus human activity may intensify the climatic effect. In other regions, the return to humid conditions triggers more intense erosion, which again may be amplified through human activity. As Bellin et al. (2013; p. 124) suggested: “improved hydrological conditions are associated with both reduced geomorphic stability and accelerated geomorphic instability dependent on the human impact on the environment”.

Perhaps one of the most complex and interesting phases of human-environment interactions from our perspective is the Roman period. Notwithstanding the evidence for Roman impact and management of the environment, the notion of forest clearance and concomitant environmental degradation is problematic in many areas, despite the claims of some writers (Harris, 2013; Hughes, 1994, 2005). There is even variation in the flooding and erosion levels associated with the Little Ice Age; one interesting signal is that derived from the analysis of OSL dates that reveals a significant peak in Spain. It is difficult to claim that soil loss or erosion was responsible for the decline or demise of any of the societies considered here, although there is little doubt that soil erosion did contribute to localized problems and even forced populations movement (for a useful discussion of general issues, see Dotterweich, 2013). The problem of causality is aggravated by the positive feedback loop between climate and erosion, in that increased rainfall in semi-arid climates can both increase agricultural productivity and thereby intensify and natural erosion rates, with the opposite being true for reductions in effective precipitation (Brown and Walsh 2017). Moreover, the fact that some of the most significant erosion signals occur during periods when we know that societies remained resilient obliges us to consider more complex socio-environmental frameworks (Butzer and Endfield, 2012), where certain societies managed soil erosion due to adaptive forms of environmental knowledge while others were less successful.

ACKNOWLEDGEMENTS

Pollen data were extracted from the European Pollen Database (EPD; <http://www.europeanpollendatabase.net/>), and the work of the data contributors and the EPD community is gratefully acknowledged”. We should also like to thank the illustrators in the Geography Department at Plymouth, Southampton and Helen Goodchild at York. The PALEOMEX-MISTRALS program of the CNRS partly supported the research of the French scientists, and Kevin Walsh wishes to acknowledge the support for a research fellowship at IMÉRA, Aix-Marseille University, Marseille (Cofund EURIAS, Labex RFIEA + and ANR-Investissements d'avenir). We would also like to thank the three anonymous reviewers for their comments, which allowed us to enhance this contribution.

Bibliography

Aldrete GS (2007) *Floods of the Tiber in Ancient Rome. Ancient Society & History*. Johns Hopkins University Press.

Amorosi A, Bini M, Giacomelli S, Pappalardo M, Ribecai C, Rossi V, et al. (2013) Middle to late Holocene environmental evolution of the Pisa coastal plain (Tuscany, Italy) and early human settlements. *Quaternary International* 303(Supplement C): 93–106. Available at: <http://www.sciencedirect.com/science/article/pii/S1040618213001638>: doi:<https://doi.org/10.1016/j.quaint.2013.03.030>.

Anthony EJ, Marriner N and Morhange C (2014) Human influence and the changing geomorphology of Mediterranean deltas and coasts over the last 6000 years: From progradation to destruction phase? *Earth-Science Reviews* 139(0): 336–361. Available at: <http://www.sciencedirect.com/science/article/pii/S0012825214001822>: doi:<http://dx.doi.org/10.1016/j.earscirev.2014.10.003>.

Arnaud-Fassetta G and Landuré C (2003) Hydroclimatic hazards, vulnerability of societies and fluvial risk in the Rhone Delta (Mediterranean France) from the Greek period to the Early Medieval Ages. In: Fouache E (ed) *The Mediterranean World Environment and History*. Paris: Elsevier, 51–76.

Arnaud-Fassetta G and Suc J-PCN-01261498 (2015) Dynamique hydrogéomorphologique et diversité végétale dans le delta du Rhône (France) de -10 000 ans à demain. In: Reynard E E-DM and Borel G édit (eds) *Le Rhône, entre nature et société*. Pôle de recherche pour l'organisation et la diffusion de l'information géographique (PRODIG) Institut des Sciences de la Terre de Paris (iSTeP): Cahiers de Vallesia, p.63-98. Available at: <https://hal.archives-ouvertes.fr/hal-01261498>.

Arnaud F, Révillon S, Debret M, Revel M, Chapron E, Jacob J, et al. (2012) Lake Bourget regional erosion patterns reconstruction reveals Holocene NW European Alps soil evolution and paleohydrology. *Quaternary Science Reviews* 51: 81–92. Available at: <http://www.sciencedirect.com/science/article/pii/S0277379112003009>: doi:<http://dx.doi.org/10.1016/j.quascirev.2012.07.025>.

Attema P (2017) Sedimentation as geomorphological bias and indicator of agricultural (un)sustainability in the study of the coastal plains of South and Central Italy in antiquity. *Journal of Archaeological Science: Reports*. Elsevier B.V. 15: 459–469. Available at: <http://dx.doi.org/10.1016/j.jasrep.2016.07.024>: doi:10.1016/j.jasrep.2016.07.024.

Baartman JEM, Veldkamp A, Schoorl JM, Wallinga J and Cammeraat LH (2011) Unravelling Late Pleistocene and Holocene landscape dynamics: The Upper Guadalentín Basin, SE Spain. *Geomorphology*. Elsevier B.V. 125(1): 172–185. Available at: <http://dx.doi.org/10.1016/j.geomorph.2010.09.013>: doi:10.1016/j.geomorph.2010.09.013.

Bagolini B (1996) Il sepolcreto e gli insediamenti eneolitici di Spilamberto - S. Cesario nel quadro culturale medio-padano. *Preistoria Alpina*. Trento 32 "[Riedi: 235–285.

Bakker J, Kaniewski D, Verstraeten G, De Laet V and Waelkens M (2012) Numerically derived evidence for late-Holocene climate change and its impact on human presence in the southwest Taurus Mountains, Turkey. *The Holocene*. SAGE Publications Sage UK: London, England 22(4): 425–438. Available at: <http://journals.sagepub.com/doi/10.1177/0959683611425546>: doi:10.1177/0959683611425546.

Bellin N, Vanacker V and De Baets S (2013) Anthropogenic and climatic impact on Holocene sediment

dynamics in SE Spain: A review. *Quaternary International*. Elsevier Ltd and INQUA 308–309: 112–129. Available at: <http://dx.doi.org/10.1016/j.quaint.2013.03.015>: doi:10.1016/j.quaint.2013.03.015.

Benito G, Rico M, Sánchez-Moya Y, Sopeña A, Thorndycraft VR and Barriendos M (2010) The impact of late Holocene climatic variability and land use change on the flood hydrology of the Guadalentín River, southeast Spain. *Global and Planetary Change*. Elsevier 70(1–4): 53–63. Available at: <https://www.sciencedirect.com/science/article/pii/S0921818109001908>: doi:10.1016/J.GLOPLACHA.2009.11.007.

Benito G, Macklin MG, Zielhofer C, Jones AF and Machado MJ (2015a) Holocene flooding and climate change in the Mediterranean. *CATENA* 130(Supplement C): 13–33. Available at: <http://www.sciencedirect.com/science/article/pii/S0341816214003464>: doi:<https://doi.org/10.1016/j.catena.2014.11.014>.

Berger J-F, Delhon C, Magnin F, Bonté S, Peyric D, Thiébault S, et al. (2016) A fluvial record of the mid-Holocene rapid climatic changes in the middle Rhone valley (Espeluche-Lalo, France) and of their impact on Late Mesolithic and Early Neolithic societies. *Quaternary Science Reviews*. Pergamon 136: 66–84. Available at: <https://www.sciencedirect.com/science/article/pii/S0277379115301839>: doi:10.1016/J.QUASCIREV.2015.11.019.

Bernabò Brea M, Cardarelli A and Cremaschi M (1997) *Le Terramare, la più antica civiltà padana*. Electa.

Bini M, Ribolini A and Baroni C (2013) *Geoarchaeology as a tool for reconstructing the evolution of the Apuo-Versilian plain (NW Italy)*. : doi:10.4461/GFDQ.2013.36.18.

Bintliff J (2002) Time, process and catastrophism in the study of Mediterranean alluvial history: A review. *World Archaeology* 33(3): 417–435. Available at: <http://www.tandfonline.com/doi/abs/10.1080/00438240120107459>: doi:10.1080/00438240120107459.

Bintliff J (1992) Erosion in the Mediterranean lands: a reconsideration of pattern , process and methodology. In: Bell M and Boardman J (eds) *Past and Present Soil Erosion, Archaeological and Geographical Perspectives*. Oxford: Oxbow, 125–131.

Borrelli P, Domdey C, Hoelzmann P, Knitter D, Panagos P and Schütt B (2014) Geoarchaeological and historical implications of late Holocene landscape development in the Carseolani Mountains, central Apennines, Italy. *Geomorphology*. Elsevier 216: 26–39. Available at: <https://www.sciencedirect.com/science/article/pii/S0169555X14001664>: doi:10.1016/J.GEOMORPH.2014.03.032.

Borrelli P, Hoelzmann P, Knitter D and Schütt B (2013) Late Quaternary soil erosion and landscape development in the Apennine region (central Italy). *Quaternary International*. Elsevier Ltd and INQUA 312: 96–108. Available at: <http://dx.doi.org/10.1016/j.quaint.2012.12.007>: doi:10.1016/j.quaint.2012.12.007.

Boyer P, Roberts N and Baird D (2006) Holocene environment and settlement on the Carsamba alluvial fan, south-central Turkey: Integrating geoarchaeology and archaeological field survey. *Geoarchaeology* 21(7): 675–698: doi:10.1002/GEA.

Brayshaw DJ, Rambeau CMC and Smith SJ (2011) Changes in Mediterranean climate during the Holocene: Insights from global and regional climate modelling. *The Holocene* 21(1): 15–31. Available at: <http://hol.sagepub.com/cgi/doi/10.1177/0959683610377528>: doi:10.1177/0959683610377528.

Brisset E, Guiter F, Miramont C, Troussier T, Sabatier P, Poher Y, et al. (2017) The overlooked human

influence in historic and prehistoric floods in the European Alps. *Geology* G38498. 1.

Brown AG (1999) Geomorphological techniques in Mediterranean landscape archaeology. *Environmental reconstruction in Mediterranean landscape archaeology* 2: 45–54.

Brown AG (Antony G (1997) *Alluvial geoarchaeology : floodplain archaeology and environmental change*. Cambridge University Press. Available at: <http://www.cambridge.org/gb/academic/subjects/archaeology/archaeological-theory-and-methods/alluvial-geoarchaeology-floodplain-archaeology-and-environmental-change?format=PB#7TLue1HfXQK3MPJ7.97>.

Brown AG and Ellis C (1996) People, climate and alluviation: theory, research design and new sedimentological and stratigraphic data from Etruria. *Papers of the British School at Rome* 63(1995): 45–73.

Brückner H, Herda A, Kerschner M, Müllenhoff M and Stock F (2017) Life cycle of estuarine islands — From the formation to the landlocking of former islands in the environs of Miletos and Ephesos in western Asia Minor (Turkey). *Journal of Archaeological Science: Reports*. The Authors 12: 876–894. Available at: <http://dx.doi.org/10.1016/j.jasrep.2016.11.024>; doi:10.1016/j.jasrep.2016.11.024.

Brückner H, Müllenhoff M, Gehrels R, Herda A, Knipping M and Vött A (2006) From archipelago to floodplain—geographical and ecological changes in Miletus and its environs during the past six millennia (Western Anatolia, Turkey). *Zeitschrift für Geomorphologie NF* 142(Supplement): 63–83.

Brückner H, Vött A, Schriever A and Handl M (2005) Holocene delta progradation in the eastern Mediterranean— case studies in their historical context. *Méditerranée*. Presses Universitaires de Provence (104): 95–106. Available at: <http://journals.openedition.org/mediterranee/2342>; doi:10.4000/mediterranee.2342.

Butzer KW and Endfield GH (2012) Critical perspectives on historical collapse. *Proceedings of the National Academy of Sciences* 109(10): 3628–3631. Available at: <http://www.pnas.org/content/109/10/3628.abstract>; doi:10.1073/pnas.1114772109.

Calmel-Avila M (2002) The Librilla “rambla”, an example of morphogenetic crisis in the Holocene (Murcia, Spain). *Quaternary International* 93–94: 101–108. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S1040618202000095>; doi:10.1016/S1040-6182(02)00009-5.

Carmona P and Ruiz JM (2011) Historical morphogenesis of the Turia River coastal flood plain in the Mediterranean littoral of Spain. *Catena*. Elsevier B.V. 86(3): 139–149. Available at: <http://dx.doi.org/10.1016/j.catena.2010.12.006>; doi:10.1016/j.catena.2010.12.006.

Carozza L, Berger JF, Burens-Carozza A and Marcigny CCN-01245488 (2015) Society and environment in Southern France from the 3rd millennium BC to the beginning of the 2nd millennium BC: 2200 BC a tipping point? *2200BC - A climatic breakdown as a cause for the collapse of the old world?*. Géographie de l'environnement (GEODE) UMR 5600 Environnement Ville Société (EVS) Civilisations atlantiques & Archéosciences (C2A). Available at: <https://halshs.archives-ouvertes.fr/halshs-01245488>.

Casana J (2008) Mediterranean valleys revisited: Linking soil erosion, land use and climate variability in the Northern Levant. *Geomorphology* 101(3): 429–442. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0169555X08000652>; doi:10.1016/j.geomorph.2007.04.031.

Castro P, Gili S, Lull V, Mico R, Rihuete C, Risch R, et al. (2000) Archaeology and Desertification in the Vera

Basin (Almeria, South-East Spain). *European Journal of Archaeology* 3(2): 147–166. Available at: <http://eja.sagepub.com/cgi/doi/10.1177/146195710000300201>: doi:10.1177/146195710000300201.

Castro P V, Chapman RW, Gili S, Lull V, Mico R, Rihuete C, et al. (1999) Agricultural production and social change in the Bronze Age of southeast Spain : the Gatas Project. 73(August): 846–856.

Certini G and Scalenghe R (2011) Anthropogenic soils are the golden spikes for the Anthropocene. *The Holocene* 21(8): 1269–1274. Available at: <http://hol.sagepub.com/cgi/doi/10.1177/0959683611408454>: doi:10.1177/0959683611408454.

Cesari J and Peche-Quilichini K (2015) L'Âge du bronze corse, l'émergence d'une élite. *Dossiers d'archéologie* 370: 24–29.

Chen L, Zonneveld K a. FF and Versteegh GJMM (2012) Short term climate variability during “Roman Classical Period” in the eastern Mediterranean. *Quaternary Science Reviews*. Elsevier Ltd 30(0): 3880–3891. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0277379111003039>: doi:10.1016/j.quascirev.2011.09.024.

Cline EH (2014) *1177 B.C.: The Year Civilization Collapsed. Turning points in ancient history.*

Columella LIM, Ash HB and Forster ES (1948) *On Agriculture: In Three Volumes. Res rustica I - IV.. 1.* Heinemann. Available at: <https://books.google.co.uk/books?id=mBYanQEACAAJ>.

Costanza R (2012) Ecosystem health and ecological engineering. *Ecological Engineering*. Elsevier 45: 24–29. Available at: <https://www.sciencedirect.com/science/article/pii/S0925857412001188>: doi:10.1016/J.ECOLENG.2012.03.023.

Crevaschi M and Nicosia C (2012) Sub-Boreal aggradation along the Apennine margin of the Central Po Plain: geomorphological and geoarchaeological aspects. *Géomorphologie : relief, processus, environnement*. Groupe français de géomorphologie 18(2): 155–174. Available at: <http://journals.openedition.org/geomorphologie/9810>: doi:10.4000/geomorphologie.9810.

Davis BAS, Zanon M, Collins P, Mauri A, Bakker J, Barboni D, et al. (2013) The European Modern Pollen Database (EMPD) project. *Vegetation History and Archaeobotany*. Springer Berlin Heidelberg 22(6): 521–530. Available at: <http://link.springer.com/10.1007/s00334-012-0388-5>: doi:10.1007/s00334-012-0388-5.

Attema P and de Haas TCA (2005) Villas and farmsteads in the Pontine region between 300 BC and 300 AD: a landscape archaeological approach. In: Santillo Frizell B and Klynne A (eds) *Roman villas around the Urbs. Interaction with landscape and environment. Proceedings of a conference held at the Swedish Institute in Rome, September 17-18, 2004.* Available at: <http://www.svenska-institutet-rom.org/villa/>.

Dearing JA and Jones RT (2003) Coupling temporal and spatial dimensions of global sediment flux through lake and marine sediment records. *Global and Planetary Change*. Elsevier 39(1–2): 147–168. Available at: <https://www.sciencedirect.com/science/article/pii/S0921818103000225>: doi:10.1016/S0921-8181(03)00022-5.

de Vente J, Poesen J, Verstraeten G, Govers G, Vanmaercke M, Van Rompaey A, et al. (2014) Corrigendum to “Predicting soil erosion and sediment yield at regional scales: Where do we stand?” [Earth-Sci. Rev. 127 (2013) 16–29]. *Earth-Science Reviews* 133: 94. Available at: <http://www.sciencedirect.com/science/article/pii/S0012825214000506>: doi:<http://dx.doi.org/10.1016/j.earscirev.2014.03.002>.

De Vincenzo S, Domdey C, Hoelzmann P, Knitter D, Moede K, Müller M, et al. (2013) Zur Archäologie und Landschaftsentwicklung im Turano-Tal (Sabiner Berge), Italien. *eTopoi* 2(2012): 45–110. Available at: <https://www.topoi.org/publication/21315/>.

Devillers B, Excoffon P, Morhange C, Bonnet S and Bertoncello F (2007) Relative sea-level changes and coastal evolution at Forum Julii (Fréjus, Provence). *Comptes Rendus Geoscience*. Elsevier Masson 339(5): 329–336. Available at: <https://www.sciencedirect.com/science/article/pii/S1631071307000387>: doi:10.1016/J.CRTE.2007.03.004.

Diamond JM (2005) Collapse. *How societies choose to fail or succeed*.

Dotterweich M (2008) The history of soil erosion and fluvial deposits in small catchments of central Europe: Deciphering the long-term interaction between humans and the environment — A review. *Geomorphology*. Elsevier 101(1–2): 192–208. Available at: <https://www.sciencedirect.com/science/article/pii/S0169555X08002213>: doi:10.1016/J.GEOMORPH.2008.05.023.

Dotterweich M (2013) The history of human-induced soil erosion: Geomorphic legacies, early descriptions and research, and the development of soil conservation—A global synopsis. *Geomorphology*. Elsevier 201: 1–34. Available at: <https://www.sciencedirect.com/science/article/pii/S0169555X1300370X>: doi:10.1016/J.GEOMORPH.2013.07.021.

Dreibrodt S, Lubos C, Lomax J, Sipos G, Schroedter T and Nelle O (2014) Holocene landscape dynamics at the tell Arslantepe, Malatya, Turkey—Soil erosion, buried soils and settlement layers, slope and river activity in a middle Euphrates catchment. *The Holocene* 24(10): 1351–1368.

Dusar B, Verstraeten G, D’haen K, Bakker J, Kaptijn E and Waelkens M (2012) Sensitivity of the Eastern Mediterranean geomorphic system towards environmental change during the Late Holocene: a chronological perspective. *Journal of Quaternary Science* 27(4): 371–382. Available at: <http://doi.wiley.com/10.1002/jqs.1555>: doi:10.1002/jqs.1555.

Dusar B, Verstraeten G, Notebaert B and Bakker J (2011a) Earth-Science Reviews Holocene environmental change and its impact on sediment dynamics in the Eastern Mediterranean. *Earth Science Reviews*. Elsevier B.V. Available at: <http://dx.doi.org/10.1016/j.earscirev.2011.06.006>: doi:10.1016/j.earscirev.2011.06.006.

Dusar B, Verstraeten G, Notebaert B and Bakker J (2011b) Holocene environmental change and its impact on sediment dynamics in the Eastern Mediterranean. *Earth-Science Reviews*. Elsevier 108(3–4): 137–157. Available at: <https://www.sciencedirect.com/science/article/pii/S0012825211000882>: doi:10.1016/J.EARSCIREV.2011.06.006.

Ellis C and Brown AG (1999) Alluvial microfabrics, anisotropy of magnetic susceptibility and overbank processes. In: Brown AG and Quine TA (eds) *Fluvial Processes and Environmental Change*, Wiley, Chichester. Chichester: Wiley, 181–206.

Evershed RP, Payne S, Sherratt AG, Copley MS, Coolidge J, Nieuwenhuys O, et al. (2008) Earliest date for milk use in the Near East and southeastern Europe linked to cattle herding. 455(September): 31–34: doi:10.1038/nature07180.

Faust D and Wolf D (2017) Interpreting drivers of change in fluvial archives of the Western Mediterranean - A critical view. *Earth-Science Reviews*. Elsevier 174: 53–83. Available at: <https://www.sciencedirect.com/science/article/pii/S0012825217304798>:

doi:10.1016/J.EARSCIREV.2017.09.011.

Fletcher WJ and Zielhofer C (2013) Fragility of Western Mediterranean landscapes during Holocene Rapid Climate Changes. *Catena*. Elsevier B.V. 103: 16–29. Available at: <http://dx.doi.org/10.1016/j.catena.2011.05.001>; doi:10.1016/j.catena.2011.05.001.

Fryirs K (2013) (Dis)Connectivity in catchment sediment cascades: a fresh look at the sediment delivery problem. *Earth Surface Processes and Landforms*. Wiley-Blackwell 38(1): 30–46. Available at: <http://doi.wiley.com/10.1002/esp.3242>; doi:10.1002/esp.3242.

Fuchs M (2007a) An assessment of human versus climatic impacts on Holocene soil erosion in NE Peloponnese, Greece. *Quaternary Research* 67(3): 349–356; doi:10.1016/j.yqres.2006.11.008.

Fuchs M (2007b) An assessment of human versus climatic impacts on Holocene soil erosion in NE Peloponnese, Greece. *Quaternary Research*. Publisher: United States : Elsevier Science B.V., Amsterdam VOL 67; NU: 349–356.

Fyfe RM, Woodbridge J and Roberts CN (2018) Trajectories of change in Mediterranean Holocene vegetation through classification of pollen data. *Vegetation History and Archaeobotany*. Springer Berlin Heidelberg 27(2): 351–364. Available at: <http://link.springer.com/10.1007/s00334-017-0657-4>; doi:10.1007/s00334-017-0657-4.

Ghilardi M, Andrieu-Ponel V, Coileu M, Sotiropoulos P, Longo F, Amato V, et al. (2018) First evidence of a lake at Ancient Phaistos (Messara Plain, South Central Crete, Greece): reconstructing paleoenvironments and differentiating the roles of human land- use and paleoclimate from Minoan to Roman times. *The Holocene*.

Ghilardi M, Fouache E, Queyrel F, Syrides G, Vouvalidis K, Kunesch S, et al. (2008a) Human occupation and geomorphological evolution of the Thessaloniki Plain (Greece) since mid Holocene. *Journal of Archaeological Science* 35(1): 111–125. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0305440307000507>; doi:10.1016/j.jas.2007.02.017.

Ghilardi M, Genç A, Syrides G, Bloemendal J, Psomiadis D, Paraschou T, et al. (2010) Reconstruction of the landscape history around the remnant arch of the Klidhi Roman Bridge, Thessaloniki Plain, North Central Greece. *Journal of Archaeological Science* 37(1): 178–191; doi:10.1016/j.jas.2009.09.030.

Ghilardi M, Istria D, Curras A, Vacchi M, Contreras D, Vella C, et al. (2017) Reconstructing the landscape evolution and the human occupation of the Lower Sagone River (Western Corsica, France) from the Bronze Age to the Medieval period. *Journal of Archaeological Science: Reports* 12: 741–754. Available at: <http://www.sciencedirect.com/science/article/pii/S2352409X16303674>; doi:<https://doi.org/10.1016/j.jasrep.2016.07.009>.

Ghilardi M, Kunesch S, Styllas M and Fouache E (2008b) Reconstruction of Mid-Holocene sedimentary environments in the central part of the Thessaloniki Plain (Greece), based on microfaunal identification, magnetic susceptibility and grain-size analyses. *Geomorphology* 97(3–4): 617–630; doi:10.1016/j.geomorph.2007.09.007.

Giguët-Covex C, Arnaud F, Enters D, Poulenard J, Millet L, Francus P, et al. (2012) Frequency and intensity of high-altitude floods over the last 3.5ka in northwestern French Alps (Lake Anterne). *Quaternary Research*. University of Washington 77(1): 12–22. Available at: <http://dx.doi.org/10.1016/j.yqres.2011.11.003>; doi:10.1016/j.yqres.2011.11.003.

Giesecke, T. *et al.* (2014) 'Towards mapping the late Quaternary vegetation change of Europe', *Vegetation History and Archaeobotany*. doi: 10.1007/s00334-012-0390-y.

Gilman A, Thornes JB and Wise S (2015) *Land-use and prehistory in south-east Spain*. Routledge. Available at:

https://books.google.co.uk/books?id=DBIcBQAAQBAJ&dq=end+of+the+argaric+culture+spain&lr=&source=gbs_navlinks_s.

Giraudi C (2005) Late-Holocene alluvial events in the Central Apennines, Italy. *Holocene* 15(5): 768–773: doi:10.1191/0959683605hl850rr.

Giesecke T, Davis B, Brewer S, Finsinger W, Wolters S, Blaauw M, *et al.* (2014) Towards mapping the late Quaternary vegetation change of Europe. *Vegetation History and Archaeobotany*: doi:10.1007/s00334-012-0390-y.

Glais A, Lespez L, Davidson R, López-Sáez JA and Birée L (2016) 3000 ans de détritisme en Grèce du Nord : entre fluctuations climatiques et anthropisation des zones humides. *Géomorphologie : relief, processus, environnement*. Groupe français de géomorphologie 22(2): 187–208. Available at: <http://journals.openedition.org/geomorphologie/11380>: doi:10.4000/geomorphologie.11380.

Goudeau MLS, Reichart GJ, Wit JC, de Nooijer LJ, Grauel AL, Bernasconi SM, *et al.* (2015) Seasonality variations in the Central Mediterranean during climate change events in the Late Holocene. *PALAEOGEOGRAPHY PALAEOCLIMATOLOGY PALAEOECOLOGY* 418(0): 304–318: doi:10.1016/j.palaeo.2014.11.004.

Granger DE and Muzikar PF (2001) Dating sediment burial with in situ-produced cosmogenic nuclides: theory, techniques, and limitations. *Earth and Planetary Science Letters*. Elsevier 188(1–2): 269–281. Available at: <https://www.sciencedirect.com/science/article/pii/S0012821X01003090>: doi:10.1016/S0012-821X(01)00309-0.

Greenfield HJ (2010) The Secondary Products Revolution: the past, the present and the future. *World Archaeology* 42(1): 29–54. Available at: <http://www.tandfonline.com/doi/abs/10.1080/00438240903429722>: doi:10.1080/00438240903429722.

Grove AT and Rackham O (2001) *The Nature of Mediterranean Europe: An Ecological History*. Yale: Yale University Press.

Guttmann EB, Simpson I a. I a., Nielsen N, Dockrill SJSJ, Bull ID, Evershed RP, *et al.* (2013) Early Anthropogenic Soil Formation at Tofts Ness, Sanday, Orkney. *Anthropocene*. Paris: Elsevier B.V. 4(1): 151–153. Available at: <http://pubs.acs.org/doi/abs/10.1021/acs.est.5b00543><http://dx.doi.org/10.1016/j.ancene.2014.07.002><http://linkinghub.elsevier.com/retrieve/pii/S2213305414000265><http://www.sciencedirect.com/science/article/pii/S2213305413000106><http://dx.doi.org/10.1002/GEA>: doi:10.1002/GEA.

Harris W V (2013) *The Ancient Mediterranean Environment between Science and History*. *Columbia Studies in the Classical Tradition*. Brill. Available at: <https://books.google.co.uk/books?id=PfGBAAAAQBAJ>.

Hilgers, A., Murray, A.S., Schlaak, N. and Radtke, U. (2001) A Comparison of quartz OSL protocols using Lateglacial and Holocene dune sands from Brandenburg, Germany. *Quaternary Science Reviews* 20, 731–736.

Hoffmann T, Thorndycraft VR, Brown AG, Coulthard TJ, Damnati B, Kale VS, *et al.* (2010) Human impact

on fluvial regimes and sediment flux during the Holocene: Review and future research agenda. *Global and Planetary Change*. Elsevier 72(3): 87–98. Available at:
<https://www.sciencedirect.com/science/article/pii/S0921818110000974>:
 doi:10.1016/J.GLOPLACHA.2010.04.008.

Horden P and Purcell N (2000) *The corrupting sea: a study of Mediterranean history*. Oxford: Blackwell.

Houben P, Schmidt M, Mauz B, Stobbe A and Lang A (2013) Asynchronous Holocene colluvial and alluvial aggradation: A matter of hydrosedimentary connectivity. *The Holocene*. SAGE PublicationsSage UK: London, England 23(4): 544–555. Available at:
<http://journals.sagepub.com/doi/10.1177/0959683612463105>: doi:10.1177/0959683612463105.

Hughes JD (1994) *Pan's Travail: Environmental Problems of the Ancient Greeks and Romans*. Baltimore and London: The Johns Hopkins University Press: doi:file:///e:\JournalArt2\reviewhughes.pdf; http://penelope.uchicago.edu/Thayer/E/Roman/Texts/Strabo/4F*.html.

Hughes JD (2005) *The Mediterranean: an environmental history. Nature and Human Societies*. Santa Barbara: ABC-CLIO.

Joannin S, Magny M, Peyron O, Vanni re B and Galop D (2014) Climate and land-use change during the late Holocene at Lake Ledro (southern Alps, Italy). *Holocene* 24(5): 591–602: doi:10.1177/0959683614522311.

Jones AF, Macklin MG and Benito G (2015) Meta-analysis of Holocene fluvial sedimentary archives: A methodological primer. *CATENA* 130(Supplement C): 3–12: doi:<https://doi.org/10.1016/j.catena.2014.11.018>.

Kayan I (1999) Holocene stratigraphy and geomorphological evolution of the Aegean coastal plains of Anatolia. *Quaternary Science Reviews* 18(4–5): 541–548: doi:10.1016/S0277-3791(98)90095-6.

Krahtopoulou A and Frederick C (2008) The stratigraphic implications of long-term terrace agriculture in dynamic landscapes: Polycyclic terracing from Kythera Island, Greece. *Geoarchaeology* 23(4): 550–585. Available at: <http://doi.wiley.com/10.1002/gea.20231>: doi:10.1002/gea.20231.

Lambrianides K and Spencer N (1998) Regional studies in the Madra Cay delta: archaeology, environment and cultural history on the Aegean coast of Turkey from the Early Bronze Age to the Ottoman Empire'in R. Matthews. *Matthews (ed.), Ancient Anatolia. Fifty Years' Work by the British Institute of Archaeology at Ankara. Exeter* 207–223.

Lespez L (2003) Geomorphic responses to long-term land use changes in Eastern Macedonia (Greece). *Catena*. Elsevier 51(3–4): 181–208. Available at:
<https://www.sciencedirect.com/science/article/pii/S0341816202001649>: doi:10.1016/S0341-8162(02)00164-9.

Lespez L (2008) L' volution des paysages du N olithique   la p riode ottomane dans la plaine de Philippes. In: Koukouli-Chryssanthaki H and Treuil R (eds) *Dikili Tash, un village pr historique en Mac doine orientale 1*. Ath nes: Soci t  Arch ologique d'Ath nes et  cole Fran aise d'Ath nes, De Boccard Diffusion.

Lespez L, Dalongeville R, Pastre JF, Darmon F, Mathieu R and Poursoulis G (2003) Late-Middle-Holocene palaeo-environmental evolution and coastline changes of Malia (Crete). *The Mediterranean World Environment and History*. Elsevier 439–452.

Lespez L and Glais A (2017) Identifying the earliest Neolithic settlements in the South-Eastern Balkans: methodological considerations based on the recent geoarchaeological investigations at Dikili Tash (Greek Eastern Macedonia). (May 2015): 27–31. Available at: https://www.academia.edu/34804506/Identifying_the_earliest_Neolithic_settlements_in_the_South-Eastern_Balkans_methodological_considerations_based_on_the_recent_geoarchaeological_investigations_at_Dikili_Tash_Greek_Eastern_Macedonia_.

Leydet M (n.d.) *The European Pollen Database*. Available at: <http://www.europeanpollendatabase.net/>.

Loeb J, Goold GP and Hooper WD (1934) *The Loeb Classical Library. No. 283, : De Re Rustica. Cato & varro*. Harvard University Press. Available at: <https://books.google.co.uk/books?id=wDXAngEACAAJ>.

Lowdermilk WC (1953) *Conquest of the land through 7,000 years*. US Department Of Agriculture; Washington.

Machado MJ, Medialdea A, Calle M, Rico MT, Sánchez-Moya Y, Sopeña A, et al. (2017) Historical palaeohydrology and landscape resilience of a Mediterranean rambla (Castellón, NE Spain): Floods and people. *Quaternary Science Reviews*. Pergamon 171: 182–198. Available at: <https://www.sciencedirect.com/science/article/pii/S0277379117301798>: doi:10.1016/J.QUASCIREV.2017.07.014.

Macklin MG, Toonen WHJ, Woodward JC, Williams MAJ, Flaux C, Marriner N, et al. (2015) A new model of river dynamics, hydroclimatic change and human settlement in the Nile Valley derived from meta-analysis of the Holocene fluvial archive. *Quaternary Science Reviews* 130: 109–123: doi:<http://dx.doi.org/10.1016/j.quascirev.2015.09.024>.

Macklin MG, Tooth S, Brewer PA, Noble PL and Duller GAT (2010) Holocene flooding and river development in a Mediterranean steep-land catchment: The Anapodaris Gorge, south central Crete, Greece. *Global and Planetary Change*. Elsevier 70(1–4): 35–52. Available at: <http://dx.doi.org/10.1016/j.gloplacha.2009.11.006>: doi:10.1016/j.gloplacha.2009.11.006.

Magny M, Joannin S, Galop D, Vannière B, Haas JN, Bassetti M, et al. (2017) Holocene palaeohydrological changes in the northern Mediterranean borderlands as reflected by the lake-level record of lake ledro, northeastern Italy. *Quaternary Research* 77(3): 382–396. Available at: <http://www.sciencedirect.com/science/article/pii/S0033589412000063>: doi:10.1016/j.yqres.2012.01.005.

Magny M, Peyron O, Gauthier E, Rouèche Y, Bordon A, Billaud Y, et al. (2009) Quantitative reconstruction of climatic variations during the Bronze and early Iron ages based on pollen and lake-level data in the NW Alps, France. *Quaternary International* 200(1–2): 102–110. Available at: <http://www.sciencedirect.com/science/article/B6VGS-4S575VN->

Magny M, Vanniere B, Zanchetta G, Fouache E, Touchais G, Petrika L, et al. (2009) Possible complexity of the climatic event around 4300–3800 cal. BP in the central and western Mediterranean. *The Holocene* 19(6): 823–833. Available at: <http://hol.sagepub.com/cgi/doi/10.1177/0959683609337360>: doi:10.1177/0959683609337360.

2/2/be4628325c70cf3bb758a7649e16ee66.

Marriner N and Morhange C (2007) Geoscience of ancient Mediterranean harbours. *Earth Science Reviews* 80(3–4): 137–194. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0012825206001541>:

doi:10.1016/j.earscirev.2006.10.003.

Mayewski P a., Rohling EE, Curt Stager J, Karlén W, Maasch K a., David Meeker L, et al. (2004) Holocene climate variability. *Quaternary Research* 62(3): 243–255. Available at: <http://linkinghub.elsevier.com/retrieve/pii/S0033589404000870>: doi:10.1016/j.yqres.2004.07.001.

Mazier F, Galop D, Brun C and Buttler A (2006) Modern pollen assemblages from grazed vegetation in the western Pyrenees, France: a numerical tool for more precise reconstruction of past cultural landscapes. *The Holocene* 16(1): 91–103.

McCormick M, Büntgen U, Cane MA, Cook ER, Harper K, Huybers P, et al. (2012) Climate Change during and after the Roman Empire: Reconstructing the Past from Scientific and Historical Evidence. *JOURNAL OF INTERDISCIPLINARY HISTORY*. MIT Press 43(2): 169–220. Available at: http://dx.doi.org/10.1162/JINH_a_00379: doi:10.1162/JINH_a_00379.

Medialdea A, Thomsen KJ, Murray AS and Benito G (2014) Reliability of equivalent-dose determination and age-models in the OSL dating of historical and modern palaeoflood sediments. *Quaternary Geochronology*. Elsevier 22: 11–24. Available at: <https://www.sciencedirect.com/science/article/pii/S1871101414000132>: doi:10.1016/J.QUAGEO.2014.01.004.

Mensing SA, Tunno I, Sagnotti L, Florindo F, Noble P, Archer C, et al. (2015a) 2700 years of Mediterranean environmental change in central Italy: A synthesis of sedimentary and cultural records to interpret past impacts of climate on society. *Quaternary Science Reviews* 116: 72–94: doi:10.1016/j.quascirev.2015.03.022.

Mensing SA, Tunno I, Sagnotti L, Florindo F, Noble P, Cifani G, et al. (2015b) 2700 years of Mediterranean environmental change in central Italy : a synthesis of sedimentary and cultural records to interpret past impacts of climate on society. 116: 72–94: doi:10.1016/j.quascirev.2015.03.022.

Montgomery DR (2007) Soil erosion and agricultural sustainability. *Proceedings of the National Academy of Sciences of the United States of America* 104(33): 13268–72. Available at: <http://www.pnas.org/cgi/content/long/104/33/13268>: doi:10.1073/pnas.0611508104.

Moody J (2000) Holocene climate change in Crete: an archaeologist's view. In: Halstead P and Frederick C (eds) *Landscape and Landuse in postglacial Greece*. Sheffield: Sheffield Academic Press, 52–61.

Morhange C, Vella C, Provansal M, Hesnard M and Laborel J (1999) Human impact and natural characteristics of the ancient ports of Marseille and Fos, in Provence, southern France. In: Leveau P, Tremant F, Walsh K and Barker G (eds) *Environmental Reconstruction in Mediterranean Landscape Archaeology*. Oxford: Oxbow, 145–153.

Notebaert B and Berger J-F (2014a) Quantifying the anthropogenic forcing on soil erosion during the Iron Age and Roman Period in southeastern France. *Anthropocene*. Elsevier 8: 59–69. Available at: <https://www.sciencedirect.com/science/article/pii/S2213305415300035>: doi:10.1016/J.ANCENE.2015.05.004.

Notebaert B and Berger JF (2014b) Quantifying the anthropogenic forcing on soil erosion during the Iron Age and Roman Period in southeastern France. *Anthropocene*. Elsevier B.V. 8(2014): 59–69. Available at: <http://dx.doi.org/10.1016/j.ancene.2015.05.004>: doi:10.1016/j.ancene.2015.05.004.

Notebaert B, Berger JF and Brochier JL (2014) Characterization and quantification of Holocene colluvial

and alluvial sediments in the Valdaine Region (southern France). *Holocene*. SAGE PublicationsSage UK: London, England 24(10): 1320–1335. Available at: <http://journals.sagepub.com/doi/10.1177/0959683614540946>: doi:10.1177/0959683614540946.

Ortega R, Vanacker V, Sanjurjo-Sánchez J and Miralles I (2017) Human–Landscape Interactions during the Early and High Medieval Period in Central Spain Based on New Estimates of Sediment Yield from the Melque Agricultural Complex. *Geoarchaeology* 32(2): 177–188. Available at: <http://dx.doi.org/10.1002/gea.21582>: doi:10.1002/gea.21582.

Palmisano A, Bevan A and Shennan S (2017) Comparing archaeological proxies for long-term population patterns: An example from central Italy. *Journal of Archaeological Science*. Elsevier Ltd 87: 59–72. Available at: <https://doi.org/10.1016/j.jas.2017.10.001>: doi:10.1016/j.jas.2017.10.001.

Penington, B., Sturt, F., Wilson, P. And Brown, A.G. (2017) The fluvial evolution of the Holocene Nile Delta. *Quaternary Science Reviews* 170, 212–231.

Pérez-Lambán F, Peña-Monné JL, Fanlo-Loras J, Picazo-Millán J V, Badia-Villas D, Rubio-Fernández V, et al. (2014) Paleoenvironmental and geoarchaeological reconstruction from late Holocene slope records (Lower Huerva Valley, Ebro Basin, NE Spain). *Quaternary Research* 81(1): 1–14. Available at: <http://www.sciencedirect.com/science/article/pii/S0033589413001233>: doi:<https://doi.org/10.1016/j.yqres.2013.10.011>.

Potter, T. (1979) *The changing face of Southern Etruria*. Elek, London.

Pope RJJ, Wilkinson KN and Millington AC (2003) Human and climatic impact on late Quaternary deposition in the Sparta Basin Piedmont: Evidence from alluvial fan systems. *Geoarchaeology*. Wiley-Blackwell 18(7): 685–724. Available at: <http://doi.wiley.com/10.1002/gea.10089>: doi:10.1002/gea.10089.

Provansal M (1995) The role of climate in landscape morphogenesis since the Bronze Age in Provence, southeastern France. *The Holocene* 5(3): 348–353. Available at: <http://hol.sagepub.com/cgi/doi/10.1177/095968369500500311>: doi:10.1177/095968369500500311.

Psomiadis D, Ghilardi M, Demory F, Delanghe-Sabatier D, Bloemendal J and Yiu C (2014) Late Pleistocene to Mid-Holocene landscape reconstruction in the western part of the Thessaloniki Plain, Greece: evidence for environmental changes, and implications for human occupation. *Zeitschrift für Geomorphologie, Supplementary Issues* 58(2): 67–87. Available at: <http://openurl.ingenta.com/content/xref?genre=article&issn=1864-1687&volume=58&issue=2&page=67>: doi:10.1127/0372-8854/2013/S-00137.

Reale O and Dirmeyer P (2000) Modeling the effects of vegetation on Mediterranean climate during the Roman Classical Period.Part I: Climate history and model sensitivity. *Global and Planetary Change* 25(3–4): 163–184.

Rixhon G, Braucher R, Bourlès D, Siame L, Bovy B and Demoulin A (2011) Quaternary river incision in NE Ardennes (Belgium)—Insights from ¹⁰Be/²⁶Al dating of river terraces. *Quaternary Geochronology*. Elsevier 6(2): 273–284.

Roberts N, Brayshaw D, Kuzucuoglu C, Perez R and Sadori L (2011) The mid-Holocene climatic transition in the Mediterranean: Causes and consequences. *The Holocene* 21(1): 3–13. Available at: <http://hol.sagepub.com/cgi/doi/10.1177/0959683610388058>: doi:10.1177/0959683610388058.

Roberts, N., Allcock, S.L., Barnett, H., Mather, A., Eastwood, W.J., Jones, M., Primmer, N., Yiğitbaşıoğlu,

H., and Vanni re, B. 2019, in press. Cause-and-effect in Mediterranean erosion: The role of humans and climate upon Holocene sediment flux into a central Anatolian lake catchment. *Geomorphology*

Runnels C, Panagopoulou E, Murray P, Tsartsidou G, Allen S, Mullen K, et al. (2005) A Mesolithic Landscape in Greece : Testing a Site-Location Model in the Argolid at Kandia. 2: 259–285.

Salomon F, Goiran J-P, Bravard J-P, Arnaud P, Djerbi H, Kay S, et al. (2014) A harbour–canal at Portus: a geoarchaeological approach to the Canale Romano: Tiber delta, Italy. *Water History*. Springer Netherlands 6(1): 31–49. Available at: <http://link.springer.com/10.1007/s12685-014-0099-1>: doi:10.1007/s12685-014-0099-1.

Schulte L (2003) River-response and terrace aggradation in the Mediterranean Iberian Peninsula during historical times. *Palaeofloods, historical data and climatic variability: applications in flood risk assessment*. CSIC, Madrid 67–72.

Sevink J, van der Plicht J, Feiken H, van Leusen PM and Bakels CC (2013) The Holocene of the Agro Pontino graben: Recent advances in its palaeogeography, palaeoecology, and tephrostratigraphy. *Quaternary International* 303: 153–162: doi:10.1016/j.quaint.2013.01.006.

Sherratt S (1998) Sea Peoples” and the economic structure of the late second millennium in the Eastern Mediterranean. *Mediterranean Peoples in Transition: Thirteenth to Early Tenth Centuries BCE*. Israel Exploration Society Jerusalem 292–313.

Sherratt A (1981) Plough and pastoralism: aspects of the secondary products revolution. Cambridge University Press.

Silva PG, Bardaj  T, Calmel-Avila M, Luis J, Zazo C, Goy JL, et al. (2008) Transition from alluvial to fluvial systems in the Guadalent n Depression (SE Spain) during the Holocene: Lorca Fan versus Guadalent n River. *Geomorphology* 100(1–2): 140–153: doi:10.1016/j.geomorph.2007.10.023.

Simonneau A, Chapron E, Vanni re B, Wirth SB, Gilli A, Di Giovanni C, et al. (2013) Mass-movement and flood-induced deposits in Lake Ledro, southern Alps, Italy: Implications for Holocene palaeohydrology and natural hazards. *Climate of the Past* 9(2): 825–840: doi:10.5194/cp-9-825-2013.

Stanley, D.J., Warne, A.G., 1997. Holocene sea-level change and early human utilization of deltas.

Styllas MN and Ghilardi M (2017) Early- to mid-Holocene paleohydrology in northeast Mediterranean: The detrital record of Aliakmon River in Loudias Lake, Greece. *The Holocene*. SAGE PublicationsSage UK: London, England 27(10): 1487–1498. Available at: <http://journals.sagepub.com/doi/10.1177/0959683617693905>: doi:10.1177/0959683617693905.

Theodorakopoulou K and Bassiakos Y (2017) Geoarchaeological studies at the cemetery of ancient Kamara, assisted by optically stimulated luminescence (OSL) dating: Insights in the post-Roman hydrological record of Eastern Crete. *Journal of Archaeological Science: Reports*. Elsevier Ltd 12: 794–804. Available at: <http://dx.doi.org/10.1016/j.jasrep.2016.07.001>: doi:10.1016/j.jasrep.2016.07.001.

Thorndycraft VR and Benito G (2006) The Holocene fluvial chronology of Spain: evidence from a newly compiled radiocarbon database. *Quaternary Science Reviews* 25(3): 223–234: doi:<https://doi.org/10.1016/j.quascirev.2005.07.003>.

Tourloukis V and Karkanias P (2012) The Middle Pleistocene archaeological record of Greece and the role of the Aegean in hominin dispersals: new data and interpretations. *Quaternary Science Reviews*.

Pergamon 43: 1–15. Available at:

<https://www.sciencedirect.com/science/article/pii/S027737911200145X>:

doi:10.1016/J.QUASCIREV.2012.04.004.

van Andel TH, Runnels C and Pope KO (1986) Five thousand years of land use and abuse in the Southern Argolid, Greece. *Hesperia* 55: 103–128.

van Andel TH and Zangger E (1990) Land use and soil erosion in prehistoric and historical Greece. *Journal of Field Archaeology* 17: 379–396.

Van Loo M, Duser B, Verstraeten G, Renssen H, Notebaert B, D’Haen K, et al. (2017) Human induced soil erosion and the implications on crop yield in a small mountainous Mediterranean catchment (SW-Turkey). *CATENA*. Elsevier 149: 491–504. Available at:

<https://www.sciencedirect.com/science/article/pii/S0341816216303344>:

doi:10.1016/J.CATENA.2016.08.023.

Vanmaercke M, Poesen J, Govers G and Verstraeten G (2015) Quantifying human impacts on catchment sediment yield: A continental approach. *GLOBAL AND PLANETARY CHANGE* 130: 22–36: doi:<http://dx.doi.org/10.1016/j.gloplacha.2015.04.001>.

Vanniere B and Jouffroy-Bapicot I ((n.d.)) *Lake Kournas - unpublished data* .

Verstraeten G (2015) Human impact on sediment dynamics across a range of spatial and temporal scales..

Verstraeten G, Broothaerts N, Van Loo M, Notebaert B, D’Haen K, Duser B, et al. (2017) Variability in fluvial geomorphic response to anthropogenic disturbance. *Geomorphology*. Elsevier 294: 20–39. Available at:

<https://www.sciencedirect.com/science/article/pii/S0169555X16311400>:

doi:10.1016/J.GEOMORPH.2017.03.027.

Vita-Finzi C (1995) Solar history and paleohydrology during the last two millennia. *Geophysical Research Letters*. Wiley-Blackwell 22(6): 699–702. Available at: <http://doi.wiley.com/10.1029/95GL00055>: doi:10.1029/95GL00055.

Walsh KJ (2013) *The Archaeology of Mediterranean Landscapes. Human-Environment Interaction from the Neolithic to the Roman Period*. Cambridge University Press. Walsh K, Attema P and De Haas T (2014) The Pontine Marshes (Central Italy): A case study in wetland historical ecology. *Babesch* 89(1): doi:10.2143/BAB.89.0.3034668.

Weiberg E, Unkel I, Kouli K, Holmgren K, Avramidis P, Bonnier A, et al. (2016) The socio-environmental history of the Peloponnese during the Holocene: Towards an integrated understanding of the past. *Quaternary Science Reviews* 136: 40–65. Available at:

<http://www.sciencedirect.com/science/article/pii/S0277379115301633>:

doi:<http://dx.doi.org/10.1016/j.quascirev.2015.10.042>.

Weiberg, E. and Finné, M., (2013) Mind or Matter? People-Environment Interactions and the Demise of Early Helladic II Society in the Northeastern Peloponnese. *American Journal of Archaeology*. Archaeological Institute of America 117(1): 1–31. Available at:

<https://www.jstor.org/stable/10.3764/aja.117.1.0001>: doi:10.3764/aja.117.1.0001.

Weninger B, Clare L, Rohling E, Bar-Yosef O, Böhner U, Budja M, et al. (2009) The Impact of Rapid Climate Change on Prehistoric Societies during the Holocene in the Eastern Mediterranean. *Documenta Praehistorica* 36: 7–59. Available at: http://arheologija.ff.uni-lj.si/documenta/abstracts36/a36_2.html:

doi:10.4312/dp.36.2.

Weninger B and Harper T (2015) The geographic corridor for rapid climate change in Southeast Europe and Ukraine. *Neolithic and Copper Age between the Carpathians and the Aegean Sea. Chronologies and Technologies from the 6th to the 4th Millennium BC (= Archäologie in Eurasien 31)*. Bonn: Habelt 474–494.

Williams AN (2012) The use of summed radiocarbon probability distributions in archaeology: a review of methods. *Journal of Archaeological Science* 39(3): 578–589. Available at: <http://www.sciencedirect.com/science/article/pii/S0305440311002482>: doi:<http://dx.doi.org/10.1016/j.jas.2011.07.014>.

Wolf D, Faust D, Guttman EB, Simpson I a. I a., Nielsen N, Dockrill SJSJ, et al. (2015) Western Mediterranean environmental changes: Evidences from fluvial archives. *Quaternary International*. Paris: Nature Publishing Group, a division of Macmillan Publishers Limited. All Rights Reserved. 6(Supplement C): 211–225. Available at: <http://hol.sagepub.com/cgi/doi/10.1177/0959683607080515>: doi:<https://doi.org/10.1016/j.quascirev.2015.04.016>.

Woodbridge J, Roberts N and Fyfe R (n.d.) Pan-Mediterranean Holocene vegetation and land-cover dynamics from synthesised pollen data. *Journal of Biogeography*. *Journal of Biogeography*.

Zanchetta G, Regattieri E, Isola I, Drysdale RN, Bini M, Baneschi I, et al. (2016) The so-called “4.2 event” in the central mediterranean and its climatic teleconnections. *Alpine and Mediterranean Quaternary* 29(1).

Zolitschka B, Behre K-E and Schneider J (2003) Human and climatic impact on the environment as derived from colluvial, fluvial and lacustrine archives—examples from the Bronze Age to the Migration period, Germany. *Quaternary Science Reviews*. Pergamon 22(1): 81–100. Available at: <https://www.sciencedirect.com/science/article/pii/S0277379102001828>: doi:10.1016/S0277-3791(02)00182-8.




Figure 1. The archaeological chronological periods referred to in the text along with key climate phases, including Rapid Climate Change 'events'. 8.2 ev - arid and cool phase, possible starting at -8.6 BP; 4.2 ev - tripartite climatic oscillation centred on 4.2 BP; 3.2 ev - mixed cool and trend towards aridity (esp. Eastern Mediterranean); EH1 - early Holadic MC - Minoan (Crete); PP - Proto Roman to LP - late Roman (Crete); Thr - Thracian; CI - Classical Ice; HRB - Helmshtet-Roman-early Byzantine; RCO - Roman Climatic Optimum; DACP - Dark Age Cold Period; WWP - Wes. Medieval Period; LIA - Little Ice Age.




Figure 2. Grey shaded areas—study areas with demographic and palynological data. Numbers refer to sites with erosion records referred to in text: 1 – Nar Lake (Turkey); 2 – Sagalassos (Turkey); 3 – Aliakmon-Axios deltas (Nth Greece); 4 – Chiklis, 5 – Angitis River and Dikili Tash (Nth Greece); 6 – Philious Basin (Sth Greece); 7 – Lake Kournas (Crete); 8 – Messara Plain (Crete); 9 – Lake Ledro (Italy); 10 – Turano Basin, Fiora, Marta and Treia rivers and Lago Lungo (Italy); 11 – Rhone Valley (France), 12 – Taravo River (Corsica), Sagone River (Corsica); 13 – Lorca alluvial fan, Librilla ‘Rambla’ (Spain).




Figure 3. Southern Turkey: pollen indicator groups, archaeological site survey (grey)(from Woodbridge et al., 2019) and erosion data (Dusar et al., 2012; Roberts et al., 2019)




Figure 4. Northern Greece pollen indicator groups, archaeological site survey (grey) and erosion data (Weiberg et al., 2019; Glais et al., 2016; Lespez and Glais, 2017). Of particular interest is the Iron Age ('Thracian' phase of erosion) that broadly coincides with the increase in the Olive, Juglans, Castanea, and Vitis curve and the start of the increase in pastoralism.




Figure 5: Southern Greece (Peloponnese and Crete) pollen indicator groups, archaeological site survey (A- SPD of radiocarbon dates (light grey = all possible sites, dark grey definite sites), B - Peloponnese sites count). (1) OSL dates from erosion data Peloponnese, OSL dates from colluvium (Fuchs 2007) (2) Relative CDP plots of geochronological dates from floods and extreme aluvial event units of eastern Mediterranean rivers (Benito et al. 2015), (3) Flood-events, Lake Kournas, Crete (Vanniere et al. in prep).




Figure 6. Phases of coastal advance on Apuo-Versilian plain (NW Italy) (after Bini et al., 2013).




Figure 7. Italy (western central zone) a. pollen indicator groups, archaeological site indices (SPDs and raw site counts) and erosion; Stoddart et al 2019 – flood events from Turano area (Borrelli et al., 2014; De Vincenzo et al., 2013). B. palynology and sedimentary data from Lago Lungo (Mensing et al., 2015)




Figure 8. South of France, including mid-Rhone Valley, pollen indicator groups, demographic indices (raw counts of sites, aoristic sum of sites, randomized duration of sites, SPD of ^{14}C dates; after Berger et al., 2019). Flooding/erosion dynamics based on 234 ^{14}C dates from the Rhone Valley (Berger et al. in press




Figure 9. North-eastern and south-eastern Spanish study zones pollen indicators, grey – demography for southern area, and combined demography for southern and northern areas (Fyfe et al 2019). Cumulative probability density plots (CPD) of radiocarbon dates from floods and extreme fluvial event units of the Iberian Peninsula (Benito et al., 2015; Thorndycraft and Benito, 2006; Wolf et al., 2015).




Figure 10. The principal erosion/flood records referred to in this paper. From east to west (left to right): Sagalassos (Turkey), synthesis of Eastern Mediterranean floods and fluvial activity, flood events Turano area (Italy). Flooding/erosion dynamics Rhone Valley (France), floods and extreme fluvial event units of the Iberian Peninsula, SPD of OSL dates from across the Mediterranean (all dates – thick black line, Greece – blue, Spain – red). Final curve is the nMDS curve for the climate data (Finné et al. this volume). This is a summary of isotope data from across the Mediterranean. Note the general trend from wetter conditions in the earlier Holocene towards drier conditions. This pattern is positive for all regions apart from northern Iberia; it is especially clearly marked in the Levant, Anatolia, the Balkans and S Iberia/Maghreb. nMDS allows the (shared) scaling of the data (e.g. $\delta^{18}\text{O}$, $\delta^{13}\text{C}$, δD) on a plot that collapses, or reduces the complex variability in these data that have been derived from 44 different sites