

2011-06-20

# Foraminifera resist ocean acidification in the Wagner Basin under conditions similar to high CO<sub>2</sub> environments of the Cretaceous-Paleogene

Hall-Spencer, Jason

<http://hdl.handle.net/10026.1/1381>

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*



## **Foraminifera resist ocean acidification in the Wagner Basin under conditions similar to high CO<sub>2</sub> environments of the Cretaceous-Paleogene**

L. Pettit (1), M.B. Hart (1), A. Medina-Sánchez (2), C.W. Smart (1), P. Collins (3), R. Rodolfo-Metalpa (1), J.M. Hall-Spencer (1), and R.M. Prol Ledesma (2)

(1) University of Plymouth, Drake Circus, Plymouth PL4 8AA, United Kingdom (riccardo@rodolfo-metalpa.com), (2) Universidad Nacional Autónoma de México, Ciudad Universitaria, Delegación Coyoacán, 04510 México D.F., Mexico, (3) Benthic Ecology Unit Zoology, National University of Ireland, Galway, Ireland

In 2007 and 2010 gas vents along the Wagner Fault (northern Gulf of California, Mexico; Canet *et al.*, 2010) were investigated, recording their effects on the carbonate chemistry of the sea water and the surrounding benthos. Over 300 massive, but diffuse seafloor gas vents were discovered causing dramatic changes in the carbonate chemistry.

The  $pH_T$  decreased from 7.98 out of the fault to 7.55 near the most active vents where the lowest saturation states of aragonite ( $\Omega_a$ ) and calcite ( $\Omega_c$ ) were 0.95 and 1.47 respectively. Large areas of the fault region had pH 7.7-7.8 suggesting continuous diffuse gas emissions.

Foraminifera (unicellular protists) present in the top 2 cm of the sediment (both living and dead individuals) had a range of mainly calcareous taxa (including *Bolivina acuminata*, *B. acutula*, *Bulimina marginata*, *Nonionella basispinata* and *Pseudoparella brayana*). This is a normal composition for these water depths and it was striking in the lack of dissolution features, damaged tests and the generally good preservation of the tests even when viewed at high magnification in a scanning electron microscope. With no evidence of breakage caused by transportation, as it is assumed that this composition is representative in terms of numbers of individuals and taxa represented. In sediments of similar pH around the island of Ischia (Italy), the assemblage of foraminifera in waters of comparable pH is much reduced, but carbonate saturation is different (Dias *et al.*, 2010). The foraminifera of the Wagner Basin appear to be surviving in high CO<sub>2</sub> environments that are comparable to those that occurred during the Cretaceous–Paleogene “greenhouse” world where atmospheric  $pCO_2$  was very much higher but with calcareous foraminifera apparently thriving (Tyrell & Zeebe 2004; Zachos *et al.*, 2005)