

2019-05-20

# Integrating local ecological knowledge, citizen science and longterm historical data for endangered species conservation: Additional records of angel sharks (Chondrichthyes: Squatinidae) in the Mediterranean Sea

Giovos, I

<http://hdl.handle.net/10026.1/14160>

---

10.1002/aqc.3089

Aquatic Conservation: Marine and Freshwater Ecosystems

Wiley

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*



**Integrating local ecological knowledge, citizen science and long-term historical data for endangered species conservation: Additional records of angel sharks (Chondrichthyes: Squatinidae) in the Mediterranean Sea**

Journal:	<i>Aquatic Conservation: Marine and Freshwater Ecosystems</i>
Manuscript ID	AQC-18-0326.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	15-Jan-2019
Complete List of Authors:	Giovas, Ioannis; iSea, Environmental Organisation for the Preservation of the Aquatic Ecosystems, Stoilas, Vasilis-Orestis; iSea, Environmental Organisation for the Preservation of the Aquatic Ecosystems, Mabruk, Sara; Omar Al-Mokhtar University, Zoology Department, Faculty of Science, Doumpas, Nikolaos; iSea, Environmental Organisation for the Preservation of the Aquatic Ecosystems Marakis, Philippos; iSea, Environmental Organisation for the Preservation of the Aquatic Ecosystems Maximiadi, Mary; iSea, Environmental Organisation for the Preservation of the Aquatic Ecosystems Moutopoulos, Dimitrios; Technological Educational Institute of Mesolonghi, Department of Aquaculture and Fisheries Management Kleitou, Periklis ; Marine and Environmental Research (MER) Lab Ltd; iSea, Environmental Organisation for the Preservation of the Aquatic Ecosystems Keramidas, Ioannis; iSea, Environmental Organisation for the Preservation of the Aquatic Ecosystems Tiralongo, Francesco ; Ente Fauna Marina Mediterranea de Maddalena, Alessandro ; Universita degli Studi di Milano-Bicocca, Adjunct Professor of Vertebrate Zoology
Broad habitat type (mandatory) select 1-2:	coastal < Broad habitat type, ocean < Broad habitat type
General theme or application (mandatory) select 1-2:	biodiversity < General theme or application, endangered species < General theme or application, monitoring < General theme or application
Broad taxonomic group or category (mandatory, if relevant to paper) select 1-2:	fish < Broad taxonomic group or category
Impact category (mandatory, if relevant to paper) select 1-2:	fishing < Impact category

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



SCHOLARONE™  
Manuscripts

1  
2  
3 **Integrating local ecological knowledge, citizen science and long-term historical data for endangered**  
4 **species conservation: Additional records of angel sharks (Chondrichthyes: Squatinidae) in the**  
5 **Mediterranean Sea**  
6  
7  
8  
9

10  
11  
12 **Abstract**  
13

- 14 1. All three species of angel sharks (genus *Squatina*) inhabiting the Mediterranean Sea are listed  
15 as Critically Endangered in the IUCN Red List of Threatened Species due to overexploitation.  
16  
17 2. New records from Cyprus, Greece, Italy and Libya were collected from citizen-scientists  
18 integrated with local knowledge obtained using structured-interviews in the four countries.  
19  
20 3. Observations and reports together with an analysis of the reconstructed fisheries data resulted  
21 in the identification of areas of interest, a review of the illegal trade of the species and a debate  
22 about the credibility of fisheries data for assessing threatened and/or protected species  
23  
24 4. Unconventional sources of information, such as social media, were identified as important  
25 tools for monitoring rare and endangered marine wildlife.  
26  
27 5. This work will contribute to promoting international cooperation for advancing angel shark  
28 conservation in line with the Eastern Atlantic and Mediterranean angel shark Conservation  
29 Strategy.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 **Keywords:** Sawback angelshark, Smoothback angelshark, Mediterranean, unconventional sources,  
41 threatened species  
42  
43  
44  
45  
46  
47

48 **1. Introduction**  
49

50 About 16% of the 465 species of sharks are threatened with extinction globally (Bräutigam *et al.*, 2015;  
51 Dulvy *et al.*, 2014). The single genus *Squatina*, includes 24 species (Froese & Pauly, 2018), 11 of which  
52 are listed as Threatened in the IUCN Red List of Threatened Species (IUCN, 2018). Angel sharks are flat-  
53 bodied coastal species, with extremely broad pectoral fins, dorsally located eyes and spiracles. They  
54 are moderately sized (average length around 1-1.5 m) and distributed over a wide geographical range,  
55  
56  
57  
58  
59  
60

1  
2  
3 from temperate to tropical marine waters. The majority of the species are restricted to small areas,  
4  
5 inhabiting the continental shelf and upper slopes down to 500 m (Compagno, 1984; Compagno, Dando,  
6  
7 & Fowler, 2005; Last & White, 2008; Stelbrink, von Rintelen, Cliff, & Kriwet, 2010). Because of their life  
8  
9 characteristics (i.e. slow growth, low reproductive rate and demersal nature), and due to the  
10  
11 intensification of fisheries, angel sharks are now the second, after sawfishes, most threatened family  
12  
13 of elasmobranchs in the world (Pristidae) (Dulvy et al., 2014).  
14

15  
16 The Mediterranean Sea is an important habitat for cartilaginous fish (Bradai, Saidi, & Enajjar, 2018),  
17  
18 and hosts three angel shark species; the sawback angelshark (*Squatina aculeata* Cuvier, 1829), the  
19  
20 smoothback angelshark (*Squatina oculata* Bonaparte, 1840) and the angelshark (*Squatina squatina*  
21  
22 Linnaeus, 1758). The Mediterranean populations of all three species are listed as “Critically  
23  
24 Endangered” in the IUCN Red List of Threatened Species (IUCN, 2018), due to their steep decline and  
25  
26 local extinctions, as a result of the historical and current overexploitation by demersal fisheries and  
27  
28 primarily trawl fishing (De Maddalena, Baensch, & Heim, 2016; Gordon *et al.*, 2017; Miller, 2016; Nieto  
29  
30 *et al.*, 2015; Walker, Cavanagh, Ducrocq, & Fowler, 2005). Currently, their distribution in the basin  
31  
32 appears scattered with several local extinctions (Gordon *et al.*, 2017; Ferretti et al., 2016; Soldo &  
33  
34 Bariche, 2016) while observations are extremely limited and the species are commercially extinct  
35  
36 (Cavanagh and Gibson, 2007; Gordon *et al.*, 2017; angel shark Sightings Map, 2018).  
37  
38

39  
40 Official fishery catch data can potentially provide valuable information and an extended time series of  
41  
42 data, regarding population trends of primarily commercial and relatively abundant species, as angel  
43  
44 sharks in the past (Cavanagh et al., 2007). However, in many cases, catch data lack accuracy (Pauly &  
45  
46 Froese, 2012) because they do not include discarded, subsistence, recreational and non-reported  
47  
48 catches, all of which are referred to as Illegal, Unreported and Unregulated catches (IUU). In addition,  
49  
50 angel sharks, although not targeted nowadays due to their scarcity, it is possible that when they are  
51  
52 caught are, often deliberately or unintentionally misreported (Dulvy *et al.*, 2014) and recorded within  
53  
54 another group of elasmobranchs (e.g. guitarfish, rays, etc.), thus jeopardizing accurate data reporting  
55  
56 about these species.  
57  
58  
59  
60

1  
2  
3 The scarcity of the observations, the deficiencies in the poor monitoring of angel sharks by the official  
4 authorities and the low economic value of their catches make the use of non-conventional information  
5 such as Local Ecological Knowledge (LEK) (Stephenson *et al.*, 2016), social media and citizen science  
6 vital. These are frequently used as alternative sources of information when conventional data are not  
7 available (Moutopoulos, Dimitriou, Katselis & Koutsikopoulos, 2017); supplementing and validating  
8 scientific knowledge, thus empowering marine scientists and managers to improve conservation and  
9 policy (e.g. Giovos, Chatzisprou, Doumpas, Stoilas, & Moutopoulos, 2018; Johannes, Freeman, &  
10 Hamilton, 2000) even in the case of the extremely rare angel sharks in the Canary Islands and the  
11 Adriatic Sea (Meyers *et al.*, 2017 and Holcer & Lazar 2017, respectively).

12  
13  
14 This study presents additional records of the three Mediterranean angel shark species from Cyprus,  
15 Greece, Italy and Libya, collected in the context of three citizen science projects, and complemented  
16 with information obtained through targeted interviews and reconstructed fisheries data (Pauly &  
17 Zeller, 2016). Through the integration and the analysis of such a multidisciplinary information will: (a)  
18 facilitate the identification of potentially important regions for the angel shark populations in the  
19 Central and Eastern Mediterranean Sea, (b) initiate the discussion on the illegal trade of the species in  
20 the basin, (c) debate about the credibility of fisheries data for threatened and/or protected species,  
21 and (d) highlight the importance to utilize unconventional sources, such as social media, for monitoring  
22 rare marine wildlife. It also aims to promote international cooperation for advancing angel shark  
23 conservation in the Mediterranean Sea.

## 24 25 26 **2. Materials and Methods**

27  
28 In the context of this study, a variety of sources were used to collect data about occurrences of angel  
29 sharks from four Mediterranean countries; namely Italy, Libya, Cyprus and Greece. Citizen science  
30 reports accompanied by photographic evidence and data on the historical and current occurrence of  
31 the species were collected, based on targeted interviews. The findings were complemented with an  
32 analysis of the reconstructed fisheries data found in the Sea Around Us catch database.

### 33 34 35 **2.1. Citizen Science Reports (CSR)**

1  
2  
3 CSR reports were collected in the context of three citizen science projects, utilizing to a large extent  
4 social media for broadcasting a call for reports. The projects were: “Sharks and Rays in Greece and  
5 Cyprus” by [iSea](#), taking place in Greece and Cyprus as part of the [Mediterranean Elasmobranch Citizens](#)  
6 [Observation \(M.E.C.O.\)](#) project, “AlienFish – alien and rare fish species” by [ENTE Fauna Marina](#)  
7 [Mediterranean](#), taking place in Italy (Tiralongo, Messina, Coco, & Lombardo, 2018), and “Marine  
8 Biology Libya” by [Marine Biology Libya](#), taking place in Libya. These are long term projects that  
9 between 1/1/2018-1/5/2018 posted an open call on their social media platforms requesting reports  
10 and pictures of angel sharks. People reporting sightings were further requested to provide  
11 photographic evidence of the observed specimens and information on the specimens’ size (total  
12 length) and/or weight, depth of the observation (when applicable), date of the observation, the exact  
13 location (if possible, with coordinates), and the number of observed individuals. All pictures reported,  
14 were checked for their authenticity and originality using the automatic image recognition tool of  
15 Google. All original images were recorded in a single Excel spreadsheet and photo-identified to the  
16 lowest possible taxonomic level. Thus, a verified citizen science model was utilized, in which  
17 observations were checked by experts (Gardiner *et al.*, 2012).

## 2.2. Targeted Interviews

18  
19 LEK of marine observers was used to complement CSR and collect additional information on the  
20 historical occurrence of angel sharks in the four countries. Credibility of interviewees is a known  
21 problem of LEK data that can compromise the quality of the information (Davis & Wagner, 2003). In  
22 the above-mentioned projects even though a large number of people participate (approximately  
23 5000), only a small fraction of them were selected based on the following criteria: (i) regularly report  
24 observations of elasmobranch species to the projects, (ii) display high success in self-identifying the  
25 elasmobranch species they report before the expert identification and (iii) appear to be highly  
26 motivated in helping and participating. Each criterion was scored either 0 or 1 based on the three  
27 criteria. Only those participants who scored 3 were included in the study independently of the spatial  
28 coverage. The structured questionnaire used during the interviews, had a skip-logic structure,  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 providing different options according to whether the respondent observed an angel shark species or  
4  
5 not (See Appendices). The aim was to retrieve **current observations (CO)** and **historical observations**  
6  
7 **(HO)** of the species. All interviews were conducted by three independent researchers, one from each  
8  
9 project (Cyprus-Greece, Italy and Libya) in situ or via skype. Before conducting the interviews, the  
10  
11 researchers were instructed to present the questionnaire in the same way.  
12  
13

### 14 **2.3. Fisheries data**

15  
16 The reconstructed Sea Around Us catch data, available at [www.seaaroundus.org](http://www.seaaroundus.org), (Pauly & Zeller,  
17  
18 2016), were used. Data were organized by fishing country and fishing sector, using the data series  
19  
20 between 1950-2014. Reconstructed catches combine official reported landings from the Food and  
21  
22 Agriculture Organization of the United Nations (FAO) (FishStat Plus, 2018:  
23  
24 <http://www.fao.org/fishery/statistics/en>) and reconstructed estimates provided by the SeaAroundUs  
25  
26 database for each studied country. The latter includes an unreported portion of the catches that is not  
27  
28 properly covered by the official national statistics, such as artisanal catches, discards, recreational and  
29  
30 subsistence fisheries that are derived from government sources, independent studies and surveys,  
31  
32 local experts and the grey literature, and followed the general catch reconstruction approach outlined  
33  
34 in Zeller & Pauly (2006). Hence, the use of such reconstructed data enhance further the integrated  
35  
36 approach to vulnerable species that are not recorded by official authorities. Reports from 11  
37  
38 Mediterranean areas of southern Europe, Northern Africa and East Asia were analysed for the  
39  
40 aforementioned period; Cyprus (Ulman *et al.*, 2015), Egypt (Mahmoud, Teh, Khalfallah, & Pauly, 2015),  
41  
42 Israel (Edelist *et al.*, 2013), Syria (Ulman, Saad, Zylich, Pauly, & Zeller, 2015b), Malta (Khalfallah,  
43  
44 Dimech, Ulman, Zeller, & Pauly, 2017), Tunisia (Halouani, Lasram, Khalfallah, Zeller, & Pauly, 2015),  
45  
46 Turkey (Marmara and Mediterranean Seas: Ulman *et al.*, 2013), Morocco (Mediterranean Sea;  
47  
48 Belhabib, Harper, Zeller, & Pauly, 2013), Algeria (Belhabib, Pauly, Harper, & Zeller, 2013), Italy (Sicily)  
49  
50 (Piroddi *et al.*, 2015) and Libya (Khalfallah, Belhabib, Zeller, & Pauly, 2015).  
51  
52  
53  
54  
55

## 56 **3. Results**

### 57 **3.1. Citizen Science Reports**



1  
2  
3 Fifteen CSR angel shark records from the four countries were obtained during this study; 14 of which  
4 were accompanied with photographic evidences (Table1; Figures 1 and 2). The only observation that  
5 was not accompanied by photographic evidence (No 10; Table 1; Figures 1 and 2) was recorded  
6 because it was reported by the same person who reported three other records of angel sharks (No 9,  
7 11 and 13 in Table1; Figures 1 and 2). Two observations could not be identified down to species level  
8 due to the poor quality of the picture. The most CSR were from Greek waters ( $\approx 43\%$ ; N=6), followed  
9 by Libya ( $\approx 36\%$ ; N=5), Italy ( $\approx 14\%$ ; N=2) and Cyprus ( $\approx 7\%$ ; N=1). Half of the specimens were identified  
10 as *S. squatina* (50%; N=7), and the rest were identified as either *S. aculeata* ( $\approx 21\%$ ; N=3) or *S. oculata*  
11 ( $\approx 14\%$ ; N=2). Unfortunately, in most cases, the individuals were already dead when reported to the  
12 projects.

### 23 **3.2. Targeted Interviews**

24  
25  
26 Thirty-nine targeted interviews were conducted with 19 ( $\approx 49\%$ ) taking place in Greece, followed by  
27 Italy ( $\approx 33\%$ ; N=13), Cyprus ( $\approx 10\%$ ; N=4) and Libya ( $\approx 8\%$ ; N=3). Almost 1/3 of the respondents were  
28 professional fishers ( $\approx 33\%$ ; N=13), followed by recreational fishers ( $\approx 31\%$ ; N=12), divers ( $\approx 28\%$ ; N=11),  
29 biologists ( $\approx 2.5\%$ ; N=1), fishmongers ( $\approx 2.5\%$ ; N=1) and other sea users ( $\approx 2.5\%$ ; N=1). Only 10  
30 interviewees reported CO of angel sharks in their areas, the majority from Italy (Table 2; Table S1),  
31 where six professional fishers reported an area (N-W off the coast of Trapani, 38°24,635 N; 11°34,270  
32 E) that angel sharks are considered relatively common, mostly caught by bottom trawlers. The other  
33 four CO were from Greece (N=2; Alexandroupoli and Kos Island) and Libya (N=2; Gulf of Sirte: Ras Lanuf  
34 and Qaminis) (Figure 1; Table S1). All who reported CO from Italy and Libya stated that they observe  
35 angel sharks frequently in their areas (N-W off the coast of Trapani and Gulf of Sirte respectively).  
36 Interestingly, six interviewees from Greece reported HO of angel sharks (Table 2; Figure 1). In Italy,  
37 professional fishers from Sicily and Calabria stated that catches of *Squatina* spp. individuals were  
38 relatively common off the south-eastern coasts of Sicily (Ionian Sea) 30 years ago, when they were  
39 caught with trammel nets and gill nets (140 mm of mesh size), and off the west coast of Calabria 20  
40 years ago.

1  
2  
3 years ago, when they were usually caught with longlines. Historical evidence on angel sharks (Figure  
4  
5 1) also overlay with the CSR observations and LEK reports.  
6

### 7 **3.3. Fisheries Data**

8  
9  
10 In four out of the 11 studied countries (12 areas) catches of angel sharks were recorded in only one or  
11  
12 two years out of the 64 years (1950 -2014) analysed (i.e. Morocco, Algeria, Italy (Sicily) and Libya; for  
13  
14 these countries separate analyses on annual catches were not included herein. In contrast, only Turkey  
15  
16 (for catches derived both from Marmara and Mediterranean Seas) had catch records covering the  
17  
18 entire study period 1950-2014 (Figure 3), followed by Malta (records for 59 years), Egypt (records for  
19  
20 57 years) and Syria (records for 54 years) (Figure 3). Turkey accounted for almost 80% of the mean  
21  
22 (1964-2014) annual reconstructed angel shark catches throughout the Mediterranean basin (Table 2),  
23  
24 with Syria, Tunisia and Egypt cumulatively contributing 20.4% and the remaining countries contributing  
25  
26 less than 1% (Table 2). In general, angel sharks represent a very small portion of each country's  
27  
28 reported catches (less than 1% in all cases) (%' in Table 2).  
29  
30

31  
32 The annual landing trends of angel shark catch per country only exhibited a long-term increasing trend  
33  
34 in Egypt, whereas a long-term decreasing trend was only exhibited for catches reported by Turkey. It  
35  
36 is worth noting that a sudden increase of angel shark was reported from Syria, from around 25t before  
37  
38 2000 to more than 100t during 2002-2006. Likewise, for the catches reported from Tunisia, angel shark  
39  
40 records appeared only after 1995, and since 2014 have fluctuated around 25 t/year (Figure 3). With  
41  
42 respect to the combined country reconstructed catch data, a declining trend was observed during  
43  
44 1963-1973 followed by an upward trend between 1973-2014, mostly due to the increased catches  
45  
46 reported from Syria and Turkey (Figure 4). More than half of the total Mediterranean reconstructed  
47  
48 angel shark catches were caught by trawlers (59.9%), mostly originating from Turkey (90% and 60% in  
49  
50 Mediterranean and Marmara Sea, respectively) and Libya (70%), whereas small-scale fisheries  
51  
52 contributed 39.7% that were mostly derived from Tunisia, Malta and Egypt (collectively more than  
53  
54 90%). Longlines represented a very small portion of the total catches (0.4%), mainly due to their  
55  
56  
57  
58  
59  
60

1  
2  
3 exclusive use in Algeria (100%), and to a lesser extent in Libya and Tunisia (less than 10% in both  
4  
5 countries).

### 6 7 **Discussion**

8  
9  
10 These additional records of all three angel shark species found in the Mediterranean Sea were obtained  
11  
12 and compared with fisheries-related information derived from LEK through in-depth targeted  
13  
14 interviews and complemented with reconstructed catches from official reported data and historical  
15  
16 information. All three species are enlisted in the Annex II (list of endangered or threatened species) of  
17  
18 the SPA/BD Protocol, which, based on Recommendation GFCM/36/2012/1, cannot be retained on  
19  
20 board, trans-shipped, landed, transferred, stored, sold or displayed or offered for sale, and must be  
21  
22 released unharmed and alive to the extent possible. In addition, *Squatina squatina* is included in  
23  
24 Appendix I and II (2017) of the Convention on the Conservation of Migratory Species of Wild Animals  
25  
26 (Appendix I-Endangered migratory species/ Appendix II-Migratory species conserved through  
27  
28 Agreements). Law enforcement is poor in all the Mediterranean countries (Beddington, Agnew, &  
29  
30 Clark, 2007) and given the fact that in most of them elasmobranchs were landed in aggregated  
31  
32 categories, misreporting, mislabeling and illegal trade is potentially taking place in the basin. However,  
33  
34 no records of illegal trade or an illegal fishery for angel sharks have been reported in the Greek Seas  
35  
36 during the last 15 years, based on the official fisheries infringement data derived from the coast guard  
37  
38 authorities of the Ministry of Mercantile Marine (period of 1999-2013: Moutopoulos, Prodromitis,  
39  
40 Mantzouni, & Koutsikopoulos, 2016).

41  
42  
43 Turkey is the only Mediterranean country with continuous landings that either implies a targeted  
44  
45 fishery (Kabasakal & Kabasakal, 2014), or that it is the only country reporting angel shark landings  
46  
47 accurately. In this context, Turkey is the only country for which discard quantities of angel sharks,  
48  
49 especially from trawls, have been also included in the reported catches with a discard/catch ratio  
50  
51 estimated of 3.4:1. This is a very high estimate when compared with the corresponding estimates from  
52  
53 the adjacent Greek fisheries (for the Aegean Sea: 0.353:1; Machias *et al.*, 2001). On the other hand,  
54  
55 Greece is the only country with no reported angel shark landings, according to the taxonomic  
56  
57  
58  
59  
60

1  
2  
3 disaggregation reported by HELSTAT (1967-2017) (Moutopoulos and Koutsikopoulos, 2014), which is  
4 probably due to the aggregated landing categories (6 elasmobranch landing categories).  
5  
6

7 Angel sharks are not currently target species in the Mediterranean due to their scarcity and normally  
8 are caught as a by catch. In both cases (i.e. targeted fishery or incidental catch) fraudulent or erroneous  
9  
10 labelling is possibly taking place in all Mediterranean countries, because of four main reasons: (i)  
11 aggregated elasmobranch landing categories, (ii) poor training of the monitoring authority staff (iii) the  
12 low awareness of the fishing communities about the legal and the conservation status of these species  
13 and (iv) spiritual reason (e.g. in Turkey angel shark meat is considered to help fighting cancer) . Illegal  
14 trade in elasmobranchs has been found in several markets around the globe (Feitosa et al., 2018) and  
15 in some Mediterranean countries (Barbuto *et al.*, 2010; Arculeo, 2015; Pazartzi et al., 2019) while  
16 Vasconcellos Bunholi *et al.* (2018) found illegal trade of angel sharks in Brazil. Angel shark landings are  
17 might aggregate with the landings of other batoid species, such as *Raja* spp. and guitarfish, as a result  
18 of misidentifications, but lately also because of the intentional misreporting due to the legal  
19 framework that protects the species. Misreporting is also helped by the morphological characteristics  
20 of these species, in this case flat body shape, that can be easily mistaken or sold as batoids once  
21 skinned. In Greece, for example, angel sharks used to be a very popular dish in the past, named “Rina”  
22 which is the common name of the species in Greek; it is still common to “Rina” in local markets and  
23 restaurants however it refers to *Dasyatis* spp. and *Raja* spp. species. Yet, fishmongers, retailers, chefs  
24 and restaurant owners, prefer to sell dishes of *Dasyatis* spp. and *Raja* spp. as “Rina” that is well known  
25 and more expensive, thus increasing their profitability (pers. obs.). In Libya, commercial fishers  
26 normally skin angel sharks and sell them as “Kulb baher” which means sea dog which is the name used  
27 for selling almost all shark species (Pers. Comm. with Sara A. A. Almabruk). In Italy, this species is very  
28 rare and usually when it is caught is eaten by fishers (Pers. Comm. with Francesco Tiralongo) a common  
29 pattern for elasmobranchs globally (Begossi, 2006), which makes impossible to estimate the total  
30 elasmobranch fishery as self-consumption is normally not included.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 In situ observations of different angel shark species in Cyclades and Dodecanese Islands confirm the  
4  
5 current presence of all three angel shark species in Greek waters. It is notable that the three  
6  
7 observations of *S. aculeata* reported in this study are the first in the Aegean Sea in the last 10 years  
8  
9 (Soldo & Bariche, 2016). The records from South Cyclades (CSR No 9, 10, 11, 13), Rhodes Island (CSR  
10  
11 No 2, 15) and the CO and HO off the coast of Rhodes and Alexandroupoli indicate interesting areas for  
12  
13 the species within the Greek waters. The statements of the Italian and Libyan fishers about relatively  
14  
15 frequent angel shark captures and observation in the Tyrrhenian Sea and the Gulf of Sirte, respectively,  
16  
17 could additionally indicate areas of interest for further actions in fisheries management and the  
18  
19 decision-making process. In the Tyrrhenian Sea, although fishers did not clearly indicate which angel  
20  
21 shark species are present, species of the genus *Squatina* are historically well-known and reported as  
22  
23 relatively common in catches with bottom trawl, as supported by the reconstructed catch data from  
24  
25 the Sicilian waters. For Cyprus, Hadjichristophoru (2006) mentioned that these species are occasionally  
26  
27 found in Cypriot waters. However, their records are almost completely absent from the reconstructed  
28  
29 fishery catch data (Ulman et al. 2015a). Nevertheless, it should be noted that close to the island is the  
30  
31 Iskenderun Bay, where recent records of juvenile *S. aculeata* were reported, suggesting a reproduction  
32  
33 or a nursery ground (Basusta, 2016).  
34  
35  
36  
37  
38

39 This work, additionally, provides evidence of the important role that citizen science, social media and  
40  
41 LEK can play in data gathering but also in the conservation for rare and endangered species. The value  
42  
43 of citizen science and social media for data gathering has already been proven for angel sharks (Holcer  
44  
45 & Lazar, 2017; Meyers *et al.*, 2017), as well as for other elasmobranch species in the Mediterranean  
46  
47 Sea (Giovos *et al.*, 2018), the same is true for LEK (Barash, Pickholtz, Pickholtz, Blaustein, & Rilov, 2018;  
48  
49 Coll *et al.*, 2014; Gonzalvo, Giovos, & Moutopoulos, 2015; Fortibuoni, Borme, Franceschini, Giovanardi,  
50  
51 & Raicevich, 2016). In this work, citizen science data accompanied with the current knowledge and the  
52  
53 official fisheries catch reports provided important information about an extremely threatened family  
54  
55 of species. However, as in any other data collection approach, the information obtained through social  
56  
57 media, CSR and LEK might include biases and uncertainty (e.g. Davis & Wagner, 2003; Katsanevakis &  
58  
59  
60

1  
2  
3 Moustakas 2018; Thurstan, Buckley, Ortiz & Pandolfi, 2016), which must be taken into account when  
4  
5 policy and conservation measures are designed. It is important to further explore in a larger, massive  
6  
7 spatial scale (e.g. the whole Mediterranean with the participation of more projects, organizations and  
8  
9 scientists) our understanding about angel sharks and other threatened species distribution in the basin  
10  
11 (Tulloch *et al.*, 2018), by designing cooperative communication campaigns asking for information  
12  
13 about such species. These efforts work towards two directions, improving data availability and  
14  
15 increasing public awareness, advancing conservation.  
16  
17

18  
19 In 2017, the Eastern Atlantic and Mediterranean Angel Shark Conservation Strategy was presented  
20  
21 (Gordon *et al.*, 2017), proposing a first conservation plan for angel sharks specifically for the  
22  
23 Mediterranean Sea, including the enhancement of our understanding of the species distribution, the  
24  
25 quantification of the incidental catches, and the enforcement of the existing management measures  
26  
27 (Gordon *et al.*, 2017). In this context, the Angel Shark Conservation Network  
28  
29 (<https://angelsharknetwork.com/>) was developed for delivering the objectives laid out in the action  
30  
31 plan and to receive updates on angel shark conservation news. This network will act as the umbrella  
32  
33 for an international effort in the Mediterranean Sea, fostering international cooperation,  
34  
35 strengthening our knowledge and influence conservation policy for the threatened angel sharks.  
36  
37 Especially in the field of the fish resources, the management of which is performed at a European scale,  
38  
39 the development of a database concerning, apart from fisheries information, socio-economic aspects  
40  
41 of fisheries will enhancing fisheries' monitoring and contribute to the definition of efficient managerial  
42  
43 measures at a regional level. The present study aims to contribute towards this effort by providing  
44  
45 additional data about angel shark occurrences in the basin, along with other recent studies (Fortibuoni  
46  
47 *et al.*, 2016; Holcer & Lazar, 2017). Further research effort, which could be carried out to reinforce our  
48  
49 findings, should be directed towards the incorporation in the official monitoring scheme of the  
50  
51 utilization of LEK in the "hot-spot" areas for angel sharks in a broader scale campaign targeting the  
52  
53 whole Mediterranean basin with the participation entities from every country, in order to explore  
54  
55 additional areas of interest. Field research should include underwater visual census surveys and/or  
56  
57  
58  
59  
60

1  
2  
3 experimental fishing depending the bathymetry of the locations. An awareness campaign is also of  
4  
5 imperative importance, for educating fishers on reporting and safely releasing angel sharks.

6  
7 Angel sharks are still fished and potentially consumed in the Mediterranean Sea despite the strict and  
8  
9 prohibitive legislation that applies in all countries. This must urgently be brought to the attention of  
10  
11 the national authorities, while NGOs and other interested parties should work more actively towards  
12  
13 the education of the public and, primarily, of professional and recreational fishers.  
14  
15

## 16 **Acknowledgements**

17  
18 The authors are grateful to all the participants of the citizen science projects, the interviewees, the  
19  
20 following fishers for sharing their information and experience: Giovanni Lo Coco, Salvo Marino, Alfonso  
21  
22 Barone, Gabriele De Leonardo, Stelios F., Lefteris Pagonis and Stamatis Kampouris and the diving  
23  
24 centers Kalymnos Diving Center, Kos Divers, Cretan Divers, and Samos Divers. Last, we would like to  
25  
26 thank Dr. Eric Glenn Haenni from Franciscan University of Steubenville, Ohio, USA for his valuable  
27  
28 comments and edits and the Editor and the two anonymous reviewers that contributed significantly  
29  
30 and improved the quality of our article.  
31  
32  
33

## 34 **References**

- 35  
36 Angel Shark Sightings Map. 2018. <http://sites.zsl.org/angelsharks/> [31 July 2018]  
37  
38  
39 Barash, A., Pickholtz, R., Pickholtz, E., Blaustein, L., & Rilov, G. (2018). Seasonal aggregations of  
40  
41 sharks near coastal power plants in Israel: an emerging phenomenon. *Marine Ecology Progress Series*,  
42  
43 590, 145-154.  
44  
45 Barbuto, M., Galimberti, A., Ferri, E., Labra, M., Malandra, R., Galli, P., Casiraghi, M. (2010). DNA  
46  
47 barcoding reveals fraudulent substitutions in shark seafood products: the Italian case of “palombo”  
48  
49 (Mustelus spp.). *Food Research International*, 43, 376-381.  
50  
51  
52 Beddington, J. R., Agnew, D. J., & Clark, C. W. (2007). Current problems in the management of  
53  
54 marine fisheries. *Science*, 316, 1713–1716.  
55  
56  
57 Begossi, A. (2006). Temporal stability in fishing spots: conservation and co-management in  
58  
59 Brazilian artisanal coastal fisheries. *Ecology and Society*, 11(1), 5.  
60

1  
2  
3 Belhabib, D., Harper, S., Zeller, D., & Pauly, D. (2013a). Reconstruction of marine culate catches  
4 from Morocco (north, central and south), 1950-2010. Pp 23-40. In: Belhabib, D., Zeller, D., Harper, S.,  
5 and Pauly, D. (eds.), Marine fisheries catches in West Africa, 1950-2010, part I. Fisheries Centre  
6  
7  
8  
9 Research Reports 20 (3). Fisheries Centre, University of British Columbia, Canada [ISSN 1198-6727].

10  
11  
12 Belhabib, D., Pauly, D., Harper, S., & Zeller, D. (2013b). Reconstruction of marine fisheries  
13 catches for Algeria, 1950-2010. Pp 1-22. In: Belhabib, D., Zeller, D., Harper, S. and Pauly, D. (eds.),  
14 Marine culate catches in West Africa, 1950-2010, part I. Fisheries Centre Research Reports 20 (3).  
15  
16  
17  
18 Fisheries Centre, University of British Columbia, Canada [ISSN 1198-6727].

19  
20  
21 Bradai, M. N., Saidi, B., & Enajjar, S. (2018). Overview on Mediterranean Shark's Fisheries: Impact  
22 on the Biodiversity. In: Turkoglu, M., Önal, U., İsmen, A., (eds), *Marine Ecology*, pp: 211-230.

23  
24  
25 Bräutigam, A., Callow, M., Campbell, I. R., Camhi, M.D., Cornish, A. S., Dulvy, N. K., ... Welch, D.  
26  
27  
28 J. (2015). Global Priorities for Conserving Sharks and Rays: A 2015–2025 Strategy.

29  
30 Cavanagh, R. D., & Gibson, C.. (2007). Overview of the Conservation Status of Cartilaginous  
31 Fishes (Chondrichthyans) in the Mediterranean Sea. IUCN. 10.2305/IUCN.CH.2007.MRA.3.en.

32  
33  
34 Coll, M., Carreras, M., Ciércoles, C., Cornax, M. J., Gorelli, G., Morote, E., & Sáez, R. (2014).  
35 Assessing fishing and marine biodiversity changes using fishers' perceptions: the Spanish  
36  
37  
38 Mediterranean and Gulf of Cadiz case study. *PLoS ONE*, 9(1), e85670.

39  
40  
41 Compagno, L., Dando, M., & Fowler, S. (2005). A field guide to the sharks of the world. Harper  
42  
43  
44 Collins Publishers, London, 496pp.

45  
46  
47 Compagno, L.J.V. (1984) FAO Species Catalogue. Vol. 4. Sharks of the world. An annotated and  
48 illustrated catalogue of shark species known to date. Part 1 – Hexanchiformes to Lamniformes. FAO  
49  
50  
51 Fish. Synop. 125(4/1):1-249. Rome, FAO.

52  
53  
54 Davis, A., & Wagner, J.R. (2003) Who knows? On the importance of identifying “experts” when  
55 researching local ecological knowledge. *Human Ecology*, 31, 463-489.

56  
57  
58 De Maddalena, A., Baensch, H., & Heim, W. (2016). Sharks of the Mediterranean. McFarland,  
59  
60  
Jefferson, 236 pp.



1  
2  
3 Dulvy, N. K., Fowler, S. L., Musick, J. A., Cavanagh, R. D., Kyne, P. M., Harrison, L. R., ... White, W.  
4  
5 T. (2014). Extinction risk and conservation of the world's sharks and rays. *eLife*, 3, e00590.  
6

7 Edelist, D., Scheinin, A., Sonin, O., Shapiro, J., Salameh, P., Rilov, G., ... Zeller, D. (2013). Israel:  
8  
9 Reconstructed estimates of total fisheries removals in the Mediterranean, 1950–2010. *Acta Adriatica*,  
10  
11 54, 253 – 264.  
12

13  
14 European Commission (2011). European Commission 2011/471. Communication from the  
15  
16 Commission to the European Parliament, the Council, the European Economic and Social Committee  
17  
18 and the Committee of the Regions. Reform of the Common Fisheries Policy. Available from:  
19  
20 <http://www.europarl.europa.eu/document/activities/cont/201109/20110916ATT26808/20110916A>  
21  
22 [TT26808EN.pdf](http://www.europarl.europa.eu/document/activities/cont/201109/20110916ATT26808/20110916A)  
23

24  
25 Feitosa, L. M., Martins, A. P. B., Giarrizzo, T., Macedo, W., Monteiro, I. L., Gemaque, R., ...  
26  
27 Carvalho-Costa, L. F. (2018). DNA-based identification reveals illegal trade of threatened shark species  
28  
29 in a global elasmobranch conservation hotspot. *Scientific Reports*, 8(1). Doi:10.1038/s41598-018-  
30  
31 21683-5  
32

33  
34 Ferretti, F., Morey, G., Serena, F., Mancusi, C., Coelho, R. P., Seisay, M., ... Buscher, E. (2016).  
35  
36 *Squatina culate*. The IUCN Red List of Threatened Species 2016: e.T61418A16570000. Downloaded on  
37  
38 15 April 2018.  
39

40  
41 Fortibuoni, T., Borme, D., Franceschini, G., Giovanardi, O., & Raicevich, S. (2016). Common, rare  
42  
43 or extirpated? Shifting baselines for common angelshark, *Squatina squatina* (Elasmobranchii:  
44  
45 Squatinidae), in the Northern Adriatic Sea (Mediterranean Sea). *Hydrobiologia*, 772, 247-259.  
46

47  
48 Froese, R., & Pauly, D. (2018). FishBase. [www.fishbase.org](http://www.fishbase.org) (02/2018).  
49

50  
51 Gardiner, M., Allee, L., Brown, P., Losey, J., Roy, H., & Smyth, R. (2012). Lessons from lady  
52  
53 beetles: accuracy of monitoring data from US and UK citizen-science programs. *Frontiers in Ecology*  
54  
55 *and the Environment*, 10, 471-476.  
56  
57  
58  
59  
60

1  
2  
3           Giovos, I., Chatzisprou, A., Doumpas, N., Stoilas, V.O., & Moutopoulos, D.K. (2018). Using  
4 unconventional sources of information for identifying critical areas for the endangered guitarfish in  
5 Greece. *Journal of the Black Sea/Mediterranean Environment*, 24, 38-50.  
6  
7

8  
9           Gonzalvo, J., Giovos, I., & Moutopoulos, D.K. (2015). Fishermen's perception on the  
10 sustainability of small-scale fisheries and dolphin–fisheries interactions in two increasingly fragile  
11 coastal ecosystems in western Greece. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 25,  
12 91-106.  
13  
14

15  
16           Gordon, C., Hood, A., Lawson, J., Dulvy, N., Barker, J., Bartolí, À., ... Meyers, E. (2017). Eastern  
17 Atlantic and Mediterranean angel shark conservation strategy. The Shark Trust.  
18  
19

20  
21           Hadjichristophoru, M. (2006): Chondrichthyes in Cyprus. In: Baştusta, N., Ç. Keskin, F. Serena &  
22 B. Seret (eds.): The Proceedings of the International Workshop on Mediterranean Cartilaginous Fish  
23 with Emphasis on Southern and Eastern Mediterranean. 14-16 October 2005, Istanbul, Turkey. Turkish  
24 Marine Research Foundation, pp. 162–168.  
25  
26

27  
28           Halouani, G., Lasram, F. B. R., Khalfallah, M., Zeller, D. & Pauly, D. (2015). Reconstruction of  
29 Marine Fisheries catches for Tunisia (1950-2010). Fisheries Centre Working Paper #2015-95, University  
30 of British Columbia, Vancouver, 12 pp.  
31  
32

33  
34           Holcer, D., & Lazar, B. (2017). New data on the occurrence of the critically endangered common  
35 angelshark, *Squatina squatina*, in the Croatian Adriatic Sea. *Natura Croatica: Periodicum Musei*  
36 *Historiae Naturalis Croatica*, 26, 313-320.  
37  
38

39           IUCN 2018. The IUCN Red List of Threatened Species. Version 2018-1.  
40 <<http://www.iucnredlist.org>>  
41  
42

43  
44           Johannes, R.E., Freeman, M.M., & Hamilton, R.J. (2000). Ignore fishers' knowledge and miss the  
45 boat. *Fish and Fisheries*, 1, 257-271.  
46  
47

48  
49           Kabasakal, H., & Kabasakal, Ö. (2014). Status of angelshark, *Squatina squatina* (Elasmobranchii:  
50 Squatiniformes: Squatinidae) in the sea of Marmara. *Annales, Series Historia Naturalis*, 24, 1.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Katsanevakis S, & Moustakas A. (2018). Uncertainty in Marine Invasion Science. *Frontiers in*  
4 *Marine Science*, 5, 38 doi: <https://doi.org/10.3389/fmars.2018.00038>

7 Khalfallah, M., Belhabib, D., Zeller, D., & Pauly, D. (2015). Reconstruction of Marine Fisheries  
8 catches for Libya (1950-2010). Fisheries Centre Working Paper #2015-47, University of British  
9 Columbia, Vancouver, 16 pp.

14 Khalfallah, M., Dimech, M., Ulman, A., Zeller, D., & Pauly, D. (2017). Reconstruction of Marine  
15 Fisheries Catches for the Republic of Malta (1950-2010). *Mediterranean Marine Science*, 18, 241-250.

18 Last, P. R., & White, W. T. (2008). Three new angel sharks (Chondrichthyes: Squatinidae) from  
19 the Indo-Australian region. *Zootaxa*, 1734, 1-26.

23 Machias, A., Vassilopoulou, V., Vatsos, D., Bekas, P., Kallianiotis, A., Papaconstantinou, C., &  
24 Tsimenides, N. (2001). Bottom trawl discards in the northeastern Mediterranean Sea. *Fisheries*  
25 *Research* 53, 181-195.

29 Mahmoud, H. H., Teh, L., Khalfallah, M., & Pauly, D. (2015). Reconstruction of marine fisheries  
30 statistics in the Egyptian Mediterranean Sea, 1950-2010. Fisheries Centre Working Paper #2015-85,  
31 University of British Columbia, Vancouver, 17 pp.

36 Meyers, E. K. M., Tuya, F., Barker, J., Jiménez Alvarado, D., Castro-Hernández, J. J., Haroun, R., &  
37 Rödder, D. (2017). Population structure, distribution and habitat use of the Critically Endangered Angel  
38 Shark, *Squatina squatina*, in the Canary Islands. *Aquatic Conservation: Marine and Freshwater*  
39 *Ecosystems*, 27, 1-12.

45 Miller, M. H. (2016) Status Review Report of 3 Species of Angel sharks: *Squatina culate*, *S. culate*,  
46 *and S. squatina*. Report to National Marine Fisheries Service, Office of Protected Resources. June 2016.  
47 74 pp.

52 Moutopoulos, D. K., Dimitriou, E., Katselis, G., & Koutsikopoulos, C. (2017). Typology of illegal  
53 fishing in transitional waters: Fisheries infringement records from Mesolonghi-Etolikon lagoons (Ionian  
54 Sea, Greece). *Ocean and Coastal Management*, 141, 20-28.

1  
2  
3 Moutopoulos D. K., Prodromitis G., Mantzouni I., & Koutsikopoulos C. (2016). Quantifying the  
4 implementation of Common Fisheries Policy: Patterns of fisheries violations and penalties imposed in  
5 Greek waters. *Marine Policy*, 70, 65-76.  
6  
7

8  
9 Moutopoulos, D. K., Koutsikopoulos, C. (2014). Fishing strange data in national fisheries statistics  
10 of Greece. *Marine Policy*, 48:114-122.  
11  
12

13  
14 Nieto, A., Ralph, G. M., Comeros-Raynal, M. T., Kemp, J., García-Criado, M., Allen, D.J., ...,  
15 Williams, J. T. (2015). European Red List of marine fishes. Publications Office of the European Union,  
16 Luxembourg, 88pp.  
17  
18

19  
20 Pauly, D., & Zeller, D. (2016). Catch reconstructions reveal that global marine fisheries catches  
21 are higher than reported and declining. *Nature Communications* 7, 10244.  
22  
23

24  
25 Pauly, D., & Froese, R. (2012). Comments on FAO's State of Fisheries and Aquaculture, or  
26 'SOFIA2010'. *Marine Policy*, 36, 746–752.  
27  
28

29  
30 Pazartzis, T., Siaperopoulou, S., Gubili, C., Maradidou, S., Loukovitis, D., Chatzisyrou, A., Griffiths,  
31 A. M., Minos, G., Imsiridou, A. (2019). High levels of mislabeling in shark meat – Investigating patterns  
32 of species utilization with DNA barcoding in Greek retailers, *Food Control* (in press).  
33  
34

35  
36 Piroddi, C., Kristina, M., Zylich, K., Greer, K., Ulman, A., Zeller, D., & Pauly, D. (2015).  
37 Reconstruction of Italy's marine fisheries removals and fishing capacity, 1950-2010. *Fisheries Research*  
38 172: 137-147.  
39  
40

41  
42 Soldo, A., & Bariche, M. (2016). *Squatina oculata*. The IUCN Red List of Threatened Species 2016:  
43 e.T61417A16569265. Downloaded on 15 April 2018.  
44  
45

46  
47 Stelbrink, B., von Rintelen, T., Cliff, G., & Kriwet, J. (2010). Molecular systematics and global  
48 phylogeography of angel sharks (genus *Squatina*). *Molecular Phylogenetics and Evolution*, 54, 395-404.  
49  
50

51  
52 Stephenson, R. L., Paul, S., Pastoors, M. A., Kraan, M., Holm, P., Wiber, M., ... Benson, A. (2016).  
53 Integrating fishers' knowledge research in science and management. *ICES Journal of Marine Science*,  
54 73, 1459-1465.  
55  
56  
57  
58  
59  
60

1  
2  
3 Thurstan R. H., Buckley S. M., Ortiz J. C., & Pandolfi J. M. (2016). Setting the record straight:  
4 assessing the reliability of retrospective accounts of change. *Conservation Letters*, 9, 98–105.

7 Tiralongo, F., Messina, G., Coco, S., & Lombardo B. M. (2018). On the presence of a well-  
8 established population of *Lobotes surinamensis* (Bloch, 1790) in the central Mediterranean Sea.  
9  
10  
11 *Annales, Series Historia Naturalis*, 28, 31–36.

14 Tulloch, A. I. T., Auerbach, N., Avery-Gomm, S., Bayraktarov, E., Butt, N., Dickman, C. R., ...  
15  
16 Watson, J. E. M. (2018). A decision tree for assessing the risks and benefits of publishing biodiversity  
17 data. *Nature Ecology & Evolution* 2, 1209–1217.

21 Ulman, A., Bekiřođlu, ř., Zengin, M., Knudsen, S., Ünal, V., Mathews, C., ... Pauly, D. (2013). From  
22 bonito to anchovy: a reconstruction of Turkey's marine fisheries catches (1950-2010). *Mediterranean*  
23 *Marine Science*, 14, 309-342.

27 Ulman, A., Çiçek, B.A., Salihoglu, I., Petrou, A., Patsalidou, M., Pauly, D., & Zeller, D. (2015a).  
28 Unifying the catch data of a divided island: Cyprus's marine fisheries catches, 1950–2010. *Environment,*  
29 *Development and Sustainability*, 17, 801-821.

34 Ulman A., Saad A., Zylich K., Pauly D., & Zeller D. (2015b). Reconstruction of Syria's fisheries  
35 catches from 1950-2010: Signs of overexploitation. *Acta Ichthyologica et Piscatoria*, 45, 259-272.

39 Vasconcellos Bunholi, I., Ferrette, B., De Biasi, J., de Oliveira M.C., Rotundo, M., Oliveira, C., ...  
40  
41 Mendonça, F., (2018). The fishing and illegal trade of the angelshark: DNA barcoding against misleading  
42 identifications. *Fisheries Research*, 206, 193-197.

45 Walker, P., Cavanagh, R., Ducrocq, M., & Fowler, S. (2005). Regional Overviews: Northeast  
46 Atlantic (including Mediterranean and Black Sea). In: Sharks, rays and chimaeras: The status of the  
47 Chondrichthyan fishes. IUCN SSC Shark specialist group (eds. By Fowler S, Cavanagh R, Camhi M,  
48 Burgess G, Cailliet G, Fordham S, Simpfendorfer C & Musick J), pp. 71–94. IUCN, Gland, Switzerland,  
49 Cambridge, UK.

52 Zeller, D., & Pauly, D. (2006). Reconstruction of marine fisheries catches for key countries and regions  
53 (1950-2005). Vancouver, *Fisheries Centre Research Reports*, 15.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


## Tables

**Table 1.** Citizen Science observations of angel sharks reported to the three citizen science projects. The location of the observations can be found in Figure 1, while the pictures of the observations are shown in Figure 2.

#	Date	Location	Species observed	N of specimen	Type of observation	Size (m)	Depth (m)	Substrate
1	29/07/2015	Sitia, Crete, GR	<i>S. oculata</i>	1	Nets	-	20	-
2	01/07/2017	Rhodes Island, GR	<i>S. squatina</i>	1	Found stranded	-	-	-
3	02/12/2017	Marzamemi, Sicily, IT	<i>S. aculeata</i>	1	Trammel net	-	70	Sandy with scattered rocks
4	05/12/2017	Marzamemi, Sicily, IT	<i>S. oculata</i>	1	Trammel net	-	55	Sandy with scattered rocks
5	14/12/2017	Ras Lanuf, LB	<i>S. squatina</i>	3	Recreational fishing	1	-	Sandy-muddy
6	02/02/2018	Brega City, LB	<i>S. squatina</i>	2	Professional trammel net	1	50	Sandy-muddy
7	04/02/2018	Brega City, LB	<i>S. squatina</i>	1	Professional trammel net	1.1	-	Sandy-muddy
8	04/02/2018	Geminis, LB	<i>S. squatina</i>	1	Boat-based Recreational fishing	0.8	-	Sandy-muddy
9	23/02/2018	Cyclades, GR	<i>S. aculeata</i>	1	Bottom trawler	1.2	165	Rocky
10	25/02/2018	Cyclades, GR	<i>Squatina</i> spp.	1	Bottom trawler	2	229	Rocky
11	11/03/2018	Cyclades, GR	<i>S. squatina</i>	1	Bottom trawler	1.5	236	Rocky-muddy
12	14/03/2018	Ras Lanuf, LB	<i>S. squatina</i>	1	Recreational fishing	1	-	Sandy-muddy
13	29/04/2018	Cyclades, GR	<i>S. aculeata</i>	1	Bottom trawler	1	130	-
14	09/05/2018	Paralimni, CY	<i>Squatina</i> spp.	1	Boat-based Recreational fishing	0.7	60	Sandy
15	13/11/2018	Rhodes Island	<i>S. aculeata</i>	1	Boat-based Recreational fishing	-	330	-

**Table 2.** Percentage contribution of the mean annual angel shark reconstructed catches per country for the Mediterranean waters during 1950-2014. %' indicated the percentage representation of angel shark to all combined country reconstructed catches.

Country	%	%'
Cyprus	0.11	0.011
Egypt (Mediterranean)	2.11	0.010
Israel (Mediterranean)	0.05	0.003
Libya	0.01	0.000
Malta	0.42	0.063
Sicily (Italy)	0.04	0.000
Syria	10.72	0.018
Tunisia	7.59	0.787
Turkey (Marmara Sea)	58.19	0.266
Turkey (Mediterranean Sea)	20.75	0.028
Algeria	0.01	0.000



**Figure 1.** The map displays the CSR observations of angel sharks collected in the context of this study (see also Table 1), the interviews of targeted marine observers (with positive and negative responses about current observations of angel sharks), the reports of historic occurrences of angel sharks collected in the context of this study and the published observation records of angel sharks available in the bibliography and the grey literature (Supporting Table 2).



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Figure 2.** Angel Shark specimens reported in the context of this study. The numbers correspond to the number of each observation as displayed in Table 1.




Figure 3. Annual reconstructed catches (in t.) of angel shark per country between 1950-2014.




Figure 4. Total reconstructed catches (in t.) of angel shark in the Mediterranean during 1950-2014.