

2019-05-21

Chronosequence of former kaolinite open cast mines suggests active intervention is required for the restoration of Atlantic heathland

Lane, M

<http://hdl.handle.net/10026.1/14228>

10.1111/rec.12983

Restoration Ecology

Wiley

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Chronosequence of former kaolinite open cast mines suggests active intervention is required for the restoration of Atlantic heathland

Lane M.^{1,2}, Hanley M.E.¹, Lunt, P.³, Knight M.E.¹,
Braungardt C.B.³, Ellis, J.S.¹,

1. School of Biological and Marine Sciences, University of Plymouth, Drake Circus, Plymouth, Devon, PL4 8AA, UK
2. SIBELCO Ltd., Headon Works, Cornwood, Ivybridge, Devon, PL21 9PW, UK
3. School of Geography, Earth and Environmental Sciences, University of Plymouth, Drake Circus, Plymouth, Devon, PL4 8AA, UK

Keywords: restoration, mining, kaolinite, vegetation, community ecology, lowland heath

Running title: Lowland heath restoration – vegetation analysis

Corresponding author: Mary Lane, School of Biological and Marine Sciences, University of Plymouth, Drake Circus, Plymouth, Devon, PL4 8AA, UK. Email: mary.lane@plymouth.ac.uk.

Author contributions: All authors conceived and designed the research, ML collected field data and performed laboratory analyses; ML, MH, JE analysed the data, ML, MH, wrote the manuscript with editorial assistance from CB, PL, MK, JE.

1 **Abstract**

2 Atlantic lowland heath (ALH) is a priority conservation habitat in Western Europe, but
3 restoration efforts have met with mixed success due to the complexity associated with
4 replicating establishment conditions. By virtue of their impoverished, often acidic soils and
5 geographic location in areas naturally occupied by ALH communities, former kaolinite mine
6 sites may offer an excellent opportunity for heathland restoration. Using a chronosequence of
7 former open cast kaolinite mines in SW England (0, 2, 27, and 150 years since mining ceased),
8 we determined the ability of ALH vegetation and soils to re-establish naturally and in addition,
9 for the three youngest sites, how the reinstatement of stored overburden affected heathland
10 regeneration. Analysis of soil characteristics revealed major differences in the levels of acidity,
11 organic content, and fertility between abandoned kaolinite sites and a nearby natural reference
12 heath. Even 150 years after mining ceased, concentrations of all major soil nutrients and
13 organic content were lower and pH higher, than undisturbed ALH. The reinstatement of
14 overburden did little to improve soil quality, since all former kaolinite sites were dominated by
15 mesotrophic grasses, rather than species characteristic of the target ALH community. We
16 conclude that to maximise the potential of former open cast kaolinite sites for ALH re-
17 establishment, changes in pre-and post-restoration management are required. These include
18 modification of how overburden is stockpiled, while the addition of organic material, microbial
19 communities, and sulphur (to reduce soil pH) to reinstated overburden are likely also essential
20 interventions to facilitate successful ALH establishment.

21

22 **Implications for Practice**

23 • Successful restoration of former quarries requires a local propagule source, but such
24 sources may be limited near large scale, open-cast mining operations.

25 • Reinstatement of stored overburden to former kaolinite sites does little to improve key soil
26 characteristics and facilitate heathland restoration.

27 • Unlike most restorations using stored overburden, a requirement for low acidity poses a
28 problem for heathland establishment and necessitates further manipulation of pH and soil
29 nutrients to prevent dominance by mesotrophic grassland species.

30

31

32

33 Introduction

34 The Atlantic lowland heaths (ALH) of NW Europe are a distinct habitat characterised by a
35 dominant heather (*Calluna* and *Erica* species), shrubby Fabaceae (i.e. *Ulex* and other Genisteeae
36 species) and distinctive graminoid (e.g. *Molinia caerulea*) community growing on low nutrient,
37 acid soils (Gimingham 1972; Loidi et al. 2010). Most heathlands are the result of anthropogenic
38 management imposed by periodic fire, grazing, or other disturbances, and as such, these
39 habitats have an important cultural, as well as biodiversity and ecosystem service value
40 (Mitchell et al. 2008; Pywell et al. 2011; Fagúndez 2012). In the UK alone, 133 conservation
41 priority (UK Post-2010 Biodiversity Framework) plant and animal species are associated with
42 ALH, including 47 invertebrates with either a restricted, or very restricted UK distribution,
43 which are also internationally rare or endangered (Webb et al. 2009). Globally, heathlands face
44 a number of threats, and due to a combination of changing management, atmospheric N
45 deposition, and habitat loss associated with building development, the ALH habitats of western
46 Europe, are particularly endangered (Fagúndez 2012; Bähring et al. 2017). For example, only
47 one sixth of the lowland heath present in England from the early 19th century remains, much
48 of this lost to urbanisation and agricultural intensification (Perrow & Davy, 2002; Webb et al.
49 2009). This dramatic decline has prompted the European Union and individual member states
50 to adopt and implement various protection and restoration strategies for the habitat (JNCC,
51 2004; Pywell et al. 2011).

52 ALH requires at least four elements to establish and persist. These include: (i) low soil pH (2.8-
53 3.9, Clarke 1997), (ii) low soil nutrient content (exchangeable calcium 80-159 $\mu\text{g Ca g}^{-1}$,
54 exchangeable phosphorus $< 10 \mu\text{g P g}^{-1}$ soil, Clarke 1997), (iii) propagule supply of dominant
55 dwarf ericoid plants, and (iv) management to prevent succession to woodland (Kleijn et al.
56 2008; Martinez-Ruiz et al. 2007; Newton et al. 2009). On this basis, many different restoration
57 techniques have been tested (Pywell et al. 1994; Clemente et al. 2016) but these can be

58 categorised into two classes: those involving soil amelioration (e.g. nutrient addition,
59 overturning), and those involving the selective addition of plants or seeds (Allison & Ausden,
60 2004; Walker et al. 2004a; Pywell et al. 2007; Glen et al. 2017). The results of many previous
61 studies indicate the most important factor for successful heathland restoration is prior land use,
62 as the most successful restorations are situated on former heathland (Walker et al. 2004b). Here,
63 the removal of invasive scrub, coupled with the distribution of heather brash and native seeds
64 (of local morphotype/genotype) cut and/or collected from local established adjacent heathland,
65 have proved the most effective strategy (Walker et al. 2004b, 2007; Diaz et al. 2008). This
66 approach not only provides the recipient community with necessary propagules and microbial
67 symbionts, it also underscores the importance of soil biogeochemistry and the major
68 contribution that the soil seed bank makes to heathland regeneration (Clarke 1993; Pywell et
69 al. 1996; Fagúndez, 2013; Nussbaumer et al. 2016). This is not, however, a sustainable or
70 practical technique to cover large areas or where there has been significant land use change,
71 such as quarrying operations over many decades.

72 In many cases, effective restoration of former open cast quarries may be achieved by ‘passive
73 restoration’ (Prach et al. 2001, 2013; Tropek et al. 2010). This approach has many advantages,
74 including relatively rapid colonisation by local ecotypes of well-adapted species, with minimal
75 economic costs to the mine operator (Prach & Hobbs 2008). Passive restoration seems to work
76 best however, when the disturbed site is small, and surrounded by natural vegetation unaffected
77 by the initial disturbance (Holl & Aide 2011; Prach et al. 2014).

78 Many OCM (Open Cast Mining) sites in the UK are located in areas naturally-dominated by
79 heathland habitats, where high concentrations of mineral deposits, such as cassiterite, ilmenite,
80 and kaolinite occur beneath the low nutrient, highly acidic, soils. This pattern is generally true
81 of European ALH, where the habitat is most commonly associated with soils originating over
82 low nutrient, sand and gravel beds (e.g. Belgium and the Netherlands), or igneous, typically

83 granitic, intrusions (Scandinavia, Western France & UK). In SW England, ALH frequently
84 coincides with deposits of the aluminosilicate mineral, kaolinite, a product of in-situ alteration
85 of the plagioclase feldspar component of the granite intrusions that surface throughout
86 Cornwall and west Devon. As a result, there has been a 300 year history of OCM kaolinite
87 extraction in the region, supporting over 5% of the global extent of ALH habitat (Devon BAP,
88 2009).

89 In general, the nature and success of any restoration depend on the planning conditions imposed
90 and the suitability of post-OCM conditions for plant establishment (Cooke & Johnson, 2002;
91 Kuter, 2013). By its nature, OCM necessitates the removal of plant communities and
92 underlying material to expose commercially-extractable minerals. The topsoil and underlying
93 sediments (overburden) are mixed and stockpiled, destroying natural soil structure. Often, there
94 is not enough original topsoil to cover the area left after extraction (Merino-Martín et al. 2017).
95 Most problematic however, is the long period (between 1-20 years) of overburden storage,
96 since over time, sub-surface soil layers in storage berms develop sub- or anoxic conditions,
97 causing changes in microbial communities and further deterioration of soil structure and quality
98 (Golos et al. 2016; Merino-Martín et al. 2017). In the specific case of ALH restoration, the
99 combination of stockpiling methods and age since removal means that the availability of donor
100 soils containing seed of the community dominant, *Calluna vulgaris*, is often limited (Pywell et
101 al. 2002). Even if seeds are present, the altered characteristics and microbial communities of
102 donor soils degraded by storage can reduce subsequent *Calluna* seedling establishment
103 (Bossuyt & Honnay 2008). *C. vulgaris* can take 25-55 years to colonize mine spoil after
104 cessation of mining operations (Roberts et al. 1982). In addition, the seeds of other component
105 plant species, including many rare heathland specialists, are poorly represented in the
106 overburden seed bank compared to their contribution in the natural ALH community (Bakker
107 & Berendse 1999). The decline of the seed bank and loss of soil structure and microbial

108 community in stored overburden underscores why the most successful attempts to restore ALH
109 have been on former heathland sites. In these cases, some vestige of pre-disturbance soil
110 propagule availability, microbial community, and soil biogeochemistry remains (Pywell et al.
111 1996; Walker et al. 2004b, 2007, Diaz et al. 2008; Wubs et al. 2018).

112 The aim of this case-study was to determine whether, and how quickly, after kaolinite mining
113 has ceased, plant communities approach those of an undisturbed, target, ALH community.
114 We also investigate temporal changes in the establishment of plant communities to see how
115 closely the vegetation of former kaolinite extraction sites followed observed changes in soil
116 quality and how quickly a post-OCM site would converge with typical ALH. In doing so, we
117 test the hypotheses that even without any active attempt to ameliorate overburden, given
118 sufficient time, it is possible to re-instate ALH following kaolinite OCM.

119

120 **Methods**

121 *Study Sites*

122 Located on the periphery of the Dartmoor National Park, Devon, SW England, commercial
123 OCM kaolin extraction has taken place at Headon China Clay Works (50.2510°N, 03.5930°W)
124 since 1855. The quarry offers a sequential series of sites where kaolinite extraction ceased in
125 1868, 1990, 2013 and 2015. As described by De Palma *et al.* (2018), this provides a space-for-
126 time substitution (where spatial comparisons are made to infer temporal change) under a
127 Control-Impact model with the associated limitations. All locations were therefore at a similar
128 altitude and experienced similar climate, although the 1990 and 2015 sites were north facing
129 and the 2013 south facing slope. The 1868 site was also south facing, but had a steeper slope
130 angle ($\pm 66\%$) compared to the younger sites ($\pm 30\%$). While the 1868 site received no known
131 post-OCM interventions, the 1990, 2013 and 2015 sites were covered in overburden stockpiled

132 outside in large mounds approximately 6 m deep for 5 years, in order to help stabilise slopes.
133 The nearby Trendlebere Down Nature Reserve (50.3641°N, 03.4424°W) was selected as a
134 typical ALH reference site, as it had no history of mining, but similar slope, aspect, altitude,
135 and geomorphology to the commercial quarry prior to kaolinite extraction. At the time the soil
136 and vegetation surveys were undertaken, these sites were 147, 27, 2 and 0 years old,
137 respectively.

138 *Soil sampling and analysis*

139 In summer 2015, ten sampling points at each restoration site were determined using the ‘W-
140 walk’ method (JNCC, 2004). In each, a starting point was selected randomly and a quadrat (0.5
141 m × 0.5 m) placed to determine the first sampling position. The sample points were 20 m apart,
142 and the total distance walked was 180 m. A 30 cm soil core was taken from the left corner of
143 each quadrat using a soil coring kit (Eijkelkamp Agrisearch Equipment BV). The O horizon
144 (~15 cm) was sampled from the cores and subsequently dried in a desiccator set to 65°C,
145 disaggregated, sieved (2 mm mesh) and stored before analysis.

146 For pH analysis, 10 g of soil in 50 ml deionised water was mixed for 15 minutes with a magnetic
147 stirrer, left to settle and determined using a Hanna 991001 pH and temperature probe (Jones Jr,
148 2001). As a proxy for organic matter, loss on ignition was used to quantify soil carbon content,
149 with (~5 g) samples dried at 105°C for 1 hour, weighed and ashed at 400°C for 2 hours in a
150 Gallenkamp hotspot furnace (Jones Jr, 2001). Mineral elements were extracted using the
151 Mehlich 3 method (Jones Jr, 2001), whereby an extraction solution (30 ml) was added to each
152 soil sample (3 g) in centrifuge tubes and mixed on a reciprocating mechanical shaker at 200
153 rpm for 5 minutes. Samples were subsequently filtered through Whatman 42 filter papers, and
154 the filtrate retained in the dark until analysis. The Na, K, Mg, Ca and P concentration of the
155 extracted solution was analysed using a Thermo Scientific iCAP7400 ICP-OES instrument.

156 To assess soil nitrate/nitrite concentrations, 3 g samples were digested in 30 ml of 0.01 M
157 calcium sulphate, shaken on a reciprocal shaker for 15 mins at 180 rpm, and filtered through a
158 Whatman 42 filter paper, followed by a cadmium reduction reaction and quantification by
159 colourimetry (HACH DR/890) (Jones Jr, 2001). Cation exchange capacity, a measure of soil
160 ability to retain key nutrients in ‘plant-available’ form, was quantified using the sodium acetate
161 method (Jones Jr, 2001). We applied One-Way ANOVA, with a Welch’s correction for unequal
162 variances, to explore how these key soil chemical parameters varied according to the factor
163 ‘time since restoration’.

164 *Vegetation sampling and analysis*

165 Within each of the ten 0.5m × 0.5m quadrats positioned along the sample transect, species
166 presence and absence was quantified (0 – absent, present – 1), and an NMDS using the Raup-
167 Crick distance used to visualise variation in community patterns between sites (Clarke, 1993;
168 Zuur, Ieno & Smith, 2007). Analysis was performed in three dimensions using metaMDS and
169 ordiellipse to highlight groupings in the ‘vegan’ (Oksanen, 2015) package in ‘R’ v.3.5.2. Once
170 the communities were plotted onto an ordination plot, the physical characteristics of the soil
171 were overlaid as vectors (for variables where $P \leq 0.001$). Lines pointed in the same direction
172 are positively correlated to each other (Zuur, Ieno & Smith, 2007). This enabled interpretation
173 of the significant physical factors and how they were aligned with the various communities.
174 An ANOSIM was performed in the ‘vegan’ (Oksanen, 2015) package in ‘R’ v.3.5.2 to examine
175 variation in plant community composition between restoration treatments.

176

177 **Results**

178 *Soil chemistry*

179 We found little evidence to support the hypothesis that even several decades after OCM
180 terminated, soils on former kaolinite sites would transition naturally towards soils favouring an
181 ALH community. Even at the oldest (147-year-old) site, key aspects of soil chemistry were
182 very different from the Trendlebere Down heathland (Table 1). Concentrations of major
183 elements (Na, Ca, K and Mg) were generally an order of magnitude lower at the 1868 site, and
184 soil P and NO₃ concentrations were 28% and 20% respectively of those in established
185 heathland. With the exception of NO₃, the restored sites had lower major element levels; less
186 than 25% of the reference site. Established heathland soil was also more acidic (pH 3.8), and
187 had considerably higher organic matter content (67.5% OM) than all former OCM sites (pH
188 4.5-4.9, <6% OM), showing that the addition of stockpiled soil to the 1990, 2013, and 2015
189 sites had minimal beneficial impact on soil chemistry (Table 1).

190

191 *Changes in vegetation community composition*

192 Multivariate analysis revealed considerable variation in the plant community characteristics
193 between each site (global R_{ANOSIM} = 0.496, P < 0.001), (Fig 1). The reference ALH community
194 at Trendlebere was tightly clustered around the major defining plant species for this habitat
195 (i.e. *Calluna*, *Molinia*, and *Erica tetralix*), these species also being more abundant here than
196 any other site. The former OCM sites were less tightly clustered around distinct species; the
197 1868 site in particular showed broad overlap across many different plants, most
198 uncharacteristic of typical ALH communities (specifically, the graminoids *Deschampsia*,
199 *Festuca*, and *Juncus*, and the forbs *Potentilla* and *Galium*; species more commonly associated

200 with acid grasslands). Nonetheless, *Calluna* and *Molinia* at the 1868 site achieved the highest
201 abundance recorded at any former kaolinite mine location.

202 The 1990 and 2013 sites were dominated by Poaceae species characteristic of acid and
203 mesotrophic grasslands (e.g. *Deschampsia flexuosa* and *Festuca rubra*), although the position
204 of the 1990 cluster in the nMDS reflects that the contribution of both *Calluna* and *Molinia* to
205 the community was much greater here, than at the younger 2013 site. Also, of note is the fact
206 that *Ulex europaeus* was considerably more abundant at the 1990 site than any other location
207 (although the presence of this N-fixing legume appeared to have little impact on soil NO₃). The
208 1990 site had the tightest cluster of all the OCM restoration sites. The 2015 site clustered
209 around *Agrostis capillaris* (Fig 1), reflecting that quadrats here were dominated by bare ground
210 and had no ALH-characteristic plants present. The most important environmental factors
211 dictating plant community composition was the addition of overburden; K and P, and time (P
212 ≥ 0.001) (Fig 1).

213

214 **Discussion**

215 Although we emphasise that this case-study lacks true replication, our results nevertheless
216 corroborate the general view that effective ALH restoration is a long-term process with little
217 or no guarantee of success (Miller et al. 2017). Indeed, even after nearly 150 years (albeit
218 with minimal additional management; i.e. grazing by livestock), soil chemistry failed to
219 approach the levels of acidity, organic content, CEC or key soil nutrients characteristic of,
220 and important in, heathland soil (Clarke 1993; Green et al. 2015). Similarly, although some
221 species typical of established ALH were abundant in the 1868 site, the community was also
222 characterised by species representative of acid or mesotrophic grasslands. There seems little
223 potential therefore, to expect long-term, natural ALH recovery on the many kaolinite open

224 cast mines located in regions where this habitat is most common, and especially where
225 restoration occurs alongside active mining. Instead, and like many OCM sites globally,
226 heathland restoration can likely only be facilitated by further interventions after mining
227 operations cease (Holmes. 2001; Benigno et al. 2013; Clemente et al. 2016; Glen et al. 2017).

228 One commonly-applied approach is to reinstate stockpiled overburden onto former OCM
229 sites, but the results suggest this practice did little to facilitate any improvement in key soil
230 characteristics, or subsequent establishment of plant species typical of the target ALH
231 community. Even on the 25-year-old (1990) site, organic content and pH of the reinstated
232 overburden had little in common with those in nearby natural ALH. In theory, the use of
233 topsoil provides a source of native seed, mycorrhizal and bacterial symbionts with which to
234 facilitate plant community restoration (Muñoz-Rojas et al. 2016; Wubs et al. 2018). In
235 practice, however, suitable topsoils are scarce and overburden (topsoil mixed with underlying
236 mineral horizons) is stockpiled into large mounds to reduce footprint on the mine site, a
237 procedure that diminishes key properties over relatively short periods (Golos et al. 2016). For
238 the most part however, recent studies reporting the impact of soil stockpiling on restoration
239 have focussed on (generally negative) changes in the soil seed bank (Dickie et al. 1988;
240 Rokich et al. 2000) or soil microbial community (Harris et al. 1989; Poncelet et al. 2014). It
241 may be the case however, that soil nutrients are less impacted by storage (Abdul-Karrem &
242 McRae. 1984; Strohmeyer 1999).

243 A deficiency in the major macronutrients (NPK) required for plant establishment and growth
244 in stored overburden and kaolinite mine waste is nonetheless well known (Marrs et al. 1981;
245 Coppin & Bradshaw 1982). Phosphorus and potassium concentrations in the sites were
246 considerably lower (i.e. less than 20%), even 25 years after overburden had been reinstated,
247 than in the adjacent target community. Soil nitrate was, however, substantially higher in sites
248 with overburden (1990, 2013, 2015) than in the 1868 site where no interventions were

249 undertaken after OCM ceased. Other important heathland macronutrients, including Mg, Na
250 and Ca (Clarke 1993, 1997), were frequently present at concentrations less than one-tenth of
251 those seen in the adjacent ALH site. Removal and mixing of thin heathland topsoils with the
252 mineral soils that underlie them before mining inevitably dilutes soil nutrients; subsequent
253 storage and leaching from a generally coarse-grained overburden, further diminishes fertility.
254 Reinstatement of a nutrient limited, mineral overburden where the symbiotic soil microflora
255 plants require to extract nutrients from low fertility heathland soils are now absent,
256 unsurprisingly limits establishment of heathland specialists, even if propagules are available
257 (Diaz et al. 2006). To compound the problem, the low water retention capacity of coarse-
258 grained, low organic content mineral overburden increases substantially the risk of plant
259 mortality and reduced growth during drought (Machado et al. 2013; Bateman et al. 2018).
260 Although relatively uncommon in SW England, future climate scenarios predict increased
261 frequency of warm, dry summers, including extreme heatwaves and drought (Guillod et al.
262 2018).

263 Unlike the majority of mine rehabilitation studies where the low pH associated with
264 overburden poses a major problem for plant community restoration (Abdul-Kareem &
265 McRae 1984; Malik & Scullion 1998), none of the former kaolinite sites studied were as
266 acidic as natural ALH. Low soil pH is critical for the establishment of the ericoid shrubs that
267 characterise lowland heaths (Pywell et al. 1994; Marrs et al. 1998). Moreover, low pH often
268 results in loss of cations from soils; Green et al. (2015) for example, reported a positive
269 correlation between pH and concentrations of extractable K, Ca & Mg, but a negative
270 association with phosphate. In-turn, soil concentrations of many elements affects the
271 bioavailability of other key nutrients and also influence greatly the growth of species that
272 might otherwise outcompete the target heathland species. Green et al. (2015) describe how at
273 higher pH (5 or above), the vegetation of restored heathland sites was dominated by *Agrostis*

274 *capillaris* and note how control of this highly competitive species is key to *Calluna* and *Erica*
275 *cinerea* establishment. Similarly, our results show how *Agrostis capillaris*, along with at least
276 one other mesotrophic grass species, was dominant on the 1868, 1990 and 2013 sites where
277 soil pH remained above 4.7.

278 The failure of key soil characteristics or plant community composition throughout our
279 chronosequence to trend towards those associated with the adjacent natural ALH strongly
280 suggest that even where stored overburden is used, further manipulation is required. Benefits
281 may accrue from reduction in overburden storage times and the depth of stockpiles (reducing
282 compaction), and regular addition of organic material to retain soil meso-fauna and microbial
283 populations and function, and water holding capacity during storage (Dickie et al. 1988;
284 Rokich et al. 2000; Ngugi et al. 2018). Following reinstatement, further addition of organic
285 matter to overburden is desirable for the same reasons (Smith & Read 2010; Muñoz-Rojas et
286 al. 2016) and the potential enhancement of nitrogen cycling rates (Van Vuuren et al. 1992),
287 while fertilizer application can also encourage plant establishment and growth, and
288 concomitant benefits to soil biota (Ngugi et al. 2018). More specific to heathland restoration,
289 soil pH is effectively reduced by the application of sulphur, with the additional benefit of
290 increasing the bioavailability of phosphate without the need for fertilizer application (Green
291 et al. 2015). Heather establishment is also strongly dependent on symbiotic interactions with
292 ericoid mycorrhizal fungi (ERM) that do not respond well to long-term soil storage (Smith &
293 Read 2010). Consequently, the introduction of essential ERM to the soil may be essential to
294 effective ALH restoration where former kaolinite OCMs are covered with overburden stored
295 for long periods. Taken together therefore, we conclude that effective restoration of ALH
296 communities on former kaolinite quarries requires multiple interventions that address the
297 limiting effects of low soil fertility, relatively high soil pH, propagule limitation, and an
298 absence of soil micro- and macro-biota. Time alone is insufficient to facilitate these changes.

299

300 **Acknowledgements**

301 We thank Ben Uphill from Sibelco for access to Headon China Clay Works. The research was
302 supported by Sibelco as part of a business education partnership. We also thank Professor Karel
303 Prach, Professor Rob Marrs and an anonymous reviewer for commenting on earlier drafts of
304 this manuscript.

305

306 **Literature Cited**

- 307 Allison M, Ausden M (2004) Successful use of topsoil removal and soil amelioration to
308 create heathland vegetation. *Biological Conservation* 120:221-228.
- 309 Abdul-Kareem AW, McRae SG (1984) The effects on topsoil of long-term storage in
310 stockpiles. *Plant and Soil* 76:357-363.
- 311 Bähring A, Fichtner A, Ibe K, Schütze G, Temperton VM, von Oheimb G, Härdtle W (2017)
312 Ecosystem functions as indicators for heathland responses to nitrogen fertilization.
313 *Ecological Indicators* 72:185-193
- 314 Bakker JP, Berendse F (1999) Constraints in the restoration of ecological diversity in grassland
315 and heathland communities. *Trends in Ecology & Evolution* 14:63-68.
- 316 Bateman A, Lewandrowski W, Stevens J, Muñoz-Rojas M (2016) Ecophysiological indicators
317 to assess drought responses of arid zone native seedlings in reconstructed soils. *Land*
318 *Degradation & Development* 29:984-993.
- 319 Benigno SM, Dixon KW, Stevens J (2013) Increasing soil water retention with native-sourced
320 mulch improves seedling establishment in postmine Mediterranean sandy soils.
321 *Restoration Ecology* 21:617-626
- 322 Bossuyt B, Honnay O (2008) Can the seed bank be used for ecological restoration? An
323 overview of seed bank characteristics in European communities. *Journal of*
324 *Vegetation Science* 19:875-884.
- 325 Clarke KR (1993) Non-parametric multivariate analyses of changes in community structure.
326 *Australian Journal of Ecology* 18:117-143.

327 Clarke CT (1997) Role of soils in determining sites for lowland heathland reconstruction in
328 England. *Restoration Ecology* 5:256-264.

329 Clemente AS, Moedas AR, Oliveira G, Martins-Loução MA, Correia O (2016) Effect of
330 hydroseeding components on the germination of Mediterranean native plant species.
331 *Journal of Arid Environments* 125:68-72.

332 Cooke JA, Johnson, MS (2002) Ecological restoration of land with particular reference to the
333 mining of metals and industrial minerals: A review of theory and practice.
334 *Environmental Reviews* 10:41-71.

335 Coppin NJ, Bradshaw AD (1982) *The Establishment of Vegetation in Quarries and Open-pit*
336 *Non-metal Mines*. Mining Journal Books: London; 112.

337 Diaz A, Green I, Benvenuto M, Tibbett M (2006) Are Ericoid Mycorrhizas a Factor in the
338 Success of *Calluna vulgaris* Heathland Restoration? *Restoration Ecology* 14:187-195.

339 Diaz A, Green I, Tibbett M (2008) Re-creation of heathland on improved pasture using top
340 soil removal and sulphur amendments: Edaphic drivers and impacts on ericoid
341 mycorrhizas. *Biological Conservation* 141:1628-1635.

342 De Palma A, Sanchez-Ortiz K, Martin PA, Chadwick A, Gilbert G, Bates AE, Börger L,
343 Contu S, Hill SL, Purvis A, (2018) Challenges with inferring how land-use affects
344 terrestrial biodiversity: Study design, time, space and synthesis. In *Advances in*
345 *Ecological Research*: 58. 163-199. Academic Press.

346 Devon, CC. (2009) *Devon Biodiversity and Geodiversity Action Plan*. Devon County
347 Council

348 Dickie JB, Gajjar KH, Birch P, Harris JA (1988) The survival of viable seeds in stored topsoil
349 from opencast coal workings and its implications for site restoration. *Biological*
350 *Conservation* 43:257–265

351 Fagúndez, J (2012) Heathlands confronting global change: drivers of biodiversity loss from
352 past to future scenarios. *Annals of Botany* 111:51-172.

353 Gimingham, CH (1972) *Ecology of heathlands*. Chapman & Hall, London, UK.

354 Glen E, Price EAC, Caporn SJM, et al (2017) Evaluation of topsoil inversion in UK habitat
355 creation and restoration schemes. *Restoration Ecology* 25:72-81.

356 Golos PJ, Dixon KW, Erickson TE (2016) Plant recruitment from the soil seed bank depends
357 on topsoil stockpile age, height, and storage history in an arid environment.
358 *Restoration Ecology* 24:53-61.

359 Green I, Evans D, Diaz A (2015) Modifying soil chemistry to enhance heathland recreation: a
360 use for sulphur captured during oil refining. *International Journal of Plant and Soil*
361 *Science* 6:272-282.

362 Guillod BP, Jones RG, Dadson SJ, Coxon G, Bussi G, Freer J, Kay AL, Massey NR, Sparrow
363 SN, Wallom DC, Allen MR, (2018) A large set of potential past, present and future
364 hydro-meteorological time series for the UK. *Hydrology and Earth System Sciences*
365 22:611-634.

366 Harris JA, Birch P, Short KC (1989) Changes in the microbial community and physico-
367 chemical characteristics of soils stockpiled during opencast mining. *Soil Use and*
368 *Management* 5:161–168.

369 Holl KD, Aide TM, (2011) When and where to actively restore ecosystems? *Forest Ecology*
370 *and Management* 261:1558-1563.

371 Holmes PM, (2001) Shrubland restoration following woody alien invasion and mining:
372 effects of topsoil depth, seed source, and fertilizer addition. *Restoration Ecology* 9:71-
373 84.

374 JNCC, (2004) *Common Standards Monitoring Guidance for Lowland heath*, Version August
375 2004, ISSN 1743-8160. Jones Jr, JB (2001) *Laboratory guide for conducting soil tests*
376 *and plant analysis*. CRC press.

377 Kleijn D, Bekker RM, Bobbink R, De Graaf MC, Roelofs JG (2008) In search for key
378 biogeochemical factors affecting plant species persistence in heathland and acidic
379 grasslands: a comparison of common and rare species. *Journal of Applied Ecology*
380 45:680-687.

381 Kuter N (2013) Reclamation of degraded landscapes due to opencast mining. In Özyavuz, M.
382 (Ed.), *Advances in landscape architecture*. Rijeka: InTech. Retrieved February 27,
383 2016, from [http://www.intechopen.com/books/advances-in-landscape-](http://www.intechopen.com/books/advances-in-landscape-architecture/reclamation-of-degraded-landscapes-due-to-opencast-mining)
384 [architecture/reclamation-of-degraded-landscapes-due-to-opencast-mining](http://www.intechopen.com/books/advances-in-landscape-architecture/reclamation-of-degraded-landscapes-due-to-opencast-mining)

385 Loidi J, Biurrun I, Campos JA, García-Mijangos I, Herrera M (2010) A biogeographical
386 analysis of the European Atlantic lowland heathlands. *Journal of Vegetation Science*
387 21:832-842.

388 Machado NDM, Leite MGP, Figueiredo MA, Kozovits AR (2013) Growing *Eremanthus*
389 *erythopappus* in crushed laterite: a promising alternative to topsoil for bauxite-mine
390 revegetation. *Journal of Environmental Management* 129:149-156.

391 Malik A, Scullion J (1998) Soil development on restored opencast coal sites with particular
392 reference to organic matter and aggregate stability. *Soil Use and Management* 14:234-
393 239.

394 Marris RH, Snow CSR, Owen KM, Evans CE (1998) Heathlands and acid grassland creation
395 on arable soils at Minsmere: Identification of potential problems and a test of
396 cropping to impoverish soils. *Biological Conservation* 85:69-82.

397 Marris RH, Roberts RD, Skeffington RA, Bradshaw AD (1981) Ecosystem development on
398 naturally-colonized china clay wastes. II. Nutrient compartmentation. *Journal of*
399 *Ecology* 69:163-169.

400 Merino-Martín L, Commander L, Mao Z, Stevens JC, Miller BP, Golos PJ, Mayence CE,
401 Dixon K (2017) Overcoming topsoil deficits in restoration of semiarid lands:
402 Designing hydrologically favourable soil covers for seedling emergence. *Ecological*
403 *Engineering* 105:102-117.

404 Miller BP, Sinclair EA, Menz MH, Elliott CP, Bunn E, Commander LE, Dalziell E, David E,
405 Davis B, Erickson TE, Golos PJ (2017) A framework for the practical science necessary
406 to restore sustainable, resilient, and biodiverse ecosystems. *Restoration Ecology*
407 25:605-617.

408 Mitchell R, Rose R, Palmer S (2008) Restoration of *Calluna vulgaris* on grass-dominated
409 moorlands: The importance of disturbance, grazing and seeding. *Biological*
410 *Conservation* 141: 2100-2111.

411 Muñoz-Rojas M, Erickson TE, Dixon KW, Merritt DJ (2016) Soil quality indicators to assess
412 functionality of restored soils in degraded semiarid ecosystems. *Restoration Ecology*
413 24:43–52.

414 Newton AC, Stewart GB, Myers G, Diaz A, Lake S, Bullock JM, Pullin AS (2009) Impacts
415 of grazing on lowland heathland in north-west Europe. *Biological Conservation*,
416 142:935-947.

417 Ngugi MR, Dennis PG, Neldner VJ, Doley D, Fechner N, McElnea A (2018) Open-cut
418 mining impacts on soil abiotic and bacterial community properties as shown by
419 restoration chronosequence. *Restoration Ecology* 26:839-850.

420 Nussbaumer Y, Cole MA, Offler CE, Patrick JW (2016) Identifying and ameliorating nutrient
421 limitations to reconstructing a forest ecosystem on mined land. *Restoration Ecology*
422 24:202-211.

423 Oksanen J (2015) Unconstrained Ordination: Tutorial with R and vegan. R package version
424 2.3-4.

- 425 Perrow MR, Davy AJ, eds, (2002) Handbook of ecological restoration (Vol. 2). Cambridge
426 University Press.
- 427 Poncelet DM, Cavender N, Cutright TJ, Senko JM (2014) An assessment of microbial
428 communities associated with surface mining-disturbed overburden. *Environmental*
429 *Monitoring and Assessment* 186:1917-1929.
- 430 Prach, K, Bartha S, Joyce CB, Pyšek, P, Van Diggelen R, Wiegand G (2001) The role of
431 spontaneous vegetation succession in ecosystem restoration: a perspective. *Applied*
432 *Vegetation Science* 4, 111-114.
- 433 Prach K, Hobbs RJ (2008) Spontaneous succession versus technical reclamation in the
434 restoration of disturbed sites. *Restoration Ecology* 16:363-366.
- 435 Prach K, Lencová K, Řehouňková K, Dvořáková H, Jírová A, Konvalinková P, Mudrák O,
436 Novák J, Trnková R (2013) Spontaneous vegetation succession at different central
437 European mining sites: a comparison across seres. *Environmental Science and*
438 *Pollution Research* 20: 7680-7685.
- 439 Prach K, Řehouňková K, Lencová K, Jírová A, Konvalinková P, Mudrák O, Študent V,
440 Vaněček Z, Tich L, Petřík P, Šmilauer P, Pyšek P, (2014) Vegetation succession in
441 restoration of disturbed sites in Central Europe: the direction of succession and
442 species richness across 19 seres. *Applied Vegetation Science* 17: 193-200.
- 443 Pywell RF, Webb NR, Putwain PD (1994) Soil fertility and its implications for the restoration
444 of heathland on farmland in southern Britain. *Biological Conservation* 70:169-181.
- 445 Pywell RF, Webb N, Putwain PD (1996) Harvested heather shoots as a resource for heathland
446 restoration. *Biological Conservation* 75:247-254.
- 447 Pywell RF, Pakeman RJ, Allchin EA, Bourn NAD, Warman EA, Walker KJ (2002) The
448 potential for lowland heath regeneration following plantation removal. *Biological*
449 *Conservation* 108:247-258.
- 450 Pywell RF, Bullock JM, Tallwin JB, Walker KJ, Warman EA, Masters G (2007) Enhancing
451 diversity of species-poor grasslands: an experimental assessment of multiple
452 constraints. *Journal of Applied Ecology* 44:81-94.
- 453 Pywell RF, Meek WR, Webb NR, Putwain PD, Bullock JM (2011) Long-term heathland
454 restoration on former grassland: The results of a 17-year experiment. *Biological*
455 *Conservation* 144:1602-1609.
- 456 Roberts RD, Marrs RH, Skeffington RA, Bradshaw AD (1981) Ecosystem development on
457 naturally- colonized china clay wastes. I. Vegetation change and overall accumulation
458 of organic matter and nutrients. *Journal of Ecology* 69: 153-161.

459 Rokich DP, Dixon KW, Sivasithamparam K, Meney KA (2000) Topsoil handling and storage
460 effects on woodland restoration in Western Australia. *Restoration Ecology* 8:196–208

461 Smith SE, Read DJ (2010) *Mycorrhizal symbiosis*. Academic press.

462 Strohmayer P (1999) Soil stockpiling for reclamation and restoration activities after mining
463 and construction. *Restoration and Reclamation Review* 4:1-4

464 Tropek R, Kadlec T, Karesova P, Spitzer L, Kocarek, P, Malenovsky I, Banar P, Tuf IH,
465 Hejda M, Konvicka M, (2010) Spontaneous succession in limestone quarries as an
466 effective restoration tool for endangered arthropods and plants. *Journal of Applied*
467 *Ecology* 47:39-147.

468 Van Vuuren MM, Aerts E, Berendse F, De Visser W (1992) Nitrogen mineralization in
469 heathland ecosystems dominated by different plant species. *Biogeochemistry* 16:151-
470 166.

471 Walker KJ, Stevens PA, Stevens DP, Mountford JO, Manchester SJ, Pywell RF (2004a) The
472 restoration and re-creation of species-rich lowland grassland on land formerly
473 managed for intensive agriculture in the UK. *Biological Conservation* 119:1-18.

474 Walker KJ, Pywell RF, Warman EA, Fowbert JA, Bhogal A, Chambers BJ (2004b) The
475 importance of former land use in determining successful re-creation of lowland heath
476 in southern England. *Biological Conservation* 116:289-303.

477 Walker KJ, Warman EA, Bhogal A, Cross RB, Pywell RF, Meek, BR, Chambers, BJ,
478 Pakeman R (2007) Recreation of lowland heathland on ex-arable land: assessing the
479 limiting processes on two sites with contrasting soil fertility and pH. *Journal of*
480 *Applied Ecology* 44:573-582.

481 Webb JR, Drewitt AL, Measures GH (2009) *Managing for species: Integrating the needs of*
482 *England's priority species into habitat management. Part 1 report. Natural England*
483 *Research Reports Number 024*

484 Wubs ERJ, Melchers PD, Bezemer TM (2018) Potential for synergy in soil inoculation for
485 nature restoration by mixing inocula from different successional stages. *Plant and Soil*
486 433:147-156

487 Zuur A, Ieno EN, Smith GM (2007) *Analysing ecological data*. Springer Science & Business
488 Media.

TABLE 1 Comparison of mean (\pm SE, n=10) soil chemical conditions at an undisturbed Atlantic lowland heathland site (Trendlebere Down, Devon, UK – TBD) and former opencast kaolinite mine sites located in an adjacent commercial mine. The date when mining ceased at each site is given, and for the 1990, 2013 & 2015 sites, the termination of operations was followed by the replacement of stored overburden. The results of a one-way ANOVA of soil param are given, with different letters in superscript indicating significant difference in variance ($P < 0.05$) between site means, following Tukey's paired comparisons. CEC = Cation Exchange Capacity

Site		Na ($\mu\text{g g}^{-1}$)	Ca ($\mu\text{g g}^{-1}$)	K ($\mu\text{g g}^{-1}$)	Mg ($\mu\text{g g}^{-1}$)	P ($\mu\text{g g}^{-1}$)	NO ₃ ($\mu\text{g g}^{-1}$)	pH	Organic Content (%)	CEC (mEq /100g)
TBD	Mean	16.5 ^A	73.2 ^A	48.2 ^A	56.4 ^A	8.1 ^A	2.0 ^A	3.8 ^A	67.5 ^A	76.7 ^A
	(SE)	1.2	5.8	4.3	4.7	0.9	0.4	0.0	5.8	3.7
1868	Mean	1.9 ^B	7.3 ^B	6.5 ^B	4.3 ^B	2.0 ^B	0.4 ^B	4.9 ^B	3.4 ^B	8.5 ^B
	(SE)	0.3	1.7	1.1	0.7	1.2	0.1	0.1	0.6	0.6
1990	Mean	2.7 ^B	7.2 ^B	9.2 ^B	3.9 ^B	2.0 ^B	1.8 ^A	4.7 ^{BC}	5.9 ^B	17.6 ^B
	(SE)	0.1	3.3	2.0	0.7	0.1	0.2	0.1	0.3	1.5
2013	Mean	1.5 ^B	16.4 ^B	4.1 ^B	3.9 ^B	2.0 ^B	1.1 ^{AB}	4.9 ^B	3.8 ^B	10.1 ^B
	(SE)	0.1	4.1	0.5	0.6	0.1	0.3	0.1	0.5	1.5
2015	Mean	1.6 ^B	9.9 ^B	5.6 ^B	3.5 ^B	0.8 ^B	1.4 ^{AB}	4.5 ^C	3.9 ^B	11.4 ^B
	(SE)	0.2	3.8	0.6	0.4	0.2	0.4	0.1	0.6	1.5
ANOVA	$F_{(4,45)}$	120.1	32.16	65.5	103.5	21.36	4.85	38.19	168.6	118.5
	P	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001




Figure 1: nMDS of the presence- absence data plant community of an undisturbed ALH site (TBD) and restored kaolinite mine sites (shown in 2 dimensions for ease of visualisation). The date labels denote the year when mining ceased. Stress =0.11. Ordellipse are present to show the overlap of the communities. The vectors are significant environmental factors ($P \geq 0.001$)

Key to environmental factors: Timenatural, TBD; Timeold, 1868; Timerecent, 1990; timenew, 2013,2015; Soiln, overburden not added; Soily, overburden added.

Key to plant species: Agr cap, *Agrostis capillaris*: Agr sto, *Agrostis stolonifera*: Ant odo, *Anthoxanthum odoratum*: Des fle, *Deschampsia flexuosa*: Fes ovi, *Festuca ovina*: Fes rub, *Festuca rubra*: Mol cae, *Molinia caerulea*: Nar str, *Nardus stricta*: Cal vul, *Calluna vulgaris*:

Eri tet, *Erica tetralix*: Eri cin, *Erica cinerea*: Rhopon, *Rhododendron ponticum* Vac myr, *Vaccinium myrtillus*: Ule eur, *Ulex europaeus*: Pot ere, *Potentilla erecta*: Ver off, *Veronica officinalis*: Gen cam, *Gentianella campestris*: Pol vul, *Polygala vulgaris*: Gal ver, *Galium verum*: Gal sax, *Galium saxatile*: Tri rep, *Trifolium repens*: Tri pra, *Trifolium pratense*: Dig pur, *Digitalis purpurea*: Dro agg, *Drosera* (agg): Jun a.a, *Juncus articulatus*: Jun bul, *Juncus bulbosus*: Jun squ, *Juncus squarrosus*: Sph agg, *Sphagnum* (agg): Cla agg, *Cladonia* (agg).