

2020-08

The cytotoxic activity of extracts of the brown alga *Cystoseira tamariscifolia* (Hudson) Papenfuss, against cancer cell lines changes seasonally

Mansur, AA

<http://hdl.handle.net/10026.1/15358>

10.1007/s10811-019-02016-z

Journal of Applied Phycology

Springer Science and Business Media LLC

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1 **The cytotoxic activity of extracts of the brown alga *Cystoseira tamariscifolia***

2 **(Hudson) Papenfuss, 1950 against cancer cell lines changes seasonally**

3 **Ahmad A. Mansur*, Murray T. Brown, Richard A. Billington**

4 School of Biological and Marine Sciences, University of Plymouth, Drake Circus,

5 Plymouth, PL4 8AA, UK

6 *Corresponding Author: ahmad.mansur@plymouth.ac.uk

7 **Abstract**

8 Brown seaweeds have been highlighted for their abundant production of bioactive
9 substances that may provide novel drugs or drug scaffolds for a range of diseases,
10 particularly cancer. Indeed, a number of promising compounds that can modulate
11 growth arrest or apoptosis have already been isolated. As previous work has
12 highlighted seasonal differences in concentrations of secondary metabolites, this study
13 aimed to evaluate seasonal variation in the cytotoxic anticancer activity of *Cystoseira*
14 *tamariscifolia* extracts. Primary and secondary metabolites were measured using
15 colourimetric techniques and extracts were exposed to human leukaemia, HL60 and
16 THP-1, and human prostate cancer PC3 cell lines *in vitro* and cell viability was
17 measured using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide
18 (MTT) assay. The results confirm a large but inconsistent seasonal variation in the
19 concentrations of biomolecules, although, generally, levels were higher in spring and
20 summer. IC₅₀ values for cytotoxicity also showed variability both seasonally and with
21 extraction protocol, but again, this was inconsistent. These data suggest that, when
22 collecting brown algae in order to isolate novel bioactive compounds, repeated
23 sampling across annual or biotic cycles of the seaweed and the use of multiple solvent
24 extraction methods could give a greater possibility of discovering bioactive-lead
25 compounds.

26

27 Keywords: Anticancer; Chemical metabolites; Seasonal variation; Seaweed.

28 Introduction

29 Cancer has become one of the major causes of premature death in an ageing
30 population. With the risk of developing cancer approaching 50% by the age of 80,
31 combined with current life expectancy figures that are increasing in developing
32 countries, cancer is likely to remain a major affliction to patients and a challenge to
33 healthcare services. Prostate cancer remains one of the leading causes of cancer-
34 related death, while 6.8 million cancer deaths reported globally in 2018, 1.8 million of
35 which were caused by prostate cancer only. Not far from that, leukaemia of various
36 types, also claiming many lives around the world every year, especially children (Siegel
37 et al. 2019; Bray et al. 2018), which makes the search for a cure for these fatal
38 diseases a noble goal of our research. The most common treatments for cancer are
39 still surgery, radiotherapy and chemotherapy, including direct and indirect hormonal
40 and immune therapies (Coffelt and de-Visser 2015; Senthebane et al. 2017).
41 Currently, most of the chemotherapeutic strategies used to treat different cancers are
42 not completely effective and are often associated with severe side effects that can
43 significantly reduce the quality of life of the patient. Thus, much research is being
44 dedicated to finding novel compounds that show clinical efficacy with reduced side
45 effects, with much of this research concentrated on natural products including those
46 from marine algae (Anand et al. 2016). Bioactive compounds from brown seaweeds
47 have been highlighted for their importance as anticancer agents (Ashwini et al. 2016)
48 and most of these compounds are generally safe and are abundant in brown
49 seaweeds, which make them a promising potential source of therapeutic compounds
50 (Zubia et al. 2009). A wide range of these bioactive ingredients has been reported to
51 have anti-tumour activity. Polyphenols have been linked with anticancer activities of
52 seaweed crude extracts, for instance, phloroglucinol and dioxinodehydroeckol from
53 brown algae display an antiproliferative activity against colon and breast tumours
54 (Lopes-Costa et al. 2017; Murphy et al. 2014). Similarly, phloroglucinol from an
55 *Ecklonia cava* extract showed apoptosis enhancing effects on the MCF-7 human
56 breast cancer cell line (Kong et al. 2009). Furthermore, crude extracts from brown
57 algae *Palmaria palmata*, *Laminaria setchellii* and *Macrocystis integrifolia* exhibited
58 cytotoxicity against the cervical cancer HeLa cell line (Yuan and Walsh 2006). The
59 anticancer or cytotoxic mechanism of polyphenols is still unclear; however, it has been

60 suggested that polyphenols may inhibit the formation of mitotic spindles by preventing
61 normal microtubule formation, decreasing angiogenesis and cell adhesion and
62 invasion (Olivares-Bañuelos et al. 2019; Zenthoefer et al. 2017; Wells et al. 2016).

63 Seaweed flavonoids such as flavones, catechins, chalcones, flavanols and
64 isoflavonoids are well recognised to exhibit a potent anticancer activity. For example,
65 flavonoids extracted in methanol, chloroform and ethyl acetate from the brown algae
66 *Padina gymnospora*, and *Sargassum wightii* reduced the cell viability of the human
67 cancer cell lines A549, HCT-15 and PC-3 (Murugan and Iyer 2013). One possible
68 mechanism of flavonoids may be exerted by altering steroid hormone production
69 through inhibition of aromatase, preventing the development of cancer cells (Sithranga
70 et al. 2010).

71 Polysaccharides from seaweeds, such as sulphated laminarin, porphyran and
72 fucoidan are naturally active as anticancer agents against human breast, gastric, liver,
73 prostate, colon, lung and urinary bladder cancers and melanoma (Yamasaki-
74 Miyamoto et al. 2009; Alekseyenko et al. 2007; Lowenthal and Fitton JH 2015). The
75 suggested mechanism is that polysaccharides cause cell cycle arrest at the G0/G1
76 phase (Senthilkumar et al. 2013). A number of proteins have been isolated from
77 seaweed such as lectins, glycoproteins and phycobiliproteins which are the most
78 studied as bioactive anticancer compounds against human lymphoma MCL, colon
79 cancer Colo201 cells and cervix cancer HeLa cell lines (Pangestuti and Kim 2015;
80 Harnedy and FitzGerald 2011). They may be able to induce cell cycle arrest and inhibit
81 the IGF-IR signalling pathway (Park et al. 2013).

82 Given the wide range of compounds that they contain and initial evidence on the
83 activities of those compounds, brown algae are a promising source of novel
84 pharmaceutical compounds. Among the brown seaweeds, the genus *Cystoseira*
85 contains about 40 species distributed widely in the Atlantic Ocean and Mediterranean
86 Sea (De Sousa et al. 2017). Crude extracts of different species have shown anti-
87 tumour activity against a wide variety of cancer cell lines. *Cystoseira* spp. provide a
88 good source of polyphenolic compounds (De Sousa et al. 2017), which are well known
89 to show cytotoxic activities against tumour cells (Gutiérrez-Rodríguez et al. 2017).
90 Polyphenols from these algae include those with unique structural features and exhibit

91 effective anticancer activities (Yong-Xin et al. 2011), for example, hexane fractions
92 extracted from *Cystoseira crinita* and *C. stricta* showed 87% and 50% reductions in
93 proliferation respectively in colorectal carcinoma (Caco2) cells after 72 h exposure with
94 200 $\mu\text{g mL}^{-1}$ extracts and the antiproliferative effect correlated with their polyphenol
95 and flavonoid contents (Alghazeer et al. 2016).

96 Several studies have demonstrated that the bioactivity properties of brown seaweeds
97 vary seasonally (e.g. Stengel et al. 2011). These studies have indicated that changes
98 in biotic and abiotic factors between seasons and locations affect the production of
99 their bioactive compounds (Celis-Plá et al. 2016; Zatelli et al. 2018). However, as far
100 as we are aware, while concentrations of secondary metabolites have been measured,
101 no assessment of seasonal variations in cytotoxicity against cancer cell lines has been
102 undertaken. Here, we investigate the seasonal variation of the chemical composition
103 and the cytotoxic activity of extracts from *C. tamariscifolia* collected from south-west
104 England. We show that extracts from *C. tamariscifolia* harvested in the winter and
105 spring contained higher levels of secondary metabolites in accordance with previous
106 work (Celis-Plá et al. 2016). Furthermore, we tested cytotoxicity of our extracts against
107 leukaemia and prostate cancer cell lines and we show seasonal variation in the
108 cytotoxic activity of four different solvent extracts against these cell lines. These
109 cytotoxic activities can range from complete cell death to undetectable depending on
110 the season and thus, we suggest that when considering collecting marine algae from
111 temperate seas in order to search for bioactive compounds, a consideration of
112 seasonality is made and samples are collected during different seasons to maximise
113 the possibility of finding bioactive compounds.

114

115 **Material and methods**

116 **Collection of *Cystoseira tamariscifolia*:**

117 *Cystoseira tamariscifolia* was collected by hand from the shallow subtidal at Hannafore
118 Point, Cornwall, UK (DD N 50.342234, DD W -4.453528) in mid-June (summer), mid-
119 October (autumn), and early March (winter) and early May (spring). The collected
120 samples were transferred to the laboratory in polythene bags kept in an icebox. On
121 arrival, the samples were rinsed with distilled water, cleaned of adhering debris and
122 epiphytes and then freeze-dried at -20°C using Edwards super Modulyo freeze dryer.

123 **Solvent extraction**

124 Extracts were made in chloroform, methanol (100% and 70%) and water. Fifty grams
125 of freeze-dried samples were mixed with solvents (1:10, w/v), and then homogenised
126 for 2 min by using an IKA T10B Ultra-Turrax disperser at 24°C . The extract was then
127 stirred for 3 min, centrifuged (5,000 g, 10 min, room temperature [RT]) and the
128 supernatants recovered. The extracts were dried at 40°C under vacuum. Solvents
129 were used to re-suspend all extracts, and then the extracts were stored at -20°C for
130 biological activity screening (Vizetto-Duarte et al. 2016).

131 **Total polyphenol content (TPC)**

132 The Folin-Ciocalteu (F-C) colourimetric technique (Velioglu et al. 1998) was used to
133 determine total phenolic concentrations (TPC). Briefly, 5 μL of the extracts at 0.1, 1
134 and 10 mg/mL were mixed with 100 μL of 10-fold diluted F-C reagent, incubated at
135 RT for 5 min and mixed with 100 μL of sodium carbonate (75 g L^{-1} , w/v). Following a
136 90 min incubation period at RT, absorbance was measured at 725 nm on a microplate
137 reader (Omega, BMG Labtech). The concentration of TPC was calculated as
138 phloroglucinol equivalents (PGE) utilising a calibration curve prepared with
139 phloroglucinol standard solutions and is expressed as mg PGE g^{-1} dry weight.

140 **Total flavonoid content (TFC)**

141 Flavonoid concentrations were quantified according to the method described by Ahn
142 et al. (2007), with modifications. Briefly, 50 μL of 2% (w/v) aluminium chloride-ethanol
143 solution was added to 50 μL of the extracts at 0.1, 1, 10 mg mL^{-1} . After one hour at RT,

144 the absorbance was measured at 420 nm on a microplate reader (Omega, BMG
145 Labtech). Quercetin was used as a standard, and results are expressed as mg of
146 quercetin equivalents (QE) g⁻¹ dry weight.

147 **Total polysaccharide content**

148 Total polysaccharide concentrations were measured by the phenol-sulfuric acid
149 method (Masuko et al. 2005). 50 µL of samples at the concentrations 0.1, 1 and 10
150 mg/mL was added to 150 µL of sulfuric acid (96% reagent grade). The mixture was
151 incubated in a 96 well plate floating on a water bath at 90°C for 5 min. 30 µL of 5%
152 phenol was added to the mixture for another 5 min in the water bath. The plate was
153 then floated on cold water for additional 5 min to cool and the absorbance was
154 measured at 490 nm on a microplate reader (Omega, BMG Labtech). Glucose was
155 used as a standard and results are expressed as mg of glucose equivalents (G) g⁻¹ dry
156 weight.

157 **Total protein**

158 The BCA assay was carried out to determine total protein concentrations using the
159 Pierce BCA Protein Assay Kit (Thermo Fisher Scientific) in a 96-well-plate format. The
160 assay mixture contained 100 µL of the reagent and 100 µL of the sample; crude
161 extracts at concentrations 0.1, 1 and 10 mg mL⁻¹. A standard curve with serial bovine
162 serum Albumin BSA solutions (ranging from 0.1 µg mL⁻¹ to 100 µg mL⁻¹) was used for
163 calibration. The mixture was incubated at 37°C and absorbance was measured at 560
164 nm on a microplate reader (Omega, BMG Labtech).

165 **Cell line maintenance and culture**

166 The culturing of the human leukaemia cancer cell lines THP-1 and HL60 was as a
167 suspension and the human prostate cancer cell line (PC3) was as a monolayer. Cells
168 were obtained from the European Collection of Authenticated Cell Cultures (ECACC),
169 and divided every 3-4 d and were used regularly between passages 10-35. The cells
170 were cultured in RPMI 1640 complete growth medium supplemented with 10% (v/v)
171 fetal bovine serum and 2 mM L-glutamine and maintained at 37°C, under 5% CO₂ in a
172 humidified incubator.

173

174 **Cell vitality assay**

175 The cells were incubated at a density of 2×10^5 cells/well in a 96-well microplate for 72
176 h at 37°C with an equal volume of assay medium containing thiazolyl blue tetrazolium
177 (2 mg mL^{-1} in PBS). The resulting formazan crystals were solubilised in 150 μL DMSO.
178 The absorbance was read at 540 nm using a microplate reader (Omega, BMG
179 Labtech). The IC_{50} values were calculated from a sigmoidal dose-response curve of
180 the data generated in SigmaPlot v. 13.0.

181 **Fluorescent staining for HL60 and THP-1 cells**

182 THP-1 and HL60 were seeded at 10^6 cells/ml in 24-well culture plates and treated with
183 100% methanol extracts of *C. tamariscifolia* for 48 h. Cells were centrifuged and
184 suspended in PBS. Propidium iodide ($5 \mu\text{g mL}^{-1}$ in PBS; Sigma Aldrich) was used for
185 staining cells for 5 min at room temperature in the dark and images were captured
186 using a Nikon fluorescence microscope.

187 **Flow Cytometry Analysis**

188 Flow cytometry was performed to determine the apoptotic effect of *C. tamariscifolia*
189 methanol extracts on human leukaemia cell lines HL 60 and THP-1. Cells were seeded
190 at a density of 10^6 mL^{-1} and exposed to extracts at a concentration of $150 \mu\text{g mL}^{-1}$
191 before being incubated at 37°C in a 5% CO_2 incubator for 24h. Cells were washed in
192 PBS and fixed in cold 70% ethanol for 1h. Cells were incubated in $50 \mu\text{g mL}^{-1}$ RNase
193 and $50 \mu\text{g mL}^{-1}$ fluorescein diacetate (FITC; Sigma Aldrich) solution for 30 min before
194 cells were analysed by Becton Dickinson FACSCalibur Flow Cytometer. All
195 experiments were performed in three replicates.

196

197

198 **Results**

199 **Effects of extraction method and season on the biochemical composition.**

200 To assess the cytotoxic bioactivity of *Cystoseira tamariscifolia* extracts from the south-
201 western coast of England, various extraction solvents were applied to material
202 collected in all four seasons, and the concentrations of primary and secondary
203 metabolites were measured (Table 1).

204 100% methanol extracts of *C. tamariscifolia* contained the highest yields of primary
205 and secondary metabolites. Polyphenol content was found to be higher in 100%
206 methanol extracts with a concentration of approximately 100 mg g⁻¹ DW, while
207 chloroform extracts showed the highest content of flavonoids with up to 45 mg g⁻¹ DW.
208 Polysaccharide content ranged from 10 and 45 mg g⁻¹ DW in the water and 100%
209 methanol extracts, respectively (Table 1).

210 The spring and summer extracts generally had higher concentrations of metabolites
211 while in autumn we recorded the lowest concentration of total polyphenols and
212 flavonoids. *C. tamariscifolia* crude extracts had low protein content compared with the
213 previous bioactive compounds. The greatest concentration of protein was in the
214 autumn in both 100% methanol and water extracts with concentrations of roughly 19
215 and 6 mg g⁻¹ DW respectively, while the chloroform extract could not be used due to
216 interference with the assay.

217 **Extraction method and seasonality impact the anticancer activity of *C.***
218 ***tamariscifolia* extracts.**

219 We were interested in seeing the effect of the extraction method and season on any
220 potential cytotoxic activity as *Cystoseira* spp. extracts have been shown to contain
221 various bioactive compounds, particularly polyphenols that show cytotoxicity against
222 cancer cell lines. Three different cell lines, HL-60, THP-1 and PC3 cells, representing
223 two leukaemia-derived lines and a prostate cancer line respectively, were treated with
224 increasing concentrations of *C. tamariscifolia* extracts and cell vitality was assessed
225 via the MTT assay. Cells were treated with extracts from all four seasons extracted via
226 all four extraction regimes separately (Figures 1, 2 and 3) and IC₅₀ values for cell
227 vitality were calculated (Table 2).

228 The solvent used to make the extracts influenced the cytotoxic activity of the extracts.
229 Generally, 100% methanol and chloroform extracts had the most cytotoxic effect on
230 the three cell lines tested with some differences between them (Table 2). The 100%
231 MeOH extract was approximately 17 times more cytotoxic in HL-60 compared to water
232 extracts, while both 100% methanol and chloroform extracts had almost identical
233 activity in HL60 and THP-1. Extracts made in water were the most cytotoxic in PC3
234 cells (Figure 2) with IC_{50} values approximately 2-5 fold higher than for the other
235 solvents whereas water extracts were much less cytotoxic than the other solvents for
236 HL-60 and THP-1 cells, being between approximately 2-130 fold less cytotoxic
237 (Figures 1, 3).

238 In addition to the extraction method effect, there was a clear seasonality in the
239 cytotoxic effects on the three different cell lines. Summer and spring extracts showed
240 the most potent cytotoxic effect on THP-1 and HL-60 cells with approximately 8 times
241 higher cytotoxicity compared to the autumn and winter extracts. There was, however,
242 an exception in that winter extracts were also highly cytotoxic against the prostate
243 cancer cells PC3.

244 HL-60 cells showed the greatest sensitivity to *C. tamariscifolia* extracts with a mean
245 IC_{50} over all seasons and extracts of $80.61 \pm 21.74 \mu\text{g mL}^{-1}$ while the similar THP-1
246 model showed a mean IC_{50} over all seasons and extracts of $199.78 \pm 37.23 \mu\text{g mL}^{-1}$
247 and PC3, $162.15 \pm 36.11 \mu\text{g mL}^{-1}$. The highest cytotoxic activity was seen in the
248 summer extracts in 100% and 70% MeOH and 100% chloroform on HL-60 cells (IC_{50}
249 values; 2.32 ± 0.21 , 7.34 ± 0.30 and $7.92 \pm 0.12 \mu\text{g mL}^{-1}$ respectively; Figure 1). These
250 values show that the cytotoxic activity of these crude extracts is very high against HL-
251 60 cells and is between approximately 17 and 62 fold higher than the corresponding
252 effects on THP-1 and PC3 cells, respectively.

253 **Methanol extracts of *C. tamariscifolia* induce apoptosis in HL60 and THP-1 Cell** 254 **lines**

255 To investigate whether the cytotoxicity activity of *C. tamariscifolia* extracts is related to
256 the induction of apoptosis, HL60 and THP-1 cells were exposed to a concentration of
257 150 mgmL^{-1} of methanol extracts of *C. tamariscifolia* for 24 h and nuclear
258 morphological changes of HL60 and THP-1 cells were observed using propidium

259 iodide staining (Fig. 4). Compared with the normal nuclear morphology of the control
260 cells, the cells treated with *C. tamariscifolia* extracts presented typical morphological
261 characteristics of apoptosis, including nuclear fragmentation. Additional confirmation
262 of apoptosis induced by methanol extracts of *C. tamariscifolia* was performed by flow
263 cytometry-based on fluorescent diacetate (FITC) staining as shown in figure 5. Control
264 cells that had not been treated with extract displayed a largely homogeneous
265 population with >99% of cells being vital. Upon treatment with the extract, this dropped
266 to 36.28 and 17.54% in HL-60 and THP-1 cells respectively and an increase in cells in
267 both necrosis and late apoptosis was observed with c. 50% of cells in late apoptosis
268 in both lines confirming the data from the cell vitality assay.

269

270 Discussion

271 In temperate seas, one would expect that seasonal differences would have a large
272 influence on the concentrations of likely bioactive compounds and thus bioactivity of
273 those extracts. Surprisingly, little work has been performed on these links. Here we
274 have investigated the effects of season and extraction method on primary/secondary
275 metabolite concentrations in extracts and their cytotoxic activity against three cancer
276 cell lines.

277 Methanol (100%) was generally the most effective solvent, extracting the highest
278 levels of the four metabolite classes, although all four solvents gave detectable levels
279 of all the investigated metabolites, the exception being chloroform that interfered with
280 the BCA assay for proteins. Concentrations of polyphenols and flavonoids were high,
281 a result in accordance with previous research showing brown algae to be good sources
282 of these two metabolite classes (Thomas et al. 2011; Alghazeer et al. 2016). A similar
283 pattern of results was obtained by Mhadhebi et al. (2011) and Yegdaneh et al. (2016)
284 who also concluded that methanol and chloroform extracts contain high concentrations
285 of metabolites, especially polyphenols. This result can be explained by the difference
286 in secondary metabolite polarity. For example, the high variation in the structures and
287 both hydrophilic and hydrophobic parts of polyphenols (Li et al. 2011) allow them to be
288 extracted typically in polar solvents including methanol and water but some can also
289 be extracted in low polarity solvents such as chloroform (Airanthi et al. 2011; Vizetto-
290 Duarte et al. 2016), which may explain the high levels of polyphenols in chloroform
291 extracts of *Cystoseira*. Protein represented the least common of the four classes of
292 metabolite as might be expected in brown algae.

293 As might be expected, there was a clear seasonality to the levels of the metabolites
294 although this varied between compound and extraction method (Celis-Plá et al. 2016;
295 Rickert et al. 2016; Cikoš et al. 2018). *C. tamariscifolia* begins to grow in late winter
296 with most growth in the spring and summer before stopping in autumn. Polyphenols
297 were highest in each extraction method in spring and summer except for chloroform
298 extracts from winter. This is consistent with observations by Abdala-Díaz et al. (2006),
299 who showed that polyphenol concentration in the tissue of *C. tamariscifolia* (in Spain
300 from June 1988 to July 2000) ranged from 2% in the winter to 8% in summer.
301 Polyphenols from species of the brown algal order Dictyotales, Fucales and

302 Laminariales, have been correlated with the tissue age and stage of the life cycle and
303 there were significantly lower levels in new branches in late winter than older, bigger
304 branches in summer (Denton et al. 1990; Mannino et al. 2014). For flavonoids,
305 polysaccharides and protein, the picture was more complex with winter and autumn
306 often offering at least one higher value than in spring/autumn. The complexity of the
307 variation of flavonoid, polysaccharide and protein levels extracted from *C.*
308 *tamariscifolia* has been highlighted in previous studies. For instance, a study by
309 Abdala-Díaz et al. (2006) found that spring and winter extracts of *C. tamariscifolia*
310 showed the highest concentration of flavonoids, however, contradictory results were
311 reported by Ramah et al. (2014) and Mannino et al. (2014) who showed that the
312 flavonoid peak occurred in summer. During the year, the highest levels of
313 polysaccharides in *C. tamariscifolia* were observed in summer, supporting the
314 observations of Teas et al. (2013) and Hurtado and Critchley (2018). These authors
315 noted that *Undaria pinnatifida*, *Ascophyllum nodosum* and *Fucus vesiculosus* showed
316 the maximum levels of polysaccharides in summer. However, other results by
317 Skriptsova (2016) revealed that *Saccharina japonica* and *Sargassum pallidum* showed
318 maximum polysaccharide content in autumn. The protein content in *C. tamariscifolia*
319 varied from 3.28 ± 0.36 to 19.57 ± 1.42 mg g⁻¹ dry weight, and it differs from season to
320 season and according to the extraction method. In general, the highest protein values
321 in *C. tamariscifolia* were found in autumn and spring, while the lowest were found
322 winter and summer. Similar results were reported by Kim et al. (2011) and Pangestuti
323 and Kim (2015), who suggested that minimum protein concentration in summer could
324 be linked with protein destruction.

325 Given the temperate nature of the collection site, autumn and winter offer lower water
326 temperatures, fewer hours of sunlight and increased wave action that will lead to
327 decreased growth and investment into primary and secondary metabolites (Fleurence
328 and Levine 2016). This may also coincide with decreased pressures of herbivory
329 during these seasons and less requirement for secondary metabolites (Duffy and Hay
330 1990). We found that new growth of *C. tamariscifolia* was already clearly apparent in
331 early March (winter) with many fresh new phosphorescent fronds present presumably
332 ready for the increased sunlight hours of spring and summer. Conversely, in October
333 (autumn) most of the thalli were showing early signs of senescence where the main

334 axis and the primary laterals had elongated while the branches were relatively short
335 and often covered in epiphytes. Despite this, autumn algae still contained high levels
336 of some metabolites although polyphenol levels were generally lowest in autumn, but
337 this depended on the extraction method. One explanation of this result could be
338 attributed to the increase in the need for the secondary metabolites as a protection
339 mechanism against increased herbivory in summer (Duffy and Hay 1990; Jormalainen
340 and Honkanen 2008). Another reason for the increase in secondary metabolites in
341 summer may be due to the photoprotective role against the high radiation dosages in
342 summer daylight (Connan et al. 2004). They also contribute to protection against
343 oxidative stress, bacterial infection and epiphytes and perform a role in algal
344 reproduction which all increase in summer (Plouguerné et al. 2006; Ferreres et al.
345 2012; Jennings and Steinberg 1997; Thomas et al. 2011).

346 In line with Duarte (2016), who demonstrated that a hexane extract of *C. tamariscifolia*
347 had a high level of cytotoxicity against AGS, HCT-15 and HepG2 cell lines with IC₅₀
348 values of 32.36, 23.59 and 13.15 µg mL⁻¹ respectively, we showed that our extracts
349 possess cytotoxic activity against cancer cell lines. In particular, our extracts were
350 considerably more cytotoxic against HL-60 cells and our evidence suggests that cell
351 death is induced via apoptosis. There is a large seasonal variation not only in the levels
352 of primary and secondary metabolites but also a significant seasonal variation in their
353 cytotoxic effects. In particular, our results clearly underline the need to test extracts
354 from different seasons and using different solvent extraction protocols. Of particular
355 interest, our data suggest that generalisations cannot be drawn with respect to season
356 or extraction method and that there is no best season or extraction method when
357 results are compared between cell types. For example, the difference between the
358 effect of the summer extracts in organic solvents for HL-60 cells and the aqueous
359 winter extracts on PC3 cells highlights this difference. Indeed, we carried out statistical
360 analyses to examine any potential interactions between metabolite composition and
361 the cytotoxicity of the extracts with season and extraction method (data not shown).
362 There was no clear interaction that could link the cytotoxic effects to any metabolite
363 group which might also suggest that various bioactive compounds may be present in
364 the extracts and that their relative amounts may vary with both season and extraction
365 method. It would be tempting to assume that summer (or an early or midpoint of the

366 growing season) might be the best time to collect material, but our results show that
367 this is not necessarily the case in all cell lines or extracts. We have not fully fractionated
368 our extracts to isolate the specific compound(s), but it is likely that different compounds
369 are responsible for the observed effects and that these change levels with season.
370 Equally, we cannot discount that there may be synergistic interactions between
371 compounds that only become apparent in different seasons. We believe that it is
372 possible that many promising extracts potentially containing novel
373 cytotoxic/chemotherapeutic agents may have been discarded in the past where
374 samples were taken in a single season, extracted using a single solvent or tested using
375 a single cell line.

376 We suggest that, where possible, samples are taken at multiple times of the year that
377 represent different phases in the growth or annual life cycle of the plant/species in
378 question. Given that many of the bioactive compounds with potential medical use are
379 probably produced in response to some type of biotic or abiotic stress, it may also be
380 good practice to include the presence of any obvious stressors (seasonal temperature,
381 irradiance, herbivores, for example) into any consideration of sampling times so as to
382 maximize the possibility of finding novel compounds.

383

384

385 **References**

- 386 Abdala-Díaz RT, Cabello-Pasini A, Pérez-Rodríguez E, Álvarez RC, Figueroa F L
387 (2006) Daily and seasonal variations of optimum quantum yield and phenolic
388 compounds in *Cystoseira tamariscifolia* (Phaeophyta). *Mar Biol* 148:459-465
- 389 Ahn MR, Kumazawa S, Usui Y, Nakamura J, Matsuka M, Zhu F, Nakayama T, (2007)
390 Antioxidant activity and constituents of propolis collected in various areas of China.
391 *Food Chem* 101:1383-1392
- 392 Airanthi MWA, Hosokawa M, Miyashita K (2011) Comparative antioxidant activity of
393 edible Japanese brown seaweeds. *J Food Sci* 76:C104-C111
- 394 Alekseyenko TV, Zhanayeva SY, Venediktova AA, Zvyagintseva TN, Kuznetsova TA,
395 Besednova NN, Korolenko TA (2007) Antitumor and antimetastatic activity of fucoidan,
396 a sulfated polysaccharide isolated from the Okhotsk Sea *Fucus evanescens* brown
397 alga. *Bull Exp Biol Med* 143:730-732
- 398 Alghazeer R, Enaeli M, Howell NK (2016) Anticancer and Antioxidant Activities of
399 Some Algae from Western Libyan Coast. *Pharmacol Toxicol. Ver. 1*, 2016090018
- 400 Anand N, Rachel D, Thangaraju N, Anantharaman P (2016) Potential of marine algae
401 (seaweeds) as a source of medicinally important compounds. *Plant Genet Resour*
402 14:303-313
- 403 Ashwini S, Babut S, Saritha MS (2016) Seaweed extracts exhibit anticancer activity
404 against Hela cell lines. *Int J Curr Pharm Res* 9:114-117
- 405 Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A (2018) Global cancer
406 statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36
407 cancers in 185 countries. *CA Cancer J Clin* 68:394-424
- 408 Celis-Plá PS, Bouzon ZL, Hall-Spencer JM, Schmidt EC, Korbee N, Figueroa FL
409 (2016) Seasonal biochemical and photophysiological responses in the intertidal
410 macroalga *Cystoseira tamariscifolia* (Ochrophyta). *Mar Environ Res* 115:89-97
- 411 Cikoš AM, Jokić S, Šubarić D, Jerković I (2018) Overview on the application of modern
412 methods for the extraction of bioactive compounds from marine macroalgae. *Mar*
413 *Drugs* 16:348
- 414 Coffelt SB, de Visser KE (2015) Immune-mediated mechanisms influencing the
415 efficacy of anticancer therapies. *Trends Immunol* 36:198-216
- 416 Connan S, Goulard F, Stiger V, Deslandes E, Gall EA (2004) Interspecific and
417 temporal variation in phlorotannin levels in an assemblage of brown algae. *Bot Mar*
418 47:410-416
- 419 De Sousa CB, Gangadhar KN, Macridachis J, Pava M, Morais, TR, Campino L, Varela
420 J, Lago JHG (2017) *Cystoseira* algae (Fucaceae): update on their chemical entities
421 and biological activities. *Tetrahedron: Asymmetry* 28:1486-1505

- 422 Denton, A (1990) Size specific concentrations of phlorotannins in three species of
423 *Fucus*. Mar Ecol Prog Ser 65:103-104
- 424 Duarte, CVG (2016) Biomedical properties of *Cystoseira* species: Insights into nutra-
425 and pharmaceutical applications. PhD Thesis, University of Algarve, Portugal.
426 <http://hdl.handle.net/10400.1/9004>
- 427 Ferreres F, Lopes G, Gil-Izquierdo A, Andrade PB, Sousa C, Mouga T, Valentão P
428 (2012) Phlorotannin extracts from Fucales characterised by HPLC-DAD-ESI-MSn:
429 approaches to hyaluronidase inhibitory capacity and antioxidant properties. Mar Drugs
430 10:2766-2781
- 431 Fleurence J, Levine I. eds (2016) Seaweed in health and disease prevention. Elsevier,
432 Amsterdam. p290
- 433 Gutiérrez-Rodríguez AG, Juárez-Portilla C, Olivares-Bañuelos T, Zepeda RC (2017)
434 Anticancer activity of seaweeds. Drug Discov Today 23:434-447
- 435 Harnedy PA, FitzGerald RJ (2011) Bioactive proteins, peptides, and amino acids from
436 macroalgae. J Phycol 47:218–232
- 437 Hay ME, Duffy JE, Fenical W (1990) Host-plant specialisation decreases predation on
438 a marine amphipod: a herbivore in plant's clothing. Ecology 71:733-743
- 439 Hurtado AQ, Critchley AT (2018) A review of multiple biostimulant and bioeffector
440 benefits of AMPEP, an extract of the brown alga *Ascophyllum nodosum*, as applied to
441 the enhanced cultivation and micropropagation of the commercially important red algal
442 carrageenophyte *Kappaphycus alvarezii* and its selected cultivars. J Appl Phycol
443 30:2859-2873
- 444 Jennings JG, Steinberg PD (1997) Phlorotannins versus other factors affecting
445 epiphyte abundance on the kelp *Ecklonia radiata*. Oecologia 109:461-473
- 446 Jormalainen V, Honkanen T (2008) Macroalgal chemical defences and their roles in
447 structuring temperate marine communities. In: Amsler CD (ed) Algal chemical ecology,
448 Springer, Berlin, Heidelberg. p57- 89
- 449 Kong CS, Kim JA, Yoon NY, Kim SK (2009) Induction of apoptosis by phloroglucinol
450 derivative from *Ecklonia cava* in MCF-7 human breast cancer cells. Food Chem
451 Toxicol 47:1653–1658
- 452 Lopes-Costa E, Abreu M, Gargiulo D, Rocha E, Ramos AA (2017) Anticancer effects
453 of seaweed compounds fucoxanthin and phloroglucinol, alone and in combination with
454 5-fluorouracil in colon cells. J Toxicol Environ Health 80:776-787
- 455 Lowenthal RM, Fitton JH (2015) Are seaweed-derived fucoidans possible future anti-
456 cancer agents? J Appl Phycol 27:2075-2077
- 457 Mannino A, Vaglica V, Oddo E (2013) Seasonal variation in total phenolic content of
458 *Dictyopteris polypodioides* (Dictyotaceae) and *Cystoseira amentacea* (Sargassaceae)
459 from the Sicilian coast. FI Med 24:39–50

460 Masuko T, Minami A, Iwasaki N, Majima T, Nishimura SI, Lee YC (2005) Carbohydrate
461 analysis by a phenol-sulfuric acid method in microplate format. *Anal Biochem* 339:69-
462 72

463 Mhadhebi L, Laroche-Clary AA, Robert J, Bouraoui A (2011) Anti-inflammatory anti-
464 proliferative and anti-oxidant activities of organic extracts from the Mediterranean
465 seaweed *Cystoseira crinita*. *Afr J Biotechnol* 10:16682-16690

466 Murphy C, Hotchkiss S, Worthington J, McKeown SR (2014) The potential of seaweed
467 as a source of drugs for use in cancer chemotherapy. *J Appl Phycol* 26:2211-2264

468 Murugan K, Iyer VV (2013) Differential growth inhibition of cancer cell lines and
469 antioxidant activity of extracts of red, brown, and green marine algae. *In Vitro Cell Dev*
470 *Biol-Animal* 49:324–334

471 Olivares-Bañuelos T, Gutiérrez-Rodríguez AG, Méndez-Bellido R, Tovar-Miranda R,
472 Arroyo-Helguera O, Juárez-Portilla C, Meza-Menchaca T, Aguilar-Rosas LE,
473 Hernández-Kelly LC, Ortega A, Zepeda RC (2019) Brown Seaweed *Egregia*
474 *menziesii*'s Cytotoxic Activity against Brain Cancer Cell Lines. *Molecules* 24:260.

475 Pangestuti R, Kim SK(2015) Seaweed proteins, peptides, and amino acids Seaweed
476 Sustainability Food and Non-Food Applications, Elsevier p.125-140

477 Park H, Kim H, Kim J, Nam T (2013) Induction of apoptosis and the regulation of ErbB
478 signalling by laminarin in HT-29 human colon cancer cells. *Int J Mol Med* 32:291–295

479 Plouguerné E, Le Lann K, Connan S, Jechoux G, Deslandes E, Stiger-Pouvreau V
480 (2006) Spatial and seasonal variation in density, reproductive status, length and
481 phenolic content of the invasive brown macroalga *Sargassum muticum* (Yendo)
482 Fensholt along the coast of Western Brittany (France). *Aquat Bot* 85:337-344

483 Ramah S, Etwarising L, Auckloo N, Gopeechund A, Bhagooli R, Bahorun T (2014)
484 Prophylactic antioxidants and phenolics of seagrass and seaweed species: a seasonal
485 variation study in a Southern Indian Ocean Island, Mauritius. *Int J Med Update* 1:27-
486 37

487 Rickert E, Wahl M, Link H, Richter H, Pohnert G (2016) Seasonal variations in surface
488 metabolite composition of *Fucus vesiculosus* and *Fucus serratus* from the Baltic Sea.
489 PLoS One 11:e0168196

490 Senthebane DA, Rowe A, Thomford NE, Shipanga H, Munro D, Al Mazeedi MA,
491 Almazyadi HA, Kallmeyer K, Dandara C, Pepper MS, Parker MI (2017) The role of
492 tumour microenvironment in chemoresistance: To survive, keep your enemies closer.
493 Int J Mol Sci 18:1586

494 Senthilkumar K, Manivasagan P, Venkatesan J, Kim SK (2013) Brown seaweed
495 fucoidan: biological activity and apoptosis, growth signalling mechanism in cancer. Int
496 J Biol Macromol 60:366-374

497 Siegel RL, Miller KD, Jemal A (2019) Cancer statistics, 2019. CA Cancer J Clin 69:7-
498 34.

499 Sithranga BN, Kathiresan K (2010) Anticancer drugs from marine flora: An overview. J.
500 Oncol 2010 :2141-6.

501 Skriptsova AV (2016) Seasonal variations in the fucoidan content of brown algae from
502 Peter the Great Bay, sea of Japan. Russ J Mar Biol 42:351–356

503 Stengel DB, Connan S, Popper ZA (2011) Algal chemodiversity and bioactivity:
504 sources of natural variability and implications for commercial application. Biotechnol
505 Adv 29:483–501

506 Teas J, Vena S, Cone D, Irhimeh M (2013) The consumption of seaweed as a
507 protective factor in the etiology of breast cancer: proof of principle. J Appl Phycol
508 25:771-779

509 Thomas NV, Kim SK (2011) Potential pharmacological applications of polyphenolic
510 derivatives from marine brown algae. Environ Toxicol Pharmacol 32:325–335

511 Velioglu YS, Mazza G, Gao I, Oomah BD (1998) Antioxidant activity and total phenolics
512 in selected fruits, vegetables, and grain products. J Agric Food Chem 46:4113–4117

513 Vizetto-Duarte C, Custodio L, Acosta G, Lago JH, Morais TR, Bruno de Sousa C,
514 Gangadhar KN, Rodrigues MJ, Pereira H, Lima RT, Vasconcelos MH, Barreira L,
515 Rauter AP, Albericio F, Varela J (2016) Can macroalgae provide promising anti-
516 tumoral compounds? A closer look at *Cystoseira tamariscifolia* as a source for
517 antioxidant and anti-hepatocarcinoma compounds. PeerJ 4:e1704

518 Wells ML, Potin P, Craigie JS, Raven JA, Merchant SS, Helliwell KE, Smith AG,
519 Camire ME, Brawley SH (2016) Algae as nutritional and functional food sources:
520 revisiting our understanding. J Appl Phycol 29: 949–982

521 Yamasaki-Miyamoto Y, Yamasaki M, Tachibana H, Yamada K (2009) Fucoidan
522 induces apoptosis through activation of caspase-8 on human breast cancer MCF-7
523 cells. J Agric Food Chem 57:8677-8682


- 524 Yegdaneh A, Ghannadi A, Dayani L (2016) Chemical constituents and biological
525 activities of two Iranian *Cystoseira* species. Res Pharm Sci 11:311-317
- 526 Yong-Xin L, Isuru W, Yong L, Se-Kwon K (2011) Phlorotannins as bioactive agents
527 from brown algae. Process Biochem 46:2219–2224
- 528 Yuan YV, Walsh NA (2006) Antioxidant and antiproliferative activities of extracts from
529 a variety of edible seaweeds. Food Chem Toxicol 44:1144–1150
- 530 Zatelli GA, Philippus AC, Falkenberg M (2018) An overview of odoriferous marine
531 seaweeds of the *Dictyopteris* genus: insights into their chemical diversity, biological
532 potential and ecological roles. Rev Bras Farmacogn 28:243-260
- 533 Zenthoefer M, Geisen U, Hofmann-Peiker K, Fuhrmann M, Kerber J, Kirchhöfer R,
534 Hennig S, Peipp M, Geyer R, Piker L, Kalthoff H (2017) Isolation of polyphenols with
535 anticancer activity from the Baltic Sea brown seaweed *Fucus vesiculosus* using
536 bioassay-guided fractionation. J Appl Phycol 29:2021-2037
- 537 Zubia M, Fabre MS, Kerjean V, Lann KL, Stiger-Pouvreau V, Fauchon M, Deslandes
538 E (2009) Antioxidant and antitumoural activities of some Phaeophyta from Brittany
539 coasts. Food Chem 116:693-701
- 540

Table 1: Metabolite concentrations in *Cystoseira tamariscifolia* as a function of season and extraction solvent. DW: dry weight. Total polyphenol Content: milligram phloroglucinol acid equivalents per gram dry weight; Total flavonoid Content: milligram Quercetin equivalents per gram dry weight. Total polysaccharide: milligram Glucose equivalents per gram dry weight; Total protein: milligram Bovine Serum Albumin equivalents per gram dry weight. Values are presented as mean \pm SD (n = 3).

Metabolite (mg g ⁻¹ DW)	Season	Extraction solvent			
		100%MeOH	70%MeOH	Water	Chloroform
Total polyphenol content	Summer	102.23 \pm 1.85	57.70 \pm 2.06	83.24 \pm 1.03	41.99 \pm 0.90
	Autumn	71.27 \pm 4.16	9.67 \pm 1.27	20.61 \pm 3.07	7.81 \pm 0.31
	Winter	71.16 \pm 4.40	31.20 \pm 1.19	14.62 \pm 0.91	48.89 \pm 1.91
	Spring	85.46 \pm 2.7	62.35 \pm 1.86	46.29 \pm 0.28	68.75 \pm 2.79
Total flavonoid content	Summer	22.87 \pm 0.80	5.55 \pm 0.46	8.56 \pm 0.71	22.27 \pm 0.66
	Autumn	27.86 \pm 1.20	3.26 \pm 0.27	5.99 \pm 0.99	16.69 \pm 0.52
	Winter	35.23 \pm 1.03	5.66 \pm 0.91	4.89 \pm 0.28	45.19 \pm 2.12
	Spring	25.54 \pm 0.5	6.69 \pm 0.97	4.80 \pm 0.12	49.21 \pm 4.83
Total polysaccharide content	Summer	48.84 \pm 3.66	8.16 \pm 0.40	14.95 \pm 1.85	42.84 \pm 3.84
	Autumn	31.10 \pm 5.80	11.19 \pm 0.96	10.25 \pm 1.58	27.81 \pm 0.71
	Winter	18.04 \pm 2.96	3.92 \pm 0.33	3.12 \pm 0.69	25.78 \pm 8.74
	Spring	39.11 \pm 1.46	19.30 \pm 2.26	16.02 \pm 0.40	26.75 \pm 0.29
Total protein content	Summer	9.28 \pm 0.28	8.10 \pm 0.59	7.18 \pm 0.63	-
	Autumn	19.57 \pm 1.42	3.31 \pm 0.81	13.51 \pm 1.08	-
	Winter	9.95 \pm 0.36	5.95 \pm 0.31	3.28 \pm 0.36	-
	Spring	11.17 \pm 0.12	5.36 \pm 0.43	3.7 \pm 0.07	-

Table 2: IC₅₀ values (µg mL⁻¹) for extracts on HL60, PC3 and THP-1 cell lines. Cells were exposed for 72 h to crude extracts of the brown alga *Cystoseira tamariscifolia* collected in all seasons. IC₅₀ values were calculated by sigmoidal dose-response of the data using SigmaPlot v. 13.0.


Season	cells	70%MeOH	100%MeOH	Chloroform	Water
Summer	HL60	7.33 ± 0.30	2.32 ± 0.21	7.9170 ± 0.12	293.54 ± 25
	PC3	452.01 ± 20	40.09 ± 3.40	112.28 ± 11	24.88 ± 0.21
	THP-1	116.48 ± 15	134.68 ± 24	60.35 ± 0.98	64.72 ± 6.2
Autumn	HL60	110.17 ± 9.6	110.43 ± 9.8	46.25 ± 39	221.51 ± 19
	PC3	469.76 ± 31	64.72 ± 5.3	263.59 ± 24	50.42 ± 5.6
	THP-1	236.74 ± 24	262.62 ± 21	396.04 ± 54	24.62 ± 4.1
Winter	HL60	39.37 ± 3.2	74.86 ± 0.51	24.63 ± 0.13	149.93 ± 0.20
	PC3	168.27 ± 15	105.60 ± 12	149.88 ± 0.41	38.69 ± 2.7
	THP-1	196.66 ± 22	342.39 ± 22	467.11 ± 27	444.29 ± 45
Spring	HL60	24.29 ± 0.27	24.28 ± 0.20	24.46 ± 0.19	128.38 ± 17
	PC3	190.75 ± 19	110.83 ± 15	99.35 ± 8.7	228.39 ± 22
	THP-1	74.98 ± 0.45	24.80 ± 0.31	74.84 ± 0.30	169.13 ± 16



544


545 Figure 1: Dose-response curves for cell vitality in HL-60 cells. HL60 cells were exposed
 546 for 72 h to crude extracts of the brown alga *Cystoseira tamariscifolia* collected in all
 547 seasons. Black circles - 70% MeOH, white circles - 100% MeOH, chloroform - black
 548 triangles, white triangles - water. (mean \pm SD; n = 9).

549



551


552 Figure 2: Dose-response curves for cell vitality in PC3 cells. PC3 cells were exposed
 553 for 72 h to crude extracts of the brown alga *Cystoseira tamariscifolia* collected in all
 554 seasons. Black circles - 70% MeOH, white circles - 100% MeOH, chloroform - black
 555 triangles, white triangles - water. (mean ± SD; n = 9).




556
 557 Figure 3: Dose-response curves for cell vitality in THP-1 cells. THP-1 cells were
 558 exposed for 72 h to crude extracts of the brown alga *Cystoseira tamariscifolia* collected
 559 in all seasons. Black circles, 70% MeOH, white circles, 100% MeOH, chloroform black
 560 triangles, white triangles water. (mean \pm SD; n = 9).

561

(A)




(B)



562

563 Figure 4: Fluorescent staining of cancer cell nuclei with propidium iodide. (A) HL60
564 and (B) THP-1 were treated with 100% methanol summer extracts at $150 \mu\text{g mL}^{-1}$ for
565 36h. Fragmented nuclei and apoptotic bodies were seen in the *C tamariscifolia* extract-
566 treated cells (b), but not in the control treatment (a). Magnification 200×, images
567 representative of three independent experiments.
568



569

570 Figure 5: The effect of *C. tamariscifolia* methanol extracts on the human leukaemia cell
 571 lines HL 60 (A) and THP-1 (B). HL 60 and THP-1 cells were resuspended in PBS and
 572 then RNase, and fluorescein diacetate (FITC) were added for 30 minutes, and cells
 573 were analysed by flow cytometry. Control cells (a) received no drug treatments. (b)
 574 Extract-treated HL60 and THP-1 cells. Early, late apoptosis and necrosis were found
 575 in treated cells but not in control. Experiments were carried out in three replicates.
 576