Faculty of Health: Medicine, Dentistry and Human Sciences

School of Health Professions

2020-07-28

The Effect of Retinal Illuminance on the Subjective Amplitude of Accommodation

Lara, F

http://hdl.handle.net/10026.1/15594

10.1097/OPX.000000000001544 Optometry and Vision Science Lippincott, Williams & Wilkins

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Manuscript title: The effect of retinal illuminance on the subjective amplitude of accommodation

Short title: Retinal illuminance and accommodation

Authors: Francisco Lara¹, PhD, Antonio J. Del Águila-Carrasco², PhD, Iván Marín-Franch^{1,3}, PhD, Resurrección Riquelme-Nicolás¹, MSc, and Norberto López-Gil¹, PhD

Affiliations:

1. Vision Sciences Research Group (CiViUM). University of Murcia, Murcia, Spain.

2. Eye and Vision Sciences Research Group, Faculty of Health: Medicine, Dentistry and Human Sciences, University of Plymouth, Plymouth, United Kingdom

3. Computational Optometry, Atarfe, Granada, Spain

Total number of words: 2750

Number of figures: 4

Number of tables: 0

Corresponding author:

Norberto López-Gil

Phone no: +44 (0)1752 588878

Faculty of Health: Medicine, Dentistry and Human Sciences, School of Health Professions, University of Plymouth, Plymouth, PL6 8BH, United Kingdom

Email: antonio.delaguila@plymouth.ac.uk

Abstract

Significance statement: We show that the amplitude of accommodation decreases with target illumination even under photopic reading conditions and a constant pupil size. This result provides a basis for clinical approaches that are not based on an optical explanation.

Purpose: We investigated the effect of retinal illuminance on the amplitude of accommodation while the pupil of the eye remained constant.

Methods: The amplitude of accommodation of 10 young subjects (from 20 to 38 years of age) and 10 presbyopic subjects (from 45 to 54 years of age) were measured subjectively through an artificial pupil of 5 mm using a Badal optometer and for four values of retinal illuminance: 222, 821, 2,138, and 5074 Td. Phenylephrine was instilled to all the subjects to ensure that their natural pupil was greater than the artificial one in all experimental runs. Two-way ANOVAs with age and log luminance as covariates were used to check whether changes in amplitude of accommodation with illumination were statistically significant.

Results: In the range of luminances tested, the amplitude of accommodation decreased on average from 6.34 D to 4.35 D in the young subjects, and from 1.69 D to 1.04 D in the presbyopic subjects. Luminance was associated with the amplitude of accommodation in both young and presbyopic groups, with p < 0.01.

Conclusions: The reduction in the amplitude of accommodation with target illumination (a phenomenon named *night presbyopia*) under photopic light conditions is not only due to a reduction in the depth of focus as a consequence of pupil dilation; it is strongly affected by the decrease of retinal illumination.

Keywords: accommodation, luminance, retinal illumination, night myopia, night presbyopia.

1 Introduction

There has been substantial evidence suggesting that the objective amplitude of 2 3 accommodation changes with object illumination since the effect was first observed in the 18th century.¹ Maskelyne in 1789² and Lord Rayleigh in 1883³ were the first to 4 describe how their eyes became short-sighted under low illumination conditions. That is, 5 6 their eyes experienced a myopic shift at night. This now well-known effect is called night 7 myopia. It took some 50 years after Rayleigh for Ferree and Rand⁴ to study systematically the changes in the near point caused by changes in light, and 10 more 8 9 years for Cabello to quantify the effect of illumination on the amplitude of accommodation (distance from the near point to far point) as a whole.⁵ Cabello coined the phrase night 10 presbyopia to capture the effect of receding near point that occurred at mesopic and 11 scotopic light conditions.⁵ Figure 1 (data from Cabello⁵ and Otero et al⁶) shows the 12 reduction of the amplitude of accommodation when the target illumination decreases. 13 Otero⁶ later demonstrated that, on average, the accommodative range decreases 14 progressively towards the point of tonic accommodation (dark focus)7-9 when the 15 stimulus gets dimmer. 16

More recent studies report both a myopic shift in distance refractions^{10,11} and reduced accommodation at low light levels.^{12,13} However, as in the studies of Cabello and Otero mentioned above,^{5,6} most of these changes occur at low mesopic and scotopic light levels, below those typically needed for reading text in the modern world.¹⁴

Given the fact that the retinal illuminance depends on target's luminance and on the pupil size¹⁵ (also depending on the large-field luminance¹⁶), the changes in near point and far point may have been affected by pupil miosis occurring as light levels change¹⁷ while accommodation takes place.^{18,19} As the pupil dilates, more peripheral optics contribute to the retinal image, and due to the change from positive to negative spherical aberration

that accompanies accommodation^{20,21} refractive measures that include the peripheral optics will tend to reveal reduced accommodation relative to measures that employ more paraxial optics.²² However, because of the impact of quantal noise,^{1,23,24} neural sensitivity declines as retinal illuminance is lowered (square root law). Because defocus primarily demodulates the higher spatial frequencies,²⁵ the lowered sensitivity to these high spatial frequencies may also contribute to a failure to accommodate at low light levels.

To separate the optical effects produced by the pupil from the neural effects, we have sought to isolate the neural effect on the depth of focus by studying the impact of retinal illuminance on the subjective amplitude of accommodation for a fixed pupil size. In this study, we constrained the light levels to those typically encountered during reading.¹⁴

36 Methods

37 Participants

Twenty-three subjects ranging from 20 to 54 years of age participated in this study. 38 39 However, three participants could not finish the experiment due to inability to accommodate, having excessive tearing, or a pupil smaller than 5 mm after the 40 instillation of phenylephrine. The remaining 20 subjects were split into two different 41 42 groups of 10 each: young subjects with ages ranging from 20 to 38 years and with a 43 mean (±standard deviation) age of 25 (±5.3) years; and 10 presbyopes, with ages 44 ranging from 45 to 54 years and with a mean age of 50 (± 3.1) years. Prior to testing, a subjective refraction²⁶ was performed by the same qualified optometrist at a luminance 45 of 108.9 (±1.7) cd m⁻². All subjects had a best-corrected visual acuity of 0.0 logMAR or 46 47 better. The study was approved by the University of Murcia Ethics Committee. All 48 participants gave written informed consent and the study adhered to the tenets of the Declaration of Helsinki. 49

50 Experimental set-up

A custom made Badal optometer with a stimulus controlled by a stepper motor²⁷ was used to vary target vergence in a range of about +10 D to -18 D. The system consisted of two achromatic doublets; one fixed, whereas the other one could be moved (i.e. Badal lens). With this automated Badal system, subjects adjusted target vergence to find their far point and near point subjectively, using a joystick control. The amplitude of accommodation was defined as the difference in vergence between these two positions. Further details about the optical system and the methodology can be found elsewhere.²⁸

The entrance pupil of the eye was chosen as the reference plane for the measurements, so that the amplitude of accommodation can be compared among subjects with different refractive errors.²⁹ For this purpose, a camera focusing at infinity and a plane mirror were used, so the iris of the participants could be focused. Amplitude of accommodation

calculations took into account that target position was fixed and its distance finite withrespect to the subject.

64 Experimental procedure

65 The left eye of all the participants was measured while the contralateral eye remained occluded. The target was a Bailey-Lovie chart placed at 6.95 m from the subject. The 66 67 target was inverted to account for the inversion introduced by the Badal system. Participants were told to put their chins on a chin rest, and then their refractive error was 68 69 compensated by placing the prescription obtained in the subjective refraction in the 70 phoropter, which was placed between the Badal lens and the eye, as close to the later 71 as possible. An artificial pupil with a diameter of 5 mm was placed in the phoropter, approximately at 10 mm from the corneal vertex. 72

73 Each participant took as many preliminary training trials as necessary until they felt they 74 were ready. After this, two drops of 10% phenylephrine were instilled on the participants' left eye, with an interval of 5 minutes. Phenylephrine does not significantly affect 75 accommodation.^{30,31} The artificial pupil was centered with respect to the natural pupil 76 before it was fully dilated. After 40 minutes, and after making sure that the pupil diameter 77 78 was greater than 5 mm and not reacting to light, the subjective measurements of the 79 amplitude of accommodation started. Different configurations of room lights and the use of two extra incandescent lamps of 500 W were used to generate four different chart 80 luminance levels (nominally, 11.3, 41.8, 108.9, and 258.4 cd m⁻², corresponding to a 81 retinal illuminance of 222, 821, 2138, 5074 Td, respectively.¹⁵) The sequence of 82 luminance conditions was random for every subject. 83

Participants were instructed to move the Badal lens up to the furthest possible position in which they were able to see the letters corresponding to 0.0 logMAR visual acuity clearly, following the "objectionable blur" criterion. The specific instructions (given in Spanish) were to search for the farthest point that sustained a "level of blur which you

88 would refuse to tolerate on a full-time basis. The blur has just reached a point at which it 89 is unacceptable."³² After obtaining the position of the far point, the subjects were 90 instructed to find their near point by bringing the Badal lens as close as possible until 91 they could perceive a maintained and unacceptable blur, with the same criterion as 92 before. Amplitude of accommodation was then obtained by averaging five far point and 93 near point measurements for each subject and luminance condition.

94 Statistical Analysis

95 Even though we designed the experiment to have two clearly distinct age groups, young 96 (amplitude of accommodation \geq 3 D) and presbyopes (amplitude of accommodation < 3 97 D), the effect of age on amplitude of accommodation was still expected to be very strong 98 within each group, especially on young subjects. We therefore incorporated age as a 99 potential confounder of the effect of illumination on amplitude of accommodation in our 100 models. Therefore, for each age group, a linear mixed-effect model for repeated-101 measures was performed with amplitude of accommodation as a linear function of age 102 (as confounder), a logarithmic function of luminance (the main explanatory variable) and subject as random effect. We used a logarithmic scale for luminance because its 103 104 association with the amplitude of accommodation is roughly linear on average (see 105 Results section). The significance level after Bonferroni correction of the repeated measures analyses for both groups was set at 0.025. The package "Ime4"33 for the R 106 107 statistical environment (https://www.R-project.org) was used to estimate the parameters of the linear mixed models. The p-values for the Kenward-Roger modification³⁴ of the F-108 statistic (an improved small sample approximation) was obtained with the function 109 110 "Anova" of the R package "car".³⁵

111 Results

112 The mean spherical equivalent obtained for the young group was -0.15 D. The standard 113 error of the mean (SEM) multiplied by 1.96 was 0.68 D. For the presbyopic group these 114 values were -0.10 and 1.11 D. The mean amplitude of accommodation (and 1.96 SEM) 115 obtained for the young group were (in descending order of luminance) 6.34 (0.35) D, 116 5.66 (0.43) D, 5.63 (0.45) D, and 4.35 (0.59) D. Figure 2 shows the amplitude of accommodation for each young subject at all four luminances. Subjects are sorted from 117 118 younger (top) to older (bottom). Overall, amplitude of accommodation decreases when 119 age increases.

120 The mean amplitude of accommodation (and 1.96 SEM) obtained for the presbyopic

121 group (Figure 3) under different luminance conditions (in increasing order of age) were

122 1.69 (0.16) D, 1.36 (0.16) D, 1.24 (0.20) D, and 1.04 (0.21) D. In general, same

tendency between amplitude of accommodation and age can be observed in this

124 group.

When averaged among each age group, our results can be summarized in Figure 4. As the illumination increased from 11.3 to 258.4 cd m⁻², the far point moved farther away from the eye approximately two times more in the young subjects (0.51 D) than in the presbyopes (0.23 D). In addition, the near point came about three times closer to the eye in young people (1.48 D) than in presbyopes (0.41 D).

For the young group, amplitude of accommodation increased with luminance by 1.36 D / \log_{10} (cd m⁻²) ($p < 4 \times 10^{-5}$) and decreased with age by 0.13 D / year (p = 0.07) as estimated with linear random-effect model for repeated measures. For the presbyope group, the estimated increase of amplitude with log luminance was more than 3 times smaller at 0.45 D / \log_{10} (cd m⁻²) ($p < 2 \times 10^{-6}$). Amplitude decrease with age was similar at 0.11 D / year (p = 0.01).

136 Discussion

137 The aim of this study was to assess the effect of changes in luminance on subjective amplitude of accommodation over the range of environmental light levels typically used 138 for reading.¹⁴ By employing a fixed 5-mm pupil diameter, these changes in stimulus 139 140 luminance altered the neural sensitivity of the visual system,¹ but avoided the confounder 141 of optical changes caused by pupil changes. Thus, the only optical changes experienced 142 by the eye were those happening during accommodation, so differences in perception of 143 the target were due to changes in neural effects produced by changes in retinal 144 illumination. In our Badal system, the retinal illuminance changed slightly during accommodation since the image nodal point of the eye changes its position,¹⁵ but that 145 change is less than 5% even for almost 7 D of accommodation.³⁶ A limitation of the 146 147 present study is the lack of objective measurements of the amplitude of accommodation, which would shed some light on whether or not the lack of light affects the response of 148 149 the ciliary muscle or if it is just a problem of photon noise.

150 Our results (an approaching far point and receding near point) are in general agreement with previous studies by Cabello,⁵ Otero⁶ and others³⁷. Unlike these previous studies, 151 152 however, in which light levels and the accommodative response both altered pupil size and therefore the optical characteristics of the image,³⁸ our use of a fixed pupil diameter 153 154 (5 mm) isolated the impact of changing neural sensitivity on amplitude of 155 accommodation. Thus, our results show that besides optical effects generated by the change in pupil size, retinal illumination plays a key role in the variation of the amplitude 156 of accommodation. 157

In addition to the main effect of presbyopia, the differences found in the magnitude of average changes of the far point with luminance could be explained by the greater transmission factor of ocular media in younger subjects producing a larger retinal illumination in the younger than the aged eye for the same object's luminance. Mean transmittance of 50 years old subjects is approximately 79% that of the 25 years old

subjects.³⁹ In relative terms, the effect of the target luminance is similar in both groups,
since the younger group accommodated between 3 and 4 times more than the
presbyopes, on average.

166 The mean age of the presbyope group was 50 years and previous studies indicate 167 average subjective amplitude of accommodation of an individual of that age is approximately 1.75 D,⁴⁰⁻⁴² which is about 0.4 D greater than the one obtained in this 168 169 study with a target luminance of 108.9 (1.7) cd m^{-2} (similar to that one used in clinical 170 measurements). This difference could be explained by the effect of the depth of focus 171 due to the difference in pupil diameters, since a typical pupil of 50 year olds at these luminance levels is approximately 4 mm,⁴³ 20% smaller than the artificial 5-mm pupil we 172 used here. A similar tendency was found for the younger subjects as for the older ones. 173 The average amplitude of accommodation for a 30-year-old individual is about 7 D,⁴⁰ 174 which is larger than the 5.66 D we observed in our younger sample. Significantly, 175 objective measures of amplitude of accommodation⁴⁴ report values of about 6 or 7 D in 176 young adults, and approximately zero in those over 50 years. This discrepancy between 177 the subjective^{40,41} and objective⁴⁴ measures of amplitude of accommodation in older eyes 178 is presumed to reflect the pseudo-accommodation or subjective depth of focus which is 179 180 incorporated into subjective measures of amplitude of accommodation as used in the present study. Because of large inter-subject differences, 40,42,45 as well as the sensitivity 181 of subjective accommodation to stimulus and instructions given to subjects.^{32,46} the small 182 differences between the current study and earlier reports is perhaps expected. 183

One main difference between a focused and a defocused image is the amplitude of the high spatial frequency content in the image.²⁵ As retinal illuminance decreases, higher spatial frequencies are affected more by photon noise.⁴⁷ Therefore, the signal indicating defocus^{48,49} becomes less visible as retinal illuminance decreases and accommodation tends to its resting state (around -2 D on average, although it varies a lot from subject to subject)¹³. Luminance reduction is far more impactful under scotopic light levels as it is

190 shown in Figure 1 with a larger jump in the transition between mesopic and scotopic ranges. The current study has shown that a decrease in amplitude of accommodation 191 192 can occur independently of the change of pupil size and is present at low photopic light levels as well, such as those used in reading, ¹⁴ which would be in agreemen with 193 Cambell's work on the minimum amount of light required to elicit the accommodation 194 reflex in humans.¹² Campbell stated that "if the luminance of the object is diminished until 195 196 it approaches the sensory threshold the perception of the less intense blurred edge of 197 the image will become impossible". Therefore, in Campbell's own words "the higher the 198 luminance of the object the easier will be the detection of this out-of-focus blurring, and the greater will become the accommodation response". 199

200 Our results highlight the importance of light levels in vision, especially in presbyopic 201 patients, when performing near vision tasks, such as reading. For instance, not having enough light could mean that a presbyopic patient may not be able to read at near 202 203 distances as a consequence of their near point getting further away from their eye due 204 to dim lightning, even when their addition has been properly calculated at photopic levels. 205 Some presbyopia treatments consist of expanding depth of focus by using an artificial small pupil in contact lenses^{50,51} or inside the cornea.⁵² However, this methodology 206 causes a decrease in retinal illuminance, which may alter the patient's near point. In a 207 208 theoretical study, Xu et al.³⁸ found that, at low light levels, visual function was generally worse when using small-pupil solutions than multifocal solutions. 209

In conclusion, our results show that the retinal illuminance changes that accompany reductions in target luminance reduce the subjective amplitude of accommodation over the stimulus range commonly encountered with text in the modern environment. It is likely therefore that lowered light levels will exacerbate any age related decline in amplitude of accommodation and therefore lower environmental lighting situations may be driving the onset of clinical presbyopia and the age at which optical aids are required.

- Lighting conditions, therefore, should be taken into account in the assessments,
- 217 diagnostics and prescriptions performed in the daily clinical practice.

218 Acknowledgments

- 219 We thank Arthur Bradley for revising the manuscript and for his useful advice in the
- 220 Discussion section.

References

- Van Nes FL, Bouman MA. Spatial Modulation Transfer in the Human Eye. J Opt Soc Am 1967;57:401–6.
- Maskelyne N. XXI. An Attempt to Explain a Difficulty in the Theory of Vision, Depending on the Different Refrangibility of Light. Philos Trans R Soc London 1789;79:256–64.
- Rayleigh L. On the Invisibility of Small Objects in a Bad Light. Proc Camb Philos Soc 1883:324.
- 4. Ferree CE, Rand G. Intensity of Light in Relation to the Near Point and the Apparent Range of Accommodation. Am J Ophthalmol 1935;18:307–18.
- Cabello J. Las Causas de la Miopía Nocturna (Causes of Night Myopia). An Fís Quím 1945:439–48.
- Otero JM, Aguilar M. Accommodation and Night Myopia. J Opt Soc Am 1951;41:1061_1-1062.
- Leibowitz H, Owens D. Anomalous Myopias and the Intermediate Dark Focus of Accommodation. Science 1975;189:646–8.
- Leibowitz HW, Owens DA. New Evidence for the Intermediate Position of Relaxed Accommodation. Doc Ophthalmol 1978;46:133–47.
- Andre JT, Owens RL, Owens DA. Dark Focus Values Measured by Retinoscopy: a Clinical Chart Review. Optom Vis Sci 1998;75:903–8.
- Epstein D, Ingelstan E, Jansson K, Tengroth B. Low-luminance Myopia as Measured with a Laser Optometer. Acta Ophthalmol 1981;59:928–43.
- 11. Arumi P, Chauhan K, Charman WN. Accommodation and Acuity Under Nightdriving Illumination Levels. Ophthalmic Physiol Opt 1997;17:291–9.

- 12. Campbell FW. The Minimum Quantity of Light Required to Elicit the Accommodation Reflex in Man. J Physiol 1954;123:357–66.
- Johnson CA. Effects of Luminance and Stimulus Distance on Accommodation and Visual Resolution. J Opt Soc Am 1976;66:138–42.
- Xu R, Gil D, Dibas M, Hare W, Bradley A. The Effect of Light Level and Small Pupils on Presbyopic Reading Performance. Invest Ophthalmol Vis Sci 2016;57:5656–64.
- Thibos LN, Lopez-Gil N, Bradley A. What is a Troland? J Opt Soc Am A 2018;35:813–6.
- Barbur JL, Harlow AJ, Sahraie A. Pupillary Responses to Stimulus Structure, Colour and Movement. Ophthalmic Physiol Opt 1992;12:137–41.
- Winn B, Whitaker D, Elliot DB, Phillips NJ. Factors Affecting Light-adapted Pupil Size in Normal Human Subjects. Invest Ophthalmol Vis Sci 1994;35:1132–7.
- Radhakrishnan H, Neil Charman W. Age-related Changes in Static Accommodation and Accommodative Miosis. Ophthalmic Physiol Opt 2007;27:342–52.
- Almutairi MS, Altoaimi BH, Bradley A. Accommodation and Pupil Behaviour of Binocularly Viewing Early Presbyopes. Ophthalmic Physiol Opt 2017;37:128–40.
- López-Gil N, Fernández-Sánchez V, Legras R, et al. Accommodation-related Changes in Monochromatic Aberrations of the Human Eye as a Function of Age. Invest Ophthalmol Vis Sci 2008;49:1736–43.
- Del Águila-Carrasco AJ, Kruger PB, Lara F, López-Gil N. Aberrations and Accommodation. Clin Exp Optom 2019; 103:95-103.
- 22. Lopez-Gil N, Fernandez-Sanchez V. The Change of Spherical Aberration During

Accommodation and its Efect on the Accommodation Response. J Vis 2010;10:12–12.

- Banks M, Geisler W, Bennett P. The Physical Limits of Grating Visibility. Vision Res 1987;27:1915–24.
- Xu R, Wang H, Thibos LN, Bradley A. Interaction of Aberrations, Diffraction, and Quantal Fluctuations Determine the Impact of Pupil Size on Visual Quality. J Opt Soc Am A 2017;34:481–92.
- Webster MA, Georgeson MA, Webster SM. Neural Adjustments to Blur. Nat Neurosci 2002;5:839–40.
- Rabbetts RB. Bennett and Rabbetts' Clinical Visual Optics. Elsevier/Butterworth Heinemann; 2007.
- Atchison DA, Bradley A, Thibos LN, Smith G. Useful Variations of the Badal Optometer. Optom Vis Sci 1995;72:279–84.
- López-Gil N, Fernández-Sánchez V, Thibos LN, Montés-Micó R. Objective Amplitude of Accommodation Computed from Optical Quality Metrics Applied to Wavefront Outcomes. J Optom 2009;2:223–34.
- Bernal-Molina P, Vargas-Martín F, Thibos LN, López-Gil N. Influence of Ametropia and Its Correction on Measurement of Accommodation. Investig Opthalmology Vis Sci 2016;57:3010-6.
- Esteve-Taboada JJ, Del Águila-Carrasco AJ, Bernal-Molina P, et al. Effect of Phenylephrine on the Accommodative System. J Ophthalmol 2016;2016:7968918.
- Del Águila-Carrasco AJ, Lara F, Bernal-Molina P, et al. Effect of Phenylephrine on Static and Dynamic Accommodation. J Optom 2019;12:30–7.
- 32. Atchison DA, Fisher SW, Pedersen CA, Ridall PG. Noticeable, Troublesome and

Objectionable Limits of Blur. Vision Res 2005;45:1967–74.

- Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using Ime4. J Stat Software 2015;67:1–48.
- Kenward M, Roger J. Small Sample Inference for Fixed Effects from Restricted Maximum Likelihood. Biometrics 1997;53:983–97.
- 35. Fox J, Weisberg S. An R Companion to Applied Regression. Sage publications;2018.
- El Hage SG, Le Grand Y. Physiological Optics. Vol 13. Berlin, Heidelberg: Springer Berlin Heidelberg; 1980.
- Lara F, Bernal-Molina P, Fernández-Sánchez V, López-Gil N. Changes in the Objective Amplitude of Accommodation with Pupil Size. Optom Vis Sci 2014;91:1215–20.
- Xu R, Wang H, Jaskulski M, et al. Small-pupil Versus Multifocal Strategies for Expanding Depth of Focus of Presbyopic Eyes. J Cataract Refract Surg 2019;45:647–55.
- Weale RA. Age and the Transmittance of the Human Crystalline Lens. J Physiol 1988;395:577–87.
- Jackson E. Amplitude of Accommodation at Different Periods of Life. Cal State J Med 1907;5:163–6.
- Duane A. An Attempt to Determine the Normal Range of Accommodation at Various Ages, Being a Revision of Donder's Experiments. Trans Am Ophthalmol Soc 1908;11:634–41.
- 42. López-Alcón D, Marín-Franch I, Fernández-Sánchez V, López-Gil N. Optical
 Factors Influencing the Amplitude of Accommodation. Vision Res 2017;141:16–
 22.

- 43. Watson AB, Yellott JI. A Unified Formula for Light-adapted Pupil Size. J Vis 2012;12.
- Anderson H, Hentz G, Glasser A, et al. Minus-lens-stimulated Accommodative Amplitude Decreases Sigmoidally with Age: a Study of Objectively Measured Accommodative Amplitudes from Age 3. Invest Ophthalmol Vis Sci 2008;49:2919–26.
- Donders F, Moore W. On the Anomalies of Accommodation and Refraction of the Eye: With a Preliminary Essay on Physiological Dioptrics. New Sydenham Society; 1864.
- 46. Stark LR, Atchison DA. Subject Instructions and Methods of Target Presentation in Accommodation Research. Invest Ophthalmol Vis Sci 1994;35:528–37.
- 47. Kotulak JC, Morse SE, Billock VA. Red-green Opponent Channel Mediation of Control of Human Ocular Accommodation. J Physiol 1995;482 (Pt 3):697–703.
- Kruger PB, Pola J. Stimuli for Accommodation: Blur, Chromatic Aberration and Size. Vision Res 1986;26:957–71.
- Esteve-Taboada JJ, Del Águila-Carrasco AJ, Bernal-Molina P, et al. Dynamic Accommodation Without Feedback Does Not Respond to Isolated Blur Cues. Vision Res 2017;136:50–6.
- Freeman E. Pinhole Contact Lenses. Am J Optom Arch Am Acad Optom 1952;29:347–52.
- Albarrán Diego C, Montés-Micó R, Pons AM, Artigas JM. Influence of the Luminance Level on Visual Performance with a Disposable Soft Cosmetic Tinted Contact Lens. Ophthalmic Physiol Opt 2001;21:411–9.
- Waring GO. Correction of Presbyopia with a Small Aperture Corneal Inlay. J Refract Surg 2011;27:842–5.

Figure Legends

Figure 1. Variation of the subjective far and near points (and thus amplitude of accommodation) as a function of stimulus luminance (open and black circles, respectively). Target vergence producing maximum visual acuity is also plotted (black squares). Luminance is given in Apostilbs (asb), being 1 asb = π^{-1} cd m⁻². Scotopic conditions correspond approximately to $\pi \cdot 10^{-3}$ asb; mesopic conditions between $\pi \cdot 10^{-3}$ asb to π asb, and values greater than π abs correspond to photopic conditions. Adapted from Cabello⁵ and Otero et al.⁶

Figure 2. Amplitude of accommodation for the young subjects for each retinal illuminance. Subjects are ranked by age with younger on top. The width of the bars indicates the amplitude of accommodation, the left border is the far point and the right border of the bar is the near point of each subject. Red vertical dashed lines show the mean far point and near point among all the young subjects. The left error bars represent the 1.96 SEM of the far point, whereas right error bars represent the 1.96 SEM of the far point, whereas for the mean among subjects.

Figure 3. Amplitude of accommodation for the presbyopic subjects for each retinal illuminance. Other details as in Figure 2.

Figure 4. Far point (solid circles) and the near point (empty circles) under different target luminance levels, for young subjects (top left panel) and for presbyopes (top right panel). Bottom panels show the relationship between the logarithm of the luminance and the amplitude of accommodation for young subjects (bottom left panel) and for presbyopes (bottom right panel). The values on the top *x*-axis in the upper panels are the retinal illuminance (in Trolands) corresponding to each log luminance level on the bottom *x*-axis. They were added to allow for direct comparisons with Figures 2 and 3. The length of the error bars represents the Gaussian estimate of the 95% confidence

intervals (\pm 1.96 SEM) around the mean values. Note that in many cases the error bars are so small in length that they are occluded by the symbols themselves.







