Faculty of Science and Engineering

School of Biological and Marine Sciences

2020-08

# Challenges in monitoring mobile populations - Applying Bayesian multi-site mark-recapture abundance estimation to the monitoring of a highly mobile coastal population of bottlenose dolphins

Ingram, Simon

http://hdl.handle.net/10026.1/16222

10.1002/aqc.3355 Aquatic Conservation: Marine and Freshwater Ecosystems Wiley

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.



#### Challenges in monitoring mobile populations - Applying Bayesian multi-site mark-recapture abundance estimation to the monitoring of a highly mobile coastal population of bottlenose dolphins

Journal:	Aquatic Conservation: Marine and Freshwater Ecosystems
Manuscript ID	AQC-19-0069.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Nykänen, Milaja; University College Cork School of Biological Earth and Environmental Sciences, Oudejans, Machiel; Kelp Marine Research Rogan, Emer; University College Cork College of Science Engineering and Food Science, School of Biological, Earth and Environmental Sciences Durban, John; Southwest Fisheries Science Center Ingram, Simon; University of Plymouth, School of Biological and Marine Sciences
Broad habitat type (mandatory) select 1-2:	coastal < Broad habitat type
General theme or application (mandatory) select 1-2:	monitoring < General theme or application, Special Area of Conservation < General theme or application, survey < General theme or application
Broad taxonomic group or category (mandatory, if relevant to paper) select 1-2:	mammals < Broad taxonomic group or category
Impact category (mandatory, if relevant to paper) select 1- 2:	

## SCHOLARONE<sup>™</sup> Manuscripts

# Challenges in monitoring mobile populations - Applying Bayesian multisite mark-recapture abundance estimation to the monitoring of a highly mobile coastal population of bottlenose dolphins

Milaja Nykänen<sup>1\*</sup>, Machiel G. Oudejans<sup>2</sup>, Emer Rogan<sup>1</sup>, John W. Durban<sup>3</sup>, and Simon N. Ingram<sup>4</sup>

<sup>1</sup> School of Biological, Earth and Environmental Sciences, University College Cork, Cork, Ireland

<sup>2</sup> Kelp Marine Research, Hoorn, The Netherlands

<sup>3</sup> Protected Resources Division, Southwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, La Jolla, California, USA

<sup>4</sup> School of Biological and Marine Sciences, University of Plymouth, Plymouth UK

\* Correspondence to be sent to: Milaja Nykänen, School of Biological, Earth and Environmental Sciences, the Butler Building, North Mall, University College Cork, Cork, Ireland, E-mail: m.nykanen@ucc.ie, Tel: +353 (0)21 4904

Peer Peurez

2		
3	1	
4 5	-	
6	2	1. Monitoring the
7 8	3	conservation manage
9 10	4	protected areas. Whe
11	5	photo-identification
12 13	6	and out of the surve
14 15	7	accuracy of resulting
16 17 18	8	2. Bayesian hierarchi
19	9	in summer 2014 to
20 21	10	dolphins, Tursiops tr
22	11	addition, the effects
23 24 25	12	in population size wa
26 27	13	3. The abundance of a
28	14	posterior density int
29 30	15	throughout the study
31 32	16	mobility. In addition
33 34	17	the six-year reporting
35 36	18	4. Given that the Spe
37	19	separate areas cover
38 39	20	approach is appropri
40 41	21	well-suited for sparse
42 42	22	of resources prevent
45 44	23	areas fail to capture th
45 46	24	multi-site approach
47 48	25	terrestrial taxa.
49 50 51 52	26	Keywords: coastal, n
53 54		
55		
56 57		
58		
59		

#### Abstract

1. Monitoring the abundance of mobile and wide-ranging cetacean populations for conservation management is challenging, especially when the management is focused on static protected areas. Where abundance estimates are derived from mark-recapture data, such as photo-identification of naturally marked individuals, unpredictable movements of animals in and out of the survey area can reduce 'capture' probabilities and affect the precision and accuracy of resulting estimates.

8 2. Bayesian hierarchical log-linear likelihood was applied to photo-identification data collected
9 in summer 2014 to derive a multi-site abundance estimate for a population of bottlenose
10 dolphins, *Tursiops truncatus*, ranging widely throughout coastal waters of western Ireland. In
11 addition, the effects of varying levels of sampling effort on the minimum detectable decrease
12 in population size was examined.

3. The abundance of dolphins was estimated as 189 (coefficient of variation: 0.11, 95% highestposterior density interval: 162–232). Over 50% of the well-marked dolphins encountered
throughout the study were sighted in more than one distinct coastal area thus displaying high
mobility. In addition, it was found that in order to detect a 25% decline in abundance within
the six-year reporting period of the EU's Habitats Directive would require biennial surveys.

4. Given that the Special Area of Conservation designated for these dolphins consists of two separate areas covering a substantial portion of the west coast of Ireland, the multi-site approach is appropriate for monitoring this population. It produces a precise estimate and is well-suited for sparse recapture data collected opportunistically at multiple sites, when the lack of resources prevent large scale surveys, or when concentrating surveys on smaller localized areas fail to capture the broad range and unpredictable occurrence of the animals. The Bayesian multi-site approach could be applied to the management of other wide-ranging marine or terrestrial taxa.

Keywords: coastal, mammals, monitoring, Special Area of Conservation, survey, modelling

#### **1. Introduction**

#### 28 1.1. General background

Bottlenose dolphins (Tursiops truncatus) are widely distributed throughout tropical and temperate seas and found in pelagic oceanic environments, on the continental shelf, as well as in coastal inshore waters (Wells and Scott, 2009). Their minimum worldwide abundance is estimated to be approximately 600,000 individuals (Wells and Scott, 2009) and numbers in European Atlantic continental shelf waters have been estimated to be around 16,000 (Hammond et al., 2013) although results from recent aerial surveys suggests that there may be strong inter-annual variation in this area (Rogan et al., 2018). Whilst the bottlenose dolphin as a species is not considered to be globally endangered, some populations, especially those inhabiting coastal areas, are small and often genetically and/or geographically isolated (e.g. Caballero et al., 2012; Fernández et al., 2011; Louis et al., 2014; Mirimin et al., 2011; Nykänen et al., 2018). This puts them at risk of losing heterozygosity and genetic resilience due to genetic drift (Lacy, 1987) placing them at greater risk to local extinctions with increased vulnerability to anthropogenic pressures. The main threats to delphinids in coastal environments include pollutants such as xenobiotic chemicals (especially PCBs and DDTs), reduced prey availability due to coastal fisheries, habitat degradation, noise and disturbance from vessel traffic, entanglement and incidental bycatch, direct hunting, marine construction and anthropogenic noise (Jepson et al., 2016; Lusseau et al., 2009; Pirotta et al., 2015; Williams et al., 2009; Williams et al., 2014). The sensitivity of bottlenose dolphins to these threats is further exacerbated by their position as apex predators in coastal ecosystems and by their low reproductive rates (Arso Civil et al., 2017; Baker et al., 2018a; Quick et al., 2014). 

*1.2. Conservation and monitoring requirements of bottlenose dolphins in Europe* 

The conservation of wild animal populations is often implemented through designation and management of protected areas that are considered to represent important habitats for foraging, breeding and other important activities (Palumbi, 2001; Reeves, 2000). This is usually followed by regular monitoring of some demographic parameters, such as abundance, survival or age structure of the individuals inhabiting these areas. In European waters, bottlenose dolphins are protected through Annexes II and IV of the European Union's Habitats Directive (European Economic Community, 1992), and the Member States are required to designate Special Areas of Conservation (SACs) as part of a European strategy to maintain or restore 'favourable conservation status' for the species. In practice, this means that in order to be classed as 

'favourable', the species (or population) should not decline from the reference level (defined by the Member States individually) by more than 25% over a six-year reporting period, alternatively, annual decline should not exceed 1%. In addition to the Habitats Directive, as top predators, bottlenose dolphins are included as one of the indicator species for 'Good Environmental Status' in European coastal waters in the Marine Strategy Framework Directive (MSFD; Council of the European Communities, 2008). 

Some bottlenose dolphin populations have a strong site-fidelity to bays and estuaries (Bearzi et al., 2008; Connor et al., 2000; Ingram and Rogan, 2002; Read et al., 2003), and their conservation monitoring has been focused on discrete, local areas such the SACs. Recent studies have found that bottlenose dolphins using the coastal waters of Ireland belong to two genetically, demographically and socially distinct populations (Mirimin et al., 2011; Nykänen et al., 2018, 2019), with a resident population using the Shannon estuary, hereafter referred to as the 'Shannon population'. Residency is determined here by individuals occurring in the area at least 50% of the months in a given year over multiple years or decades, adapted from Rosel et al. (2011). The second population, hereafter the 'west coast population', is more widely distributed using other coastal areas of western Ireland (Ingram et al., 2003; Oudejans et al., 2010). Consequently, two discrete SACs have been designated to ensure the protection of these populations; the Lower River Shannon Estuary SAC and the West Connacht Coast SAC (see inset map in Figure 1). While the area-based monitoring of the Shannon population has been successful, capturing majority of the individuals inhabiting the estuary (based on discovery curves) and producing precise abundance estimates (Englund et al., 2008; Ingram and Rogan, 2003), this approach may not be suitable for more mobile and dispersed populations that have spatially and temporally variable use of large areas of habitat, presenting a challenge for monitoring. For example, on the east coast of Scotland, the effectiveness of the Moray Firth SAC, designated based on high site-fidelity exhibited by a population of bottlenose dolphins to the area, has recently been questioned due to the population extending their range to areas outside the SAC over the past decade (Wilson, 2016; Wilson et al., 2004). Similarly, in Irish waters, the range of the west coast population extends beyond the designated SAC. Conservation efforts may thus need to move away from area-based management and instead focus on populations whilst considering population dynamic processes such as dispersal (genetic and demographic) that affect the dynamics and the overall viability of populations. Moreover, in order to achieve efficient monitoring appropriate for the MSFD and to ensure that 

http://mc.manuscriptcentral.com/aqc

91 effective conservation of dispersed coastal populations is achieved, it is crucial to design an92 appropriate monitoring strategy suitable for the population in question.

Compared to the Shannon estuary population that has been the focus of most research on bottlenose dolphins in Ireland (Baker et al., 2018a; 2018b; Berrow et al., 2012; Englund et al., 2008; Foley et al., 2010; Ingram and Rogan, 2002; 2003), much less is known about the west coast population. Preliminary studies identified a significant number of bottlenose dolphins inhabiting the waters off the west coast of Ireland (Ingram et al., 2001; 2003) with an estimated mean abundance of 171 dolphins using the waters around Connemara, Co. Galway (see Figure 1) (Ingram et al., 2009). However, this estimate was based on surveys over a limited length of coast within the West Connacht Coast SAC and was relatively imprecise with a coefficient of variation (CV) of 0.28. Moreover, despite multi-annual re-sightings of individuals, it appears that these animals are highly mobile and have a large home range with encounters occurring throughout the west coast (Ingram et al., 2001; 2003). The widespread distribution together with unpredictable movements make monitoring the abundance of this population especially challenging. Therefore, one of the aims of this study was to provide an abundance estimate which could be used as a baseline for long-term monitoring. Further, the distribution and the rate of individual movements were investigated and some of the possible underlying factors driving the distribution explored and discussed. Finally, the effect of different levels of survey effort on the precision of the abundance estimate was quantified using a power analysis. This will help inform a cost-effective strategy for future monitoring that is sufficiently sensitive to changes in abundance to reliably detect population decline in a timely manner. 

-1 112

## **2. Methods**

## 5 114 2.1. Data collection and photograph analysis

Mark-recapture is widely applied in ecological studies to estimate the number of individuals in a population or the density of animals within a surveyed area (Otis et al., 1978). Individual bottlenose dolphins can be identified from naturally occurring markings (Würsig and Würsig, 1977). These marks mostly consist of scars and nicks from interactions with conspecifics and they can be permanent, such as deep nicks or scars on the dorsal fin, or temporary, such as superficial scratches (Appendix 1). Heavily marked animals can be identified over periods of many years, whereas more superficial markings, such as tooth rake scars, may fade within a 

period of about a year reducing inter-annual re-sighting probabilities of less heavily marked individuals. In this study, identification photographs were taken of individual bottlenose dolphins encountered in schools during dedicated and opportunistic boat-based surveys. Here, a school is defined as "all dolphins within a 100m radius of each other" after (Irvine et al., 1981). Boat-based surveys were conducted along a 250km stretch of coastal waters in western Ireland (Figure 1) during the summer months (May-September) of 2014. Efforts were made to photograph the dorsal fins of all members of each encountered school. 

The best quality photograph of each identifiable dolphin was selected from each encounter and assigned an image quality score of 1 to 4 (1 being the highest quality and 4 the poorest, see Appendix 2) with no consideration of the degree of marking of the individual dolphin. Each photographed individual was then assigned one of three grades of mark-severity (Appendix 1), and visually matched against the archived catalogue of dolphins identified during previous encounters. To minimise bias in capture probability resulting from identification errors, photographs of quality grade 4 were excluded from subsequent abundance estimation. Further, only the "well-marked" dolphins (M1, see Appendix 1), easily distinguishable and identifiable from both the left and the right side, were included in the analyses. For this study, the wider study area was divided into three discrete and geographically separated blocks where survey effort had been concentrated (see Figure 1). Photographs from encounters were compared within and between the blocks to establish whether individuals were seen across the whole study area during the study period. 

#### 2.2. Abundance analysis

Mark-recapture models that assume population closure (zero net migration and births and deaths) within a single defined area, are typically used in abundance estimation of dolphins with strong site-fidelity to specific areas (Berrow et al., 2012; Louis et al., 2015; Read et al., 2003; Wilson et al., 1999). However, when the animals are moving non-randomly into and out of the area within the sampling (survey) period, and the effective number of animals available for re-capture therefore changes, closed models become less applicable as the violation of population closure assumption can result in biased abundance estimates (Kendall, 1999). Bayes' theorem, as opposed to traditional frequentist maximum likelihood (ML) based estimation, has recently become more widely applied in mark-recapture abundance estimation (Mäntyniemi and Romakkaniemi, 2002; Michielsens et al., 2006). It has been applied to a range of cetacean species (Beck et al., 2014; Cheney et al., 2013; Durban et al., 2010; Durban et al., 2005; Fearnbach et al., 2012; Moore and Barlow, 2011) due to its utility with sparse data and/or

opportunistic data collection. In this study, due to the large combined coastal area surveyed, Bayesian inference was applied to a model of hierarchical log-linear likelihood of counts of identified dolphins across three discrete blocks, and a combined abundance estimate of bottlenose dolphins using the entire survey area extending from Connemara to Donegal Bay was derived (Figure 1). This method, developed by Durban et al. (2005), is well-suited for data sets with low number of individual re-sightings and for situations when it is unfeasible to do systematic surveys covering the entire population's range. The model also takes into account different ranging patterns of individuals and geographical dependencies between multiple sites, enabling the estimation of movement rates of animals between sampling locations. An advantage of using Bayesian inference instead of traditional frequentist statistics is that prior knowledge of the parameter (prior) distribution can be incorporated into the model to produce a joint posterior distribution for the parameter in question. An example of this would be setting a realistic maximum value to the prior for the abundance of all well-marked animals in an area. This informative prior is then incorporated into the model to facilitate the convergence of Markov Chain Monte Carlo (MCMC) chains. 

A contingency table of sighting histories of well-marked (M1) bottlenose dolphins was created based on their presence or absence in each of the study blocks during a single survey season (Table 1). The re-sightings of individuals among multiple sites therefore represented spatial, rather than temporal, capture-recapture events (Durban et al., 2005). The model predicts the number of animals not captured at any of the survey sites and incorporates this value to compute the estimate of the overall abundance of well-marked animals across the entire study area. The model also incorporates the proportion of well-marked individuals as a binomial sample of the total number of animals seen; therefore, it predicts the total number of individuals (including unmarked animals) in the study area (see Cheney et al., 2013). The model averaging and prediction (Durban et al., 2005) were performed using MCMC sampling in WinBUGS software (Lunn et al., 2000) with 100,000 burn-in followed by 100,000 iterations. Three independent chains were run to confirm consistency between runs and inspected visually for convergence. 

# 50 51 182 2.3. Range of bottlenose dolphins encountered in 2001-2014

In order to describe the extent of movements of bottlenose dolphins sighted on the west coast of Ireland, the photographs taken in 2014 were supplemented with data collected over a longer time period, in 2001-2013, and the range of the sighting latitudes were plotted for the 39 most sighted ( $\geq$ 5 times) well-marked dolphins. These were the same individuals that were included in the social structure analyses. In addition, the dependency of the sighting latitude range (the

difference between the maximum and the minimum latitude) and the encounter frequency was
determined by calculating Pearson's correlation coefficient, *r*.

#### 190 2.4. Analyses of social structure

To investigate whether this population of coastal dolphins could be divided into social clusters reflecting site-fidelity to their sighting locations and/or geographic range, analyses of social structure were performed in SOCPROG 2.4 compiled version (Whitehead, 2009a; 2009b). Applying the 'gambit of the group' concept (Whitehead and Dufault, 1999), the rate at which individuals were photographed within the same schools, was used as a proxy for social association. Daily sampling periods were used to ensure the independence between the sampling periods (Whitehead, 2008). The dataset was restricted to good quality photographs (Q1-3, Appendix 2) of individuals with permanent and obvious markings (mark severity grade M1, Appendix 1) in order to identify individuals over multi-annual periods, and only dolphins photographed in at least five sampling periods (days) were included to reduce bias caused by rarely seen individuals (Whitehead, 2009a). Social analyses included entire sighting histories from 2001 up until 2014, the duration of photo-ID surveys of this population. 

The strength of association between pairs of individuals (i.e. dyads) was measured using the half-weight association index (HWI). This index of co-occurrence takes values between 0 (never seen together) and 1 (always observed together), and is appropriate when not all associates within a group have been identified (Cairns and Schwager, 1987). Standard deviation (SD) and CV of the HWI were also calculated. A Monte Carlo permutation test (Bejder et al., 1998; Whitehead, 1999) was used to test whether the observed association patterns (real data) were different than expected from randomly associating individuals (permutated data). The permutations were performed using 20,000 iterations with 1000 trials per iteration. A higher SD of the observed association indices compared to the SD of permutated data is considered as an indication of preferred and/or avoided associations between the sampling periods (Whitehead, 2009a). 

The power of the analysis to capture a true representation of the social system was estimated as the correlation of the observed and estimated association indices using the maximum likelihood estimator (Pearson's correlation coefficient, r). A measure of social differentiation, S, calculated as the CV of real association indices, was used to describe the variability in the social system, with values >0.5 indicating a well differentiated society (Whitehead, 2009a). Standard errors (SEs) for r and S were calculated by bootstrapping with 100 replications. In 

order to determine whether the population of bottlenose dolphins could be divided into clusters where association indices are higher among members of the same cluster than expected by chance, an eigenvector-based maximum modularity coefficient, Q (Lusseau, 2007; Newman, 2004; 2006; Whitehead, 2009a), was calculated. This method accounts for different levels of gregariousness between the individuals (i.e. the average number of associates) with modularities greater than  $\sim 0.3$  considered to represent effective community divisions (Newman, 2004; Whitehead, 2009a). NetDraw (Borgatti et al., 2002) was used to visualize a social network diagram using the network statistics calculated in SOCPROG. 

#### $\frac{1}{8}$ 228 2.5. Power to detect change in abundance

Program TRENDS (Gerrodette, 1987, 1991) was used to conduct a power analysis in order to estimate the annual rate of decline in population abundance within a six-year period (as mentioned previously, six years is the reporting interval set in the Habitats Directive) that could be detected with the level of precision (here, the CV) achieved in this study. The precision that would be required to detect an annual decline of 1% in population size over the six-year period was also estimated, as identifying this rate of annual decline is one of the requirements of the Habitats Directive.

Further, the effect of different amounts of sampling effort (here, number of years between surveys) on the minimum detectable overall decline in population size was examined using a longer theoretical study period of 25 years and a range of CVs varying from 0.01 (very high precision) to 0.30 (low precision). Specifically, scenarios were tested when abundance surveys were conducted every six years (five years between surveys), every three years (two years between surveys), every two years (one year between surveys) or every year, over the 25-year period. In all the power analyses, the desired power was set to 80%, the probability of Type I and II errors to 0.05, and a one-tailed test was used, as the purpose was to detect a decrease and not a general change in abundance. A linear population model was used for a non-recovering population as in Fruet et al. (2015). 

**3. Results** 

#### *3.1. Data collection*

In 2014, 146 survey hours yielded six encounters with bottlenose dolphin schools around
 Connemara, seven around Mullet peninsula and eight in Donegal Bay (Figure 1). School size
 ranged from 9 to 95 with the largest schools encountered in Donegal Bay (median school size

of 36). In total, nearly 10,000 photographs were analysed. From these, 169 new dolphin identifications from photographs obtained from either the left, the right, or both sides of the animal were added to the archive of dolphin images collected since 2001. Note that due to the fact that bottlenose dolphin markings can change over time and that some individuals are known only from one side, the number of identifications in the archive does not equal the number of individuals in the population, especially when considering the gaps in the years when photo-ID surveys were conducted. Nevertheless, 71 animals were matched to individuals identified from encounters made in previous years with seven identifications dating back to 2001.

20 260 3.2. Abundance and movements

From the photographs taken during May–September 2014, a total of 91 well-marked dolphins, identified or identifiable from both sides, were included in the abundance analysis (Table 1). Forty-nine (54%) of these animals were seen in more than one study block, and eight (9%) were encountered in all the study sites. The highest overlap of individuals occurred between Mullet peninsula and Donegal Bay with 28 dolphins (31%) sighted in both of these areas. Donegal also had the highest number (n = 23) of animals seen in only one of the three study sites. The average proportion of well-marked dolphins (to all dolphins, marked and unmarked) was 0.57 across all encounters in 2014. The Bayesian multi-site median abundance estimate of the total number of dolphins for the whole study area for the summer 2014 was 189 (CV =0.11, 95% HPDI = 162-232). The non-significant P-value (P = 0.158) from the closure test of Otis et al. (1978) suggested that the closure assumption was not violated. 

<sup>41</sup><sub>42</sub> 272 3.3. Range of bottlenose dolphins encountered in 2001-2014

The range of sighting latitudes of the most sighted ( $\geq 5$  times) well-marked dolphins is presented in Figure 2a; it appears that while most of these animals were sighted from Donegal Bay to Connemara, with the distance between the areas of more than 250km (over water), there were four animals (IDs 1056, 1094, 1038 and 1049) that had even wider distribution having been sighted from Co. Cork to Donegal Bay between 2001 and 2014 with >500km between these sites. In contrast, there were also a number of individuals with much narrower latitudinal range, that were encountered only in two of the sites; two individuals were only encountered in Connemara and around the Mullet peninsula (IDs 1099 and 1244, Figure 2a), and 12 individuals were only recorded around the Mullet peninsula and in Donegal Bay (for example, IDs 1444 and 1468) during 2001-2014. The range of the sighting latitudes was not dependent 

on the number of times the animal was encountered (Pearson's r = 0.180, P = 0.125, see Appendix 3).

#### *3.4. Social structure*

When including only good quality photographs of well-marked (M1) individuals encountered in at least five sampling periods, 39 bottlenose dolphins were included in the analyses of social structure. These data were collected during 51 encounters over 48 days in 2001-2014. The mean number of observations per dolphin was 7.21 (SE = 1.95) and the maximum number of times that an individual was encountered was 13. The individuals had, on average, 63 associations with other individuals.

The mean HWI was 0.226, which did not differ significantly from the permuted random data (mean = 0.226, P > 0.05). However, the SD (0.206) and CV (0.910) in the real data were significantly higher than in the random data (SD = 0.203, P <0.001; CV = 0.898, P <0.001), suggesting that individuals did not associate completely randomly but that short- or long term preferred companionships exist within the community (Whitehead, 2009a). Moreover, the proportion of non-zero elements was significantly larger (P <0.01) in the permuted data (proportion = 0.732) compared to the real data (proportion = 0.729) which suggests that some individuals may avoid others (Whitehead, 2009a). 

The correlation coefficient (r) between the true and estimated HWIs was 0.695 (SE = 0.042), indicating that the estimated association indices adequately represented the underlying social structure (Whitehead, 2009a). The estimate of social differentiation, S, was 0.633 (SE = 0.091). which indicates a well differentiated social system. However, a cophenetic correlation coefficient of 0.787 (less than the threshold of 0.8 for an effective social structure representation), combined with a maximum modularity (Q) of 0.264 (below the cut-off value of 0.3), shows a lack of evidence for the existence of social clusters within the community (Whitehead, 2009a) and therefore insufficient evidence for spatial segregation between individuals (Figure 2b). 

#### 51 309 *3.5. Power to detect trends in abundance*

According to the power analysis, detecting an annual decline of 1% in the population abundance with 80% certainty over a six-year period could only be achieved with CV of  $\leq 0.01$ whilst surveying every year. With the CV of 0.11 (the precision achieved in this study), on the other hand, an annual decline of 6% could be detected but only if surveys were conducted every year.

When considering a longer theoretical sampling period of 25 years and using a CV of 0.11, abundance surveys would have to be conducted every other year in order to detect an overall 25% decline (threshold in the Habitats Directive) in abundance (Figure 3). Surveying every three years with this level of precision would enable the detection of a decline of 26% and a survey frequency of every six years would only enable the detection of a larger 35% decrease in the population. On the other hand, if surveys were taking place every six years and the target was to detect the 25% decline, the CV around the estimate would have to be as low as 0.07 (Figure 3).

4. Discussion

#### 4.1. Abundance and movements of bottlenose dolphins on the west coast of Ireland

During summer 2014, the number of bottlenose dolphins using a 250km stretch of coastal waters between Connemara, Co. Galway and Co. Donegal on Ireland's west coast was estimated as 189 individuals (95% HPDI: 162-232). This estimate makes this the largest bottlenose dolphin population known to use Irish coastal waters, exceeding the numbers of animals estimated to inhabit the Shannon estuary (Berrow et al., 2012). Over 70 animals were matched with an existing catalogue with seven dolphins identified as far back as 2001. Such long-term re-identifications indicate that at least some of the animals using the coastal waters off the west and north-west of Ireland show a degree of site-fidelity, and it appears that the combined area between Connemara, Mullet peninsula and Donegal Bay form an important part of the home-range for a large number of bottlenose dolphins. While some of the members of this population were seen in only one of the coastal sites in 2014, several individuals exhibited high levels of mobility undertaking movements of over 250km during a single summer season. This high mobility presents challenges to the monitoring of the population, as wide-scale habitat use results in patchy temporal site occupancy with individuals and schools ranging freely over considerable distances around the Irish coast and further afield. Overall, the estimate derived in this study is remarkably similar to the cumulative number of animals (n = 179) identified around the Mullet peninsula in 2008-2009 (Oudejans et al., 2010) and to a previous abundance estimate of 171 (95% CI: 100-294) for dolphins using the waters around Connemara in 2009 (Ingram et al., 2009). However, the precision reached in this study (CV = (0.11) far exceeds the precision around the previous abundance estimate (CV = 0.28), making the 2014 estimate more robust for monitoring purposes, as shown by the power analysis. Nevertheless, biennial surveys would be required to detect the 25% overall decline in the 

population, and even in this case, the precision would have to remain at or below the CV of0.11 which may not be realistic year after year.

The impacts of anthropogenic habitat degradation on coastal dolphins require detailed understanding of the demographic parameters of the populations and the ranging behaviour and site-fidelity of individuals within the populations. Efficient and regular long-term monitoring of abundance is thus a vital part of the management of protected areas designated for bottlenose dolphin conservation. Studies in some other areas around the British Isles appear to show a high degree of site-fidelity to a single confined area (e.g. Shannon estuary and Sound of Barra) simplifying conservation management planning, but in other areas, such as the Moray Firth, changes in habitat use and distribution of bottlenose dolphins have been reported over the past 15 years (Arso Civil et al., 2019; Wilson et al., 2004). Similarly, the high degree of mobility of the coastal population in this study presents challenges in designing effective spatial management plans and implementing robust monitoring strategies. This study provides a benchmark for long-term monitoring of the population and its use of the West Connacht Coast SAC and illustrates how methods need to be adapted for monitoring more mobile populations. 

It is essential that bespoke monitoring strategies are designed to provide accurate and precise data on the status of populations that are sensitive to changes in abundance, population viability and survival rates. The Bayesian multi-site approach used here suits the transient behaviour of the west coast bottlenose dolphin population and provides a precise and comprehensive estimate of the abundance of animals in this large and variable habitat. A multi-site estimate is likely to better reflect the true abundance of the population than previous localised estimates due to the wider-scale sampling over a larger coastal area which increases the probability of encountering more of these animals as reflected in the lower CV value obtained in this study. Furthermore, it accounts for pseudoreplication of individuals sampled at different sites and is robust to unpredictable and unknown inter-annual variability in the distribution or occupancy of the animals. In contrast, a single site approach to monitoring this population could produce biased and highly variable abundance estimates if sections of the population were not encountered within a single site during a survey season. Further, it would be unfeasible to survey the entire known coastal range used by these animals as part of a routine monitoring strategy. With unpredictable and wide ranging movements of the animals, multi-site analysis of data enables simultaneous surveys of coastal areas by multiple research teams, and photo-identification surveys could be done opportunistically with help from a citizen science sightings network whilst maximising weather windows and keeping the costs low. 

The Bayesian multi-site approach assumes population closure with no births, deaths, immigration or emigration occurring in the area during the study period (Durban et al., 2005) as does more conventional closed maximum likelihood estimation frequently used in cetacean abundance studies (Bearzi et al., 2008; Brown et al., 2014; Gnone et al., 2011; Vermeulen and Cammareri, 2009). It is likely that although this assumption may be susceptible to violation due to the large scale of the animals' ranges, the inclusion of multiple sites over a broad geographical area should improve this model's performance. Furthermore, the short duration of the annual survey season (May-September in 2014) likely reduces the probability of migration of individuals out of the wider study area thus increasing the likelihood of effective closure of the sampled population. This is supported by the non-significant result of the closure test.

The bottlenose dolphins used the entire study area during 2014, with over half (54%) of all well-marked animals sighted in more than one of the three survey blocks, and 9% sighted in all of the study blocks with over 250km between the furthest sighing locations. Similarly, Cheney et al. (2013) found a large percentage of dolphins (58%) using more than one study site on the east coast of Scotland, however, the percentage of animals photographed in all of the sites was much smaller (only up to 1%) compared to this study, despite similar distances between the sites in both studies. In addition, up to 44% of the dolphins on the west coast of Scotland had similar long-range movements, with individuals ranging between the north and south of Skye (Cheney et al., 2013), even though the dolphins in the Sound of Barra did not exhibit movements outside this area. Some of the bottlenose dolphins in the present study that were encountered during surveys in 2014 had previously been recorded as far south as Co. Cork and appear to range widely around the west coast of Ireland and possibly beyond (Figure 2a). For example, a dolphin that was encountered in Donegal Bay in the summer of 2014 had previously been photographed in the Moray Firth in 2001 and around the Scottish Hebrides in 2004 (but is not one of the individuals regularly inhabiting these areas) (Robinson et al., 2012), thus providing further evidence of the long distance movements and transient behaviour of at least some of these animals. However, despite the large scale movements, the Irish west coast population appears to be genetically differentiated from the individuals sampled in east or west of Scotland (Nykänen et al., 2019). 

#### 56 410 *4.2. Social structure*

Even though different individuals showed varied ranging patterns with some dolphins ranging
over 500km and others encountered more locally, there was no evidence that the bottlenose

Page 15 of 39

dolphins occupying the waters of western Ireland form spatially segregated social clusters, unlike the social segregation previously documented between the coastal and offshore bottlenose dolphins (Oudejans et al., 2015). In fact, the west coast dolphins in this study seem to lack social groupings altogether, and it appears that the community consists of fluid social ties where individuals have a large number of associates, even though evidence of some short and long term companions and non-preferred associates was found. Bottlenose dolphins generally live in fluid "fission-fusion" societies (Connor et al., 2000), which means that animals usually form small social groups whose composition can change rapidly within the scale of a few hours. However, division into social clusters is common in some bottlenose dolphin societies (e.g. Chilvers and Corkeron, 2002), and this clustering has been linked to sex (Connor et al., 2011; Connor and Krützen, 2015; Frère et al., 2010; Smolker et al., 1992), specialized foraging techniques (Chilvers and Corkeron, 2001; Daura-Jorge et al., 2012; Krützen et al., 2005; 2014; Mann and Sargeant, 2003; Mann et al., 2008; Simões-Lopes et al., 1998; Smolker et al., 1997) and differential ranging patterns and spatial segregation (Louis et al., 2015; Lusseau et al., 2006). For example, Louis et al. (2015) found that individuals belonging to a social cluster were mainly observed within a specific area of the wider Normano-Breton Gulf, France. However, this clustering did not reflect genetic structuring as these dolphins were part of the same genetic population (Louis et al., 2014), so at least some spatial overlap is required to prevent genetic differentiation. In contrast, social separation was accompanied by genetic isolation between two adjacent populations of bottlenose dolphins occupying the Shannon estuary and the Irish west coast waters outside the estuary in a recent study (Nykänen et al., 2018). However, the lack of social clustering found in the present study may also be an artefact of the low number of re-sightings  $(n \ge 5)$  compared to some other studies; for example Frère et al. (2010) used a minimum of 30 identifications to estimate the social system of female Indo-Pacific bottlenose dolphins, Tursiops aduncus. 

4.3. Monitoring populations rather than protected areas? 

Most current marine conservation management requires the designation of some form of fixed marine protected area (MPA). However, MPAs have been criticised for being too small and failing to incorporate much of the range of the animals that they were designated for (Agardy et al., 2011; Hooker and Gerber, 2004; Wilson, 2016). Furthermore, the size, distribution and ranging behaviour of wild animal populations can alter as a consequence of changes in prey density and distribution (Angerbjorn et al., 1999; Friedlaender et al., 2006; Walton et al., 2001), habitat degradation or changes in environmental conditions linked to anthropogenic climate 

change (Harley et al., 2006; MacLeod et al., 2005; Parmesan and Yohe, 2003; Walther et al.,
2002). Therefore, MPAs with static boundaries advocated for the conservation of marine top
predators may not be the most appropriate method to protect mobile species. However, there
are a few examples where designation of static MPAs have been linked to improved survival
probability and increased population growth rate (Cheney et al., 2019; Gormley et al., 2012),
or the MPA has been large enough to encompass the majority of the range of most of the
animals (White et al., 2017).

There have been calls for more dynamic MPAs where the boundaries can be adjusted in response to changing species distributions or site use (Hartel et al., 2015; Hooker et al., 2011; Hooker and Gerber, 2004). However, it is likely that shifting of MPA boundaries would present such logistical and economic difficulties that the managing bodies and stakeholders may be reluctant to adopt this strategy. An alternative strategy could be to protect multiple clearly defined areas within a population's range where specific anthropogenic threats represent 'risk hot-spots' where impacts can be closely monitored and mitigated. Another proposed approach has been the development of more comprehensive marine spatial plans and ecosystem based management (Agardy et al., 2011; Halpern et al., 2010; MacLeod et al., 2005; Wilson, 2016) emphasizing integrated protection of the ecosystem as a whole while acknowledging connectivity among systems (MacLeod et al., 2005). In this context, the Great Barrier Reef Marine Park in Australia has been described as a success story of a large scale network of MPAs with its integrated and adaptive management (McCook et al., 2010). In Europe, the MSFD, where bottlenose dolphins are listed as one of the indicator species of good environmental status of coastal habitats, and the Natura 2000 network of SACs designated for the species seem to be a step in the right direction due to the potential of a network of MPAs enhancing connectivity among populations. However, transnational co-operation in the monitoring of these areas is required since the individual Member States are responsible for reporting on the status of species only in their own national SACs, and mobile populations can have ranges extending beyond country boundaries. 

If protection were focused on a population instead of a protected area, this protection could extend over the population's entire range (Reeves, 2000) rather than arbitrary portions of the population's range lying within an MPA. SACs designated for bottlenose dolphins in Irish coastal waters were designated based on limited spatial data from wide-scale and patchy surveys, and it is highly likely that the current SAC designations do not encompass the entire ecological needs of this species in the coastal waters of western Ireland. Moreover, monitoring

a population only within a designated area that covers only a portion of a population or species' habitat could give a biased view of the status of the population, if its range has expanded to other areas (Wilson et al., 2004) and a considerable part of the population is using areas outside the SAC (Arso Civil et al., 2019). However, the authorities responsible for the assessment of the conservation status of the Moray Firth SAC bottlenose dolphins have taken the recent range expansion and high mobility of the individuals into account in the monitoring of the population and with the advice that planned developments will need to be considered in assessments if they have potential impacts on bottlenose dolphins anywhere within the population's range, as they are likely to have a significant effect on the conservation objectives of the SAC (Arso Civil et al., 2019; SNH Natura Casework Guidance, 2019). This provides a good example of how governments can adjust previously set restricted management schemes under changing conditions, and that successful monitoring of mobile populations requires data collection across the population's range (Arso Civil et al., 2019). 

#### 492 4.4. Recommendations for future monitoring

The current approach to species conservation in the EU and other parts of the world is largely reliant on fixed area based protection and monitoring. While this approach may be applicable to some species in certain areas, this study provides an interesting contrast to studies of populations with higher site fidelity to coastal sites and illustrates how monitoring methods need to be adapted for more mobile populations.

Regular monitoring with at least biennial surveying on multiple known key sites across the population's range is recommended as the most appropriate monitoring strategy for the highly mobile bottlenose dolphins in this study, and a similar approach could be applied to other mobile marine and terrestrial populations worldwide. The multi-site approach maximises sighting probabilities at selected high-use sites throughout a large part of the population's range and produces accurate and precise estimates that are robust to temporary changes in ranging behaviour whilst delivering a cost-efficient way to monitor the population. Moreover, it offers great potential to be used as a tool for monitoring abundance in networks of connected MPAs, such as the Natura 2000 network of SACs designed to protect species across the EU. Indeed, evidence suggests that these coastal bottlenose dolphins have large ranges (Ingram et al., 2001) extending beyond national boundaries (O'Brien et al., 2009; Robinson et al., 2012). Even though genetic dispersal may be limited between the populations (Nykänen et al., 2019), 

510 increased transnational cooperation may be necessary for wide-scale monitoring in order to511 deliver on shared obligations under the MSFD and to some extent also the Habitats Directive.

Clearly, further research is required to uncover the entire ranging patterns and year-round habitat use of the bottlenose dolphins inhabiting coastal areas of western Ireland. Since unpredictable weather conditions on the west coast make surveying difficult and even unfeasible in the winter leading to data gaps in the populations' seasonal habitat use, photo-ID surveys could be supplemented with other methods, such as passive acoustic monitoring (PAM), to monitor temporal and spatial habitat use. In fact, preliminary PAM data suggests that bottlenose dolphins use key sites (outside and within the SAC) on the Irish west coast year round (Garagouni, 2019; Nykänen, 2016). Regular assessment of small mobile populations, such as the one in this study, is imperative to ensure that any deterioration in the conservation status of the population will be detected early, allowing for responsive mitigation measures to be put in place in a timely manner.

#### 523 Acknowledgements

This project was funded by Thomas Crawford Hayes Fund and National Parks and Wildlife Service (NPWS), Department of Arts, Heritage and Gaeltacht, Ireland. Special thanks to Oliver Ó Cadhla and Ferdia Marnell (NPWS) for their support over the course of the project. The authors would like to acknowledge Barbara Cheney for the advice in implementing the multi-site mark-recapture code, and everyone who contributed to the data collection, including Anneli Englund, Ailbhe Kavanagh, Rúadhán Ó Ceallaigh, Marie Kearns, Ross Culloch, Barry McGovern, Lochie O'Kelly, Damien Haberlin, Róisín Pinfield and Claudia Melville. We would also like to thank the Irish Whale and Dolphin Group, Brian and Cyndi Graham from St John's Point, Killary Cruises, skippers of the Pirate Queen, Shane Bisgood, John Britain, Máirtín Ó Meallaigh, Selkie sailing, Killybegs Coast Guard, Paddy Byrne, Anthony Irwin, Vincent Sweeney and Simon Sweeney for the help in providing sightings, and all the lovely people of Connemara, Mayo and Donegal. Thank you also to John Baxter and one anonymous referee for the useful feedback when revising the manuscript. 

53 537

#### **Conflict of interest**

539 The authors declare no conflict of interest.

2 3	541	References
4 5	• · -	
6 7	542	Agardy, T., di Sciara, G. N., and Christie, P. (2011). Mind the gap: Addressing the
8 9 10	543	shortcomings of marine protected areas through large scale marine spatial p
10 11 12	544	Marine Policy, 35(2), 226-232. https://doi.org/10.1016/j.marpol.2010.10.00
13 14	545	Angerbjorn, A., Tannerfeldt, M., and Erlinge, S. (1999). Predator-prey relationship
15 16	546	foxes and lemmings. Journal of Animal Ecology, 68(1), 34-49.
17 18	547	https://doi.org/10.1046/j.1365-2656.1999.00258.x
19 20 21	548	Arso Civil, M., Cheney, B., Quick, N. J., Thompson, P. M., and Hammond, P. S. (2
22 23	549	new approach to estimate fecundity rate from inter-birth intervals. Ecosphere
24 25	550	e01796. https://doi.org/10.1002/ecs2.1796
26 27 28	551	Arso Civil, M., Quick, N. J., Cheney, B., Pirotta, E., Thompson, P. M., and Hamme
28 29 30	552	(2019). Changing distribution of the east coast of Scotland bottlenose dolph
31 32	553	population and the challenges of area-based management. Aquatic Conserve
33 34 35	554	Marine and Freshwater Ecosystems, 29(S1), 178–196.
36 37	555	https://doi.org/10.1002/aqc.3102
38 39	556	Baker, I., O'Brien, J., McHugh, K., and Berrow, S. (2018a). Female reproductive p
40 41 42	557	and population demographics of bottlenose dolphins (Tursiops truncatus) in
42 43 44	558	Shannon Estuary, Ireland. Marine Biology, 165(1). https://doi.org/10.1007/s
45 46	559	017-3265-z
47 48	560	Baker, I., O'Brien, J., McHugh, K., Ingram, S. N., and Berrow, S. (2018b). Bottlen
49 50 51	561	dolphin (Tursiops truncatus) social structure in the Shannon Estuary, Irelan
52 53	562	distinguished by age- and area-related associations. Marine Mammal Science
54 55	563	458-487. https://doi.org/10.1111/mms.12462
56 57	564	Bearzi, G., Agazzi, S., Bonizzoni, S., Costa, M., and Azzellino, A. (2008). Dolphin
50 59		

shortcomings of marine protected areas through large scale marine spatial planning.
Marine Policy, 35(2), 226-232. https://doi.org/10.1016/j.marpol.2010.10.006
erbjorn, A., Tannerfeldt, M., and Erlinge, S. (1999). Predator-prey relationships: Arctic
foxes and lemmings. Journal of Animal Ecology, 68(1), 34-49.
https://doi.org/10.1046/j.1365-2656.1999.00258.x
Civil, M., Cheney, B., Quick, N. J., Thompson, P. M., and Hammond, P. S. (2017). A
new approach to estimate fecundity rate from inter-birth intervals. Ecosphere, 8(4),
e01796. https://doi.org/10.1002/ecs2.1796

- Civil, M., Quick, N. J., Cheney, B., Pirotta, E., Thompson, P. M., and Hammond, P. S. (2019). Changing distribution of the east coast of Scotland bottlenose dolphin population and the challenges of area-based management. Aquatic Conservation:
  - Marine and Freshwater Ecosystems, 29(S1), 178–196.
- https://doi.org/10.1002/aqc.3102
- r, I., O'Brien, J., McHugh, K., and Berrow, S. (2018a). Female reproductive parameters and population demographics of bottlenose dolphins (Tursiops truncatus) in the Shannon Estuary, Ireland. Marine Biology, 165(1). https://doi.org/10.1007/s00227-017-3265-z
- r, I., O'Brien, J., McHugh, K., Ingram, S. N., and Berrow, S. (2018b). Bottlenose dolphin (Tursiops truncatus) social structure in the Shannon Estuary, Ireland, is distinguished by age- and area-related associations. Marine Mammal Science, 34(2), 458-487. https://doi.org/10.1111/mms.12462
- zi, G., Agazzi, S., Bonizzoni, S., Costa, M., and Azzellino, A. (2008). Dolphins in a bottle: Abundance, residency patterns and conservation of bottlenose dolphins 565 60

2		
3 4 5 6 7 8	566	Tursiops truncatus in the semi-closed eutrophic Amvrakikos Gulf, Greece. Aquatic
	567	Conservation: Marine and Freshwater Ecosystems, 18(2), 130–146.
	568	https://doi.org/10.1002/aqc.843
10 11	569	Beck, S., Foote, A. D., Kötter, S., Harries, O., Mandleberg, L., Stevick, P. T., Whooley, P.,
12 13	570	and Durban, J. W. (2014). Using opportunistic photo-identifications to detect a
14 15	571	population decline of killer whales (Orcinus orca) in British and Irish waters. Journal
16 17 18 19 20	572	of the Marine Biological Association of the United Kingdom, 94(06), 1327–1333.
	573	https://doi.org/10.1017/S0025315413001124
21 22	574	Bejder, L., Fletcher, D., and Brager, S. (1998). A method for testing association patterns of
23 24 25	575	social animals. Animal Behaviour, 56, 719–725.
23 26 27	576	https://doi.org/10.1006/anbe.1998.0802
28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52	577	Berrow, S., O'Brien, J., Groth, L., Foley, A., and Voigt, K. (2012). Abundance Estimate of
	578	Bottlenose Dolphins (Tursiops truncatus) in the Lower River Shannon candidate
	579	Special Area of Conservation, Ireland. Aquatic Mammals, 38(2), 136-144.
	580	https://doi.org/10.1578/AM.38.2.2012.136
	581	Borgatti, S. P., Everett, M. G., and Freeman, L. C. (2002). Ucinet for Windows: Software for
	582	Social Network Analysis.
	583	Brown, A. M., Bejder, L., Pollock, K. H., and Allen, S. J. (2014). Abundance of coastal
	584	dolphins in Roebuck Bay, Western Australia. Report to WWF-Australia. Murdoch
	585	University Cetacean Research Unit.
	586	Caballero, S., Islas-Villanueva, V., Tezanos-Pinto, G., Duchene, S., Delgado-Estrella, A.,
	587	Sanchez-Okrucky, R., and Mignucci-Giannoni, A. A. (2012). Phylogeography,
53 54	588	genetic diversity and population structure of common bottlenose dolphins in the
55 56 57	589	Wider Caribbean inferred from analyses of mitochondrial DNA control region
58 59	590	sequences and microsatellite loci: Conservation and management implications.
60		

2 3	501	Animal Conservation 15(1) 95-112 https://doi.org/10.1111/j.1469-
4 5	551	minut Conservation, 15(1), 95-112. https://doi.org/10.1111/j.1409-
6 7	592	1795.2011.00493.x
, 8 9	593	Cairns, S. J., and Schwager, S. J. (1987). A comparison of association indexes. Animal
10 11	594	Behaviour, 35, 1454-1469. https://doi.org/10.1016/s0003-3472(87)80018-0
12 13	595	Cheney, B. J., Thompson, P. M., and Cordes, L. S. (2019). Increasing trends in fecundity and
14 15 16	596	calf survival of bottlenose dolphins in a marine protected area. Scientific Reports,
17 18	597	9(1), 1767. https://doi.org/10.1038/s41598-018-38278-9
19 20	598	Cheney, B., Thompson, P. M., Ingram, S. N., Hammond, P. S., Stevick, P. T., Durban, J. W.
21 22 23	599	et al. (2013). Integrating multiple data sources to assess the distribution and
23 24 25	600	abundance of bottlenose dolphins Tursiops truncatus in Scottish waters: Abundance
26 27	601	of bottlenose dolphins around Scotland. Mammal Review, 43(1), 71-88.
28 29 20	602	https://doi.org/10.1111/j.1365-2907.2011.00208.x
31 32	603	Chilvers, B. L., and Corkeron, P. J. (2001). Trawling and bottlenose dolphins' social
33 34	604	structure. Proceedings of the Royal Society of London Series B-Biological Sciences,
35 36	605	268(1479), 1901-1905. https://doi.org/10.1098/rspb.2001.1732
37 38 39	606	Chilvers, B. L., and Corkeron, P. J. (2002). Association patterns of bottlenose dolphins
40 41	607	(Tursiops aduncus) off Point Lookout, Queensland, Australia. Canadian Journal of
42 43	608	Zoology-Revue Canadienne De Zoologie, 80(6), 973–979.
44 45 46	609	https://doi.org/10.1139/z02-075
47 48	610	Connor, R. C., Watson-Capps, J. J., Sherwin, W. B., and Krützen, M. (2011). A new level of
49 50	611	complexity in the male alliance networks of Indian Ocean bottlenose dolphins
51 52	612	(Tursiops sp.). Biology Letters, 7(4), 623-626. https://doi.org/10.1098/rsbl.2010.0852
53 54 55 56 57 58 59 60	613	

3 4	614	Connor, R.C., Wells, R. S., Mann, J., and Read, A. J. (2000). The Bottlenose Dolphin: Social
5 6	615	Relationships in a Fission-Fusion Society. In: J. Mann (Ed.), Cetacean Societies:
/ 8 0	616	Field Studies of Dolphins and Whales. University of Chicago Press.
9 10 11	617	Connor, Richard C., and Krützen, M. (2015). Male dolphin alliances in Shark Bay: Changing
12 13	618	perspectives in a 30-year study. Animal Behaviour, 103, 223-235.
14 15	619	https://doi.org/10.1016/j.anbehav.2015.02.019
16 17 18	620	Daura-Jorge, F. G., Cantor, M., Ingram, S. N., Lusseau, D., and Simoes-Lopes, P. C. (2012).
19 20	621	The structure of a bottlenose dolphin society is coupled to a unique foraging
21 22	622	cooperation with artisanal fishermen. Biology Letters, 8(5), 702-705.
23 24 25	623	https://doi.org/10.1098/rsb1.2012.0174
25 26 27	624	Durban, J., Ellifrit, D., Dahlheim, M., Waite, J., Matkin, C., Barrett-Lennard, L. et al. (2010).
28 29	625	Photographic mark-recapture analysis of clustered mammal-eating killer whales
30 31 22	626	around the Aleutian Islands and Gulf of Alaska. Marine Biology, 157(7), 1591–1604.
32 33 34	627	https://doi.org/10.1007/s00227-010-1432-6
35 36	628	Durban, J. W., Elston, D. A., Ellifrit, D. K., Dickson, E., Hammond, P. S., and Thompson, P.
37 38	629	M. (2005). Multisite mark-recapture for cetaceans: Population estimates with
39 40 41	630	Bayesian model averaging. Marine Mammal Science, 21(1), 80–92.
42 43	631	https://doi.org/10.1111/j.1748-7692.2005.tb01209.x
44 45	632	Englund, A., Ingram, S. N., & Rogan, E. (2008). An updated population status report for
46 47 48	633	bottlenose dolphins using the Lower River Shannon SAC in 2008. Final Report to the
49 50	634	National Parks and Wildlife Service.
51 52	635	Fearnbach, H., Durban, J., Parsons, K., and Claridge, D. (2012). Photographic mark-recapture
53 54 55	636	analysis of local dynamics within an open population of dolphins. Ecological
55 56 57 58 59 60	637	Applications, 22(5), 1689–1700.

2		
3 4	638	Fernández, R., Santos, M. B., Pierce, G. J., Llavona, Á., López, A., Silva, M. A. et al. (2011).
5 6	639	Fine-scale genetic structure of bottlenose dolphins, Tursiops truncatus, in Atlantic
/ 8 0	640	coastal waters of the Iberian Peninsula. Hydrobiologia, 670(1), 111-125.
9 10 11	641	https://doi.org/10.1007/s10750-011-0669-5
12 13	642	Foley, A., McGrath, D., Berrow, S., and Gerritsen, H. (2010). Social Structure Within the
14 15	643	Bottlenose Dolphin (Tursiops truncatus) Population in the Shannon Estuary, Ireland.
16 17 18	644	Aquatic Mammals, 36(4), 372-381. https://doi.org/10.1578/AM.36.4.2010.372
19 20	645	Frère, C. H., Krützen, M., Mann, J., Watson-Capps, J. J., Tsai, Y. J., Patterson, E. M. et al.
21 22	646	(2010). Home range overlap, matrilineal and biparental kinship drive female
23 24 25	647	associations in bottlenose dolphins. Animal Behaviour, 80(3), 481-486.
25 26 27	648	https://doi.org/10.1016/j.anbehav.2010.06.007
28 29	649	Friedlaender, A., Halpin, P., Qian, S., Lawson, G., Wiebe, P., Thiele, D. et al. (2006). Whale
30 31	650	distribution in relation to prey abundance and oceanographic processes in shelf waters
32 33 34	651	of the Western Antarctic Peninsula. Marine Ecology Progress Series, 317, 297–310.
35 36	652	https://doi.org/10.3354/meps317297
37 38	653	Fruet, P. F., Daura-Jorge, F. G., Möller, L. M., Genoves, R. C., and Secchi, E. R. (2015).
39 40 41	654	Abundance and demography of bottlenose dolphins inhabiting a subtropical estuary in
41 42 43	655	the Southwestern Atlantic Ocean. Journal of Mammalogy, 96(2), 332-343.
44 45	656	https://doi.org/10.1093/jmammal/gyv035
46 47	657	Garagouni, M. (2019). Habitat preferences and movement patterns of bottlenose dolphins at
48 49 50	658	various spatial and temporal scales [PhD Thesis, University College Cork].
50 51 52	659	http://hdl.handle.net/10468/8618
53 54	660	Gerrodette, T. (1987). A power analysis for detecting trends. <i>Ecology</i> , 68(5), 1364–1372.
55 56 57 58	661	https://doi.org/10.2307/1939220
59 60		

2 3 4	662	Gerrodette, T. (1991). Models for Power of Detecting Trends: A Reply to Link and Hatfield.
5 6 7	663	Ecology, 72(5), 1889-1892. https://doi.org/10.2307/1940986
7 8	664	Gnone, G., Bellingeri, M., Dhermain, F., Dupraz, F., Nuti, S., Bedocchi, D. et al. (2011).
9 10 11	665	Distribution, abundance, and movements of the bottlenose dolphin (Tursiops
12 13	666	truncatus) in the Pelagos Sanctuary MPA (north-west Mediterranean Sea). Aquatic
14 15	667	Conservation: Marine and Freshwater Ecosystems, 21(4), 372–388.
16 17 18	668	https://doi.org/10.1002/aqc.1191
19 20	669	Gormley, A. M., Slooten, E., Dawson, S., Barker, R. J., Rayment, W., du Fresne, S. et al.
21 22	670	(2012). First evidence that marine protected areas can work for marine mammals:
23 24 25	671	Protected areas for marine mammals. Journal of Applied Ecology, 49(2), 474–480.
26 27	672	https://doi.org/10.1111/j.1365-2664.2012.02121.x
28 29	673	Halpern, B. S., Lester, S. E., and McLeod, K. L. (2010). Placing marine protected areas onto
30 31 32	674	the ecosystem-based management seascape. Proceedings of the National Academy of
32 33 34 35 36 37 38 39 40 41	675	Sciences, 107(43), 18312-18317. https://doi.org/10.1073/pnas.0908503107
	676	Hammond, P. S., Macleod, K., Berggren, P., Borchers, D. L., Burt, L., Cañadas, A. et al.
	677	(2013). Cetacean abundance and distribution in European Atlantic shelf waters to
	678	inform conservation and management. Biological Conservation, 64, 107–122.
42 43	679	Harley, C. D. G., Randall Hughes, A., Hultgren, K. M., Miner, B. G., Sorte, C. J. B.,
44 45	680	Thornber, C. S. et al. (2006). The impacts of climate change in coastal marine
46 47 48	681	systems. Ecology Letters, 9(2), 228–241. https://doi.org/10.1111/j.1461-
48 49 50	682	0248.2005.00871.x
51 52	683	Hartel, E. F., Constantine, R., and Torres, L. G. (2015). Changes in habitat use patterns by
53 54 55	684	bottlenose dolphins over a 10-year period render static management boundaries
56 57	685	ineffective. Aquatic Conservation: Marine and Freshwater Ecosystems, 25(5), 701-
58 59 60	686	711. https://doi.org/10.1002/aqc.2465

2		
3 4	687	Hooker, S., Cañadas, A., Hyrenbach, K., Corrigan, C., Polovina, J., and Reeves, R. (2011).
5 6 7	688	Making protected area networks effective for marine top predators. Endangered
7 8 9	689	Species Research, 13(3), 203–218. https://doi.org/10.3354/esr00322
10 11	690	Hooker, S. K., and Gerber, L. R. (2004). Marine reserves as a tool for ecosystem-based
12 13	691	management: The potential importance of megafauna. <i>Bioscience</i> , 54(1), 27–39.
14 15 16	692	http://bioscience.oxfordjournals.org/content/54/1/27.short
17 18	693	Ingram, S., Englund, A., and Rogan, E. (2001). An extensive survey of bottlenose dolphins
19 20	694	(Tursiops truncatus) on the west coast of Ireland. Heritage Council Report no.
21 22	695	WLD/2001/42.
23 24 25	696	Ingram, S., Englund, A., and Rogan, E. (2003). Habitat use, abundance and site-fidelity of
26 27	697	bottlenose dolphins (Tursiops truncatus) in Connemara coastal waters, Co. Galway.
28 29	698	Report to the Heritage Council, Wildlife Grant #12314.
30 31 32	699	Ingram, S., Kavanagh, A., Englund, A., and Rogan, E. (2009). Site assessment of the waters
33 34	700	of northwest Connemara. A survey of bottlenose dolphins (Tursiops truncatus). Report
35 36	701	for the National Parks and Wildlife Service of Ireland.
37 38 30	702	Ingram, S., and Rogan, E. (2002). Identifying critical areas and habitat preferences of
39 40 41	703	bottlenose dolphins Tursiops truncatus. Marine Ecology-Progress Series, 244, 247-
42 43	704	255. https://doi.org/10.3354/meps244247
44 45 46	705	Ingram, S., and Rogan, E. (2003). Bottlenose dolphins (Tursiops truncatus) in the Shannon
40 47 48	706	Estuary and selected areas of the west-coast of Ireland. Report to the National Parks
49 50	707	and Wildlife Service.
51 52	708	Irvine, A. B., Scott, M., Wells, R., and Kaufmann, J. H. (1981). Movements and activities of
53 54 55 56 57	709	the Atlantic bottlenose dolphin, Tursiops truncatus, near Sarasota, Florida (Vol. 79).
58 59		
60		

2 3	710	Jepson, P. D., Deaville, R., Barber, J. L., Aguilar, À., Borrell, A., Murphy, S. et al. (2016).
4 5 6	711	PCB pollution continues to impact populations of orcas and other dolphins in
7 8	712	European waters. Scientific Reports, 6, 18573. https://doi.org/10.1038/srep18573
9 10 11	713	Kendall, W. L. (1999). Robustness of Closed Capture-Recapture Methods to Violations of the
12 13	714	Closure Assumption. <i>Ecology</i> , 80(8), 2517. https://doi.org/10.2307/177237
14 15 16	715	Krützen, M., Kreicker, S., MacLeod, C. D., Learmonth, J., Kopps, A. M., Walsham, P., et al.
10 17 18	716	(2014). Cultural transmission of tool use by Indo-Pacific bottlenose dolphins
19 20	717	(Tursiops sp.) provides access to a novel foraging niche. Proceedings of the Royal
21 22 22	718	Society B: Biological Sciences, 281(1784), 20140374–20140374.
23 24 25	719	https://doi.org/10.1098/rspb.2014.0374
26 27	720	Krützen, M., Mann, J., Heithaus, M. R., Connor, R. C., Bejder, L., and Sherwin, W. B.
28 29	721	(2005). Cultural transmission of tool use in bottlenose dolphins. Proceedings of the
30 31 32	722	National Academy of Sciences of the United States of America, 102(25), 8939–8943.
33 34	723	http://www.pnas.org/content/102/25/8939.short
35 36	724	Lacy, R. C. (1987). Loss of Genetic Diversity from Managed Populations: Interacting Effects
37 38 39	725	of Drift, Mutation, Immigration, Selection, and Population Subdivision. Conservation
40 41	726	Biology, 1(2), 143–158. https://doi.org/10.1111/j.1523-1739.1987.tb00023.x
42 43	727	Louis, M., Gally, F., Barbraud, C., Béesau, J., Tixier, P., Simon-Bouhet, B., et al. (2015).
44 45 46	728	Social structure and abundance of coastal bottlenose dolphins, Tursiops truncatus, in
40 47 48	729	the Normano-Breton gulf, English Channel. Journal of Mammalogy, 96(3), 481-493.
49 50	730	Louis, M., Viricel, A., Lucas, T., Peltier, H., Alfonsi, E., Berrow, S. et al. (2014). Habitat-
51 52	731	driven population structure of bottlenose dolphins, Tursiops truncatus, in the North-
53 54 55	732	East Atlantic. <i>Molecular Ecology</i> , 23(4), 857–874. https://doi.org/10.1111/mec.12653
56 57	733	Lunn, D. J., Thomas, A., Best, N., and Spiegelhalter, D. (2000). WinBUGS-a Bayesian
58 59 60	734	modelling framework: Concepts, structure, and extensibility. Statistics and

2 3	705	$C_{\text{commuting}} = 10(4) = 225 = 227$
4	/35	<i>Computing</i> , 10(4), 525–557.
6	736	http://link.springer.com/article/10.1023/A:1008929526011
/ 8 9	737	Lusseau, D. (2007). Why are male social relationships complex in the Doubtful Sound
10 11	738	bottlenose dolphin population? Plos One, 2(4), e348.
12 13	739	https://doi.org/10.1371/journal.pone.0000348
14 15 16	740	Lusseau, D, Bain, D., Williams, R., and Smith, J. (2009). Vessel traffic disrupts the foraging
17 18	741	behavior of southern resident killer whales Orcinus orca. Endangered Species
19 20	742	Research, 6, 211–221. https://doi.org/10.3354/esr00154
21 22 22	743	Lusseau, D., Wilson, B., Hammond, P. S., Grellier, K., Durban, J. W., Parsons et al. (2006).
23 24 25	744	Quantifying the influence of sociality on population structure in bottlenose dolphins.
26 27	745	The Journal of Animal Ecology, 75(1), 14–24.
28 29	746	MacLeod, C. D., Bannon, S. M., Pierce, G. J., Schweder, C., Learmonth, J. A., Herman, J. S.
30 31 32	747	et al. (2005). Climate change and the cetacean community of north-west Scotland.
33 34	748	Biological Conservation, 124(4), 477–483.
35 36	749	https://doi.org/10.1016/j.biocon.2005.02.004
37 38 20	750	Mann, J., and Sargeant, B. L. (2003). Like mother, like calf: The ontogeny of foraging
40 41	751	traditions in wild Indian Ocean bottlenose dolphins (Tursiops sp.). In: D. Fragaszy
42 43	752	and S. Perry (Eds.), The biology of traditions: Models and evidence (pp. 236–266).
44 45	753	Cambridge University Press.
46 47 48	754	Mann, J., Sargeant, B. L., Watson-Capps, J. J., Gibson, Q. A., Heithaus, M. R., Connor, R. et
49 50	755	al. (2008). Why do dolphins carry sponges? PLoS One, 3(12), e3868.
51 52	756	http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0003868
53 54 55	757	Mäntyniemi, S., and Romakkaniemi, A. (2002). Bayesian mark-recapture estimation with an
56 57	758	application to a salmonid smolt population. Canadian Journal of Fisheries and
58 59 60	759	Aquatic Sciences, 59, 1748-1758. https://doi.org/10.1139/f02-146

http://mc.manuscriptcentral.com/aqc

2		
3 4	760	McCook, L. J., Ayling, T., Cappo, M., Choat, J. H., Evans, R. D., De Freitas, D. M. et al.
5 6	761	(2010). Adaptive management of the Great Barrier Reef: A globally significant
7 8	762	demonstration of the benefits of networks of marine reserves. Proceedings of the
9 10 11	763	National Academy of Sciences, 107(43), 18278–18285.
12 13 14 15 16 17 18 19 20	764	https://doi.org/10.1073/pnas.0909335107
	765	Michielsens, C. G., McAllister, M. K., Kuikka, S., Pakarinen, T., Karlsson, L.,
	766	Romakkaniemi, A. et al. (2006). A Bayesian state space mark recapture model to
	767	estimate exploitation rates in mixed-stock fisheries. Canadian Journal of Fisheries
21 22	768	and Aquatic Sciences, 63(2), 321–334.
23 24 25	769	http://www.nrcresearchpress.com/doi/abs/10.1139/f05-215
25 26 27 28 29	770	Mirimin, L., Miller, R., Dillane, E., Berrow, S. D., Ingram, S., Cross, T. F. et al. (2011). Fine-
	771	scale population genetic structuring of bottlenose dolphins in Irish coastal waters.
30 31	772	Animal Conservation, 14(4), 342-353. https://doi.org/10.1111/j.1469-
32 33 34	773	1795.2010.00432.x
35 36	774	Moore, J. E., and Barlow, J. (2011). Bayesian state-space model of fin whale abundance
37 38	775	trends from a 1991-2008 time series of line-transect surveys in the California Current:
39 40 41	776	Bayesian trend analysis from line-transect data. Journal of Applied Ecology, 48(5),
42 43	777	1195–1205. https://doi.org/10.1111/j.1365-2664.2011.02018.x
44 45	778	Newman, M. E. J. (2004). Analysis of weighted networks. Physical Review E, 70, 056131.
46 47 48	779	https://doi.org/056131 10.1103/PhysRevE.70.056131
49 50	780	Newman, M. E. J. (2006). Modularity and community structure in networks. Proceedings of
51 52	781	the National Academy of Sciences, 103(23), 8577–8582.
53 54	782	https://doi.org/10.1073/pnas.0601602103
55 56		
57 58		
59		

2 3	783	Nykänen, M. (2016). Phylogeography, population structure, abundance and habitat use of				
4 5 6	784	bottlenose dolphins, Tursiops truncatus, on the west coast of Ireland [University				
7 8	785	College Cork]. https://cora.ucc.ie/handle/10468/3828				
9 10 11	786	6 Nykänen, M., Dillane, E., Englund, A., Foote, A. D., Ingram, S. N., Louis, M. et al. (				
12 13	2 3 787 Quantifying dispersal between marine protected areas by a highly mobile					
14 15 16	788bottlenose dolphin, Tursiops truncatus. Ecology and Evolution.					
17 18	789 https://doi.org/10.1002/ece3.4343					
19 20	19 20 20 Nykänen, M., Louis, M., Dillane, E., Alfonsi, E., Berrow, S., O'Brien, J. et al. (20					
21 22 23	791	1 Fine-scale population structure and connectivity of bottlenose dolphins, <i>TURSIOPS</i>				
24 25	792	TRUNCATUS, in European waters and implications for conservation. Aquatic				
26 27	793	3 <i>Conservation: Marine and Freshwater Ecosystems</i> , 29(S1), 197–211.				
28 29 30	794	https://doi.org/10.1002/aqc.3139				
31 32	795	O'Brien, J., Berrow, S., Ryan, C., McGrath, D., O'Connor, I., Giovanna, P. et al. (2009). A				
33 34	796	note on long-distance matches of bottlenose dolphins (Tursiops truncatus) around the				
35 36 37	797	Irish coast using photo-identification. Journal of Cetacean Research and Managment.				
38 39	798	Otis, D. L., Burnham, K. P., White, G. C., and Anderson, D. R. (1978). Statistical inference				
40 41	799	from capture data on closed animal populations. <i>Wildlife Monographs</i> , 62, 7–135.				
42 43 44	800	Oudejans, M. G., Ingram, S. N., Englund, A., and Rogan, E. (2010). Bottlenose dolphins in				
45 46	801	Connemara & Mayo in 2008 and 2009; population structure and movement patterns				
47 48	802	between two coastal sites in northwest Mayo. Report to the National Parks and				
49 50 51	803	Wildlife Service.				
52 53	804	Oudejans, M. G., Visser, F., Englund, A., Rogan, E., and Ingram, S. N. (2015). Evidence for				
54 55	805	Distinct Coastal and Offshore Communities of Bottlenose Dolphins in the North East				
56 57 58 59 60	806	Atlantic. PLOS ONE, 10(4), e0122668. https://doi.org/10.1371/journal.pone.0122668				

2					
3 4	807	Palumbi, S. R. (2001). The ecology of marine protected areas. In: M. Bertness, S. Gaines, and			
5 6 7	808	M. Hay (Eds.), Marine community ecology (pp. 509–530). Sinauer Associates Inc.			
7 8 9	809	Parmesan, C., and Yohe, G. (2003). A globally coherent fingerprint of climate change			
10 11	810	impacts across natural systems. Nature, 421(6918), 37-42.			
12 13	811	https://doi.org/10.1038/nature01286			
14 15 16	812	Pirotta, E., Merchant, N. D., Thompson, P. M., Barton, T. R., and Lusseau, D. (2015).			
16 17 18	813	Quantifying the effect of boat disturbance on bottlenose dolphin foraging activity.			
19 20	<sup>19</sup> 20 814 <i>Biological Conservation</i> , 181, 82–89. https://doi.org/10.1016/j.biocon.201				
21 22	815 Quick, N. J., Cheney, B., Islas-Villanueva, V., Janik, V. M., Thompson, P. M., and				
23 24 25	816	Hammond, P. S. (2014). The east coast of Scotland bottlenose dolphin population:			
26 27	817	Improving understanding of ecology outside the Moray Firth SAC. Report produced			
28 29	818	as part of the UK Department of Energy and Climate Change's offshore energy			
30 31 32	819	Strategic Environmental Assessment programme (No. 14D-086; OESEA2 Supporting			
32 33 34	820	Documents, p. 87). Department of Energy and Climate Change.			
35 36	821	Read, A. J., Urian, K. W., Wilson, B., and Waples, D. M. (2003). Abundance of bottlenose			
37 38	822	dolphins in the bays, sounds and estuaries of North Carolina. Marine Mammal			
39 40 41	823	Science, 19(1), 59-073. https://doi.org/10.1111/j.1748-7692.2003.tb01092.x			
42 43	824	Reeves, R. R. (2000). The Value of Sanctuaries, Parks, and Reserves (Protected Areas) as			
44 45	825	Tools for Conserving Marine Mammals. Final Report to the Marine Mammal			
46 47 48	826	Commission, contract number T74465385. Marine Mammal Commission.			
49 50	827	Robinson, K. P., O'Brien, J., Berrow, S., Cheney, B., Costa, M., Elsfield S. M. et al. (2012).			
51 52	828	Discrete or not so discrete: Long distance movements by coastal bottlenose dolphins			
53 54	829	in UK and Irish waters. Journal of Cetacean Research and Management.			
55 56 57 58 59 60	830	http://cual.openrepository.com/cual/handle/10759/560746			

2 3 4	831	Rogan, E., Breen, P., Mackey, M., Cañadas, A., Scheidat, M., Geelhoed, S. et al. (2018).
5 6	832	Aerial surveys of cetaceans and seabirds in Irish waters: Occurrence, distribution
7 8 9 10 11 12 13 14 15 16 17 18 19 20	833	and abundance in 2015-2017. Department of Communications, Climate Action &
	834	Environment and National Parks and Wildlife Service (NPWS), Department of
	835	Culture, Heritage and the Gaeltacht.
	836	Rosel, P. E., Mullin, K. D., Garrison, L., Schwacke, L., Adams, J., Balmer, B. et al. (2011).
	837	Photo-identification capture-mark-recapture techniques for estimating abundance of
	838	bay, sound and estuary populations of Bottlenose dolphins along the U.S. East Coast
21 22	839	and Gulf of Mexico: A workshop report. 38.
23 24	840	Simões-Lopes, P. C., Fabián, M. E., and Menegheti, J. O. (1998). Dolphin interactions with
26 27	841	the mullet artisanal fishing on Southern Brazil: A qualitative and quantitative
28 29	842	approach. Revista Brasileira de Zoologia, 15(3), 709–726.
30 31 32 33 34 35 36	843	https://doi.org/10.1590/S0101-81751998000300016
	844	Smolker, R. A., Richards, A. F., Connor, R. C., and Pepper, J. W. (1992). Sex-differences in
	845	patterns of association among Indian Ocean bottlenose dolphins. Behaviour, 123, 38-
37 38	846	69. https://doi.org/10.1163/156853992x00101
39 40 41	847	Smolker, R., Richards, A., Connor, R., Mann, J., and Berggren, P. (1997). Sponge Carrying
42 43	848	by Dolphins (Delphinidae, Tursiops sp.): A Foraging Specialization Involving Tool
44 45	849	Use? Ethology, 103(6), 454–465. https://doi.org/10.1111/j.1439-0310.1997.tb00160.x
46 47 48	850	SNH Natura Casework Guidance. (2019). Natura casework guidance-How to consider
48 49 50	851	plans and projects affecting Special Areas of Conservation (SACs) and Special
51 52	852	Protection Areas (SPAs): Version 9.0.
53 54 55 56 57 58 59	853	Vermeulen, E., and Cammareri, A. (2009). Residency Patterns, Abundance, and Social
	854	Composition of Bottlenose Dolphins (Tursiops truncatus) in Bahía San Antonio,
00		

2					
3 4	855	Patagonia, Argentina. Aquatic Mammals, 35(3), 378-385.			
5 6	856	https://doi.org/10.1578/AM.35.3.2009.378			
7 8 0	857	Walther, G. R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C. et al. (2002).			
9 10 11	858	Ecological responses to recent climate change. Nature, 416(6879), 389-395.			
12 13	859	https://doi.org/10.1038/416389a			
14 15	860	Walton, L. R., Cluff, H. D., Paquet, P. C., and Ramsay, M. A. (2001). Movement Patterns o			
10 17 18	861	Barren-Ground Wolves in the Central Canadian Arctic. Journal of Mammalogy,			
19 20	862	82(3), 867–876. http://www.jstor.org/stable/1383622			
21 22	863	Wells, R., and Scott, M. D. (2009). Common bottlenose dolphin (Tursiops truncatus). In: W.			
23 24 25	864	Perrin, B. Wursig, and J. Thewissen (Eds.), Encyclopedia of Marine Mammals.			
26 27	865	Second Edition (pp. 249–255). Elsevier.			
28 29	866	White, T. D., Carlisle, A. B., Kroodsma, D. A., Block, B. A., Casagrandi, R., De Leo, G. A.			
30 31 32	867	et al. (2017). Assessing the effectiveness of a large marine protected area for reef			
33 34	868	shark conservation. Biological Conservation, 207, 64–71.			
35 36	869	https://doi.org/10.1016/j.biocon.2017.01.009			
37 38 30	870	Whitehead, H. (1999). Testing association patterns of social animals. Animal Behaviour, 57,			
40 41	871	26–29.			
42 43	872	Whitehead, H. (2009a). Programs for analyzing social structure. SOCPROG2.4 (for			
44 45	873	MATLAB7.7.0.471, release 2008b).			
46 47 48	874	Whitehead, H. (2009b). SOCPROG programs: Analysing animal social structures.			
49 50	875	Behavioral Ecology and Sociobiology, 63(5), 765–778.			
51 52	876	https://doi.org/10.1007/s00265-008-0697-y			
53 54 55	877	Williams, R., Bain, D., Smith, J., and Lusseau, D. (2009). Effects of vessels on behaviour			
56 57	878	patterns of individual southern resident killer whales Orcinus orca. Endangered			
58 59 60	879	Species Research, 6, 199–209. https://doi.org/10.3354/esr00150			

1 ว		
2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25	880	Williams, R., Erbe, C., Ashe, E., Beerman, A., and Smith, J. (2014). Severity of killer whale
	881	behavioral responses to ship noise: A dose-response study. Marine Pollution Bulletin,
	882	79(1-2), 254-260. https://doi.org/10.1016/j.marpolbul.2013.12.004
	883	Wilson, B. (2016). Might marine protected areas for mobile megafauna suit their proponents
	884	more than the animals? Aquatic Conservation: Marine and Freshwater Ecosystems,
	885	26(1), 3-8. https://doi.org/10.1002/aqc.2619
	886	Wilson, B., Hammond, P. S., and Thompson, P. M. (1999). Estimating size and assessing
	887	trends in a coastal bottlenose dolphin population. Ecological Applications, 9(1), 288-
	888	300. http://onlinelibrary.wiley.com/doi/10.1890/1051-
	889	0761(1999)009%5B0288:ESAATI%5D2.0.CO;2/full
26 27	890	Wilson, B., Reid, R. J., Grellier, K., Thompson, P. M., and Hammond, P. S. (2004).
28 29 30 31 32 33 34	891	Considering the temporal when managing the spatial: A population range expansion
	892	impacts protected areas-based management for bottlenose dolphins. Animal
	893	Conservation, 7(4), 331-338. https://doi.org/10.1017/S1367943004001581
35 36	894	Würsig, B., and Würsig, M. (1977). Photographic determination of group size, composition
37 38	895	and stability of coastal porpoises (Tursiops truncatus). Science, 198(4318), 755–756.
39 40 41 42 43	896	https://doi.org/10.1126/science.198.4318.755
	897	
44 45 46	898	
47 48		
49 50		
51		
52 53		
54 55		
56		
57 58		
59 60		
00		

#### Tables

Table 1. Contingency table of the counts of well-marked (M1) bottlenose dolphins present (Y) or absent (N) in each of the study blocks Connemara, Mullet peninsula and Donegal Bay in 2014.

	Block			
Count	Connemara	Mullet peninsula	Donegal Bay	
8	Y	Y	Y	
2	Y	Y	Ν	
13	Ν	Y	Ν	
6	Y	Ν	Ν	
28	Ν	Y	Y	
11	Y	Ν	Y	
23	Ν	Ν	Y	
NA†	Ν	N	N	

<sup>†</sup> The missing value (NA) represents the number of individuals that were not seen in any of the study blocks (i.e., "missed" well-marked dolphins) RZ. RZ

# **Figure legends**

907 Figure 1. The three coastal 'blocks' (circled areas), Donegal Bay, Mullet Peninsula and
908 Connemara surveyed during summer 2014. The red areas in the inset map show the location of
909 the two bottlenose dolphin SACs in Irish waters, and the hatched area in the large map shows
910 the extent of the West Connacht SAC. Triangles denote the location of encounters with
911 bottlenose dolphin schools during the study period.

Figure 2. The 39 most sighted well-marked bottlenose dolphins and a) the geographic range of their sighting locations, and b) their social network diagram. The bottlenose dolphins were encountered at least on five occasions during the data collection period 2001-2014 on the west coast of Ireland. The individual ID numbers are given on the x-axis in figure a) and next to the circles in figure b). The outline of Ireland has been scaled to correspond with the sighting latitudes. The centre line in the boxplot and the bottom and top of the box represent the 50<sup>th</sup>. 25<sup>th</sup> and 75<sup>th</sup> percentiles, respectively, and the whiskers the 5<sup>th</sup> and 95<sup>th</sup> percentile in figure a). The dots represent rarely visited latitudes. The data has been arranged by increasing median latitude. The length of the line in the network diagram in figure b) inversely represents the strength of the association between a dyad calculated as half-weight association index. 

Figure 3. The effect of coefficient of variation (CV) on the minimum detectable decline in the
abundance of a theoretical population with different survey frequency; surveys conducted
every six years (five years between surveys), every three years (two years between surveys),
every other year or annually, over a theoretical 25-year period.

#### Appendices

**Appendix 1.** Examples of bottlenose dolphin fins showing the three grades of mark severity used in photograph analysis. Each dolphin was graded from one to three as follows: (A) grade M1 marks, consisting of significant fin damage or deep scarring that were considered permanent; (B) grade M2 marking that consist of deep tooth rakes and lesions, with only minor cuts present; (C) fin with grade M3 marks, having only superficial rakes and lesions. Grade M1 (and to some extent, M2) are considered to last many years, enabling long-term identification of these dolphins. In contrast, 'superficial' markings (grade M3), such as tooth rakes may fade and heal within a relatively short period of time and inter-annual re-sighting probabilities of these animals are likely to be reduced. 

Appendix 2. Scoring criteria for the quality of bottlenose dolphin identification photographs. 

**Appendix 3.** Range of sighting latitude (difference between maximum and minimum latitude) plotted against the number of times each individual well-marked bottlenose dolphin was sighted on the west coast of Ireland 2001–2014. A Loess smooth curve is fitted through the observations.

Pere.

http://mc.manuscriptcentral.com/aqc





Figure 1. The three coastal 'blocks' (circled areas), Donegal Bay, Mullet Peninsula and Connemara surveyed during summer 2014. The red areas in the inset map show the location of the two bottlenose dolphin SACs in Irish waters, and the hatched area in the large map shows the extent of the West Connacht SAC. Triangles denote the location of encounters with bottlenose dolphin schools during the study period.

268x201mm (300 x 300 DPI)





Figure 2. The 39 most sighted well-marked bottlenose dolphins and a) the geographic range of their sighting locations, and b) their social network diagram. The bottlenose dolphins were encountered at least on five occasions during the data collection period 2001-2014 on the west coast of Ireland. The individual ID numbers are given on the x-axis in figure a) and next to the circles in figure b). The outline of Ireland has been scaled to correspond with the sighting latitudes. The centre line in the boxplot and the bottom and top of the box represent the 50th, 25th and 75th percentiles, respectively, and the whiskers the 5th and 95th percentile in figure a). The dots represent rarely visited latitudes. The data has been arranged by increasing median latitude. The length of the line in the network diagram in figure b) inversely represents the strength of the association between a dyad calculated as half-weight association index.

208x201mm (300 x 300 DPI)



Figure 3. The effect of coefficient of variation (CV) on the minimum detectable decline in the abundance of a theoretical population with different survey frequency; surveys conducted every six years (five years between surveys), every three years (two years between surveys), every other year or annually, over a theoretical 25-year period.

172x119mm (300 x 300 DPI)

http://mc.manuscriptcentral.com/aqc