

2020-12-01

Accelerometers for the Assessment of Recovery, AFAR. Patient-centred measurement of recovery from day surgery using wrist worn accelerometers; a pilot and feasibility study

Ratcliffe, A

<http://hdl.handle.net/10026.1/16236>

10.1111/anae.15267

Anaesthesia

Wiley

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Type of article: Original article

Submitting author: Professor J Robert Sneyd, Peninsula Medical School, University of Plymouth,
John Bull Building, Plymouth Science Park, Research Way, Plymouth PL6 8BT, UK.

Email: robert.sneyd@pms.ac.uk

**Patient-centred measurement of recovery from day-case surgery using wrist worn accelerometers:
a pilot and feasibility study***

A.M. Ratcliffe,¹ B. Zhai,² Y. Guan,³ D.G. Jackson,⁴ South West Anaesthetic Research Matrix (SWARM)⁵
and J.R. Sneyd⁶

*1 Academic Clinical Fellow and Specialist Registrar, Department of Anaesthesia, University Hospitals
Plymouth NHS Trust, Plymouth, UK.*

*2 Doctoral Trainee, 3 Lecturer, 4 Senior Research Associate, Open Lab, School of Computing,
Newcastle University, Newcastle upon Tyne, UK*

5 South West Anaesthesia Research Matrix (SWARM), <https://www.ukswarm.com/>

6 Emeritus Professor, Peninsula Medical School, University of Plymouth, UK.

For collaborators please see Appendix 1.

*Interim results presented at the Anaesthesia Research Society, London, 2019.

Correspondence to: Professor J Robert Sneyd, Peninsula Medical School, University of Plymouth,
John Bull Building, Plymouth Science Park, Research Way, Plymouth PL6 8BT, UK.

Email: robert.sneyd@pms.ac.uk Telephone: 07870 271531

Keywords: Postoperative recovery, day case surgery, accelerometer, activity monitor

Twitter handles: AMR @AnnaMRatcliffe; YG @YuGuan14; DGL @danielgjackson; SWARM
@ukswarm; JRS @robsneyd

Short title: Accelerometer assessment of postoperative recovery

Summary

This pilot and feasibility study evaluated wrist-worn accelerometers to measure recovery from day-case surgery in comparison to daily quality of recovery-15 scores. The protocol was designed with extensive patient and public involvement and engagement, and delivered by a research network of anaesthesia trainees. Forty-eight patients recruited through pre-operative assessment clinics wore wrist accelerometers for seven days before (pre-operative) and immediately after elective surgery (early postoperative), and again at three months (late postoperative). Validated activity and quality of recovery questionnaires were administered. Raw accelerometer data were archived and analysed using open-source software. The mean (SD) number of valid days of accelerometer wear per participant in the pre-operative, early and late postoperative periods were 5.4 (1.7), 6.6 (1.1) and 6.6 (1.0) days respectively. On the day after surgery, Euclidian norm minus one (a summary measure of raw accelerations), step count, light physical activity and moderate/vigorous physical activity were depressed to 57%, 47%, 59% and 35% of baseline values respectively. Activity progressively increased on a daily basis but had not returned to baseline values by 7 days. Patient questionnaires suggested subjective recovery by postoperative day 3 to 4; however, accelerometry data showed that activity levels had not returned to baseline at this point. All activity measures had returned to baseline by 3 months. Wrist-worn accelerometry is acceptable to patients and feasible as a surrogate measure for monitoring postoperative recovery from day-case surgery. Our results suggest that patients may overestimate their rate of recovery from day-case surgery, which has important implications for future research.

Introduction

The identification and measurement of outcomes from surgery and anaesthesia is an established priority for research in peri-operative medicine [1]. Progress and setbacks after discharge are monitored commonly by telephone calls, although these are labour intensive and contingent on the patient being contactable [2]. Further, even daily inquiries lack granularity at a time when the patient's condition may be changing rapidly. Wearable activity monitors are in common use for self-monitoring of exercise and might be useful to describe recovery at home following discharge from hospital. Accelerometers have been used previously in cohort studies including 100,000 participants in the Biobank study [3] and 4000 participants in the Whitehall 2 study [4]. Typically, a wearable activity monitor uses a tri-axial accelerometer to measure acceleration along three axes, which can then be analysed using computer algorithms and represented as activities. These can be count-based and device specific (e.g. step count), describe sedentary or active time in terms of energy expenditure (e.g. vigorous/moderate/low intensity activity), measure sleep quality and quantity, or characterise complex activity using machine learning techniques [5]. Notably, time to mobilisation and sleep quality are endpoints recommended by the standardised endpoints in peri-operative medicine (StEP) initiative for the measurement of patient comfort in peri-operative research [6]. Whilst the activity of hospital in-patients is known to be low [7,8], mobilisation after discharge is not well characterised.

Accelerometers for the assessment of recovery (AFAR) is a pilot and feasibility study designed and run by the trainee South West Anaesthesia Research Matrix (SWARM) [9]. This study was undertaken in collaboration with OpenLab, a laboratory specialising in human computer interaction and ubiquitous computing at Newcastle University. We used wrist-worn accelerometers to record raw acceleration data and open source software for analysis. This approach allows: reproducibility of research; re-analysis; data aggregation [10]; researcher control of the raw data; and can reduce costs [11]. An example is the development of an open source algorithm for sleep analysis using raw accelerometer data from the Whitehall 2 study [4], applied later to archived data from the UK Biobank [12].

We asked the following research questions: first, are wrist-worn accelerometers an acceptable and feasible way to measure recovery from day-case surgery, and could this method be deployed at scale across a research network?; and second, do available open source algorithms show utility in representing the activity (including sleep) of patients recovering from day-case surgery, and is this comparable with results of the quality of recovery-15 score [13]? We also aimed to demonstrate the

capability of SWARM to conduct a clinical trial whilst giving research experience to trainees in anaesthesia.

Methods

Prospective ethical approval for the study protocol was gained. Written informed consent was provided by all patients and included agreement to archiving of their de-identified raw accelerometry data in a secure data repository. This was a pilot study run at two SWARM sites: University Hospital Plymouth NHS Trust, a large regional teaching hospital and tertiary referral centre; and Royal Cornwall Hospitals NHS trust, a busy district general hospital. We aimed to recruit approximately 50 patients between the two sites; this sample size was based on a previous similar study [2], group experience, and guidelines for pilot study size [14].

Our pilot study was developed with patient and public involvement and engagement including a questionnaire-based scoping survey (see supplementary material, Appendix 2) distributed at the Royal Cornwall Hospitals NHS Trust pre-assessment clinic with the aim to identify: the level of technology awareness and use amongst our patient cohort; the acceptability of wearing a device; and how important movement and activity is to our local patient cohort in the (anticipated) recovery period. All participating patients were invited to complete a feedback questionnaire (see supplementary material, Appendix 2) during their involvement with the study. Finally, a semi-structured focus group session was held with a convenience sample of five study participants and three members of the research team to explore in-depth patient experience of the study, and to discuss plans for a further study.

Adult patients (aged > 18 years) scheduled for day-case surgery under general or neuraxial anaesthesia were eligible for inclusion in the study. Patients were still included in the study if they had a one-night planned admission for social reasons or a one-night unplanned admission. We did not study patients who were unable to consent and/or having surgical procedures or other factors likely to limit mobility (e.g. knee arthroscopy).

Participants wore an AX3 tri-axial accelerometer (Axivity; Newcastle upon Tyne, UK) on their dominant wrist for three wear periods (Fig. 1). These were: seven consecutive days pre-operatively (pre-operative period); the first seven postoperative days (early postoperative period); and a further seven consecutive days approximately three months after surgery (late postoperative period). The logistical processes for distribution and return of the accelerometers are detailed in Table 1. We aimed to acquire activity profiles at baseline, during an initial recovery period and following full

recovery from surgery. The AX3 device was pre-programmed to record accelerations between -8 g and $+8\text{ g}$ for 7 days at a sampling frequency of 100 Hz to give optimal sampling frequency to battery time ratio. These are the default settings, have appropriate sensitivity to movement (accelerations) of the study population and have been used in other studies with these devices [3,15]. Raw accelerometry data were downloaded and de-identified by the research team before transfer to a secure file-sharing area hosted by Newcastle University.

Demographic data and operation details were collected to define the group characteristics. The general practice physical activity questionnaire [16] (GPPAQ) and a modified version of the Duke activity status index (DASI) [17] were completed at the end of each wear period (see supplementary material, Appendix 2). Completion of the questionnaires was supervised directly in the pre-assessment clinic, encouraged by telephone call in the early postoperative period and without reminder in the late postoperative period. Modifications to the Duke activity status index included minor language adjustments for a UK patient population and the omission of the question relating to sexual activity. During the early postoperative period the quality of recovery-15 score [13] was measured daily by a questionnaire completed by telephone. Figure 1 depicts the timeline of data collection and study outcome measures recorded during the peri-operative journey.

Summary statistics and graphs were used to describe the data; inferential statistics were not used as this was a pilot study. A flowchart was developed from the consolidated standards of reporting trials (CONSORT) guidelines for pilot studies [18] to display feasibility results and recommendations for best practice in accelerometer-based research [19]. The post-participation focus group session was transcribed verbatim and analysed thematically using NVivo version 12 software (QSR International (UK) Limited, Daresbury, UK). Summarisation of free-text comments on feedback questionnaires was conducted manually.

Accelerometry data were processed in R using the software package GGIR (Vincent T van Hees, version 1.11-0, CRAN Archive, <https://cran.r-project.org/web/packages/GGIR/>). Python 3.8 libraries (www.python.org) 'numpy', 'pandas', 'statsmodels', 'sklearn', 'matplotlib' and 'seaborn' were used for analyses and graphs. Data extracted between the start and the end of recording were retained for analysis, yielding a maximum of seven consecutive 24-h recording periods. If any records exceeded 168 h, we discarded the surplus data. Calibration error was estimated based on static periods in the data and corrected if necessary (calibration correction range = 0.8 mg to 10.0 mg , mean correction = 2.5 mg [20]).

Non-wearing time was detected by a tuned classifier using established methods [21,22]. The classifier aimed to distinguish genuine non-wear (i.e. device removal for showering or sleep) from immobility during rest or sleep. Subject to contextual information from a wider 60 min window, a 15 min block of data was classified as a non-wear period if two out of three axes absolute value was < 50 mg or their SD < 13.0 mg (1 mg = 0.00981 m.s⁻²) [20]. We excluded these non-wearing data blocks for all wear periods and did not impute the accelerometer data. Days were defined as up to seven consecutive 24 h periods from the beginning of the accelerometer recording. A valid day was defined as having a minimum of wearing time of 10 h, a criterion which is used commonly in physical activity and health assessment studies [23,24]. Patients with at least one valid day per wear period were included in the analysis of accelerometry.

The Euclidean norm minus one, where r_i is the vector magnitude at the time point i , is shown below:

$$r_i = \sqrt{x_i^2 + y_i^2 + z_i^2} - 1000$$

This was used to quantify the acceleration related to the movement registered and is expressed in mg [21]. The Euclidean norm minus one subtracted a fixed offset value of 1 g at each time point to remove gravity [21] and was averaged over 5-s epochs. Negative values were rounded to zero to reduce bias and error. Using the dominant wrist, periods of physical activity were classified into: sedentary (< 50 mg); light (50-110 mg); and moderate/vigorous physical activity (>110 mg) (see supplementary material, Appendix 2). Sleep periods were detected using a validated algorithm [11]. For sleep period analysis, a valid day was defined as having at least 16 h of wearing time. Sleep duration (the interval between sleep onset time and end of sleep awakening time) and sleep efficiency (the proportion of that time actually asleep) were calculated. Eighty-five percent is considered normal sleep efficiency. To calculate the number of steps we resampled the accelerometer data to 15 Hz in R (version 3.6) using the software package GGIR. Step counts were then calculated using an algorithm suitable for wrist-worn accelerometers [25] [26].

Results

We distributed 100 scoping questionnaires of which 56 were completed and returned. Full results are available in the supplementary material (Appendix 2). The majority of respondents (53/56) were willing to wear an activity monitor for research purposes and expressed a preference for wearing it on their dominant wrist; therefore, this site was used in the study. Activity-related concerns ranked second among aspects of recovery that were of greatest importance to respondents.

Screening was pragmatic and based on the availability in the pre-assessment clinic of a SWARM investigator. Eighty-nine eligible patients were approached between November 2018 and May 2019 of whom 63 agreed to take part and 48 were studied (Fig. 2). The baseline characteristics of participating patients are summarised in Table 2.

Wear compliance was good, although 10 patients were unable to complete the full seven-day pre-operative wear period as their surgery fell within a week of recruitment. The mean (SD) number of valid days of accelerometer wear per participant in the pre-operative, early and late postoperative periods were 5.4 (1.7), 6.6 (1.1) and 6.6 (1.0) days respectively. No serious adverse events occurred. One participant developed a localised rash under the watch strap. Usable accelerometer data were collected during the majority of wear periods (Fig. 2). Some data were lost due to battery failure affecting several devices. Defective devices were subsequently replaced by the manufacturer and the problem resolved.

Forty-seven patients returned their early postoperative feedback questionnaires. Patients were asked to score the AX3 device (minimum 0, maximum 10) for comfort and appearance. The mean (SD) scores for comfort and appearance were 9.0 (1.3) and 8.8 (1.7) respectively. We were interested to learn if patients were worried about having their activity monitored (e.g. feeling that it was an invasion of their privacy), but no concerns in this regard were reported. Practical concerns reported included the need to keep the device clean and dry at work or when washing. The most common complaints were that the strap could catch clothing (reported five times) and that skin could become itchy/sweaty under the strap (reported three times). Explanations for temporary removal of the device included: two patients who couldn't wear the device at work; two patients preferred not to wear it at night; one patient took the device off when re-admitted overnight to hospital during the first postoperative week; and one patient removed the device due to the development of a rash under the strap.

Postoperative median (IQR [range]) quality of recovery-15 scores showed an upward trend during the first postoperative week, increasing from 123 (105-138 [76-150]) at 24 h to 145 (132-150 [100-150]) at 7 days (Fig. 3). The proportion of patients achieving the highest possible score at 24 h, 48 h and seven days was 1/44, 2/41, and 11/43 respectively, which is a pattern consistent with other studies [27].

Activity profiles during the three wear periods are shown in Figures 4 and 5. Euclidean norm minus one and step counts showed similar profiles during the pre-operative and late postoperative wear periods with a recovery trajectory during the first seven post-operative days. Physical activity did not

return to baseline during the week after surgery but had done so by three months.

Moderate/vigorous physical activity was reduced markedly after surgery; light physical activity was also reduced but not to the same degree (Fig. 5).

Scores for the quality of recovery-15 sleep question were low after surgery and then normalised over several days (Fig. 6). However, this trend was not reflected in the accelerometer-derived metrics for sleep duration and efficiency (Fig. 7).

The GPPAQ and modified DASI were completed by all participants at the end of the pre-operative period, usually under direct supervision on the day of surgery. Despite a telephone reminder, 12/96 questionnaires were not returned or were unusable after the early postoperative period. This increased to 22/96 in the late postoperative period when the reminder was included in the accelerometer packaging. Self-reported activity scores decreased after surgery and had returned to baseline by three months (Table 3).

Our post-participation focus group included five study participants (mean (SD) age 63 (12) years), three male) of which two were in full-time employment, one was semi-retired and two fully retired. Wearing the accelerometer was reported to be acceptable by all, although some were concerned they might damage the devices whilst working during the pre-operative phase. Reported anxieties predominantly related to surgery and the underlying diagnosis. Suggestions for extending the scope of accelerometry included: improved postoperative monitoring at home; encouraging patients to increase their activity; and as an aid to weight loss. Our thematic analysis is summarised in Table 4.

Discussion

We have demonstrated that wearable accelerometry is acceptable to patients and feasible as a surrogate measure for monitoring postoperative recovery from day-case surgery. As an objective measure of activity it may have advantages over questionnaires, especially if these have not been validated [28]. Our attempts to involve patients were successful and useful. Their input informed the design of the protocol (e.g. device worn on the dominant wrist), explained missing data (e.g. fear of device damage, workplace issues) and gave context (e.g. prevalence of cancer fears and anxiety about surgery). Patients also made helpful suggestions for routinely integrating accelerometry into peri-operative recovery (e.g. self-monitoring, motivation and remote support).

This study was delivered by the SWARM network, which offers anaesthesia trainees participation in meaningful research projects. Our trainees achieved the National Institute of Academia

Anaesthesia's (NIAA) research engagement objective of becoming 'research experienced' [29], undertaking all stages of the project from funding application, patient recruitment, data collection and roles of principal and chief investigator, thereby preparing them to become engaged participants in a research-active speciality.

We have demonstrated that derivatives of unsupervised home accelerometry such as the Euclidean norm minus one and step count, describe a recovery profile consistent with the resource-intensive daily application of a validated recovery measure (quality of recovery-15). The greatest impact of surgery on activity was the reduction in moderate/vigorous physical activity to 35% of baseline, which is biologically plausible and intuitive. The lesser effect on light physical activity (reduced to 59% of baseline) may reflect that those activities are less painful or physiologically disturbing in the context of recent surgery. However, all activity measures were below baseline values throughout the first postoperative week, even though patients reported themselves recovered after three days. This implies that accelerometry can detect degrees of functional impairment that patients do not consider important, or that the quality of recovery-15 score is unable to detect. Alternatively, patients may have been feeling fully recovered but obeying physician or self-imposed limits on activity. Similar overestimation of self-reported activity has been observed in patients having bariatric surgery [30].

Each wear period was one week in duration and we observed changes in the accelerometry derivatives throughout the early postoperative period suggesting that this is the optimal measurement time frame. In the pre-operative and late postoperative periods, the initial 2 or 3 days was consistent with the remainder of the 7 day period; therefore, is potentially not necessary to collect data for an entire week to characterise a patient's baseline function or to confirm their return to normal.

The inconsistency between responses to the quality of recovery-15 sleep question and our quantification of night-time sleep duration and efficiency was unexpected and requires further investigation. Movement-derived indices of sleep may not reflect what patients perceive as sleep quality. Advanced signal processing of accelerometry data is a development field with inconsistent performance of candidate analyses [31]. Extraction of multiple mobility indices and their subsequent combination may be useful [32]. Emerging analytics may yield specialist data, for example in orthopaedic surgery, where bouts of shortened stepping may indicate joint pain. However, the quantification of posture (sitting, lying, standing) and purposeful activity remains experimental with variable results [31].

Others have employed postoperative accelerometry to: describe recovery of mobility after lung surgery [33]; demonstrate the ineffectiveness of interventions designed to improve patient mobilisation after prolapse surgery [34]; and quantify the utility of a behavioural support tool after abdominal cancer surgery [35]. Bisgaard et al. evaluated hip-worn accelerometry postoperatively for quantifying physical activity and sleep in a small group of patients and volunteer controls, and reported a clinically acceptable correlation with physical activity and an unvalidated self-assessment of sleep [36]. However, methodology remains heterogeneous. A recent review noted that “...*future research would benefit from consistency in measurement methods and agreement on the most crucial types of outcomes to measure*”. [37]

Commercial activity monitors for domestic use typically report a limited range of accelerometer-derived activities using proprietary and non-disclosed algorithms which are subject to change. Raw data are discarded and there are ethical and data protection issues [38]. Further, there are a plethora of devices available which is a limiting factor; a systematic review of accelerometer use to measure physical activity in hospital (including 10 studies in the postoperative setting) identified that 17 different devices were used, and concluded it was not possible to aggregate data for meta-analysis due to the heterogeneity of the measured outcomes [39]. Sharing raw data permits pooled analysis, an approach pioneered with pharmacokinetic data shared through the open TCI initiative (www.opentci.org). Our commitment to open-source software complies with earlier recommendations [40] and our patients agreement to open-data archiving allows this to be explored with future novel algorithms.

Accelerometers have seen increasing use to document mobilisation after surgery and evaluate measures intended to improve it. Standardisation of outcome measures is recommended [41] (although the methods of measurement are not specified) and we used a validated recovery measure compliant with that initiative [13]. Possible roles for accelerometry in the peri-operative period are summarised in Table 5.

Moving beyond the descriptive, activity monitoring could be integrated into care pathways to identify poor progress or as part of an intervention. Full mobilisation after laparoscopic cholecystectomy takes over a week and may be modestly accelerated by combining motivational accelerometry with personalised advice on exercise [42]. Recovery from major surgery is enhanced by ambulation, and step counts on the first postoperative day are inversely correlated with duration of hospital stay [43]. This suggests a testable hypothesis that feedback to patients of their objectively quantified activity might reduce their time in hospital. However, patient engagement with technology derived advice cannot be taken for granted. When a cohort of 79,953 patients were

invited to share their fitness-tracker data with their healthcare provider, less than 1% did so. Further, the investigators concluded “...*patients most at risk for poor health outcomes are least likely to share personal fitness-tracker data*” [44]. Our focus group analysis offers some insights into the complex personal world in which individual surgical episodes are embedded and suggests the importance of continued attention to the social science and qualitative dimensions of future interventions. Patients at a Dutch academic medical centre were reluctant to participate in accelerometer monitoring after surgery and in some cases sleep monitoring was limited by discomfort from the hip accelerometer [24]. We avoided this type of problem with public engagement in the design of our protocol.

We have demonstrated effective peri-operative deployment of accelerometry by a trainee research network. Our experience suggests accelerometry has the potential to support and monitor patient mobilisation. The information generated could evaluate activity-related interventions either by motivating patients to exercise more or by identifying a subset of patients who progress less well than expected. In either case the experimental objective would be to improve patient outcomes. Our library of raw accelerometry files, questionnaire results and demographic data will be analysed further to explore potential activity metrics.

Acknowledgments and competing interests

This study was funded by the National Institute of Academic Anaesthesia (Association of Anaesthetists/Anaesthesia grant). AM is an academic clinical fellow funded by the National Institute for Health Research. No competing interests declared.

Appendix 1. List of SWARM collaborators.

G Minto, University Hospitals Plymouth NHS Trust, Plymouth, UK; S Howell, University Hospitals Plymouth NHS Trust, Plymouth, UK; F Miller, University Hospitals Plymouth NHS Trust, Plymouth, UK; JL Retief, University Hospitals Plymouth NHS Trust, Plymouth, UK; DA Webster, University Hospitals Plymouth NHS Trust, Plymouth, UK; P Veeralakshmanan, University Hospitals Plymouth NHS Trust, Plymouth, UK; JF Graterol, Royal Cornwall Hospitals NHS Trust, Truro, UK; DP Kotwinski, Royal Cornwall Hospitals NHS Trust, Truro, UK; S Maxwell, Royal Cornwall Hospitals NHS Trust, Truro, UK; B Parish, Royal Cornwall Hospitals NHS Trust, Truro, UK; S Spinney, Royal Cornwall Hospitals NHS Trust, Truro, UK.

Appendix 2.

Scoping Survey questionnaire

1. Do you own a Smartphone? (please delete appropriate) Yes/No If “yes”, please write below the type of phone and operating system if known: (Eg iPhone, 6s, iOS 10) If you answered “no” to question 1 please proceed to question 3.
2. Do you use applications (apps) to track your activities, health, or weight, for example Strava, iSki, My Fitness Pal (please list)
3. Do you own an activity monitor or pedometer for example FitBit, Garmin.. (please delete appropriate) Yes/ No. If yes please state which:
4. Would you be willing to wear an activity sensing device (like a wrist watch) continuously for a week following your surgery? (please delete appropriate) Yes / No
If no, please explain why below:
5. Would you prefer to wear the device on: a. Ankle, b. Wrist, c. Waist, d. Back or e. Hip.
6. Would you see any problems with returning a device by post, in a stamped addressed envelope, after a wearing it for one week? (please delete appropriate) Yes/ No. If yes, please describe:
7. Would you mind being contacted by email or text message every day to complete a questionnaire or to remind you to wear you device? (please delete appropriate) Yes/ No. If yes, please explain why:
8. What concerns do you have about your recovery from anaesthesia/ surgery- what worries you the most?
9. How important are the following aspects of recovery from an operation to you? : Please circle as appropriate (1 = very important, 5= not important)

Getting back to usual sleep patterns	1	2	3	4	5
Getting back to usual activities at work or home	1	2	3	4	5
Getting back to usual exercise routine	1	2	3	4	5
Energy levels	1	2	3	4	5
Post-operative pain	1	2	3	4	5
Post-operative nausea/vomiting	1	2	3	4	5
Being able to breathe easily	1	2	3	4	5

Sense of wellbeing/ feeling anxious or depressed

1

2

3

4

5

Scoping Survey results. Data from 56 patients surveyed in the pre-assessment clinic. Values are number (proportion) or mean (SD).

Question	Number (%)		
1. Own a smartphone	31 (56%)		
2. Use an activity app	10 (18%)		
3. Have an activity monitor	16 (29%)		
4. Willing to wear activity monitor	54 (95%)		
5. Wearing preference			
• Wrist	38 (68%)		
• Waist	8 (14%)		
• Ankle	3 (5%)		
• Hip	2 (4%)		
• Back	2 (4%)		
• Not answered	3 (5%)		
6. Willing to return device by post	52 (93%)		
7. Willing to be contacted at home	45 (80%)		
8.			
	Score	Rank	
Important aspects of recovery			
• Sleep	2.2 (1.3)	5	
• Resuming activity	1.7 (1.2)	=2	
• Exercise routine	2.1 (1.3)	4	
• Energy levels	1.8 (1.3)	=3	
• Post-op pain	1.8 (1.2)	=3	
• Post-op nausea	1.8 (1.1)	=3	
• Breathing easily	1.6 (1.2)	1	
• Wellbeing	1.7 (1.2)	=2	

Physical activity questionnaires.

(To be completed at the end of each wear period)

Study Number.....

(please tick which wear period this questionnaire responds to i.e. has just been completed)

- Pre-operative period** ☐ **(COMPLETE ON DAY OF OPERATION)**
- Early postoperative period** ☐ **(VIA PHONE DAY 7 POST-OP)**
- Late postoperative period** ☐ **(SEND WITH AX3 DEVICE)**

Part 1 (Modified Duke activity status index, DASI).

Please think back over your last week. Have you been able to do the following?

1. Take care of yourself, that is eating, dressing, bathing and using the toilet?	<input type="checkbox"/>
2. Walk indoors, such as around your house?	<input type="checkbox"/>
3. Walk 100m on flat ground?	<input type="checkbox"/>
4. Walk up one flight of stairs?	<input type="checkbox"/>
5. Walk up two flights of stairs, or up a hill	<input type="checkbox"/>
6. Can you run a short distance?	<input type="checkbox"/>
7. Can you do light work around the house, such as dusting or washing up?	<input type="checkbox"/>
8. Can you do moderate work around the house such as hovering, sweeping the floor, carrying the shopping?	<input type="checkbox"/>
9. Can you do heavy work around the house, such as scrubbing floors, lifting or moving heavy furniture	<input type="checkbox"/>

10. Can you do gardening such as raking, weeding, or lawn mowing?	<input type="checkbox"/>
11. Can you do moderate exercise, eg walking on the beach or coast path, yoga, golf, bowling	<input type="checkbox"/>
12. Can you do strenuous exercise, eg sports such as rugby, football, netball, running or jogging, road or mountain biking, rowing, surfing, aerobics/cross fit/circuit training.	<input type="checkbox"/>

Part 2. General Practice Physical Activity Questionnaire (GPPAQ). Available at:

<https://www.gov.uk/government/publications/general-practice-physical-activity-questionnaire-gppaq>

1. Please tell us the type and amount of physical activity involved in your work. Please tick one box that is closest to your present work from the following five possibilities:

		Please mark one box only
a	I am not in employment (e.g. retired, retired for health reasons, unemployed, full-time carer etc.)	
b	I spend most of my time at work sitting (such as in an office)	
c	I spend most of my time at work standing or walking. However, my work does not require much intense physical effort (e.g. shop assistant, hairdresser, security guard, childminder, etc.)	
d	My work involves definite physical effort including handling of heavy objects and use of tools (e.g. plumber, electrician, carpenter, cleaner, hospital nurse, gardener, postal delivery workers etc.)	
e	My work involves vigorous physical activity including handling of very heavy objects (e.g. scaffolder, construction worker, refuse collector, etc.)	

2. During the last week, how many hours did you spend on each of the following activities?
Please answer whether you are in employment or not

Please mark one box only on each row

		None	Some but less than 1 hour	1 hour but less than 3 hours	3 hours or more
a	Physical exercise such as swimming, jogging, aerobics, football, tennis, gym workout etc.				
b	Cycling, including cycling to work and during leisure time				
c	Walking, including walking to work, shopping, for pleasure etc.				
d	Housework/Childcare				

e	Gardening/DIY				
---	---------------	--	--	--	--

3. How would you describe your usual walking pace? Please mark one box only.

Slow pace (i.e. less than 3 mph)	<input type="checkbox"/>	Steady average pace	<input type="checkbox"/>
Brisk pace	<input type="checkbox"/>	Fast pace (i.e. over 4mph)	<input type="checkbox"/>

Exercise intensity thresholds, exemplar activities, METs and Euclidean norm minus one ranges.

Intensity Threshold	Example physical activities [45]	MET*	Euclidean norm minus one (milligravity, <i>mg</i>)
Sedentary	Resting	1	<50
Light Physical Activity	Watering plants, standing, slow walking (e.g. Less than 2.0 mph, level ground, strolling, very slow), dusting (light effort), washing dishes, making the bed	<3	50-110
Moderate Physical Activity	Mopping, tailoring, weaving, dancing (e.g. disco, folk, square, line dancing, irish step dancing, polka, contra, and country dancing.) Mowing lawn (e.g. operating power mower)	3-6	>110-440
Vigorous Physical Activity	Track and field (high jump, long jump, triple jump, javelin, pole vault) walking at 5.0 mph, water jogging. Bicycling (> 10mph with moderate effort)	>6	>440

* Metabolic Equivalent of Task, MET. For reference, when measured by a wrist worn Actigraph accelerometer, 100.6 *mg* = 3 METs and 428.8 *mg* = 6 METs [46].

Feedback questionnaire administered at the end of the early postoperative period (day 7).

1. How did you find wearing the accelerometer; please score out of 10:

Comfort (/10)

Appearance (/10)

2. Did you have concerns about wearing the AX3 device?

If yes please describe in as much detail as you can:

3. Did you complete the 7 days pre-operation wear period?

If not please explain in as much detail as you wish why this was:

4. Did you complete the 7 days post-operation wear period?

If not please explain in as much detail as you wish why this was:

5. Were there any times when you found it was difficult to wear the accelerometer, (for example when performing particular tasks, work policy, uniform restrictions)?

If you would like us to contact you to discuss your experience in more detail, please give your preferred contact details below:

References

1. Boney O, Bell M, Bell N, et al. Identifying research priorities in anaesthesia and perioperative care: final report of the joint National Institute of Academic Anaesthesia/James Lind Alliance Research Priority Setting Partnership. *BMJ Open* 2015; **5**: 1-5.
2. Benton JI, Vickery PJ, Davies P, Langton JA, Sneyd JR Recovery profiles after day-case diagnostic gynaecological laparoscopy: comparison of two anaesthetic techniques over 1 week follow-up. *British Journal of Anaesthesia* 1999; **83**: 523P-4P.
3. Doherty A, Jackson D, Hammerla N, et al. Large scale population assessment of physical activity using wrist worn accelerometers: The UK Biobank Study. *PLoS One* 2017; **12**: e0169649.
4. Menai M, Van Hees VT, Elbaz A, Kivimaki M, Singh-Manoux A, Sabia S Accelerometer assessed moderate-to-vigorous physical activity and successful ageing: results from the Whitehall II study. *Scientific Reports* 2017; **7**: 45772.
5. Troiano RP, McClain JJ, Brychta RJ, Chen KY Evolution of accelerometer methods for physical activity research. *British Journal of Sports Medicine* 2014; **48**: 1019-23.
6. Myles PS, Boney O, Botti M, et al. Systematic review and consensus definitions for the Standardised Endpoints in Perioperative Medicine (StEP) initiative: patient comfort. *British Journal of Anaesthesia* 2018; **120**: 705-11.
7. Mudge AM, McRae P, McHugh K, et al. Poor mobility in hospitalized adults of all ages. *Journal of Hospital Medicine* 2016; **11**: 289-91.
8. Jonker LT, Hendriks S, Lahr MM, van Munster BC, de Bock GH, van Leeuwen BL Postoperative recovery of accelerometer-based physical activity in older cancer patients. *European Journal of Surgical Oncology* 2020 (in-press) <https://doi.org/10.1016/j.ejso.2020.06.012>.
9. SWARM (South West Anaesthesia Research Matrix) www.ukswarm.com (accessed 20th August 2020)
10. Migueles JH, Rowlands AV, Huber F, Sabia S, van Hees VT GGIR: A research community-driven open source R package for generating physical activity and sleep outcomes from multi-day raw accelerometer data. *Journal for the Measurement of Physical Behaviour* 2019; **2**: 188-96.
11. van Hees VT, Sabia S, Jones SE, et al. Estimating sleep parameters using an accelerometer without sleep diary. *Scientific Reports* 2018; **8**: 1-11.
12. Jones SE, Lane JM, Wood AR, et al. Genome-wide association analyses of chronotype in 697,828 individuals provides insights into circadian rhythms. *Nature Communications* 2019; **10**: 1-11.
13. Stark PA, Myles PS, Burke JA Development and psychometric evaluation of a postoperative quality of recovery score: the QoR-15. *Anesthesiology* 2013; **118**: 1332-40.
14. Whitehead AL, Julious SA, Cooper CL, Campbell MJ Estimating the sample size for a pilot randomised trial to minimise the overall trial sample size for the external pilot and main trial for a continuous outcome variable. *Statistical Methods in Medical Research* 2016; **25**: 1057-73.
15. Baldwin C, van Kessel G, Phillips A, Johnston K Accelerometry shows inpatients with acute medical or surgical conditions spend little time upright and are highly sedentary: systematic review. *Physical Therapy* 2017; **97**: 1044-65.
16. General Practice Physical Activity Questionnaire (GPPAQ). www.gov.uk/government/publications/general-practice-physical-activity-questionnaire-gppaq (accessed 20/8/2020)
17. Hlatky MA, Boineau RE, Higginbotham MB, et al. A brief self-administered questionnaire to determine functional capacity (the Duke Activity Status Index). *Am J Cardiol* 1989; **64**: 651-4.
18. Eldridge SM, Chan CL, Campbell MJ, et al. CONSORT 2010 statement: extension to randomised pilot and feasibility trials. *BMJ* 2016; **355**: i5239.

19. Matthews CE, Hagströmer M, Pober DM, Bowles HR Best practices for using physical activity monitors in population-based research. *Medicine and science in sports and exercise* 2012; **44**: S68.
20. Van Hees VT, Fang Z, Langford J, et al. Autocalibration of accelerometer data for free-living physical activity assessment using local gravity and temperature: an evaluation on four continents. *J Appl Physiol (1985)* 2014; **117**: 738-44.
21. Van Hees VT, Gorzelniak L, Leon ECD, et al. Separating movement and gravity components in an acceleration signal and implications for the assessment of human daily physical activity. *PLoS One* 2013; **8**: e61691.
22. Syed S, Morseth B, Hopstock LA, Horsch A evaluating the performance of raw and epoch non-wear algorithms using multiple accelerometers and electrocardiogram recordings. *Scientific Reports* 2020; **10**: 1-18.
23. Wennman H, Pietilä A, Rissanen H, et al. Gender, age and socioeconomic variation in 24-hour physical activity by wrist-worn accelerometers: the FinHealth 2017 Survey. *Scientific Reports* 2019; **9**: 1-9.
24. Van der Meij E, Van der Ploeg HP, Van Den Heuvel B, et al. Assessing pre-and postoperative activity levels with an accelerometer: a proof of concept study. *BMC Surgery* 2017; **17**: 1-10.
25. Patterson MR. https://github.com/ShimmerEngineering/Verisense-Toolbox/blob/master/Verisense_step_algorithm/validation_paper.pdf (accessed 1st August 2020)
26. Gu F, Khoshelham K, Shang J, Yu F, Wei Z Robust and accurate smartphone-based step counting for indoor localization. *IEEE Sensors Journal* 2017; **17**: 3453-60.
27. Chazapis M, Walker E, Rooms M, Kamming D, Moonesinghe S Measuring quality of recovery-15 after day case surgery. *BJA: British Journal of Anaesthesia* 2016; **116**: 241-8.
28. Sliepen M, Lipperts M, Tjur M, Mechlenburg I Use of accelerometer-based activity monitoring in orthopaedics: benefits, impact and practical considerations. *EFORT Open Reviews* 2019; **4**: 678-85.
29. The NIAAs three point scale of research engagement. <https://www.niaa.org.uk/The-NIAAs-three-point-scale-of-research-engagement> (accessed 28th October 2020)
30. Possmark S, Sellberg F, Willmer M, Tynelius P, Persson M, Berglind D Accelerometer-measured versus self-reported physical activity levels in women before and up to 48 months after Roux-en-Y Gastric Bypass. *BMC Surgery* 2020; **20**: 39.
31. Anderson JL, Green AJ, Yoward LS, Hall HK Validity and reliability of accelerometry in identification of lying, sitting, standing or purposeful activity in adult hospital inpatients recovering from acute or critical illness: a systematic review. *Clinical Rehabilitation* 2018; **32**: 233-42.
32. Buchman AS, Dawe RJ, Leurgans SE, et al. Different Combinations of Mobility Metrics Derived From a Wearable Sensor Are Associated With Distinct Health Outcomes in Older Adults. *The Journals of Gerontology: Series A* 2019; **75**: 1176-83.
33. Kaplan SJ, Trotman PA, Porteous GH, et al. Functional recovery after lung resection: a before and after prospective cohort study of activity. *Ann Thorac Surg* 2019; **107**: 209-16.
34. Arunachalam D, Heit MH Impact of postoperative instructions on physical activity following pelvic reconstructive surgery: a randomized controlled trial. *International Urogynecology Journal* 2020: 1-9.
35. Porserud A, Aly M, Nygren-Bonnier M, Hagströmer M Objectively measured mobilisation is enhanced by a new behaviour support tool in patients undergoing abdominal cancer surgery. *European Journal of Surgical Oncology* 2019; **45**: 1847-53.
36. Bisgaard T, Kjærsgaard M, Bernhard A, Kehlet H, Rosenberg J Computerized Monitoring of Physical Activity and Sleep in Postoperative Abdominal Surgery Patients. *Journal of Clinical Monitoring and Computing* 1999; **15**: 1-8.

37. Baldwin CE, Parry SM, Norton L, Williams J, Lewis LK A scoping review of interventions using accelerometers to measure physical activity or sedentary behaviour during hospitalization. *Clin Rehabil* 2020; 269215520932965.
38. Watson DS, Krutzinna J, Bruce IN, et al. Clinical applications of machine learning algorithms: beyond the black box. *BMJ* 2019; **364**.
39. Migueles JH, Cadenas-Sanchez C, Rowlands AV, et al. Comparability of accelerometer signal aggregation metrics across placements and dominant wrist cut points for the assessment of physical activity in adults. *Scientific Reports* 2019; **9**: 1-12.
40. Gurchiek RD, Choquette RH, Beynnon BD, et al. Open-Source Remote Gait Analysis: A Post-Surgery Patient Monitoring Application. *Scientific Reports* 2019; **9**: 17966.
41. Moonesinghe SR, Jackson AIR, Boney O, et al. Systematic review and consensus definitions for the Standardised Endpoints in Perioperative Medicine initiative: patient-centred outcomes. *British Journal of Anaesthesia* 2019; **123**: 664-70.
42. Wasowicz-Kemps DK, Slootmaker SM, Kemps HM, Borel-Rinkes IH, Biesma DH, van Ramshorst B Resumption of daily physical activity after day-case laparoscopic cholecystectomy. *Surgical Endoscopy* 2009; **23**: 2034-40.
43. Daskivich TJ, Houman J, Lopez M, et al. Association of wearable activity monitors with assessment of daily ambulation and length of stay among patients undergoing major surgery. *JAMA Network Open* 2019; **2**: e187673-e.
44. Pevnick JM, Fuller G, Duncan R, Spiegel BM A large-scale initiative inviting patients to share personal fitness tracker data with their providers: initial results. *PLoS One* 2016; **11**: e0165908.
45. Ainsworth BE, Haskell WL, Herrmann SD, et al. 2011 Compendium of Physical Activities: a second update of codes and MET values. *Medicine & Science in Sports & Exercise* 2011; **43**: 1575-81.
46. Hildebrand M, VAN VH, Hansen BH, Ekelund U Age group comparability of raw accelerometer output from wrist-and hip-worn monitors. *Medicine & Science in Sports & Exercise* 2014; **46**: 1816-24.

Table 1. Procedure for distribution and return of accelerometers and the retrieval of data for each of the three seven-day wear periods. Data from each wear period were downloaded locally, de-identified, labelled with a study number, and uploaded to the Newcastle University secure file-sharing site for analysis.

Wear Period	Activity	Procedure
Pre-operative	Distribution and deployment	Accelerometer either: - given directly to the participant at recruitment if their surgery fell within 2 weeks of recruitment; or - sent by post.
	Return and data retrieval	Participant brings the accelerometer with them on day of surgery. A member of the research team meets them, downloads accelerometer data and recharges device using designated laptop.
Early postoperative	Distribution and deployment	Accelerometer returned to the patient with instructions to re-commence wearing on leaving the hospital.
	Return and data retrieval	Stamped addressed envelope supplied for return after seven days wear. Data downloaded by researcher in Plymouth onto designated laptop.
Late postoperative	Distribution and deployment	Accelerometer sent by post (may not be the same device).
	Return and data retrieval	Stamped addressed envelope supplied for return after seven days wear. Data downloaded by researcher in Plymouth using designated laptop.

Table 2. Characteristics of 48 patients undergoing elective day-case surgery and enrolled to wear wrist-based accelerometers pre- and postoperatively. Data are mean (SD), median (IQR [range]) or number. Self-assessed baseline physical activity was categorised by the general practice physical activity questionnaire (GPPAQ).

Age; years	55 (17)
Sex; female	28
BMI; kg.m ⁻²	29 (27-34 [21-48])
ASA physical status	
- 1	11
- 2	27
- 3	10
Baseline physical activity	
- Active	10
- Moderately active	10
- Moderately inactive	6
- Inactive	22
Surgical speciality	
- General	17
- Urology	12
- Gynaecology	7
- Breast	6
- Head and neck	4
- Orthopaedic	1
- Plastics	1
Duration of surgery; min	86 (37)
Duration of stay in recovery area; min	57 (36)
Overnight stay	
- Yes (planned)	1
- Yes (unplanned)	4*
- No	43

*Causes of unplanned admission were: surgical complication (n=1); slow recovery (n=1); and pain (n=2).

Table 3. Results of physical activity questionnaires in patients (n=48) undergoing day-case surgery assessed using modified Duke activity status index (DASI) and general practice physical activity questionnaire (GPPAQ). Values are number and median (IQR [range]).

	Pre-operative	Early Postoperative	Late Postoperative
Modified DASI			
- Valid	48/48	43/48	39/48
- Missing	-	5*	9*
- Score	10 (8-12 [4-12])	8 (6-9 [2-12])	11 (7-9 [4-12])
GPPAQ			
- Valid	48/48	41/48	35/48
- Missing	-	7	13
- Active	10	5	8
- Moderately active	10	10	4
- Moderately inactive	6	6	6
- Inactive	22	20	17

*Includes two participants who inserted numeric values rather than crosses when completing the modified Duke activity status index

Table 4. Results of qualitative analysis of post-participation feedback focus group, organised into “overarching themes”, “sub-themes” and “codes”, including quotes as examples of text within each code.

Overarching themes	Sub-themes	Codes	Quotes
AX3 wrist-worn accelerometer	Practicality and wearability	<ul style="list-style-type: none"> - Comfort - Appearance - Occupational considerations - Concerns: getting device wet/dirty/damaged 	<p><i>“Initially I was conscious that I had it on, not to get it too dirty, because you have to keep it quite tight on your wrist otherwise it slips around and what have you, and I can suggest some of the places that I put my hand on a livestock farm”</i></p> <p><i>“When I was dealing with patients I was very much aware that it was there. Not that it was not a big deal, it wouldn’t cause me to stop wearing it in the future”</i></p>
	Current model functionality and potential future modifications	<ul style="list-style-type: none"> - Current state functionality and seeking clarification of what it can measure (sleep, pain, blood pressure) - To view activity count - Heart rate analysis - Wireless connectivity 	<p><i>“I wasn’t totally sure what the watch was measuring whilst it was on my wrist...could it register pain? Is it like a pedometer when you go walking and it counts your paces?”</i></p>
	Burden and awareness	<ul style="list-style-type: none"> - Burden - Awareness - “Big Brother” effect - Effect on activity 	<p><i>“I can honestly say for myself there was nothing to it to be honest, you put it on and forget it was there; I wore it 99% of the time”</i></p> <p><i>“I have to say I did find it a little bit irritating, well just because the strap-the tongue of it kept undoing itself”</i></p>

	Accuracy and misrepresentation	<ul style="list-style-type: none"> - Seasonal differences in activity 	<i>"I'd like to point that out just in case it hadn't been noticed, in the winter you do less, come the summer, if we had the operation in the summer, I'm sure we'd all get out and about a lot more."</i>
Peri-operative experience	Relating to having an operation	<ul style="list-style-type: none"> - Apprehension about the operation - Activity levels and attitude to activity around an operation - State of mind and recovery - Coping with peri-operative anxiety 	<i>"I think the only thing you think of before your op, is your op."</i> <i>"So let me say that you're not in the same state of mind before an operation"</i> <i>"But don't forget, well I guess we've all had cancer... I wake up first thing in the morning and well you don't think about it but it never leaves you, so you have got that shall I say, that in your head, you carry around with you most of the time."</i>
	Related to the underlying diagnosis	<ul style="list-style-type: none"> - Anxiety/awareness about underlying health condition - Coping strategies - Effect on activity 	<i>"I must say that I told nobody. I closed down"</i> <i>"XXX is always saying I could do anything, even though I was an old man. I could beat anyone. And then you get that letter come in, and you think.. and it does drag you down. And so joining different things, it takes a bit of moving on"</i>
Changing times	Science and technology	<ul style="list-style-type: none"> - Internet - Telephone signal coverage - Wireless connectivity 	<i>"They basically they look at the index of what the regard as people require to live (referring to measuring poverty) , and that now includes people mobile phones, it now includes the internet, these are standard, whereas we wouldn't have dreamt of that 20 years ago"</i>
	Society and culture		<i>"What you're saying is there are far more pansies in the world these days than there used to be"</i>
	Health system	<ul style="list-style-type: none"> - NHS Systems - Health advice 	<i>"You've got no convalescent homes as was"</i>

Differences and inequalities		<ul style="list-style-type: none"> - Different personalities - Generational differences - Geographical differences and inequality - Rural urban differences 	<p><i>"I'm a country boy so you think that was a problem in the rural population, but probably I see and know more people because I know everyone around me. Whereas if you live in a street in town you don't make that same social contact, do you?"</i></p> <p><i>"I know technology is always improving but connectivity is important, and you've got to remember where we are in this country, I mean I can't even get mobile phone signal where I was"</i></p> <p><i>"For the younger generation, it's always a competition...to do your 10,000 steps or whatever it is."</i></p>
The NHS, people and systems		<ul style="list-style-type: none"> - Whose responsibility is it to follow up patients postoperatively in the community? - Saving time for doctors - Burden or workload of the NHS - Relieving pressure on the NHS 	<p><i>"As I understand, with the NHS, that basically you're good at what you do, and you get the operation done, and it's the bit between actually having the op, having a bed in the hospital, and then going home.."</i></p> <p><i>"If you have, or you need to have medical care afterwards and you're told to do things and you don't then you're putting the burden back on the NHS aren't you?"</i></p>

Table 5. Possible roles for accelerometry during the peri-operative period. CPET, cardiopulmonary exercise testing; ERAS, enhanced recovery after surgery.

Phase	Time	Task	Rationale
Pre-operative	When scheduled for surgery	Assess baseline activity	Accelerometry derivatives correlate with CPET data. Perhaps avoid CPET. Potential for advising on surgical risk and stratifying care.
		Assess baseline activity	Context for interpretation of later assessments
	Between scheduling and surgery	Monitoring progress of pre-habilitation	Possibly intervene to improve adherence
		As a therapeutic component of pre-habilitation	Provide feedback to motivate patient
Postoperative	Immediate	Track mobilisation	Facilitate interventions to enhance patient progress if they are falling off track
		Remote monitoring	Patient safety
		Provide feedback to motivate patient	Enhance effectiveness of ERAS programme
	Late	Remote monitoring	Patient safety
		Confirm return to baseline	Facilitate interventions to enhance patient progress if they are falling off track
All phases	All times	Research	Evaluate interventions and devices

Figures and Legends




Figure 1. Schedule of data collection for 48 patients undergoing elective day-case surgery who wore the AX3 tri-axial accelerometer on their dominant wrist over three seven-day wear periods: pre-operative (baseline); early postoperative (first 7 postoperative days); and late postoperative (3 months). In addition to recording of demographic and surgical data, questionnaires were administered on the day of surgery, daily for seven days after surgery and at the end of the study.




Figure 2. Recruitment, retention, and accelerometer data collection for 48 patients undergoing day-case surgery including outcomes required in the CONSORT extension for reporting pilot studies [18] and following recommendations for best practice in reporting research using physical activity monitors [19].




Figure 3. Daily quality of recovery-15 scores of 48 patients after elective day-case surgery. Boxes represent median and interquartile range, whiskers are 5th and 95th centiles. Values outside 1.5 times the interquartile range above the upper quartile or below the lower quartile are shown as outliers.




Figure 4. Accelerometry derivatives (a) total accelerations expressed as Euclidean norm minus one ($\text{milligravity} \cdot \text{day}^{-1}$) and (b) step count. Data were collected over three seven-day wear periods: pre-operative (baseline); early postoperative (first 7 postoperative days); and late postoperative (3 months). Thin lines represent the mean value for the group, the shaded area shows the 95%CI of the population mean and the thick regression line highlights the trend in group results with time.




Figure 5. Accelerometry derivatives (a) duration of light physical activity and (b) moderate or vigorous physical activity. Data were collected over three seven-day wear periods: pre-operative (baseline); early postoperative (first 7 postoperative days); and late postoperative (3 months). Thin lines represent the mean value for the group, the shaded area shows the 95%CI of the population mean and the thick regression line highlights the trend in group results with time.




Figure 6. Sleep score of 48 patients after elective day surgery. Boxes represent median and interquartile range, whiskers are 5th and 95th centiles. Values outside 1.5 times the interquartile range above the upper quartile or below the lower quartile are shown as outliers.




Figure 7. Accelerometry derivatives (a) sleep efficiency and (b) night-time sleep duration. Data were collected over three seven-day wear periods: pre-operative (baseline); early postoperative (first 7 postoperative days); and late postoperative (3 months). Thin lines represent the mean value for the group, the shaded area shows the 95%CI of the population mean and the thick regression line highlights the trend in group results with time.