

2021-08

COVID-19: UK frontline intensivists' emerging learning

Montgomery, H

<http://hdl.handle.net/10026.1/16403>

10.1177/1751143720931731

Journal of the Intensive Care Society

SAGE Publications

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1
2
3
4
5
6 Corresponding Author
7 Niran Rehill N.Rehill@ucl.ac.uk
8
9

10
11
12 Covid-19: UK Front Line Intensivists' Emerging
13 Learning
14
15
16
17

18 Authors
19

- 20
21
22 **Hugh Montgomery***, Whittington Hospital, Intensive Care Society [Chair]
23
24 **Niran Rehill**, NIHR Applied Research Collaboration North Thames
25
26 **Luigi Camporota***, Guy's and St Thomas' Hospital NHS Foundation Trust
27
28 **Nicki Credland***, British Association of Critical Care Nurses (BACCN)
29
30 **Mike Grocott***, University Hospital Southampton NHS Foundation Trust
31
32 **Mark Hamilton***, St George's Hospital
33
34 **Daniel Martin***, Royal Free Hospital
35
36 **Brijesh Patel***, Royal Brompton and Harefield NHS Foundation Trust
37
38 **Ganesh Suntharalingam***, Northwick Park Hospital, Intensive Care Society
39
40 **Tamas Szakmany***, Royal Gwent Hospital
41
42 **Shahana Uddin***, Kings College London
43
44 **Andre Vercueil***, King's College Hospital
45
46 **Mike Roberts**, UCL Partners & Queen Mary University of London*
47
48 *members of the expert panel
49
50

51 Conflicts of Interest
52 The Authors declare that there are no conflicts of interest.
53

54 Financial Support
55 This research received no specific grant from any funding agency in the public, commercial, or not-
56 for-profit sectors.
57
58
59
60

Acknowledgements

With thanks to Dr Sandy Mather & the Intensive Care Society for hosting the webinar, Dr Amanda Begley & Morgan McKean, UCL Partners, for coordinating and supporting the rapid synthesis of findings, & Professor Rosalind Raine, NIHR ARC North Thames for reviewing an early draft.

Dr Niran Rehill is seconded to NIHR Applied Research Collaboration North Thames. This paper presents independent research supported by the National Institute for Health Research (NIHR). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Keywords

Coronavirus, COVID-19, SARS CoV-2, Critical care,

This is NOT a clinical guideline. The suggestions for practice outlined represent professional opinion only.

Introduction:

At the time of writing, 51,608 people have tested positive for COVID-19 in the UK in less than three months; 5,373 have died. (1) These figures will have risen by the time you read this. Around 5% (2)² of COVID-19 infections appear to require intensive care unit (ICU) admission. As the number of people contracting the illness increases rapidly, so does ICU clinical experience. Hundreds of intensivists will by now have had to manage multiple complex COVID-19 patients.

A variety of guidelines have been rapidly produced (e.g. NICE guidance on critical care escalation (3), Surviving Sepsis guidelines (4) and more), and NHS England with the Faculty of Intensive Care Medicine plan release of UK management guidelines shortly. However, solid data upon which to base these are sparse. As a result, optimal management depends on rapid dissemination of experiential learning.

To address this need, the Intensive Care Society held a webinar on April 3rd 2020 at which representatives from 11 of the most COVID-19 experienced hospital trusts in England & Wales shared learning around 5 specific topic areas in an open forum. Further webinars will be arranged over the next few weeks. Details can be obtained from info@ics.ac.uk ([subject] Knowledge-sharing).

This paper summarises the emerging learning and practice shared by those frontline clinicians. It represents their professional opinions and should not be used as a clinical guideline document. Key emerging knowledge and suggestions for practice are presented in **bold**. *Bullets in italics outline modifications to clinical practice made by individual trusts.*

1. Covid-19 underlying patho-physiology & presentation of respiratory failure

 - Covid-19 appears to have several phases (5). Management should be guided by timing of presentation in relation to the onset of symptoms to understand where in the trajectory of the disease the patient is.
 - Early phase respiratory failure seems predominantly due to impaired pulmonary perfusion. Pro-coagulation leading to micro-vascular pulmonary thrombosis has been observed. Lung compliance is generally good.
 - Later respiratory failure can involve ARDS and bacterial pneumonia

2. Mechanical Ventilation

- **Aggressive ventilation in the early phase may adversely affect later outcomes. Starting positive end-expiratory pressure (PEEP) should be lower than previously recommended. PEEP 10cm H₂O or less appears satisfactory for many: higher PEEP can impair pulmonary perfusion, and gains in oxygenation are often limited. CO₂ clearance is often not greatly improved by aggressive ventilation, given the high shunt fraction/increase in physiological deadspace which is a core driver.**
- **Consider early proning of patients to improve VQ matching.**
 - Examples of clinical practice:*
 - *proning on admission to ICU if in early phase (predominantly perfusion) disease. It can be done irrespective of PF ratio, and if response is positive, this may avoid aggressive ventilation.*
 - *using cut off PF ratio ≤ 16 for proning*
 - *using 'proning teams' to manage turning, utilising non-ICU staff from around the hospital*
- **Pulmonary vasodilatation may provide short-term benefit.**
 - Examples of clinical practice:*
 - *Using nitric oxide in early stages – it can help but may become refractory after 96 hours or so*
 - *Using nebulised or IV prostacyclin – this may be helpful as part of therapeutic trial, if you are using wet ventilation circuits*
- **Cast formation and plugging can affect dry circuits: wet circuits *may* be beneficial.**
 - Examples of clinical practice:*
 - *Using wet circuits for all-Covid areas where full personal protective equipment is in use*
 - *Using checklists to monitor HMEs e.g. 12-hourly as these can fill with water rapidly – this is important in context of reduced nursing ratio. Routinely change every 24 hours in any event if not needing a change before that.*
 - *Use of mucolytics (e.g. N-acetyl cysteine) may be considered.*
 - *Managing patients on anaesthetic machines with higher flow rates may help limit need for soda lime changes (but beware total O₂ use limitations), zoning the machines together, educating nursing staff on use of the machines, asking anaesthetists familiar with machines to support, transferring patients out for weaning*
- **Severe upper airway swelling in some patients may make extubation difficult.**
 - Examples of clinical practice:*
 - *Using dexamethasone prior to extubation, having nebulised adrenaline available, with surgical airway expertise (e.g. ENT) on site and on standby.*
 - *Cohorting patients ready for extubation to areas with relevant expertise and extubation protocols*
 - *A mobile airway team might be an alternative*
- **Re-intubation rates within 24-48 hours seem higher than expected (up to 60%) so delaying initial extubation for longer than usual may be sensible.**
 - Examples of clinical practice*
 - *Waiting 48 hours past fever resolution and monitoring the inflammatory markers to ensure hyperinflammatory state is improving before attempted extubation*

3. Antibiotics

- **Antibiotic usage should be judicious. There are some reports of later aspergillosis and candida infections.**

Examples of clinical practice:

- *Stopping antibiotics in Covid patients unless clearly indicated, using procalcitonin (PCT) and other inflammatory markers to monitor for bacterial infection and restarting as required**
- *Using procalcitonin as a 'stop' signal to guide when to stop antibiotic use**

**False negative PCTs seem less of an issue than false positives in determining antibiotic use - anecdotally, rising procalcitonin has been seen in patients without evidence of bacterial infection, perhaps in relation to 'cytokine storm', and so a low PCT may be more helpful (true negative) than a high PCT (false positive)*

4. Fluid Balance & Renal Support

- **Profound pyrexia, followed by use of CPAP, may mean that patients are dehydrated.**
- **Careful use of fluids aiming for euvolaemia may be beneficial in early phase disease - *provided that* the ventilation effort is well controlled to avoid oedema – as perfusion is a problem and prior to intensive care patients tend to have been dry.**
- **Renal injury has been more common in UK cases than anticipated (20-35% of ICU patients). Careful attention to adequate hydration, and use of lower PEEP, may help.**
- **Consider commencing therapeutic anticoagulation prior to haemofiltration, as filter life appears very short in its absence.**

Examples of clinical practice:

- *Using systemic unfractionated or treatment dose low molecular weight heparin and using aPTT and anti-factor Xa levels for monitoring*
- *Close partnership with renal team to manage resources*
- *Using shorter sharper diafiltrating to service machines to more than 1 person & manage filter supply*

5. Workforce & Infection Control

- **Ensure staff are comfortable in their Personal Protective Equipment (PPE).**

Example of clinical practice:

- *Buddying for donning & doffing, potentially using medical students to observe & confirm effective practice*

- **Promote effective use of the Teams model (e.g. for resus, intubation, proning).**

Example of clinical practice:

- *Developing standard operating procedures for tasks for helpers. Resources are available including the [NHS England website](#) & [BACCN](#) which are updated daily and can be adapted for local use*

- **Beware of 'cognitive overload' which can detract from getting basic ITU care right.**

Example of clinical practice:

- *Having a tactical commander on site so the clinical leads can focus on clinical tasks & provide support to the nursing and allied staff*

References

1. Public Health England. COVID-19 Dashboard.
<https://www.arcgis.com/apps/opsdashboard/index.html#/f94c3c90da5b4e9f9a0b19484dd4bb14>.
(2020, accessed 07 April 2020)
2. Wu, Z, McGoogan, J. Characteristics of and Important Lessons from the Coronavirus Disease 2019 (COVID-19) Outbreak in China. JAMA - Journal of the American Medical Association (2020).
doi:10.1001/jama.2020.2648.
3. National Institute for Health and Care Excellence. COVID-19 rapid guideline: critical care in adults. NICE 2020 (NICE guideline [NG159]). <https://www.nice.org.uk/guidance/ng159>
4. Alhazzani W, Møller M, Arabi Y, Loeb M, Gong M, Fan E et al. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). *Intensive Care Medicine*. 2020; pp. 1-41. doi:10.1007/s00134-020-06022-5.
5. Gattinoni L. et al. COVID-19 pneumonia: different respiratory treatment for different phenotypes? *Intensive Care Medicine* (2020); doi: 10.1007/s00134-020-06033-2