

2021-03

# Galleria mellonella larvae exhibit a weight-dependent lethal median dose when infected with methicillin-resistant *Staphylococcus aureus*

Hesketh-Best, PJ

<http://hdl.handle.net/10026.1/16889>

---

10.1093/femspd/ftab003

Pathogens and Disease

Oxford University Press (OUP)

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

1 ***Galleria mellonella* larvae exhibit a weight-dependent lethal**  
2 **median dose when infected with Methicillin-resistant**  
3 ***Staphylococcus aureus***

4 **Authors**

5 Poppy J. Hesketh-Best<sup>1\*</sup>, Michelle V. Mouritzen<sup>2</sup>, Kayleigh Shandley-Edwards<sup>1</sup>,  
6 Richard A. Billington<sup>1</sup>, and Mathew Upton<sup>2</sup>

7 <sup>1</sup>School of Biological and Marine Sciences, and <sup>2</sup>School of Biomedical Sciences,  
8 University of Plymouth, PL4 8AA UK.

9 \* Corresponding author: Poppy J. Hesketh Best

10 ([poppy.heskethbest@plymouth.ac.uk](mailto:poppy.heskethbest@plymouth.ac.uk))

11 **Abstract**

12 *Galleria mellonella* is a recognised model to study antimicrobial efficacy; however,  
13 standardisation across the scientific field and investigations of methodological  
14 components are needed. Here we investigate the impact of weight on mortality  
15 following infection with Methicillin-resistant *Staphylococcus aureus* (MRSA). Larvae  
16 were separated into six weight groups (180-300 mg at 20 mg intervals) and infected  
17 with a range of doses of MRSA to determine the 50% lethal dose (LD<sub>50</sub>), and the  
18 'lipid weight' of larvae post-infection was quantified. A model of LD<sub>50</sub> values  
19 correlated with weight was developed. The LD<sub>50</sub> values, as estimated by our model,  
20 were further tested *in vivo* to prove our model.

21 We establish a weight-dependent LD<sub>50</sub> in larvae against MRSA and demonstrate that  
22 *G. mellonella* is a stable model within 180-260 mg. We present multiple linear  
23 models correlating weight with: LD<sub>50</sub>, lipid weight, and larval length. We demonstrate  
24 that the lipid weight is reduced as a result of MRSA infection, identifying a potentially  
25 new measure in which to understand the immune response. Finally, we demonstrate  
26 that larval length can be a reasonable proxy for weight. Refining the methodologies  
27 in which to handle and design experiments involving *G. mellonella*, we can improve  
28 the reliability of this powerful model.

## 29 **Key Words**

30 Methicillin-resistant *Staphylococcus aureus*; *Galleria mellonella*; antibiotic testing;  
31 LD<sub>50</sub>; pre-clinical model; fat body

## 32 **Introduction**

33 *Galleria mellonella* (Greater wax moth) larvae are widely utilised for toxicity  
34 screening (Desbois and Coote 2012; Maguire, Duggan and Kavanagh 2016; Coates  
35 *et al.* 2019) and to study host-pathogen interactions (Peleg *et al.* 2009; Olsen *et al.*  
36 2011; Junqueira 2012; Wojda and Taszłow 2013). Unlike many insect models, *G.*  
37 *mellonella* can be incubated at 37°C, which facilitates the investigation of human  
38 pathogens. This has included most of the ESKAPE pathogens: *Enterococcus*  
39 *faecium* (Chibebe Junior *et al.* 2013; Luther *et al.* 2014); *Staphylococcus aureus*  
40 (Brackman *et al.* 2011; Ramarao, Nielsen-Leroux and Lereclus 2012; Sheehan,  
41 Dixon and Kavanagh 2019); *Klebsiella pneumoniae* (Wand *et al.* 2013; Diago-  
42 Navarro *et al.* 2014); *Acinetobacter baumannii* (Peleg *et al.* 2009); and  
43 *Pseudomonas aeruginosa* (Jander, Rahme and Ausubel 2000; Seed and Dennis  
44 2008). Additionally, *Escherichia coli* (Leuko and Raivio 2012; Alghoribi *et al.* 2014;  
45 Jønsson *et al.* 2017; Guerrieri *et al.* 2019), *Bulkholderia mallei* (Schell, Lipscomb and  
46 DeShazer 2008) and several fungi (Cotter, Doyle and Kavanagh 2000; Reeves *et al.*  
47 2004; Mylonakis *et al.* 2005) have also been studied using *G. mellonella*. Crucially, a  
48 positive correlation between the virulence and immune responses between  
49 mammalian models and *G. mellonella* has been established for *P. aeruginosa*  
50 (Jander, Rahme and Ausubel 2000), *Cryptococcus neoformans* (Mylonakis *et al.*  
51 2005), and *S. aureus* (Sheehan, Dixon and Kavanagh 2019), demonstrating the  
52 powerful potential of this invertebrate model.

53 Antibiotic efficacy at dosages recommended for human use can be tested in *G.*  
54 *mellonella*, in addition to their toxicity correlating with toxicity observed in murine  
55 models (Ignasiak and Maxwell 2017). This has been shown with both natural and  
56 synthetic compounds (Gibreel and Upton 2013; Smitten *et al.* 2019), opening up the  
57 possibility of a rapid and cheap model for the early stages of discovery and  
58 development of natural and synthetic products, without the challenges of ethical  
59 approval, specialist training and the difficulties of using mice-models in early-stage  
60 drug development. Infections caused by antibiotic-resistant *S. aureus* are of global

61 concern and it is listed as a high priority pathogen for which new antibiotics are  
62 urgently needed (The World Health Organisation 2017). Methicillin-resistant *S.*  
63 *aureus* (MRSA) has been utilised with *G. mellonella* for the study of virulence  
64 (Mannala *et al.* 2018), pathogenicity (Ebner *et al.* 2016), antimicrobial efficacy of  
65 existing antimicrobials (Ba *et al.* 2015; Ferro *et al.* 2016), and for novel candidates  
66 (Gibreel and Upton 2013; Jacobs *et al.* 2013; Dong *et al.* 2017) (Table S1).

67 Despite the increased popularity of *G. mellonella*, there is much variability in method  
68 application (Andrea, Krogfelt and Jenssen 2019). This includes differences in larval  
69 size, storage, infective dose, and injection intervals. In this study, we address larval  
70 size and its potential impact in experimental design. In antibiotic efficacy studies,  
71 typically the model is infected with a pathogen shortly before the candidate treatment  
72 is presented. This has not been standardised with respect to the parameters  
73 previously mentioned for *G. mellonella*. In our preliminary experimentation in  
74 determining a 50% lethal dose (LD<sub>50</sub>) for MRSA in *G. mellonella*, it was noted that  
75 smaller larvae were more susceptible to infection than larger larvae. This was when  
76 using a broad range of larval weights (~200-300 mg), as previously reported (Jacobs  
77 *et al.* 2013). Furthermore, the larval weight has been demonstrated to positively  
78 correlate with the larval liquid volume, leading to recommendations on how *in vivo*  
79 concentrations of injected compounds and pathogens should be calculated (Andrea,  
80 Krogfelt and Jenssen 2019). This led us to hypothesise that the larvae LD<sub>50</sub> for a  
81 pathogen, in our case here MRSA, is directly proportional to the larvae weight and  
82 that larvae weight is an essential parameter in experimental design that must be  
83 tightly controlled.

84 When physical and anatomical barriers are breached, the wax moth larvae have an  
85 innate immune response relying on germline-encoded factors for the detection and  
86 clearance of microbial pathogens (Trevijano-Contador and Zaragoza 2019). There  
87 are two branches, cellular and humoral immunity. Cellular immunity is conducted by  
88 haemocytes, which are present in an open circulatory system called the  
89 haemolymph, which is analogous to vertebrate blood. There are at least six  
90 subpopulations of haemocytes which perform similar roles to those of the myeloid  
91 lineage in vertebrates (Boman and Hultmark 1987; Lavine and Strand 2002), and  
92 they are also associated with digestive system, trachea and fat body (Ratcliffe 1985).  
93 Five types of haemocytes were identified in fifth larval instar of *G. mellonella*;

94 prohaemocytes, plasmatocytes, granulocytes, oenocytoids and spherulocytes  
95 (Salem *et al.* 2014). The main immune processes include coagulation, phagocytosis  
96 and encapsulation (Tojo *et al.* 2000). Circulating haemocyte density increases during  
97 pathogenesis due to the release of suspended cells from the fat body (Tojo *et al.*  
98 2000). Haemocyte density and subpopulation variations changes with time of  
99 exposure to pathogen and pathogen virulence (Arteaga Blanco *et al.* 2017).

100 Melanisation additionally occurs in the haemolymph, the process of melanin  
101 production resulting in the darkened appearance of the larvae (Tojo *et al.* 2000). The  
102 humoral branch is involved in the production of lytic enzymes (Vogel *et al.* 2011),  
103 and antimicrobial peptides (AMPs) that are active against bacterial pathogens  
104 (Cytryńska *et al.* 2007; Tsai, Loh and Proft 2016). These molecules are mostly  
105 produced by the larval 'fat body', analogous to the mammalian liver, and are  
106 released into the haemolymph (Zasloff 2002).

107 A proteomic investigation has shown *S. aureus* infections lead to an increase in  
108 production of proteins such as AMPs and peptidoglycan recognition proteins  
109 (Sheehan, Dixon and Kavanagh 2019). Critically, the same study identified  
110 similarities between *G. mellonella* and mammal immune response to *S. aureus*  
111 infections. What has not been investigated is the physiological change in *G.*  
112 *mellonella* lipid as a result of *S. aureus* infections. For this investigation, we were  
113 motivated to quantify the lipid weight, a proxy for the fat body, of the larvae to  
114 observe how the fat body might have been affected as a result of MRSA infection.

115 The aim of the work here is to investigate methodological adjustments which may  
116 improve the reproducibility and reliability of using pet-food grade *G. mellonella* as an  
117 experimental model. This was achieved by (i) examining the effect of larval weight on  
118 the LD<sub>50</sub> to MRSA infection, and (ii) characterising physiological changes occurring  
119 to lipid weight as a result of the larval immune response to MRSA.

## 120 **Materials and Methods**


### 121 *Cultivation of MRSA*

122 A single colony of Methicillin-resistant *Staphylococcus aureus* (MRSA) NTCT 12493  
123 was streaked onto fresh Luria broth (LB, Fischer Scientific, UK; Tryptone 10 g/L,  
124 yeast extract 5 g/L, and sodium chloride 10 g/L) solidified with 1.5% agar (Acros

125 Organics, UK) 24 h before experimentation. Single colonies were suspended in  
126 Dulbecco A Phosphate buffered saline (PBS, Oxoid UK) to a range of optical density  
127 (OD) read at 600 nm (Eppendorf BioPhotometer, Netherlands). These dilutions were  
128  $OD_{600} = 0.1 - 1.0$  in 0.1 increments. Viable cell counts were made of each dilution.

### 129 *Determining a weight-based LD<sub>50</sub> for Galleria mellonella larvae*

130 Larvae were purchased commercially from Livefoods UK Ltd. (Somerset, UK;  
131 [www.livefoods.co.uk](http://www.livefoods.co.uk)). On receipt, larvae were individually weighed using an  
132 accurate scale and grouped into the following weight bands: 180-200, 201-220, 221-  
133 240, 241-260, 261-280, and 281-300 mg. Larvae were stored at 4°C for up to 7 days  
134 in the dark with no food and water. Healthy larvae were identified by a uniform cream  
135 colour, with no indications of melanisation such as spots or markings (Fig. 1A) (Li *et*  
136 *al.* 2018). Larvae were euthanised by chilling them at 4°C for 1 h, before freezing  
137 them at -20°C for a minimum of 24 h.



138

139 **Figure 1:** *Galleria mellonella* larvae. (A) Melanisation is a visual indication of the health of  
140 the larvae, as larvae progress from none to complete melanisation as a result of stress  
141 and/or infection. (B) Larvae pupation. (C) Route of infection for larvae is by intra-  
142 haemocoelic injection at the penultimate pro-leg (arrow). Larvae diagram was adapted from  
143 Singkum *et al.* (2019). (D) Larvae are divided up into six weight groups.

144 Larvae ( $n = 10$ ) from each weight band were infected by injecting 10  $\mu$ l of one of the  
145 10 dilutions of MRSA into the left penultimate pro-leg (Fig. 1C), using a 50  $\mu$ l  
146 Hamilton 750 syringe (Hamilton Company, UK) with a removable needle. Injected  
147 larvae were placed into Petri dishes lined with tissue paper (KIMTECH, UK). Three

148 independent replicates of this experiment were carried out. Syringes were cleaned  
149 before and after each bacterial dilution. Cleaning consisted of taking up and  
150 discarding of each wash solution thrice before progressing to the next wash solution.  
151 Wash solution order was as follows: distilled H<sub>2</sub>O (dH<sub>2</sub>O), 70% ethanol, and dH<sub>2</sub>O.

152 After infection, the larvae were maintained at 37°C in the dark without food or water.  
153 A placebo control of sterile PBS was used to account for the effect of the physical  
154 trauma of injection, along with a non-manipulation (NM) control. After 24 h the  
155 live/dead counts were recorded. Larvae were recorded as dead when they met the  
156 following: (i) complete melanisation (Fig. 1A), (ii) did not respond to touch, and (iii)  
157 could not correct itself when rolled onto its back.

158 Determining the weight-dependent LD<sub>50</sub>, live/dead counts were converted into  
159 percentage mortality at 24 h for each group. For this investigation we have defined  
160 LD<sub>50</sub> as CFU of MRSA per mg of organism resulting in 50% mortality. To model the  
161 dose-response and describe the relationship between increasing the infection dose  
162 on survival for each weight group, a non-linear sigmoidal regression curve was  
163 plotted. The infection dose, represented as CFU/ mg of total weight of larva, was log-  
164 transformed. A non-linear regression curve was calculated to fit best the data  
165 generated from three independent replicas. From the equation generated from this  
166 curve, the theoretical LD<sub>50</sub> was calculated along with the standard deviation (SD).  
167 Estimated LD<sub>50</sub> from each weight groups were plotted against the mean larvae  
168 weight. A regression line was drawn, and the coefficient of determinant  $R^2$  was  
169 calculated.

#### 170 *Correlating larval size with rate of pupation*

171 On the day of receipt, larvae were placed into weight groups in Petri dishes. They  
172 were immediately placed at 37°C, in the dark with no food or water and permitted to  
173 pupate over 15 days. Larvae were observed daily and pupation events recorded.

#### 174 *Quantifying lipid weight of G. mellonella*

175 Following investigation of the LD<sub>50</sub> for MRSA, the lipid weight for all living and dead  
176 larvae was quantified. Live larvae from treatments, the NM and PBS controls were  
177 ethically euthanised. Dead larvae were stored at -20°C until needed. Larvae were left  
178 to thaw at room temperature for 24 h and were weighed and individually placed in

179 Eppendorf tubes to be dried over 7 days at 55°C, and re-weighed to reveal their dry  
180 weight. Larvae were then submerged in ≥99.9% diethyl ether (Sigma-Aldrich, UK)  
181 and left for 3 days at 4°C to dissolve lipid. Diethyl ether was utilised as the lipid  
182 extraction solvent (Tzompa-Sosa *et al.* 2014). After, ether was left to evaporate in a  
183 fume hood for 24 h. Once dried, larvae were weighed again to acquire the post-ether  
184 weight. Quantities are then presented as followed: ‘total weight’ is the weight of the  
185 larvae pre-experimentation; ‘water weight’ ( $water\ weight =$   
186  $pre-experimentation\ weight - dry\ weight$ ); ‘lipid weight’ ( $lipid\ weight =$   
187  $dry\ weight - post-ether\ weight$ ).

### 188 *Statistical analysis*


189 All statistical analysis was performed using PRISM GraphPad 8.4.2 (GraphPad  
190 Software, San Diego, CA, USA). One-Way ANOVA (two-tailed), Two-Way ANOVA,  
191 and Pearson’s correlation coefficients were used when applicable to compare  
192 treatment groups. Log-rank Mantel-Cox tests compared survival curves for  
193 antimicrobial efficacy tests and pupation. A *p*-value of: < 0.05 (\*), < 0.01 (\*\*) or <  
194 0.001 (\*\*\*), < 0.0001 (\*\*\*\*) was considered to be significantly different.

## 195 **Results**

### 196 *Larval weight affects LD<sub>50</sub>*

197 To begin testing our hypothesis, LD<sub>50</sub> values were determined for each weight group.  
198 A sigmoidal non-linear model best fit the dose-dependent response of the data,  
199 resulting in an LD<sub>50</sub> calculated for each weight group (Fig. 2). When adjusted to the  
200 number of cells injected into each larvae per one unit of body weight (CFU/mg), the  
201 resulting LD<sub>50</sub> ranged from 1.19 x10<sup>7</sup>CFU/mg, for the 180-200 mg group, to the  
202 highest LD<sub>50</sub> which was 8.97 x10<sup>7</sup> CFU/mg for the 261-280 mg group (Table 1). The  
203 LD<sub>50</sub> increased across weight groups except for the 281-300 mg group, which had a  
204 lower LD<sub>50</sub> than the 261-280 mg weight group. Throughout this experiment, we  
205 encountered some difficulties when handling larvae from the two higher weight-  
206 bands (261-280 and 281-300 mg), such as high variation in mortality at the lowest  
207 infective dosages (0-40% mortality) and highest dosages (60-100% mortality).  
208 Nevertheless, we were able to calculate an LD<sub>50</sub> with the final data.





209

210

211 **Figure 2.** Sigmoidal non-linear logistic regressions best fit the dose-dependent  
 212 response observed when calculating an LD<sub>50</sub> for MRSA. LD<sub>50</sub> was calculated for each weight  
 213 group with 10 larvae/group. Data are shown as mean ± SD (n = 10) of three independent  
 214 replicas.

215 **Table 1.** Summary of the LD<sub>50</sub>s as calculated by non-linear models for each weight group (N,  
 216 number of replicas; R<sup>2</sup>, coefficient of determination).

Weight group (mg)	LD <sub>50</sub> (CFU/mg) <sup>(a)</sup>	SD (CFU/mg)	N	R <sup>2</sup>
180-200	1.19 x10 <sup>7</sup>	1.47	30	0.85
201-220	1.26 x10 <sup>7</sup>	2.45	30	0.77
221-240	2.34 x10 <sup>7</sup>	1.57	30	0.80
241-260	4.40 x10 <sup>7</sup>	3.35	30	0.76
261-280	8.97 x10 <sup>7</sup>	1.46	30	0.69
281-300	4.19 x10 <sup>7</sup>	1.81	30	0.78


217

<sup>(a)</sup>Example of how this is calculated can be found in Table S3

218

219 We observed a positive correlation between weight of the larvae and LD<sub>50</sub>, as  
 220 calculated by Pearson correlation test (r = 0.87, p = 0.025, n = 18). A linear  
 221 regression model arriving at an equation (y = 0.007966x + 5.548) was used to  
 222 estimate LD<sub>50</sub> (Fig. 3A). The LD<sub>50</sub> values, as estimated by our model, were tested *in*

223 *vivo*, demonstrating an approximate 50-56% ( $\pm 5.7 - 10\%$ ) survival for four of the  
 224 weight groups (Fig. 3B). Survival at 24 h for the weight groups 261-280 and 281-300  
 225 mg was 30% ( $\pm 0\%$ ) and 43% ( $\pm 15.3\%$ ), respectively.




226

227 **Figure 3.** LD<sub>50</sub> as calculated by non-linear regression models positively correlated with weight and  
 228 was validated *in vivo* for all but the two highest weight groups. (A) Calculated LD<sub>50</sub> by non-linear  
 229 models correlation positively with total weight. (B) LD<sub>50</sub> value as calculated by the model was  
 230 validated by injecting into larvae and observing mortality. Data are shown as mean  $\pm$  SD (n=10) of  
 231 three independent replicas.

### 232 *MRSA infection leads to a reduction in lipid weight*

233 With the non-manipulated (NM) group, we assessed the overall relationship between  
 234 total weight, dry weight, and lipid weight and length of the larvae (Fig. 4). Determined  
 235 by Pearson's correlation test, we found a positive correlation between the total and  
 236 dry weight, ( $r = 0.972$ ,  $p < 0.0001$ ,  $n = 83$ ) (Fig. 4A), and total and water weight ( $r =$   
 237  $0.989$ ,  $p < 0.0001$ ,  $n = 83$ ) (Fig. 4B). These two results support the findings of  
 238 previous research (Andrea, Krogfelt and Jenssen 2019). Two additional positive  
 239 correlations were observed between total weight and lipid ( $r = 0.788$ ,  $p < 0.0001$ ,  $n =$   
 240  $83$ ) (Fig. 4C), and total weight and length ( $r = 0.9944$ ,  $p < 0.0001$ ,  $n = 252$ ) (Fig. 4D).




241

242 **Figure 4.** Multiple correlations observed between larvae total weight and dry weight, water  
 243 weight, lipid weight, and larvae length. Non-manipulated (NM) larvae were used to analyse  
 244 the relationships between (A) total weight and dry weight, (B) total weight and the lipid  
 245 weight after here presented as lipid weight, (C) total weight and lipid weight as proportional  
 246 to the total weight, water weight, and (D) total weight and larvae length where data is  
 247 presented as mean  $\pm$  SD (n = 252).

248 We also investigated the effect of infection on the lipid weight of all the larvae used in  
 249 determining the LD<sub>50</sub> for MRSA (Fig. 5). As calculated by one-way ANOVA, injection  
 250 with MRSA resulted in an overall decrease in the lipid weight for both dead (18.7 mg  
 251  $\pm$  8.541,  $p < 0.0001$ , n = 573) and live larvae (22.4 mg  $\pm$  6.556,  $p < 0.0001$ , n = 524),  
 252 when compared to the NM control (31.92 mg  $\pm$  8.815, n = 83) (Fig 5A). When  
 253 compared to one another, live larvae had a significantly greater lipid weight  
 254 compared to dead larvae ( $p < 0.0001$ ). There was no significant reduction in the lipid  
 255 weight between NM and PBS control (27.81 mg  $\pm$  5.825,  $p > 0.999$ , n = 50) (Fig 5A  
 256 and Table S2).

257




258

259 **Figure 5.** Injection of the larvae with MRSA results in an overall decreased in the lipid weight  
 260 of the larvae. (A) Statistical results from a one-way ANOVA are illustrated above the bars as  
 261 compared to the NM control. Summary of multiple analysis can be found in Table S2. (B)  
 262 Box-plots above and to the right of the scatter plot are to illustrate the distribution of the data.  
 263 Colours are as follows: black, NM control; purple, PBS control; blue, live larvae; and red,  
 264 dead larvae 24 h post-MRSA infection. Correlations of infective dose and lipid weight for (C)  
 265 living and (D) dead larvae. Data is presented as Log[CFU], as the infective doses are not  
 266 adjusted for larvae weight. Data presented as mean  $\pm$  SD ( $n = 10$ ) of three independent  
 267 replicas.

268 Finally, we observed that at a high infective dosage of MRSA, the larvae had a lipid  
 269 weight close to the mean of the NM and PBS control compared to the lower dosages  
 270 (Fig. 5C-D). This was supported by a positive correlation between lipid weight and  
 271 infective dose for both live ( $r = 0.778$ ,  $p = 0.008$ ) (Fig. 5C) and dead larvae ( $r =$   
 272  $0.669$ ,  $p = 0.035$ ) (Fig. 5D).

273 *Pupation is unaffected by weight*

274 To explore whether larger larvae were closer to the final instar stage (pupae) in  
275 which they begin to pupate into adult moths, an observational experiment was  
276 performed. NM larvae were left to pupate at 37°C, and it was observed that 80-100%  
277 of larvae pupated within the 15 day incubation period, independent on their weight  
278 grouping, as calculated by Log-rank (Mantel-Cox) test ( $X^2(5, N = 60) = 4.004, p =$   
279  $0.549$ ) (Fig. 6).



280

281 **Figure 6.** The weight did not influence the probability of pupation of NM larvae. NM larvae  
282 were incubated at 37°C for 15 days and observed daily for pupation events. No significant  
283 difference was found between the weight group and the probability of pupation as calculated  
284 by a Log-rank Mantel-Cox test ( $p = 0.5489$ ). Data are shown as mean  $\pm$  SD ( $n = 10$ )  
285 repeated twice.

## 286 Discussion

### 287 *MRSA exhibits a weight-dependent LD<sub>50</sub>*

288 In this study, we have demonstrated it is possible to develop a model in which a LD<sub>50</sub>  
289 can be predicted based on the weight of the larvae, and that the prediction can be  
290 experimentally validated (Fig. 3). The linear model correlating total and water weight  
291 (Fig. 4B) imply that in increasingly larger larvae, the *in vivo* dilution of MRSA  
292 increases requiring a greater density of pathogen to reach the LD<sub>50</sub>. Likewise for the

293 positive correlation confirmed with total and lipid weight (Fig. 4C), the presence of a  
294 larger fat body that can be degraded for the production of immune factors, may well  
295 be why we observe the weight-dependent effect on LD<sub>50</sub>. The LD<sub>50</sub>s (1.19 – 8.97  
296 x10<sup>7</sup> CFU/mg) for the MRSA strain was not within range of infective dosages utilised  
297 in previously investigated MRSA and Methicillin-sensitive *S. aureus* (MSSA) strains  
298 (0.8 – 5.0 x10<sup>6</sup> CFU) (Table S1). However, a direct comparison may not be  
299 appropriate given the variation in reporting densities as in our study the LD<sub>50</sub> was  
300 adjusted to account for *in vivo* dilution in the larvae as described in Andrea, Krogfelt  
301 and Jenssen (2019), but this is not always done.

302 During the process of this investigation, we found two of the largest weight groups  
303 (261-280 and 281-300 mg) to be unreliable, which hindered progress. This was  
304 consistent across multiple batches of larvae orders. LD<sub>50</sub>, as calculated by our model  
305 for 261-280 and 281-300 mg larvae, resulted in less than 50% survival at 24 h (Fig.  
306 3B), indicating that our model for a weight-dependent LD<sub>50</sub> had overestimated the  
307 LD<sub>50</sub>. Our first assumptions were that larger larvae were older and closer to pupation  
308 than the smaller weight groups, as larvae increase in size until pupation (Jorjão *et al.*  
309 2018), which might somehow impact on survival. Given the difficulty in identifying an  
310 age for each larva, it is a difficult hypothesis to test beyond quantifying the number of  
311 days it took for NM larvae from each weight group to pupate.

312 When this was conducted, we found that larval size did not influence the probability  
313 of pupation (Fig. 6), and we conclude that the larvae received from the supplier had  
314 an 80-100% probability of pupating within 15 days if kept at 37°C, regardless of  
315 weight. It would appear that larger larvae were not likely to be closer to pupation than  
316 smaller ones, so the reason for our observed decrease in LD<sub>50</sub> for large larvae  
317 remains unknown. Since larvae were kept without food, this may be a reason for the  
318 observed similar pupation times across all weight groups as lack of food source may  
319 be forcing the larvae into pupation. Feeding regimes are not the standard protocol  
320 when investing antibiotic efficacy, as such we feel this best represented the  
321 conditions larvae would be exposed to at the start of experimentation.

322 Using *G. mellonella* does have drawbacks, one such being the functional equivalent  
323 of adaptive immunity termed 'immune priming' (Little and Kraaijeveld 2004; Sadd  
324 and Schmid-Hempel 2006). Individual larvae that survive infection or exposure to a

325 particular pathogen may exhibit increased immune resistance against the same or  
326 similar pathogens. Priming with heat-killed pathogens was observed to result in  
327 increased larval survival (Wu *et al.* 2014). Ultimately there will be no control over the  
328 immune history of the larvae and this should always be recognised when working  
329 with pet-food grade *G. mellonella*. Across the literature, a wide range of weight  
330 bands have been utilised: 150-200 mg (Mannala *et al.* 2018); 300-700 mg (Ebner *et*  
331 *al.* 2016); 200-300 mg (Jacobs *et al.* 2013); and in other studies this is not declared  
332 (Ba *et al.* 2015; Jorjão *et al.* 2018). Our results suggest that choosing weight ranges  
333 as wide as 300-700 mg and 200-300 mg could result in inconsistent data. While a  
334 weight range of only 20 mg is likely a conservative approach, ranges such as 100  
335 mg or greater in our weight-dependent LD<sub>50</sub> model for MRSA indicates that there  
336 would be significant differences in survival (Fig. 3A).

337 Weighing individual larvae is a time-consuming procedure. This study also  
338 demonstrated that larvae length is reasonable proxy for the weight (Fig. 4D). Larval  
339 length has been previously used to characterise larvae for experimentation where  
340 larvae of 15-25 mm were utilised (Bazaid *et al.* 2018). Like total weight, a large  
341 length grouping may also encounter similar challenges. A 20 mg weight grouping  
342 would equate to roughly 1 mm, for example, 180-200 mg would be 20-21 mm.  
343 Measuring length may be a preferred alternative to accurately weighing all larval.  
344 When sourcing larvae from our supplier, we frequently found that larvae belonging to  
345 the weight groups 201-220 and 221-240 mg were most abundant, which will  
346 inevitably be the practical determining factor in weight group selection. Our findings  
347 would support selection of larvae in this range.

#### 348 *Lipid metabolism occurs in response to MRSA infection*

349 MRSA infection leads to a decreased lipid weight in the larvae after 24 h, whether  
350 they died or survived the infection (Fig. 5A). The reduction in lipid weight is likely the  
351 result of lipolysis during an immune response. This is to be expected, the fat body of  
352 the larvae produce many defence compounds essential to the larvae's immune  
353 response (Cytryńska *et al.* 2007; Tsai, Loh and Proft 2016). This reaction can be  
354 rapid, in some models showing production of AMPs within the first 4 to 6 h post-  
355 infection (Sheehan, Dixon and Kavanagh 2019; Trevijano-Contador and Zaragoza  
356 2019). This is supported by proteomic work, which demonstrated that at 6 and 24 h

357 post-*S. aureus* infection larvae had increased expression of AMPs (Sheehan, Dixon  
358 and Kavanagh 2019).

359 On exposure to the infecting pathogen, there may be a rapid metabolism of the fat  
360 body to provide the required energy to fight the infection. Larvae with larger lipid  
361 weight before infection might be more likely to survive, as seen with the surviving  
362 larvae having a greater lipid weight than dead larvae (Fig. 5A). Within this  
363 experimental design, the larvae are not fed before or during the experiment, and  
364 therefore they cannot be acquiring more lipid. Where lipid weight was seen as closer  
365 to the NM and PBS control baseline, as observed in the trend of lipid weight  
366 positively correlating with infective dose (Fig. 5C-D), it is more likely that lipid  
367 metabolism has been compromised.

368 What could reasonably be expected is that lipolysis of the fat body occurs to  
369 increase the production of AMPs and additional defence compounds. When  
370 *Drosophila* are stimulated by a systemic infection with *S. aureus*, signalling from the  
371 Toll receptor increases, which leads to increased production of AMPs and reduced  
372 accumulation of lipids (Liu *et al.* 2016; Lee and Lee 2018). This could suggest that  
373 for larvae surviving high infective dosages, there are additional immune responses  
374 that do not deplete the fat body.

375 We intended to quantify the larval lipid weight to aid in understanding the weight-  
376 dependent LD<sub>50</sub> effect and the observed unreliability of the two largest weight groups  
377 (261-280 and 281-300 mg). We report several observations regarding the lipid  
378 weight and MRSA infection; however, none can fully explain the irregularity we  
379 encountered for the largest weight groups. Analysing larval lipid weight has proved  
380 some insight, but would benefit from further investigation, though alternative  
381 methods to estimate lipid mass would be required.

### 382 *Overall assessment of G. mellonella as a model*

383 There remains a lack of widely available and cheap standardised stocks of larvae  
384 reared under controlled conditions. Temperature (Mowlds and Kavanagh 2008), diet  
385 (Banville, Browne and Kavanagh 2012; Jorjão *et al.* 2018), past infections (Fallon,  
386 Kelly and Kavanagh 2012), and antibiotics and hormones in the feed (Büyükgüzel  
387 and Kalender 2008) are all reported to influence laboratory experimentation. Most



388 larvae currently used are acquired from commercial insect food providers (Andrea,  
389 Krogfelt and Jenssen 2019), where it is understood that use of antibiotics and  
390 hormones in the culture medium is common practice, and acquiring accurate  
391 information regarding the conditions in which the larvae are reared is challenging. All  
392 of which may vary between larvae suppliers, which is a challenge that warrants  
393 further investigation.

394 Ultimately from our investigation, it would appear that lipid deposits are essential in  
395 *G. mellonella* response to MRSA. Prior investigation has evaluated the effect of  
396 nutrient deprivation on larvae (Banville, Browne and Kavanagh 2012), and the  
397 selection of diet (Jorjão *et al.* 2018), which both influence susceptibility to *S. aureus*  
398 infection. This emphasises the issues associated with a having lack of knowledge of  
399 rearing conditions used by suppliers and how they will influence experimental  
400 results. TruLarv™ (BioSystems Technology, UK) currently provide the only  
401 standardised *G. mellonella* in the UK. While cheap compared to murine models, it is  
402 considerably costlier (£1.20 per larvae) than purchasing larvae from commercial pet  
403 food providers.

404 However for pet-food grade larvae to be reliably used in research, more significant  
405 consideration should be taken over the parameters that can be controlled, and in this  
406 study, we emphasise that such experiments can be reproducible and reliable. We  
407 recommend that investigators consider the potential variability associated with using  
408 different larval weight as we have shown herein. We would recommend using weight  
409 groupings as a means to control this. . Our data suggests that all larvae used should  
410 be within 10 mg of the mean weight of all larvae to provide consistency. Additionally,  
411 larvae of >260 mg should not be avoided.

412 In this work, we present several linear regression curves that could be used as tools  
413 to aid in experimental design, such as the linear model for LD<sub>50</sub> (Fig. 3A), weight and  
414 lipid content (Fig. 4C), and length (Fig. 4D). Finally, we demonstrate that the lipid  
415 weight is reduced as a result of MRSA infection, identifying a potentially new  
416 measure in which to understand the immune response. Similarities between *G.*  
417 *mellonella* and mammals in response to *S. aureus* infections can be used to study  
418 the efficacy and interactions of novel antimicrobials, even at early development  
419 stages. By refining and standardising methodologies in which to handle and select

420 *G. mellonella* for study, we can improve the reliability of this powerful model for  
421 multiple purposes.

## 422 **Author Contributions**

423 Conceptualisation, PJHB and MVM; Methodology, PJHB, MVM, KSE, RAB, and MU;  
424 Validation, PJHB, MVM, and KSE; Formal Analysis, PJHB; Investigation, PJHB and  
425 MVM; Resources, PJHB and MVM; Data Curation, PJHB and MVM; Writing –  
426 Original Draft Preparation, PJHB; Writing – Review & Editing, PHB, MVM, KSE,  
427 RAB, and MU; Visualization, PJHB; Supervision, RAB, and MU; Funding Acquisition,  
428 MU.

## 429 **Funding**

430 This study was funded in part by the University of Plymouth, School of Biology and  
431 Marine Science postgraduate research studentship and Innovate UK Antibiotic  
432 Development grant (Innovate UK 103358).

## 433 **Transparency declaration**

434 The authors declare no conflict of interests. The funders had no role in the design of  
435 the study; in the collection, analyses, or interpretation of data; in the writing of the  
436 manuscript, or in the decision to publish the results.

## 437 **References**

438 Algoribi MF, Gibreel TM, Dodgson AR *et al.* *Galleria mellonella* infection model  
439 demonstrates high lethality of ST69 and ST127 uropathogenic *E. coli*. *PLoS*  
440 *One* 2014;**9**:e101547–e101547.

441 Andrea A, Krogfelt KA, Jenssen H. Methods and Challenges of Using the Greater  
442 Wax Moth (*Galleria mellonella*) as a Model Organism in Antimicrobial  
443 Compound Discovery. *Microorganisms* 2019;**7**:85.

444 Arteaga Blanco LA, Crispim JS, Fernandes KM *et al.* Differential cellular immune  
445 response of *Galleria mellonella* to *Actinobacillus pleuropneumoniae*. *Cell Tissue*  
446 *Res* 2017;**370**:153–68.

447 Ba X, Harrison EM, Lovering AL *et al.* Old drugs to treat resistant bugs: Methicillin-  
448 resistant *Staphylococcus aureus* isolates with *mecC* are susceptible to a  
449 combination of penicillin and clavulanic acid. *Antimicrob Agents Chemother*

450 2015;**59**:7396–404.

451 Banville N, Browne N, Kavanagh K. Effect of nutrient deprivation on the susceptibility  
452 of *Galleria mellonella* larvae to infection. *Virulence* 2012;**3**:497.

453 Bazaid AS, Forbes S, Humphreys GJ *et al.* Fatty Acid Supplementation Reverses  
454 the Small Colony Variant Phenotype in Triclosan-Adapted *Staphylococcus*  
455 *aureus*: Genetic, Proteomic and Phenotypic Analyses. *Sci Rep* 2018;**8**, DOI:  
456 10.1038/s41598-018-21925-6.

457 Boman HG, Hultmark D. Cell-free immunity in insects. *Annu Rev Microbiol*  
458 1987;**41**:103–26.

459 Brackman G, Cos P, Maes L *et al.* Quorum sensing inhibitors increase the  
460 susceptibility of bacterial biofilms to antibiotics in vitro and in vivo. *Antimicrob*  
461 *Agents Chemother* 2011;**55**:2655–61.

462 Büyükgüzel E, Kalender Y. *Galleria mellonella* (L.) Survivorship, Development and  
463 Protein Content in Response to Dietary Antibiotics. *J Entomol Sci* 2008;**43**:27–  
464 40.

465 Chibebe Junior J, Fuchs BB, Sabino CP *et al.* Photodynamic and Antibiotic Therapy  
466 Impair the Pathogenesis of *Enterococcus faecium* in a Whole Animal Insect  
467 Model. *PLoS One* 2013;**8**:55926.

468 Coates CJ, Lim J, Harman K *et al.* The insect, *Galleria mellonella*, is a compatible  
469 model for evaluating the toxicology of okadaic acid. *Cell Biol Toxicol*  
470 2019;**35**:219–32.

471 Cotter G, Doyle S, Kavanagh K. Development of an insect model for the in vivo  
472 pathogenicity testing of yeasts. *FEMS Immunol Med Microbiol* 2000;**27**:163–9.

473 Cytryńska M, Mak P, Zdybicka-Barabas A *et al.* Purification and characterization of  
474 eight peptides from *Galleria mellonella* immune hemolymph. *Peptides*  
475 2007;**28**:533–46.

476 Desbois AP, Coote PJ. Chapter 2 - Utility of Greater Wax Moth Larva (*Galleria*  
477 *mellonella*) for Evaluating the Toxicity and Efficacy of New Antimicrobial Agents.  
478 In: Laskin AI, Sariaslani S, Gadd GM (eds.). *Advances in Applied Microbiology*.  
479 Vol 78. Academic Press, 2012, 25–53.

480 Diago-Navarro E, Chen L, Passet V *et al.* Carbapenem-resistant *Klebsiella*  
481 *pneumoniae* exhibit variability in capsular polysaccharide and capsule  
482 associated virulence traits. *J Infect Dis* 2014;**210**:803–13.

483 Dong CL, Li LX, Cui ZH *et al.* Synergistic effect of pleuromutilins with other  
484 antimicrobial agents against *Staphylococcus aureus* in vitro and in an  
485 experimental *Galleria mellonella* model. *Front Pharmacol* 2017;**8**:553.

486 Ebner P, Rinker J, Nguyen MT *et al.* Excreted cytoplasmic proteins contribute to  
487 pathogenicity in *Staphylococcus aureus*. *Infect Immun* 2016;**84**:1672–81.

488 Fallon J, Kelly J, Kavanagh K. *Galleria mellonella* as a Model for Fungal  
489 Pathogenicity Testing. *Methods Mol Biol* 2012;**845**:469–85.

490 Ferro TAF, Araújo JMM, dos Santos Pinto BL *et al.* Cinnamaldehyde Inhibits  
491 *Staphylococcus aureus* Virulence Factors and Protects against Infection in a  
492 *Galleria mellonella* Model. *Front Microbiol* 2016;**7**:2052.

493 Gibreel TM, Upton M. Synthetic epidermicin NI01 can protect *Galleria mellonella*  
494 larvae from infection with *Staphylococcus aureus*. *J Antimicrob Chemother*  
495 2013;**68**:2269–73.

496 Guerrieri CG, Pereira MF, Galdino ACM *et al.* Typical and Atypical  
497 Enterococcal *Escherichia coli* Are Both Virulent in the *Galleria mellonella*  
498 Model. *Front Microbiol* 2019;**10**:1791.

499 Ignasiak K, Maxwell A. *Galleria mellonella* (greater wax moth) larvae as a model for  
500 antibiotic susceptibility testing and acute toxicity trials. *BMC Res Notes* 2017;**10**,  
501 DOI: 10.1186/s13104-017-2757-8.

502 Jacobs AC, DiDone L, Jobson J *et al.* Adenylate kinase release as a high-  
503 throughput-screening-compatible reporter of bacterial lysis for identification of  
504 antibacterial agents. *Antimicrob Agents Chemother* 2013;**57**:26–36.

505 Jander G, Rahme LG, Ausubel FM. Positive correlation between virulence of  
506 *Pseudomonas aeruginosa* mutants in mice and insects. *J Bacteriol*  
507 2000;**182**:3843–5.

508 Jønsson R, Struve C, Jenssen H *et al.* The wax moth *Galleria mellonella* as a novel  
509 model system to study Enterococcal *Escherichia coli* pathogenesis.

- 510 *Virulence* 2017;**8**:1894–9.
- 511 Jorjão AL, Oliveira LD, Scorzoni L *et al.* From moths to caterpillars: Ideal conditions  
512 for *Galleria mellonella* rearing for in vivo microbiological studies. *Virulence*  
513 2018;**9**:383–9.
- 514 Junqueira JC. *Galleria mellonella* as a model host for human pathogens: Recent  
515 studies and new perspectives. *Virulence* 2012;**3**, DOI: 10.4161/viru.22493.
- 516 Lavine MD, Strand MR. Insect hemocytes and their role in immunity. *Insect*  
517 *Biochemistry and Molecular Biology*. Vol 32. Pergamon, 2002, 1295–309.
- 518 Lee KA, Lee WJ. Immune–metabolic interactions during systemic and enteric  
519 infection in *Drosophila*. *Curr Opin Insect Sci* 2018;**29**:21–6.
- 520 Leuko S, Raivio TL. Mutations that impact the enteropathogenic *Escherichia coli* Cpx  
521 envelope stress response attenuate virulence in *Galleria mellonella*.  
522 *Infect Immun* 2012;**80**:3077–85.
- 523 Li Y, Spiropoulos J, Cooley W *et al.* *Galleria mellonella* - a novel infection model for  
524 the *Mycobacterium tuberculosis* complex. *Virulence* 2018;**9**:1126–37.
- 525 Little TJ, Kraaijeveld AR. Ecological and evolutionary implications of immunological  
526 priming in invertebrates. *Trends Ecol Evol* 2004;**19**:58–60.
- 527 Liu B, Zheng Y, Yin F *et al.* Toll Receptor-Mediated Hippo Signaling Controls Innate  
528 Immunity in *Drosophila*. *Cell* 2016;**164**:406–19.
- 529 Luther MK, Arvanitis M, Mylonakis E *et al.* Activity of daptomycin or linezolid in  
530 combination with rifampin or gentamicin against biofilm-forming *Enterococcus*  
531 *faecalis* or *E. faecium* in an in vitro pharmacodynamic model using simulated  
532 endocardial vegetations and an *in vivo* sur. *Antimicrob Agents Chemother*  
533 2014;**58**:4612–20.
- 534 Maguire R, Duggan O, Kavanagh K. Evaluation of *Galleria mellonella* larvae as an in  
535 vivo model for assessing the relative toxicity of food preservative agents. *Cell*  
536 *Biol Toxicol* 2016;**32**:209–16.
- 537 Mannala GK, Koettwitz J, Mohamed W *et al.* Whole-genome comparison of high and  
538 low virulent *Staphylococcus aureus* isolates inducing implant-associated bone

539 infections. *Int J Med Microbiol* 2018;**308**:505–13.

540 Mowlds P, Kavanagh K. Effect of pre-incubation temperature on susceptibility of  
541 *Galleria mellonella* larvae to infection by *Candida albicans*. *Mycopathologia*  
542 2008;**165**:5–12.

543 Mylonakis E, Moreno R, El Khoury JB *et al.* *Galleria mellonella* as a model system to  
544 study *Cryptococcus neoformans* pathogenesis. *Infect Immun* 2005;**73**:3842–50.

545 Olsen RJ, Ebru Watkins M, Cantu CC *et al.* Virulence of serotype M3 group A  
546 *Streptococcus* strains in wax worms (*Galleria mellonella* larvae). *Virulence*  
547 2011;**2**:111.

548 Peleg AY, Jara S, Monga D *et al.* *Galleria mellonella* as a model system to study  
549 *Acinetobacter baumannii* pathogenesis and therapeutics. *Antimicrob Agents*  
550 *Chemother* 2009;**53**:2605–9.

551 Ramarao N, Nielsen-Leroux C, Lereclus D. The insect *Galleria mellonella* as a  
552 powerful infection model to investigate bacterial pathogenesis. *J Vis Exp* 2012,  
553 DOI: 10.3791/4392.

554 Ratcliffe NA. Invertebrate immunity - A primer for the non-specialist. *Immunol Lett*  
555 1985;**10**:253–70.

556 Reeves EP, Messina CGM, Doyle S *et al.* Correlation between gliotoxin production  
557 and virulence of *Aspergillus fumigatus* in *Galleria mellonella*. *Mycopathologia*  
558 2004;**158**:73–9.

559 Sadd BM, Schmid-Hempel P. Insect Immunity Shows Specificity in Protection upon  
560 Secondary Pathogen Exposure. *Curr Biol* 2006;**16**:1206–10.

561 Salem HM, Hussein MA, Hafez SE *et al.* Ultrastructure changes in the haemocytes  
562 of *Galleria mellonella* larvae treated with gamma irradiated *Steinernema*  
563 *carpocapsae* BA2. *J Radiat Res Appl Sci* 2014;**7**:74–9.

564 Schell MA, Lipscomb L, DeShazer D. Comparative genomics and an insect model  
565 rapidly identify novel virulence genes of *Burkholderia mallei*. *J Bacteriol*  
566 2008;**190**:2306–13.

567 Seed KD, Dennis JJ. Development of *Galleria mellonella* as an alternative infection

568 model for the *Burkholderia cepacia* complex. *Infect Immun* 2008;**76**:1267–75.

569 Sheehan G, Dixon A, Kavanagh K. Utilization of *Galleria mellonella* larvae to  
570 characterize the development of *Staphylococcus aureus* infection. *Microbiol*  
571 *(United Kingdom)* 2019;**165**:863–75.

572 Singkum P, Suwanmanee S, Pumeesat P *et al.* A powerful in vivo alternative model  
573 in scientific research: *Galleria mellonella*. *Acta Microbiol Immunol Hung*  
574 2019;**66**:31–55.

575 Smitten KL, Southam HM, de la Serna JB *et al.* Using Nanoscopy To Probe the  
576 Biological Activity of Antimicrobial Leads That Display Potent Activity against  
577 Pathogenic, Multidrug Resistant, Gram-Negative Bacteria. *ACS Nano*  
578 2019;**13**:5133–46.

579 The World Health Organisation. *Global Priority List of Antibiotic-Resistant Bacteria to*  
580 *Guide Research, Discovery, and Development of New Antibiotics.*, 2017.

581 Tojo S, Naganuma F, Arakawa K *et al.* Involvement of both granular cells and  
582 plasmatocytes in phagocytic reactions in the greater wax moth, *Galleria*  
583 *mellonella*. *J Insect Physiol* 2000;**46**:1129–35.

584 Trevijano-Contador N, Zaragoza O. Immune response of *Galleria mellonella* against  
585 human fungal pathogens. *J Fungi* 2019;**5**, DOI: 10.3390/jof5010003.

586 Tsai CJ-YY, Loh JMS, Proft T. *Galleria mellonella* infection models for the study of  
587 bacterial diseases and for antimicrobial drug testing. *Virulence* 2016;**7**:214–29.

588 Tzompa-Sosa DA, Yi L, van Valenberg HJF *et al.* Insect lipid profile: Aqueous versus  
589 organic solvent-based extraction methods. *Food Res Int* 2014;**62**:1087–94.

590 Vogel H, Altincicek B, Glöckner G *et al.* A comprehensive transcriptome and  
591 immune-gene repertoire of the lepidopteran model host *Galleria mellonella*.  
592 *BMC Genomics* 2011;**12**, DOI: 10.1186/1471-2164-12-308.

593 Wand ME, McCowen JWI, Nugent PG *et al.* Complex interactions of *Klebsiella*  
594 *pneumoniae* with the host immune system in a *Galleria mellonella* infection  
595 model. *J Med Microbiol* 2013;**62**:1790–8.

596 Wojda I, Taszłow P. Heat shock affects host-pathogen interaction in *Galleria*

597 *mellonella* infected with *Bacillus thuringiensis*. *J Insect Physiol* 2013;**59**:894–  
598 905.

599 Wu G, Zhao Z, Liu C *et al.* Priming *Galleria mellonella* (Lepidoptera: Pyralidae)  
600 Larvae With Heat-Killed Bacterial Cells Induced an Enhanced Immune  
601 Protection Against *Photobacterium luminescens* TT01 and the Role of Innate  
602 Immunity in the Process. *J Econ Entomol* 2014;**107**:559–69.

603 Zasloff M. Antimicrobial peptides of multicellular organisms. *Nature* 2002;**415**:389–  
604 95.

605



## Supplementary

**Table S1.** Summary of experimental design utilised for *G. mellonella* experiments. Table adapted from Andrea, Krogfelt and Jenssen (2019) to include details of strains and a brief summary of their characteristics. (LP, left pro-leg; NA, not available, P-, privately purchased larvae from country specified).

<b>S. aureus strain</b>	<b>Strain characteristics</b>	<b>Larval weight/size</b>	<b>Larval origin</b>	<b>Storage</b>	<b>Injection site</b>	<b>Infection</b>	<b>Reference</b>
EDCC 5455-5461	clinical isolates	150–200 mg	Reared	30°C	Not given	1.0 x10 <sup>6</sup>	(Mannala <i>et al.</i> 2018)
EDCC 5464	clinical isolate, MRSA strain						
RN4220		NA	P-UK	4°C	Between segments	~1.3 x10 <sup>6</sup>	(Ba <i>et al.</i> 2015)
LGA251	veterinary isolate, <i>mecC</i> positive						
02.5099.D	clinical isolate, <i>mecC</i> positive						
ATCC 43300	MRSA, type culture	15–25 mm long	P-UK	7 days	LP	0.8–2.6 x10 <sup>6</sup>	(Bazaid <i>et al.</i> 2018)
Newman							
NCTC 13277	MRSA, type culture						
2x clin. isol.							
ATCC 29213	MSSA	~250 mg	P-China	NA	LP	~1.0 x10 <sup>6</sup>	(Dong <i>et al.</i> 2017)
ATCC 43300	MRSA						
N54	MSSA, clinical isolate						
MRSA N9	MRSA, clinical isolate						
USA300 JE2	MRSA	300–700 mg	P-Netherlands	NA	LP	1.0 x10 <sup>6</sup>	(Ebner <i>et al.</i> 2016)
ATCC 25923	type culture	~200 mg	NA	NA	LP	1.0 x10 <sup>3</sup>	(Ferro <i>et al.</i> 2016)
ATCC 6538	type culture						
SA01-04	clinical isolates						
USA300-0114		200–300 mg	P-USA	4°C, 14 days	LP	5.0 x10 <sup>6</sup>	(Jacobs <i>et al.</i> 2013)
UAMS-1							
UAMS-1112							
RN4220							
ATCC 6538	type culture	NA	Reared	28°C	NA	10 <sup>8</sup> CFU/ml	(Jorjão <i>et al.</i> 2018)
ATCC 11195	MSSA, type culture	NA	P-UK	4°C, 14 days	NS	~2.5 x10 <sup>6</sup>	(Gibreel and Upton 2013)

**Table S2.** Multiple comparison results for change in lipid weight 24 h post-MRSA infection described in Figure 5. (PBS; phosphate-buffered saline injection; NM, no manipulation control)

Interaction	Mean rank diff.	Adjusted p-value	Summary
NM vs. PBS	78.87	>0.9999	ns
NM vs. Live	339.5	<0.0001	****
NM vs. Dead	490.7	<0.0001	****
PBS vs. Live	260.6	<0.0001	****
PBS vs. Dead	411.8	<0.0001	****
Live vs. Dead	151.2	<0.0001	****

**Table S3.** Editable excel table that can be used in order to calculate the LD50 for MRSA based on weight-grouping.

### Calculating the LD50 for MRSA NCTC 12493

Equation as given by linear model:  $y = 0.007966x + 5.548$

Larvae weight (mg): $x =$	180	<i>Insert here the median weight for the selected weight group</i>
LD <sub>50</sub> (Log[CFU/mg]): $y =$	9.61E+06	This is the LD <sub>50</sub> for your selected weight group
LD <sub>50</sub> (CFU):	2.88E+09	This is the total density of MRSA 12493 that must be injected into larva of the selected weight-group in order to kill 50% of the population