

2021-03-24

# Cracked Tooth Syndrome: Assessment, Prognosis and Predictable Management Strategies.

Gill, T

<http://hdl.handle.net/10026.1/17598>

---

10.1922/EJPRD\_2232Gill10

European Journal of Prosthodontics and Restorative Dentistry

Dennis Barber

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

## Title of Article: Cracked Tooth Syndrome: Assessment, Prognosis and Predictable Management Strategies

### Abstract:

Cracked tooth syndrome (CTS) is a common presentation in general practice. The diagnosis and management of teeth with CTS may be difficult due to the unknown extent of the crack. This article reviews the aetiology, diagnosis, management and prognosis of teeth with CTS. A thorough examination is required to effectively assess CTS. Intervention should aim to relieve symptoms and brace the remaining tooth structure effectively against further flexion. Restored teeth with CTS have a guarded prognosis due to the risk of further crack propagation, but the chances of survival at 5-years is acceptable (74.1-96.8%).

### Dental Descriptors:

- Cracked Tooth Syndrome / therapy\*
- Dental Restoration, Permanent / methods
- Dental Bonding
- Dental Onlay

### Introduction

Cracked Tooth Syndrome (CTS) often presents as an incomplete tooth fracture. This can be defined as a fracture plane of unknown depth and direction passing through tooth structure which, if not already involving, may progress to communicate with the pulp and or periodontal ligament.<sup>1</sup> Cracks in teeth can be grouped broadly into: craze lines, fractured cusp, cracked tooth, split tooth and vertical root fracture (Table 1).<sup>2</sup> Talim and Gohil (1974) developed a more detailed (or topographical/structure-based) classification which divided fractures into: fractures of enamel, fractures involving enamel and dentine without pulp involvement, fractures involving enamel and dentine with pulpal involvement and fractures of the root.<sup>3</sup>

Diagnosis and management of cracked tooth syndrome (CTS) can present a significant clinical challenge. This frequently occurring clinical presentation is most commonly associated with patients over 40 years old, and there is no clear evidence of gender being a related factor.<sup>4</sup> CTS is relatively common with 4.4 posterior teeth per 100 adults experiencing cracks each year, these are approximately 2.4 times more likely to occur in molars than premolars.<sup>5</sup> The most commonly involved teeth are mandibular second molars, mandibular first molars and maxillary first molars.<sup>6</sup>

CTS is three times more likely to occur in teeth where a preparation involving the marginal ridge has been necessary, compared to teeth with an occlusal restoration.<sup>7</sup> Treatment of CTS can range from minor, requiring simple treatment to resolve, to severe requiring root canal treatment or possibly extraction. However, the decision as to which treatment option to select has often been a dilemma for dentists. It is the purpose of this paper to help practitioners in making diagnostic and treatment decisions which may result in more predictable outcomes in the management of CTS.

### Aetiology of Cracked tooth syndrome.

The aetiology of CTS is multifactorial and can be difficult to determine clinically, occurring in unrestored and restored teeth. Lynch et al., (2002) described the causes as restorative procedures, occlusal factors, or other factors such as developmental conditions.<sup>8</sup> One possible mechanism for CTS occurring more commonly in the ageing dentition is the structures of the tooth becoming more brittle and less elastic, making them more prone to cracks through exceeding the elastic potential of dentine.<sup>9,10</sup> In unrestored teeth, cracks can occur due to excessive occlusal loads due to bruxism, or masticatory accidents such as biting on a hard object. It has also been proposed that affected teeth may have developmental weaknesses due to incomplete fusion of calcifying areas leading to a propensity to form cracks.<sup>11</sup>

If wedging forces, for example from the mesio-palatal cusp of an upper tooth, are placed on both buccal and lingual cuspal inclines the resultant crack may occur in the midline of the tooth and propagate towards the pulp, especially in unrestored teeth. A tooth which has a relatively large restoration will tend to fracture more superficially and produce fewer or no symptoms<sup>7</sup>.

The most commonly affected tooth is the mandibular second molar<sup>12</sup> This is thought to be due to higher loading as the result of the class II lever effect and their proximity to the fulcrum (temporomandibular joint). In addition, a plunger cusp is a common anatomical feature of the opposing maxillary second molar, which can create a central wedging force. Premature contacts or interferences can also result in larger forces being placed on these teeth. Other natural predisposing features include extensive attrition or abrasion, or anatomical factors such as the lingual inclination of the lingual cusps of mandibular molars and steep cusp angle of maxillary premolars.<sup>13</sup> The susceptibility of posterior teeth to increased forces can be due to toothwear reducing the steepness of excursive guidance of the anterior teeth. Additionally, the under-contouring of restorations in posterior teeth can lead to over-eruption and result in deeper intercuspatation of the posterior teeth, which fail to disclude in lateral excursion of the mandible. During mastication, forces placed on teeth are comparatively low (70.6-146N)<sup>14</sup> compared to maximal biting forces (98-715N).<sup>15</sup> The force ratio of molars, premolars and incisors is thought to be 4:2:1.<sup>16</sup> However, individuals can create much higher forces during nocturnal bruxism than conscious function, due to the cortical inhibitors being suppressed during sleep.<sup>17</sup>

Preparation of teeth for caries removal has been shown to reduce tooth rigidity.<sup>7</sup> In addition, un-restored carious teeth have also been shown to be more prone to fracture.<sup>18</sup> Lab based studies have shown that cuspal flexion under load becomes greater with increasing restorative preparation from sound tooth to MO to MOD restoration.<sup>19</sup> As the cusp height doubles, due to the relative apical movement of the cavity floor during caries removal, the deflection will increase 8-fold.<sup>7</sup> This results in microcracks which can propagate and lead to the development of cracked tooth syndrome. Choice of restorative technique can also be an important factor, for instance differences in the coefficient of thermal expansion between the restorative material and tooth can induce fractures.<sup>13</sup> Restorative procedures that have been shown to be linked with crack formation include the placing of self-threading dentine pins, non-incremental use of composite resin, excessive hydraulic pressure when luting tight fitting indirect intra-coronal restorations, or placing amalgam with excessive condensation pressure.<sup>20,21</sup>

### Presentation and Diagnosis

Cracked tooth syndrome commonly presents as a short sharp pain on biting that worsens on release, but cracked teeth may be asymptomatic leading to a delayed diagnosis. If this is the case, cracks can propagate to the pulp causing pulpitis, pulpal necrosis or catastrophic failure. Diagnosis is typically made from a detailed dental examination and pain history, and by excluding other causes.<sup>22</sup>

Characteristic signs of cracked tooth syndrome include patients who clench, grind, or press their teeth together, especially in teeth with a wear facet into dentine. Additional signs of CTS include visible cracks on the distal surface of a tooth, and cracks that block transilluminated light.<sup>23</sup> Diagnosis of CTS is likely when patients report clenching or grinding their teeth, have pain symptoms consistent with CTS, and clinical examination reveals a molar with crack that blocks transilluminated light. In addition, a localized periodontal pocket in an otherwise healthy dentition is highly suggestive of a crack.<sup>24</sup> Kanamaru et al. (2017) found presence of a non-working side interference was commonly found in vital molars with CTS (38 out of 44 teeth).<sup>25</sup>

A thorough examination (See Figure 1) should always be carried out but often cracks are not visible. Inspection of the tooth using magnification and rubber dam isolation can aid diagnosis. Vitality (sensitivity) of the tooth should be confirmed through two methods, typically cold testing and electric pulp tester (EPT). If the patient is symptomatic an attempt to elicit these symptoms should be made to confirm diagnosis. Pressure should be applied to the tooth with a crack finding tool. Utilisation of a crack finding tool is the most specific method to identify the symptomatic tooth as it can be applied to a single cusp and if symptoms are induced, specifically on release, it is a strong indicator of cracked tooth syndrome. Transillumination is another useful method to detect cracks. A curing light is shone towards the tooth from different angles, light will not usually be propagated beyond the fracture line. It should be noted that teeth with deep fissures can also prevent light propagation so the anatomy of the tooth should be considered. If cracked tooth syndrome is suspected, it is advisable to remove any restorations under rubber dam and high magnification to assess the remaining tooth structure and the floor of the cavity for cracks. At this stage an assessment of the tooth's restorability can be made (Figure 2).

### Management of Cracked Teeth

Once a diagnosis of a CTS is made, the management of these teeth should be based clinically on pulp status, restorability of the remaining tooth structure, and the patient's wishes.

A thorough occlusal examination should be undertaken (as part of the diagnosis stage), which should include checking of the static and dynamic occlusion, as well as slides from retruded contact position (RCP) to the position of maximum intercuspation (ICP). When restoring teeth ICP contacts should be maintained to prevent an unplanned change in the occlusal scheme. However, damaging shearing forces can be avoided by planning restorations on non-guiding teeth to disclude on lateral and protrusive excursions of the mandible. The use of articulated study casts will help with planning restorations in cracked teeth.

In CTS, the aims of treatment should be to stabilise the remaining tooth structure to prevent further crack propagation which could result in pulpal communication or catastrophic failure of the tooth. In the first instance, the tooth should undergo a 'provisionalisation' treatment

stage where the effect of the intervention on vitality and clinical symptoms are monitored. If the tooth is necrotic or has irreversible pulpitis, and has been checked under high magnification to exclude the possibility of a crack having propagated over the pulpal floor or into the roots, then the transition to the definitive restorative stage can be immediate (Figure 3, Figure 4) after completing root canal treatment.

In a cracked tooth, some cusps may be undermined leading to weakness. Lateral pressure can be applied to the cusp to see if it fractures away. Cusps should be reduced so that they are approximately 2mm at their thinnest point.<sup>26</sup> Selective enamoplasty can be used to modify plunger cusps, or steep inclines of guiding cusps, prone to excursive forces, in order to minimise the forces on the cracked tooth. However, careful planning and consent is necessary if any occlusal modification is being considered, especially if this involves adjustments to teeth not immediately effected by pathology.

In an unrestored tooth it may be advisable to place an orthodontic band<sup>27</sup> or composite splint.<sup>28</sup> The direct composite splint (DCS) is a directly bonded overlay, in supraocclusion, applied to a tooth with suspected CTS as a diagnostic tool<sup>29</sup>. A published audit of 151 CTS teeth treated with DCS showed that approximately 87% of the splints were successful at 3-months, and that the restorations were surprisingly well tolerated.<sup>29</sup> If placement of an orthodontic band or DCS resolves the patient's symptoms then this can be considered confirmation of a diagnosis of CTS. Further, if the tooth remains asymptomatic it can then be predictably restored with a cuspal coverage restoration. However, if vitality remains unclear the tooth can be prepared, and a long-term provisional crown can be constructed (Composite or Polymethyl Methacrylate [PMMA]) to be placed with a temporary cement. It is essential to avoid eugenol containing temporary cements for this phase as the obtundent (sedative) effect on the pulp may mask underlying symptoms of pulpitis. After a period of monitoring nerve sensibility (the authors recommend 3-months asymptomatic whilst maintaining nerve vitality) a definitive restoration can be provided, or if indicated root canal treatment (RCT) can be completed prior to definitive restoration.

The choice of definitive treatment will depend on crack position, remaining tooth structure, type of restoration to be provided and patient preference. In a tooth with sufficient ferrule, a crown transfers the stresses of occlusal forces to the tooth structure circumscribed by the margin of the crown. If the crown is biomechanically stable this should increase the stability of the crack.<sup>30</sup> Whilst the 'chasing out' of cracks is debated in the literature, there is no convincing evidence that this practice improves outcomes,<sup>31</sup> but where possible crown margins should be placed on sound tooth tissue.<sup>32</sup>

The definitive restoration should provide cuspal coverage<sup>25</sup> and can be direct<sup>31</sup> (Figure 5) or indirect (Figure 6, Figure 7).<sup>33</sup> In teeth where symptoms are due to a cusp base fracture the tooth can be restored with a partial coverage cuspal onlay restoration. However, in teeth where there is a central crack, we would advocate a full coverage restoration due to the previously mentioned need to increase the tooth's biomechanical stability. The authors recommend gold as the material of choice due to superior biocompatibility and structural properties, along with minimal preparation being required. However, several tooth coloured materials also exist, and the decision will be patient and dentist derived after discussion of the risks and benefits.

Following restoration, patients with CTS will need their occlusion monitored and protected from parafunctional loads and an appropriately constructed splint (Figure 8) should be provided after delivery of the final restoration. Particularly if there is any concern regarding parafunction. It is not the remit of this paper to discuss splint design but Jagger and King (2018) give a considered overview of considerations on this topic.<sup>34</sup>

### Prognosis of cracked teeth

Teeth diagnosed with CTS and reversible pulpitis, in the absence of a cuspal coverage are associated with over 8 times higher risk of pulp complications. Therefore, it is essential to brace cracked teeth to minimise flexion and crack propagation. When treated early, these teeth have been shown to maintain vitality in 71% cases at 3 years.<sup>35</sup> However, research methodologies are variable and other studies have found much lower maintenance of vitality in treated teeth, some as low as 20%.<sup>36</sup> The literature shows that the proportion of teeth requiring root canal treatment is greater when there is an associated deep periodontal probing depth corresponding to a crack. Krell and Caplan (2018) found the key prognostic factors negatively associated with survival to be: distal marginal ridge crack, periapical diagnosis such as chronic periapical periodontitis, and pocketing depth over 5mm in the fracture line.<sup>6</sup>

Teeth with cracks extending onto the pulpal floor often require extraction (odds ratio = 4.5, P = 0.07), and multivariable analyses found that extension of cracks onto the pulpal floor independently increased the odds of tooth loss 11-fold (odds ratio = 11, P = 0.033), with other factors being held constant.<sup>37</sup> The 5-year survival estimate in the absence and presence of crack extension onto the pulpal floor was 99% and 88%, respectively.<sup>37</sup> Chen et al. (2017) found that visual cues can help to predict crack severity and showed that the larger the length of crack on the occlusal surface the longer the length of crack on the proximal surface and concluded that both length and width of crack on the occlusal surface are useful predictors of whether the crack involves the root.<sup>38</sup> However, the factor most consistently related to failure was having an isolated probing pocket depth of more than 5 mm in the line of suspected fracture.<sup>36,39,40</sup> Clearly, probing pocket depths must be considered in the context of the patient's general periodontal status.

There is debate regarding whether root canal therapy or extraction is the most appropriate treatment when a diagnosis of pulpal necrosis is considered. Berman and Kuttler (2010) have argued that extraction should be recommended in cracked teeth with necrotic pulps and minimal restorations, because of variability in outcomes.<sup>41</sup> However, the authors suggest that if there are not probing pocket depths  $\geq 5$  mm and a full coverage restoration is placed during and after root canal treatment, these teeth have a reasonable (74.1-96.8%) chance of survival at 5 years<sup>40</sup>. Krell and Caplan (2018) suggested a staging based on clinical findings which gave a prognostic outcome of success at 12 months (Table 2).<sup>6</sup>

### Conclusion

Cracked tooth syndrome can be difficult to diagnose. Unfortunately, in dentistry no two presentations are the same and the diagnosis may not be entirely apparent. Understandably, clinicians may be concerned about undertaking irreversible, as well as biologically and financially costly steps for the management of such teeth. However, teeth

with CTS have been shown to have encouraging success rates when treated early. In cases where a diagnosis is unclear, it is sensible to remove any interferences and place either an orthodontic band or direct composite splint. Then, once a diagnosis is confirmed the tooth should be prepared for direct or indirect cuspal coverage to prevent further flexion of the tooth. The superiority of full coverage versus more conservative treatments such as adhesive onlays is not clear in the literature. A balance between conservation of structure, maintenance of vitality, and the ability to brace and prevent flexion in the remaining tooth structure needs to be achieved.

In CTS, if a crack runs from the mesial to distal, especially in minimally restored tooth, the authors of this paper would give strong consideration to full coverage crown restoration. In cases where symptoms are due to a fractured cusp then a partial onlay restoration is more appropriate, depending on the restorative status of the tooth. The associated risks of loss of tooth vitality will need to be discussed with the patient. Patients who have experienced CTS should have a thorough occlusal assessment and if appropriate establishment of protective anterior guidance and provision of a hard Michigan-type splint may be indicated.

Teeth diagnosed with cracked tooth syndrome do have a guarded prognosis but have been shown to have relatively high survival/success rates at 5-years when managed appropriately. However, in the presence of probing depths over 5mm and cracks across both marginal ridges these teeth have a poorer prognosis. In such cases, the patient should be made aware that it may not be possible to predictably restore this tooth, and a conversation regarding the management of the resultant space may be required.

#### References:

1. Ellis SG. Incomplete tooth fracture--proposal for a new definition. *Br Dent J* 2001 Apr 28;190(8):424-428.
2. American Association of Endodontics. *Cracking the Cracked Tooth Code: Detection and Treatment of Various Longitudinal Tooth Fractures*. 2008.
3. Talim ST, Gohil KS. Management of coronal fractures of permanent posterior teeth. *J Prosthet Dent* 1974 Feb;31(2):172-178.
4. Roh B, Lee Y. Analysis of 154 cases of teeth with cracks. *Dent Traumatol* 2006 Jun;22(3):118-123.
5. Bader J, Martin J, Shugars D. Incidence rates for complete cusp fracture. *Community Dentistry and Oral Epidemiology* 2001;29(5):346-353.
6. Krell KV, Caplan DJ. 12-month Success of Cracked Teeth Treated with Orthograde Root Canal Treatment. *J Endod* 2018 Apr;44(4):543-548.
7. Homewood CI. Cracked tooth syndrome--incidence, clinical findings and treatment. *Aust Dent J* 1998 Aug;43(4):217-222.
8. Lynch CD, McConnell RJ. The cracked tooth syndrome. *J Can Dent Assoc* 2002 Sep;68(8):470-475.
9. Yahyazadehfar Mobin, Zhang Dongsheng, Arola Dwayne. On the Importance of Aging to the Crack Growth Resistance of Human Enamel. *Acta biomaterialia* 2016 March;32:264-274.
10. Nazari A, Bajaj D, Zhang D, Romberg E, Arola D. Aging and the reduction in fracture toughness of human dentin. *J Mech Behav Biomed Mater* 2009 Oct;2(5):550-559.

11. Hasan S, Singh K, Salati N. Cracked tooth syndrome: Overview of literature. *International Journal of Applied & Basic Medical Research* 2015 Sep-Dec;5(3):164-168.
12. Hiatt WH. Incomplete crown-root fracture in pulpal-periodontal disease. *J Colo Dent Assoc* 1975 May;53(3):4-15.
13. Lubisich EB, Hilton TJ, Ferracane J. Cracked teeth: a review of the literature. *J Esthet Restor Dent* 2010 Jun;22(3):158-167.
14. Anderson DJ. Measurement of stress in mastication. I. *J Dent Res* 1956 Oct;35(5):664-670.
15. Helkimo E, Carlsson GE, Helkimo M. Chewing efficiency and state of dentition. A methodologic study. *Acta Odontol Scand* 1978;36(1):33-41.
16. Arnold M. Bruxism and the occlusion. *Dent Clin North Am* 1981 Jul;25(3):395-407.
17. Attanasio R. Nocturnal bruxism and its clinical management. *Dent Clin North Am* 1991 Jan;35(1):245-252.
18. Goel VK, Khera SC, Gurusami S, Chen RC. Effect of cavity depth on stresses in a restored tooth. *J Prosthet Dent* 1992 Feb;67(2):174-183.
19. Hood JA. Biomechanics of the intact, prepared and restored tooth: some clinical implications. *Int Dent J* 1991 Feb;41(1):25-32.
20. Trushkowsky R. Restoration of a cracked tooth with a bonded amalgam. *Quintessence Int* 1991 May;22(5):397-400.
21. Nguyen V, Palmer G. A review of the diagnosis and management of the cracked tooth. *Dent Update* 2009 Jul-Aug;36(6):338-346.
22. Kahler W. The cracked tooth conundrum: terminology, classification, diagnosis, and management. *Am J Dent* 2008 Oct;21(5):275-282.
23. Hilton TJ, Funkhouser E, Ferracane JL, Gilbert GH, Baltuck C, Benjamin P, et al. Correlation between symptoms and external characteristics of cracked teeth: Findings from The National Dental Practice-Based Research Network. *J Am Dent Assoc* 2017 04;148(4):246-256.e1.
24. Mathew S, Thangavel B, Mathew CA, Kailasam S, Kumaravadivel K, Das A. Diagnosis of cracked tooth syndrome. *J Pharm Bioallied Sci* 2012 -8;4(Suppl 2):S242-S244.
25. Kanamaru J, Tsujimoto M, Yamada S, Hayashi Y. The clinical findings and managements in 44 cases of cracked vital molars. *Journal of Dental Sciences* 2017 Sep;12(3):291-295.
26. Elayouti A, Serry MI, Geis-Gerstorfer J, Löst C. Influence of cusp coverage on the fracture resistance of premolars with endodontic access cavities. *Int Endod J* 2011 Jun;44(6):543-549.
27. Ehrmann EH, Tyas MJ. Cracked tooth syndrome: diagnosis, treatment and correlation between symptoms and post-extraction findings. *Aust Dent J* 1990 Apr;35(2):105-112.
28. Banerji S, Mehta SB, Millar BJ. Cracked tooth syndrome. Part 2: restorative options for the management of cracked tooth syndrome. *Br Dent J* 2010 Jun;208(11):503-514.
29. Banerji S, Mehta SB, Kamran T, Kalakonda M, Millar BJ. A multi-centred clinical audit to describe the efficacy of direct supra-coronal splinting--a minimally invasive



- approach to the management of cracked tooth syndrome. *J Dent* 2014 Jul;42(7):862-871.
30. Mamoun JS. On the ferrule effect and the biomechanical stability of teeth restored with cores, posts, and crowns. *European Journal of Dentistry* 2014 Apr;8(2):281-286.
  31. Mamoun JS, Napoletano D. Cracked tooth diagnosis and treatment: An alternative paradigm. *European Journal of Dentistry* 2015 Apr-Jun;9(2):293-303.
  32. Sharma A, Rahul GR, Poduval ST, Shetty K. Short clinical crowns (SCC) - treatment considerations and techniques. *Journal of clinical and experimental dentistry* 2012 Oct;4(4):230.
  33. Lin C, Chang Y, Hsieh S, Chang W. Estimation of the failure risk of a maxillary premolar with different crack depths with endodontic treatment by computer-aided design/computer-aided manufacturing ceramic restorations. *J Endod* 2013 Mar;39(3):375-379.
  34. Jagger R, King E. Occlusal splints for bruxing and TMD - A balanced approach? *Dental Update* 2018 Nov;45(10):912-918.
  35. Wu S, Lew HP, Chen NN. Incidence of Pulpal Complications after Diagnosis of Vital Cracked Teeth. *J Endod* 2019 May;45(5):521-525.
  36. Kim S, Kim S, Cho S, Lee G, Yang S. Different treatment protocols for different pulpal and periapical diagnoses of 72 cracked teeth. *J Endod* 2013 Apr;39(4):449-452.
  37. Sim IGB, Lim T, Krishnaswamy G, Chen N. Decision Making for Retention of Endodontically Treated Posterior Cracked Teeth: A 5-year Follow-up Study. *J Endod* 2016 Feb;42(2):225-229.
  38. Chen M, Fu K, Qiao F, Zhang X, Fan Y, Wang L, et al. Predicting extension of cracks to the root from the dimensions in the crown: A preliminary in vitro study. *J Am Dent Assoc* 2017 10;148(10):737-742.
  39. Tan L, Chen NN, Poon CY, Wong HB. Survival of root filled cracked teeth in a tertiary institution. *Int Endod J* 2006 Nov;39(11):886-889.
  40. Kang SH, Kim BS, Kim Y. Cracked Teeth: Distribution, Characteristics, and Survival after Root Canal Treatment. *J Endod* 2016 Apr;42(4):557-562.
  41. Berman LH, Kuttler S. Fracture necrosis: diagnosis, prognosis assessment, and treatment recommendations. *J Endod* 2010 Mar;36(3):442-446.