

2020-03-01

# Microplastics and seafood: lower trophic organisms at highest risk of contamination

Walkinshaw, C

<http://hdl.handle.net/10026.1/17612>

---

10.1016/j.ecoenv.2019.110066

Ecotoxicology and Environmental Safety

Elsevier

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

1 [Disclaimer](#)

2 ['This is a copy of the accepted paper as submitted for publication. Readers are advised to refer to](#)  
3 [the final version of the paper which can be found at \*Ecotoxicology and Environmental Safety\*, 190,](#)  
4 [110066. doi:10.1016/j.ecoenv.2019.110066](#)

5  
6 **Microplastics and seafood: lower trophic organisms at highest risk of**  
7 **contamination**

8  
9 Chris Walkinshaw<sup>1,2</sup>, Penelope K. Lindeque<sup>1</sup>, Richard Thompson<sup>3</sup>, Trevor Tolhurst<sup>2</sup> & Matthew Cole<sup>1\*</sup>

10  
11 <sup>1</sup> Marine Ecology and Biodiversity group, Plymouth Marine Laboratory, Plymouth PL1 3DH, UK

12 <sup>2</sup> School of Environmental Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4  
13 7TJ, UK

14 <sup>3</sup> School of Biological and Marine Sciences, University of Plymouth, Drake Circus, Plymouth, Devon PL4  
15 8AA, UK

16 \* Corresponding author: MC; email: mcol@pml.ac.uk; telephone: +44(0)1752 633100

17  
18 **Abstract**

19 Microplastic debris is a prevalent global pollutant that poses a risk to marine organisms and ecological  
20 processes. It is also suspected to pose a risk to marine food security; however, these risks are currently  
21 poorly understood. In this review, we seek to understand the current knowledge pertaining to the  
22 contamination of commercially important fished and farmed marine organisms with microplastics,  
23 with the aim of answering the question “Does microplastic pollution pose a risk to marine food  
24 security?”. A semi-systematic review of studies investigating the number of microplastics found in  
25 commercially important organisms of different trophic levels suggests that microplastics do not  
26 biomagnify, and that organisms at lower trophic levels are more likely to be contaminated by  
27 microplastic pollution than apex predators. We address the factors that influence microplastic  
28 consumption and retention by organisms. This research has implications for food safety and highlights  
29 the risks of microplastics to fisheries and aquaculture, and identifies current knowledge gaps within  
30 this research field.

31  
32 **Keywords**

33 Plastic; Food security; Aquaculture; Trophic transfer; Biomagnification

34

## 35 **Highlights**

- 36 • Microplastic contamination of commercially important marine and aquaculture species was  
37 assessed
- 38 • We provide evidence that microplastics do not biomagnify
- 39 • Microplastics are more prevalent in lower trophic organisms
- 40 • There is currently insufficient evidence to consider risks to human health

41

## 42 **1. Introduction**

43 Microplastics are a ubiquitous global contaminant, identified throughout the marine environment,  
44 including seawater, sediment and biota (Cole *et al.*, 2011; Law and Thompson, 2014). Microplastics  
45 describe tiny plastic particulates, although a coherent definition remains under debate, especially in  
46 terms of their size (Frias and Nash, 2019; Hartmann *et al.*, 2019). For the purposes of this review, we  
47 refer to microplastics and nanoplastics as synthetic solid particles or polymer matrices, with at least  
48 one dimension ranging 0.1  $\mu\text{m}$ –1 mm. The literature describes microplastic shapes in a myriad of  
49 different ways, from spheres, beads and fragments, to films, filaments and fibres; for consistency, we  
50 here opt for using the terms “bead” (any spherical plastic), “fibre” (plastic threads such as those used  
51 in clothing), or “fragment” (irregularly shaped particulates). Microplastics can be further classified  
52 based on their origin: primary microplastics are manufactured in the micro size range, and include  
53 cosmetic microbeads, pre-production pellets and industrial scrubbers; secondary microplastics are  
54 formed by the breakdown of macroplastics within the environment (Andrady, 2017). Microplastic  
55 fibres have been identified as a particular concern for the environment, owing to their abundance and  
56 bioavailability, with research suggesting that microplastic fibres can contribute up to 91% of all plastics  
57 collected in global seawater samples (Barrows, Cathey and Petersen, 2018).

58


59 Plastic production has increased rapidly since its inception, with an estimated 8.3 billion metric tonnes  
60 of virgin plastic produced to date. Approximately 4.6 billion metric tonnes of this (55%) has been  
61 produced since 2000 (Geyer, Jambeck and Law, 2017). Microplastics enter the marine ecosystem  
62 through many different pathways, including riverine transport, sewage and wastewater effluent,  
63 direct release (e.g. from shipping and ports) and atmospheric deposition (Boucher and Friot, 2017).  
64 Plastics are incredibly durable,, and rather than undergoing a straightforward process of  
65 mineralization in the marine environment, plastics first degrade into smaller and smaller pieces,  
66 eventually forming micro- and nanoplastics (Andrady, 1998, 2011). Microplastic debris can travel vast  
67 distances via oceanic currents and winds, impinging on remote habitats including mid-oceanic islands  
68 and the polar ice caps (Barnes *et al.*, 2009; Peeken *et al.*, 2018). Sinks of microplastics include the

69 ocean gyres, sediments, shorelines, polar sea ice, and biota, including animals destined for human  
 70 consumption (Hardesty *et al.*, 2017; Peeken *et al.*, 2018). Whilst there are efforts to remove  
 71 microplastics from the marine environment, it is widely accepted that once released, it is practically  
 72 and economically infeasible to recapture marine microplastics for recycling or responsible disposal.

73

74 Microplastics pose a risk to marine life and ecological processes (Galloway, Cole and Lewis, 2017), and  
 75 it has been suggested they may further impact on food security (Barboza *et al.*, 2018a), socio-  
 76 economic wellbeing (Beaumont *et al.*, 2019) and human health (Galloway, 2015). The perceived risks,  
 77 pathways, effects, and consequences arising from microplastic pollution on food security and  
 78 ecosystem health in the marine environment are displayed in Fig. 1.

79



80

81 Fig. 1. Perceived impact pathways of microplastics on food security and ecosystem health.

82

### 83 1.1 Marine food security

84 Fisheries and aquaculture provide a critical proportion of the world's food supply, providing over 4.5  
 85 billion people with at least 15% of their average per capita intake of animal protein (Béné *et al.*, 2015),  
 86 and production is predicted to grow in the future, from 171 million tonnes in 2016 to approximately  
 87 201 million tonnes in 2030, an increase of 17.5% (FAO, 2018). Global fish exports in 2017 were valued  
 88 at 152 billion USD (FAO, 2018). Total capture from fisheries has remained fairly constant since the  
 89 1990s and is not expected to increase considerably, with growth instead expected from aquaculture,  
 90 predominantly in Asia, which as a continent accounts for almost two thirds of global fish consumption

91 (Béné *et al.*, 2015). The FAO predicts that aquaculture production will reach 109 million tonnes in 2030  
92 (FAO, 2018).

93

94 Food security is defined by the Food and Agriculture Organisation as "a situation that exists when all  
95 people, at all times, have physical, social and economic access to sufficient, safe and nutritious food  
96 that meets their dietary needs and food preferences for an active and healthy life" (FAO *et al.*, 2017).  
97 Current identified risks to food security include climate variability due to both short-term events and  
98 climate change, eutrophication, ocean acidification, oxygen depletion, conflict, economic recession,  
99 pathogens, and pollution (Chakraborty and Newton, 2011; Wollenberg *et al.*, 2016). Larger plastic  
100 debris, particularly derelict fishing gear (i.e. abandoned or lost nets, lines, pots), has been shown to  
101 pose a substantial risk to food security. For example, in Chesapeake Bay the removal of 34,408 derelict  
102 fishing pots led to the harvest of an additional 13,504 metric tonnes in blue crab (*Callinectes sapidus*)  
103 valued at 21.3 million USD (Scheld, Bilkovic and Havens, 2016). However, whilst there has been  
104 considerable research into the effects of microplastics on marine organisms, evidence is lacking on  
105 the effect of microplastics on food security and food safety. We hypothesise that in marine ecosystems  
106 already affected by a multitude of environmental stressors, microplastics may represent a significant  
107 additional risk to food security.

108

109 In this review, we critically assess microplastics research with relevance to fishing and aquaculture,  
110 the health of commercially exploited organisms, and food security; to understand the current state of  
111 microplastics research and evaluate whether microplastics pose a risk to food security. Several marine  
112 pollutants are known to biomagnify, causing heightened risk to higher trophic organisms, however,  
113 very little research is available to show whether this may occur with microplastics, with current  
114 research giving opposing viewpoints (GESAMP, 2016; Akhbarizadeh, Moore and Keshavarzi, 2019;  
115 Hantoro *et al.*, 2019). We evaluate currently available data regarding microplastic content within  
116 organisms of different trophic levels to assess whether biomagnification is likely to be a risk with  
117 microplastic contamination. Current research gaps will also be discussed to highlight areas where  
118 unknown risks may threaten marine food security and human health.

119

## 120 **2. Methods**

### 121 **2.1 Sourcing reference material**

122 In order to investigate the prevalence of microplastics in commercially exploited marine organisms,  
123 including fish, shellfish, crustaceans and macroalgae, we undertook a semi-systematic review of the  
124 scientific literature, performed by using a specific set of search terms separated by Boolean operators

125 (Table 1), utilising the academic literature search engines *Web of Science*, *ScienceDirect*, *Pubmed* and  
 126 *PLOS ONE*. This search method was supplemented by use of a snowballing method, where further  
 127 literature was identified in the references of the articles reviewed to encompass the broadest set of  
 128 literature. Only articles published up to the end of 2018 were included in the data analysis in this  
 129 review. See Table 2 for a summary of the number of articles found from each search engine. These  
 130 articles were considered for relevant information and subjected to a quality control step (see below);  
 131 literature that passed this stage was utilised in this review.

132

133 Table 1. Search terms and Boolean operators used in the identification of scientific literature.

Search term	Boolean operator	Search term
Microplastic	AND OR	Food security
Microplastic pollution		Food
Marine microplastic		Marine
		Health
		Fish (including individual species searches)
		Effect
		Shellfish (including individual species searches)
		Bivalve (including individual species searches)
		Organism

134

135 Table 2. Relevant literature identified through searches of different academic literature search engines

Academic search engine	Results retrieved
Web of Science	955
ScienceDirect	1516
PubMed	668
PLOS ONE	46

136

## 137 2.2 Quality control

138 The primary literature from which data was extracted for analysis had been peer-reviewed prior to  
 139 publication, providing a base level of quality assurance. We additionally conducted a quality  
 140 assessment to verify that: (1) experimental replication was performed for statistical analysis; and (2)  
 141 suitable controls were implemented in the study protocol (e.g. negative controls in toxicity testing,  
 142 procedural blanks, and contamination controls in environmental analyses). If any of these quality  
 143 control parameters was not met, the literature was not included in this review. After these steps, the

144 identified literature was cross-referenced with available data showing organisms of global importance  
 145 to aquaculture and fisheries. Following further narrowing of studies to select those that analysed  
 146 organisms of commercial importance, 32 pieces of literature were selected to ascertain the data  
 147 presented in this review.

148

149 **2.3 Data analysis**

150 In the literature data is typically presented as the number of microplastics per individual  
 151 (MP/individual) for fish, or microplastics per gram (wet weight, w. w.) (MP/gram) for shellfish. For  
 152 assessing whether microplastics biomagnify within lower trophic level organisms it was necessary to  
 153 convert MP/individual values by ascertaining mean wet weights for individual species, drawn from  
 154 primary and grey literature. MP/gram w. w. values were subsequently estimated by dividing average  
 155 microplastics per organism by the average mass of that organism as reported in the literature (see  
 156 Table S1 for further information).

157

158 **3. Results**

159

160 **3.1 Risks to food security**

161

162 **3.1.1 Prevalence of microplastics in commercially exploited species**

163 Microplastics can be ingested by a wide range of marine life, and the presence of microplastics in  
 164 marine organisms destined for human consumption has been widely reported. Tables 3 and 4 below  
 165 show the 10 most caught marine species and 10 most farmed aquaculture species in 2016 (FAO, 2018),  
 166 alongside evidence of their capacity to ingest microplastic debris. 60% of the most farmed aquaculture  
 167 species have been investigated for the presence of microplastics, and 80% of the most caught marine  
 168 species have been investigated. The organisms that are not mentioned in any microplastic ingestion  
 169 studies up to the end of 2018 represented a total of approximately 22.5 million tonnes of food in 2016.

170

171 Table 3. 10 most cultured aquaculture species in 2016 (data from FAO, 2018). NIF = no information found.

Common name	Species name	Production (thousand tonnes, 2016)	Habitat	Feeding strategy	Microplastic ingestion reference
Grass carp	<i>Ctenopharyngodon idellus</i>	6 068	Freshwater	Herbivorous	NIF

Silver carp	<i>Hypophthalmichthys molitrix</i>	5 301	Freshwater	Planktivorous	Jabeen <i>et al.</i> , 2017
Cupped oysters NEI	<i>Crassostrea spp.</i>	4 864	Estuarine	Filter feeder	Van Cauwenberghe and Janssen, 2014; Rochman <i>et al.</i> , 2015; Phuong <i>et al.</i> , 2018; Waite, Donnelly and Walters, 2018
Common carp	<i>Cyprinus carpio</i>	4 557	Freshwater	Omnivorous	Jabeen <i>et al.</i> , 2017
Japanese carpet shell	<i>Ruditapes philippinarum</i>	4 229	Seawater and estuarine	Filter feeder	Li <i>et al.</i> , 2015
Nile tilapia	<i>Oreochromis niloticus</i>	4 200	Freshwater	Omnivorous	Rochman <i>et al.</i> , 2015; Biginagwa <i>et al.</i> , 2016
Whiteleg shrimp	<i>Penaeus vannamei</i>	4 156	Seawater	Planktivorous (plus more: detritus, worms, bivalves and crustaceans)	NIF
Bighead carp	<i>Hypophthalmichthys nobilis</i>	3 527	Freshwater	Planktivorous	NIF
Crucian carps	<i>Carassius spp.</i>	3 006	Freshwater	Omnivorous	Jabeen <i>et al.</i> , 2017; Yuan <i>et al.</i> , 2019
Catla	<i>Catla catla</i>	2 961	Freshwater	Planktivorous	NIF



Table 4. 10 most caught marine species in 2016 (data from FAO, 2018). NIF = no information found.

Common name	Species name	Production (thousand tonnes, 2016)	Habitat	Feeding strategy	Microplastic ingestion reference
Alaska pollock	<i>Theragra chalcogramma</i>	3 476	Demersal	Fish and invertebrates	NIF
Peruvian anchovy	<i>Engraulis ringens</i>	3 192	Pelagic	Planktivorous	Ory <i>et al.</i> , 2018
Skipjack tuna	<i>Katsuwonus pelamis</i>	2 830	Pelagic	Fish, crustaceans, molluscs	Rochman <i>et al.</i> , 2015; Choy and Drazen, 2013; Markic <i>et al.</i> , 2018
Sardinellas NEI	<i>Sardinella spp.</i>	2 290	Pelagic	Planktivorous	NIF
Jack and horse mackerels NEI	<i>Trachurus spp.</i>	1 744	Pelagic/ demersal	Fish and plankton	Neves <i>et al.</i> , 2015; Foekema <i>et al.</i> , 2013; Lusher, McHugh and Thompson, 2013; Murphy <i>et al.</i> , 2017; Markic <i>et al.</i> , 2018; Güven <i>et al.</i> , 2017
Atlantic herring	<i>Clupea harengus</i>	1 640	Pelagic	Planktivorous	Ogonowski <i>et al.</i> , 2017; Foekema <i>et al.</i> , 2013; Rummel <i>et al.</i> , 2016; Hermsen <i>et al.</i> , 2017
Pacific chub mackerel	<i>Scomber japonicus</i>	1 599	Pelagic	Fish and plankton	Neves <i>et al.</i> , 2015; Rochman

					<i>et al.</i> , 2015; Güven <i>et al.</i> , 2017; Ory <i>et al.</i> , 2018
Yellowfin tuna	<i>Thunnus albacares</i>	1 463	Pelagic	Fish, crustaceans, molluscs	Choy and Drazen, 2013; Markic <i>et al.</i> , 2018
Atlantic cod	<i>Gadus morhua</i>	1 329	Demersal	Fish and crustaceans	Foekema <i>et al.</i> , 2013; Bråte <i>et</i> <i>al.</i> , 2016; Liboiron <i>et al.</i> , 2016; Rummel <i>et</i> <i>al.</i> , 2016
Japanese anchovy	<i>Engraulis japonicus</i>	1 304	Pelagic	Planktivorous	Tanaka and Takada, 2016

175

### 176 **3.1.2 Fish**

177 Many species of edible demersal, pelagic and reef fish, sampled from across the globe, have been  
178 found to ingest microplastics (Bellas *et al.*, 2016; Rummel *et al.*, 2016; Bråte *et al.*, 2016; Lusher,  
179 McHugh and Thompson, 2013; Ory *et al.*, 2018; Tanaka and Takada, 2016; Rochman *et al.*, 2015; Neves  
180 *et al.*, 2015; Critchell and Hoogenboom, 2018). Of the seven most farmed aquaculture species which  
181 are fish (Table 3), all are freshwater species, and their feeding strategies are mostly planktivorous or  
182 omnivorous, with the exception of the grass carp which is herbivorous and feeds mostly on aquatic  
183 weeds. These fish may be likely to consume microplastics due to their prey being within a similar size  
184 range. However, microplastic ingestion investigations have only been performed on Common carp,  
185 Crucian carps, Nile tilapia and Silver carp, and no data is available for the other three species, even  
186 though they represent a combined 12.5 million tonnes of farmed fish (as of 2016). These studies gave  
187 a combined average amount of microplastics per organism of  $2.5 \pm 1.3$  MP/individual (Common carp),  
188  $1.9 \pm 1.0$  MP/individual (Crucian carps), and  $3.8 \pm 2.0$  MP/individual (Silver carp). Nile tilapia data was  
189 presented by the authors as the number of individuals which had consumed microplastics, which was  
190 an average of 16% (Rochman *et al.*, 2015; Biginagwa *et al.*, 2016). Where it is possible to view the  
191 morphology of plastic particles ingested, fibres are the most common microplastic shape seen and  
192 make up 57.6-86.5% of the plastic shapes observed.

193

194 Of the ten most caught species (Table 4), all are marine fish; the majority are pelagic species that  
195 consume mostly plankton and small fish, with three exceptions (pollock, tuna and cod). The  
196 microplastic content of these fish are much more studied than common aquaculture species, with 80%  
197 of the top ten most fished species included in at least one microplastic study. Collating all available  
198 literature on these organisms gives the following percentages of each species that were seen with  
199 microplastics in their gastrointestinal tract (GIT): 0.9% Peruvian anchovy; 9.4% Skipjack tuna; 24.5%  
200 Jack and Horse mackerels; 8.8% Atlantic herring; 23.3% Pacific chub mackerel; 23.4% Yellowfin tuna;  
201 2.8% Atlantic cod, and 76.6% Japanese anchovy (Neves *et al.*, 2015; Foekema *et al.*, 2013; Lusher,  
202 McHugh and Thompson, 2013; Murphy *et al.*, 2017; Güven *et al.*, 2017; Ogonowski *et al.*, 2017;  
203 Rummel *et al.*, 2016; Hermsen *et al.*, 2017; Rochman *et al.*, 2015; Ory *et al.*, 2018; Choy and Drazen,  
204 2013; Markic *et al.*, 2018; Bråte *et al.*, 2016; Liboiron *et al.*, 2016; Tanaka and Takada, 2016). Other  
205 species of commercial importance that have been included in several pieces of literature (plus  
206 percentages seen with microplastics in their GIT) include Scads (*Decapterus spp.*, 46%), European  
207 pilchards (*Sardina pilchardus*, 26%), Blue whiting (*Micromesistius poutassou*, 29.8%), and Atlantic  
208 mackerel (*Scomber scombrus*, 23.2%). As with aquacultured species, fibres are the most common  
209 microplastic shape seen, forming 30-87.6% of the plastic shapes observed. Unfortunately it is not  
210 possible to view in detail the most common size of microplastics observed in each species due to how  
211 the data is reported, however this information may not be reliable due to constraints in minimum  
212 observable size in the methodology used (e.g. choice of filters, sensitivity of analytical techniques,  
213 Lusher *et al.*, 2017). Notable by its absence in the literature is the Alaska Pollock (*Theragra*  
214 *chalcogramma*) and members of *Sardinella spp.*, neither of which were found to have been analysed  
215 to investigate microplastic ingestion in the literature. Both species are an extremely important food  
216 source, with more than 3.47 million tonnes of Pollock and 2.29 million tonnes *Sardinella spp.* fished in  
217 2016.

218

### 219 **3.1.3 Shellfish**

220 Cupped oysters (*Crassostrea spp.*) and Japanese carpet shell (*Ruditapes philippinarum*) are among the  
221 most prevalently aquacultured shellfish species worldwide. Microplastic ingestion in shellfish is  
222 generally reported as the number of microplastics per gram of wet tissue. In Cupped oysters, the  
223 average result reported ranged from 0.18 to 3.84 microplastics gram<sup>-1</sup> w. w., and in the Japanese  
224 carpet shell, the average reported result ranged from 0.9 to 2.5 microplastics gram<sup>-1</sup> w. w.

225

226 By far the most studied shellfish are mussels of the family *Mytilidae*. 9 pieces of literature were  
227 identified that studied the amount of microplastics found in sea mussels in their natural environments,  
228 with ingestion ranges varying from 0.2-5.36 microplastics g<sup>-1</sup> w. w. (Bråte et al., 2018; Catarino et al.,  
229 2018; De Witte et al., 2014; Li et al., 2015, 2016; Phuong et al., 2018; Qu et al., 2018; Van  
230 Cauwenberghe and Janssen, 2014; Van Cauwenberghe et al., 2015). Whilst ingestion values look  
231 different when analysing the number of microplastics ingested per individual, when normalised for  
232 soft tissue weight, the values for all three species overlap, seemingly showing that microplastic  
233 ingestion in shellfish is not species-specific. Though shellfish can show selective feeding, rejecting  
234 particles based on size or lack of organic material (Newell and Jordan, 1983; Defosse and Hawkins,  
235 1997), they are found to ingest microplastics. Whilst these species all ingest similar amounts of  
236 microplastics, it is possible that they selectively ingest different size microplastics due to organism  
237 size, with for example oysters being able to ingest larger particles than mussels. Data from the analysis  
238 of mussels and oysters taken from the French Atlantic coast (Phuong et al., 2018) suggests this, as  
239 both organisms ingested a majority of microplastics in the 50-100 µm size range, but mussels ingested  
240 a higher proportion of 20-50 µm particles than oysters (37% and 15 %, respectively), and oysters  
241 ingested a higher proportion of > 100 µm particles than mussels (32% and 11%, respectively).

242

#### 243 **3.1.4 Crustaceans**

244 Crustaceans form a very large and diverse group of organisms including many that are important for  
245 worldwide food security, such as crabs, lobsters, crayfish and prawns. Many edible species of  
246 crustaceans have been shown to ingest microplastics (Devriese et al., 2015; Welden and Cowie, 2016a;  
247 Abbasi et al., 2018). Organisms such as copepods and krill are also critically important as a food for  
248 organisms which are consumed by humans, and have been reported to ingest microplastics (Botterell  
249 et al., 2019). No studies have been performed to investigate microplastic ingestion in the Whiteleg  
250 shrimp, one of the top ten most farmed aquatic species with 4.2M tonnes farmed in 2016 (Table 3),  
251 however, investigations have taken place with other commercially important species. Brown shrimp,  
252 *Crangon crangon*, a commercially important crustacean fished in the eastern Atlantic and  
253 Mediterranean Sea, were found with an average of 0.68 ± 0.55 microplastics gram<sup>-1</sup> w. w. and 63% of  
254 the 165 shrimp analysed containing microplastics (Devriese et al., 2015). Green tiger prawn, *Penaeus*  
255 *semisulcatus*, an organism of commercial important in East Africa and Asia, was found to have ingested  
256 an average of 7.8 particles per individual (1.5 particles gram<sup>-1</sup>, n=12) in the Musa estuary, Persian Gulf  
257 (Abbasi et al., 2018). Nylon fibres were observed in the stomachs of 5.93% *Plesionika narval* (narwhal  
258 shrimp), an important fishery in the Aegean Sea, although it is hypothesised by the authors that these  
259 fibres may result from the fishing method (Bordbar et al., 2018). Other commercially important

260 species that have been observed to contain microplastics include *Eriocheir sinensis* (Wójcik-  
261 Fudalewska, Normant-Saremba and Anastácio, 2016), *Carcinus maenas* (Watts *et al.*, 2014, 2015), and  
262 *Nephrops norvegicus* (Murray and Cowie, 2011; Welden and Cowie, 2016b).

263

### 264 **3.1.5 Macroalgae**

265 Seaweeds have been consumed as a traditional food around the globe; however, consumption of  
266 seaweed has been increasing in recent years with much of this increase from farming of seaweed  
267 rather than from harvesting wild crops. Statistics from the Food and Agriculture Organisation of the  
268 United Nations state that aquatic plant production grew from 13.5 million tonnes to over 30 million  
269 tonnes from 1995 to 2016, with 96.5% of the 31.2 million tonnes produced in 2016 from aquaculture  
270 (FAO, 2018). Seaweeds for consumption are generally classified into three groups: red algae  
271 (Rhodophyta) such as Dulse and Nori, brown algae (Phaeophyceae) such as kelp and green algae (an  
272 informal group containing Chlorophyta, Charophyta, Mesostigmatophyceae, Chlorokybophyceae and  
273 Spirotaenia) such as sea lettuce. *Fucus vesiculosus* is a common seaweed in the British Isles and  
274 Atlantic coastlines, in the class of brown algae, and is often consumed as a health supplement. Recent  
275 studies have shown the ability for 20 µm polystyrene microparticles to sorb to *F. vesiculosus* (Sundbæk  
276 *et al.*, 2018). Trophic transfer via this macroalgae has also been observed; Gutow *et al.* (2016)  
277 demonstrated the ability for the common periwinkle *Littorina littorea* to ingest microplastics via *Fucus*  
278 *vesiculosus*. Algal pieces were exposed to polystyrene microbeads (10 µm), fragments (1-100 µm), and  
279 polyacrylic fibres (90 to 2200 µm), followed by a washing step. Feeding assays with the three types of  
280 microplastic-contaminated algal pieces showed that *Littorina littorea* did not show a feeding  
281 preference between contaminated and non-contaminated algal pieces, and microplastics were found  
282 in the stomach content, gut and faecal pellets, with 89% of *L. littorea* faecal pellets containing  
283 microplastics.

284

## 285 **3.2 Factors influencing microplastic consumption**

286

### 287 **3.2.1 Feeding strategy**

288 Broadly speaking, there are two main ways for marine organisms to ingest microplastics: direct  
289 ingestion from the natural environment; or indirect ingestion, including trophic transfer from prey and  
290 consumption of contaminated aquaculture feedstock. Furthermore, there is some indication that  
291 microplastics can be taken up via the gills (Watts *et al.*, 2014). Dietary strategy may be a defining  
292 characteristic influencing microplastic ingestion in fish, with planktivores more likely to consume

293 microplastics direct from the natural environment, while piscivores (e.g. tuna) would be expected to  
294 consume microplastics mainly through trophic transfer via prey or accidental ingestion while feeding.

295

296 Direct ingestion of microplastics is often a consequence of feeding strategy. Indiscriminate feeders  
297 show no selection in the matter that they ingest, ingesting prey in proportion to their availability in  
298 the environment, whilst discriminate feeders select based on preferential feeding factors (colour, size  
299 etc.). Filter feeders such as some bivalves can be considered as indiscriminate feeders as they feed by  
300 filtering water through their gills, capturing particulate matter such as plankton and microalgae. This  
301 is generally in a non-selective manner; however some of the filtered matter can be rejected. This has  
302 been shown recently by Ward *et al.* (2019), who demonstrated that the bivalves *Crassostrea virginica*  
303 and *Mytilus edulis* selectively ingested microplastics preferentially, based on the physical  
304 characteristics of the plastic. In this way, microplastics are ingested if they resemble the properties of  
305 the organic matter these organisms feed on, such as in size and shape. Discriminate feeders may  
306 directly ingest microplastics either when they resemble prey items, or incidentally whilst feeding, e.g.  
307 in contaminated feedstock; this feeding strategy is generally utilised by higher trophic-level organisms.  
308 Discriminate feeders such as fish may therefore ingest microplastics that resemble their prey.  
309 Amberstripe scad (*Decapterus muroadsi*) appear to ingest blue microplastics preferentially as they  
310 resemble their copepod prey in both colour and size (Ory *et al.*, 2017). Evidence of selective feeding  
311 on the blue copepods *Pontella sinica* and *Sapphirina spp.* was seen, as was selectivity for blue  
312 microplastics.

313

314 Indirect ingestion, or “trophic transfer” occurs when organisms consume prey that have already  
315 consumed microplastics. Trophic transfer from blue mussels *Mytilus edulis* to the shore crab *Carcinus*  
316 *maenas* has been observed in laboratory conditions (Farrell and Nelson, 2013; Watts *et al.*, 2014).  
317 Farrell and Nelson (2013) fed 0.5 µm fluorescent polystyrene microspheres to *M. edulis*, with *C.*  
318 *maenas* subsequently being fed one mussel per crab. Microspheres were subsequently detected in  
319 the stomach, hepatopancreas, ovary, gills and haemolymph of the crabs. Results from Nelms *et al.*  
320 (2018) suggest the ability for microplastics to be ingested by grey seals (*Halichoerus grypus*) through  
321 trophic transfer from Atlantic mackerel (*Scomber scombrus*). Detritivores may also be prone to  
322 indirectly consuming microplastics present in faeces of contaminated organisms; for example  
323 coprophagous copepods can ingest microplastics present in other copepods’ egests (Cole *et al.*, 2016).  
324 Feedstock contaminated with microplastics may be a risk to aquaculture, as fishmeal is a commonly  
325 used fish feed manufactured from whole fish, therefore any microplastics within the fish may pass  
326 into the processed fishmeal (Karbalaee *et al.*, 2019).

327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355

### 3.2.2 Trophic level

The percentage of planktivorous and piscivorous fish populations contaminated with microplastics might suggest that trophic level and feeding strategy alone are not indicative of microplastic ingestion, however, this may be due to a difference in how microplastics data are usually presented (See Table 5). For example, Markic *et al.* (2018), saw no significant difference in their study on plastic ingestion rate in 23 species of fish in the South Pacific based on their trophic level, with the only significant difference in ingestion rates seen between benthic predators and omnivores. However, while similar proportions of the total population of marine organisms with different dietary strategies contained microplastics, the number of microplastics per gram of tissue may be very different. For example, data presented in this review shows a similar percentage of *S. japonicus* (23.3%) and *T. albacares* (23.4%) contained microplastics, but the average weight of *T. albacares* caught by Markic *et al.* (2018) is 5228.7 g, whereas the average caught weight for *S. japonicus* by Güven *et al.* (2017) was 28.86 g. Using these weights, the average amount of plastic particles per gram (wet weight) for *Scomber japonicus* from Güven *et al.* (2017) is estimated as 0.33 particles gram<sup>-1</sup> and the maximum number of microplastics found per gram in *Thunnus albacares* from Markic *et al.* (2018) is estimated at 5.9x10<sup>-4</sup> particles gram<sup>-1</sup>, a 1000-fold difference.

In order to investigate this further, 11 commercially exploited taxa, including bivalves, crustaceans and fish, were selected for analysis from a variety of trophic levels. Taxa were selected that had either a wide range of literature available for analysis (e.g. *Mytilus spp.*, *Scomber japonicus*), or were at a trophic level not covered by other data (e.g. *Thunnus albacares*, *Katsuwonus pelamis*). The data was normalized to give the number of microplastics ingested per gram wet weight of these organisms. Table 5 lists the fish, crustaceans and bivalves in which the number of microplastics per gram wet weight of tissue has been calculated.

Table 5. Number of microplastics per gram wet weight marine organisms.

Species	Common name	Family	Diet	Microplastics per gram wet weight	Raw data references
<i>Katsuwonus pelamis</i>	Skipjack tuna	Fish	Largely piscivorous	0.000249	Markic <i>et al.</i> , 2018

<i>Thunnus albacares</i>	Yellowfin tuna	Fish	Largely piscivorous	0.00059	Markic <i>et al.</i> , 2018
<i>Clupea harengus</i>	Atlantic herring	Fish	Planktivorous	0.01	Foekema <i>et al.</i> , 2013
<i>Engraulis ringens</i>	Peruvian anchovy	Fish	Planktivorous	0.057	Ory <i>et al.</i> , 2018
<i>Trachurus spp.</i>	Jack and horse mackerels NEI	Fish	Planktivorous	0.000126-0.14	Foekema <i>et al.</i> , 2013; Neves <i>et al.</i> , 2015; Güven <i>et al.</i> , 2017; Markic <i>et al.</i> , 2018
<i>Scomber japonicus</i>	Pacific chub mackerel	Fish	Planktivorous	0.0025-0.33	Neves <i>et al.</i> , 2015; Güven <i>et al.</i> , 2017; Ory <i>et al.</i> , 2018
<i>Crangon crangon</i>	Brown shrimp	Crustacean	Planktivorous/ herbivorous	0.13-1.23	Devriese <i>et al.</i> , 2015
<i>Penaeus semisulcatus</i>	Green tiger prawn	Crustacean	Planktivorous/ herbivorous	1.5	Abbasi <i>et al.</i> , 2018
<i>Ruditapes philippinarium</i>	Japanese carpet shell	Shellfish	Filter feeder	0.9-2.52	Li <i>et al.</i> , 2015; Davidson and Dudas, 2016
<i>Crassostrea spp.</i>	Cupped oysters	Shellfish	Filter feeder	0.18-3.84	Foekema <i>et al.</i> , 2013; Van Cauwenberghe and Janssen, 2014; Phuong <i>et al.</i> , 2018; Waite, Donnelly and Walters, 2018
<i>Mytilus spp.</i>	Sea mussels	Shellfish	Filter feeder	0.2-5.36	De Witte <i>et al.</i> , 2014; Van Cauwenberghe and Janssen, 2014; Li <i>et al.</i> , 2015,



					2016; Van Cauwenberghe <i>et al.</i> , 2015; Bråte <i>et al.</i> , 2018; Catarino <i>et al.</i> , 2018; Phuong <i>et al.</i> , 2018; Qu <i>et al.</i> , 2018
--	--	--	--	--	--------------------------------------------------------------------------------------------------------------------------------------------------------------------

356

357


358 There is up to four magnitudes of difference between microplastics per gram present in shellfish  
359 compared to higher trophic level fish. The data presented above therefore suggests that trophic level  
360 and feeding strategy may play a key role in the level of microplastic contamination within marine  
361 organisms; though similar percentages of the total population of organisms at different trophic levels  
362 contain microplastics within their body tissues, lower trophic level organisms have a higher proportion  
363 of microplastic comparatively with body weight, which may be more indicative of risks from  
364 microplastics. Fig. 2 displays a comparison of microplastics per gram wet weight of the organisms in  
365 Table 5 with the amount of mercury in tissues of similar organisms reported by Plessi, Bertelli and  
366 Monzani (2001; *Mytilus spp.*) and the FDA (FDA, 2017; all other species). Mercury is well known to  
367 biomagnify, and values are inversely proportional with the microplastic data presented here, which  
368 shows a decrease in microplastic concentration with increasing trophic level. Based on this data, we  
369 conclude that unlike other contaminants such as organochlorines (Borgå, Gabrielsen and Skaare,  
370 2001) or mercury (Lavoie *et al.*, 2013), microplastics do not biomagnify. This is likely because the  
371 evidence currently suggests that microplastics do not, in most cases, translocate from the digestive  
372 system into tissues or circulatory fluid, therefore it is a more transitory contaminant with a limited  
373 residence time within organisms.

374

375

376

377



378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

Fig. 2. A comparison of the number of microplastics (MP) per gram wet weight of organisms of different trophic levels to the amount of Mercury (ppm) reported in the tissues of similar organisms as listed by the FDA (FDA, 2017) and Plessi, Bertelli and Monzani (2001). Trophic level shows general increase with direction of arrow. Line of best fit added to show trend in data. \*Value is average value of ranges shown in Table 5 and Table S1, with error bars displaying the range of the results.

### 3.2.3 Environmental concentrations

It is possible that another variable such as habitat may have a pronounced effect on the amount of microplastic ingestion. Markic *et al.* (2018) saw a significant difference in the vertical habitat of a species and their plastic ingestion rates. Although they did not see a significant difference with respect to horizontal distribution (Neritic/Neritic-oceanic/Oceanic), it may be expected that for example fish caught in an oceanic gyre or other area of high microplastic load may have a higher incidence of ingestion than those caught in other areas. In fact, this is observed in the study in question; significantly higher ingestion of microplastic debris was observed in a sampling area within the South Pacific 'garbage patch' than in fish from other locations. This was seen with for example *Thunnus albacares*, where ingestion was seen in 70% of individuals within the garbage patch, and 24% and 15% at two locations outside of this area. In juvenile fish, there was an increased incidence of microplastic ingestion and increasing concentrations of microplastic in seawater with proximity to the coast, with

397 higher encounter rates where microplastic concentrations exceeded those of fish larvae (Steer *et al.*,  
398 2017).

399

400 Environmental concentrations may be a particularly important variable for microplastic ingestion in  
401 crustaceans and molluscs (Li *et al.*, 2019). As bivalves are filter feeders, any differences in microplastic  
402 ingestion are likely due to microplastic distribution in their habitat. Li *et al.* (2016) investigated  
403 microplastic abundance in mussels in 22 sites along the coast of China, and significant differences in  
404 microplastic ingestion were seen at different sites. Wild mussels contained on average 2.7 items/g  
405 (4.6 items/individual) and farmed mussels contained on average 1.6 items/g (3.3 items/individual). In  
406 heavily contaminated areas, mussels contained an average of 3.3 items/g (5.3 items/individual),  
407 whereas in less contaminated areas, microplastic abundance in mussels was significantly lower (1.6  
408 items/g or 3.3 items/individual). Gut content of individuals of the crustacean *Nephrops norvegicus*  
409 collected from three sites in North and West Scotland had significantly different microplastic  
410 ingestion; 84.1%, 43% and 28.7% of *N. norvegicus* individuals ingested microplastic in the Clyde Sea  
411 Area, North Minch and North Sea, respectively (Welden and Cowie, 2016a), suggesting crustaceans  
412 may also ingest microplastics relative to environmental availability.

413

### 414 **3.3 Risks of microplastics to marine organisms**

415

#### 416 **3.3.1 Retention in the digestive system (gut blockages)**

417 Following ingestion, microplastics may be rejected by the organism through pseudofaeces or post-  
418 ingestion rejection, egested through faeces, transferred across the GIT epithelium, or be retained in  
419 the GIT. Microplastic retention in the digestive system may adversely affect organism health through  
420 physical perforation of the gut or by giving the organism a feeling of false satiety, decreasing feeding  
421 activity and nutrient intake.

422

423 Shore crabs fed with 10 µm polystyrene microspheres had plastic detected in the foregut 5 days after  
424 exposure to microplastic-containing mussels (Watts *et al.*, 2014). In this feeding experiment, crabs  
425 were fed with mussels that had been exposed to microplastics and subsequently sampled over a 21-  
426 day period, and n=6 crabs were analysed for microplastics in the foregut at each time point post-  
427 ingestion. Polystyrene microspheres were detected in all six crabs after 24hrs; decreasing to 50-66%  
428 of the crabs from days 2-5. Microplastics were then not detected in the crab faecal pellets after 7 and  
429 22 days post-exposure (but were on day 14).

430

431 Blue mussels (*Mytilus edulis*) were shown to ingest 9% of all available microplastic fibres (approx. 450  
432  $\mu\text{m}$  length) in an ingestion study where microplastic fibres were ingested alongside the microalgae  
433 *Rhodomonas salina* (Woods *et al.*, 2018). Mussel filtration rate decreased when exposed to  
434 microplastic fibres in addition to *R. salina*, and though most fibres (71%) were rejected as  
435 pseudofaeces, 9% were ingested, and < 1% were excreted in faeces. Microplastics were identified in  
436 the gills, digestive gland and other soft tissues at all time points over a 72 hr exposure period. In  
437 another experimental study, 2 of 31 Palm Ruff (*Seriolella violacea*) fish were shown to retain  
438 microplastics after a 49-day exposure period (Ory *et al.*, 2018). The transitory nature of microplastics  
439 within the digestive system of organisms may explain why microplastics do not appear to biomagnify.  
440 If microplastics pass through the GIT of organisms and are not retained within the GIT or tissues, it is  
441 much less likely that organisms at higher trophic levels will ingest significant amounts of microplastics  
442 through a carnivorous diet.

443

444 Research by Welden and Cowie (2016) suggests whilst Norway lobster (*Nephrops norvegicus*) are seen  
445 to retain microplastics within their foregut for extended periods of time, the main route by which they  
446 are removed is by ecdysis, whereby the individual moults and sheds its gut lining. This gut lining was  
447 found to contain microplastics which were removed from the individual during moulting.

448

### 449 **3.3.2 Growth rate, reproduction or function affected?**

450 Any changes to growth rate, reproduction, mortality or behaviour due to external factors may  
451 significantly alter population dynamics. In the case of commercially important organisms, this may  
452 significantly affect the efficiency and profitability of fishing and aquaculture. Lower growth rates may  
453 mean that fewer organisms can be harvested in a season, or lower reproduction rates may cause  
454 population decreases in following seasons, both of which would have a negative effect on food  
455 security. A similar concept is discussed by Galloway, Cole and Lewis (2017), who propose that, though  
456 chronic exposure to microplastic is not usually lethal, it is associated with reductions in energy,  
457 growth, fecundity and reproductive output. These individual and population-level effects can as a  
458 consequence cause ecosystem level effects, such as community shifts and changes to ecosystem  
459 function, which would result in risks to food security.

460

461 Several articles have shown reduction of growth rates and reproductive function (Cole *et al.*, 2015;  
462 Sussarellu *et al.*, 2016), and behavioural changes (Cole *et al.*, 2015; Sussarellu *et al.*, 2016; Ribeiro *et al.*  
463 *et al.*, 2017; Woods *et al.*, 2018) in marine organisms as a result of exposure to microplastics. Significant  
464 effects from microplastic exposure were observed in laboratory exposure studies with the Pacific

465 oyster (*Crassostrea gigas*) (Sussarellu *et al.*, 2016). Significantly higher algal consumption was  
466 observed for oysters exposed to microplastics, possibly in an attempt for the oyster to compensate  
467 for lower nutrient intake. Significant reproductive effects were observed; exposed female oysters had  
468 fewer, smaller oocytes and a reduction in D-larval yield; exposed male oysters had lower sperm  
469 velocity. *C. gigas* larval growth was significantly slower, with a reduction in mean size of 18.6% at 17  
470 days post-fertilization and a 6-day lag time to metamorphosis.

471

472 Behavioural changes are observed in clams; 20 µm polystyrene microplastics also induced effects on  
473 antioxidant capacity, DNA damage, neurotoxicity and oxidative damage in *Scrobicularia plana* (Ribeiro  
474 *et al.*, 2017), and reduced clearance rate in *Atactodea striata* (Xu *et al.*, 2017). Behaviour may also be  
475 affected in the presence of nanoplastics. For example, Wegner *et al* (2012) observed no pseudofaeces  
476 production in *Mytilus edulis* exposed to microalgae alone, but found heightened pseudofaeces  
477 production in *Mytilus edulis* exposed to microalgae (*Pavlova lutheri*) and 30 nm polystyrene, along  
478 with a decrease in filtering activity.

479

### 480 **3.3.3 Risk of disease**

481 Once in the marine environment, microplastics are quickly colonised by a variety of organisms termed  
482 the plastisphere (Zettler, Mincer and Amaral-Zettler, 2013). The plastisphere is a risk to the marine  
483 environment, aquaculture and food security as it has the potential to support pathogenic  
484 microorganisms, and allow them to become more bioavailable to the organisms consuming  
485 microplastics. Recent research has identified hazardous microorganisms present on microplastics,  
486 along with microorganisms usually found in sewage and gut-associated pathogens (Oberbeckmann,  
487 Löder and Labrenz, 2015). The microbial biofilms discussed here affect the physical characteristics of  
488 the plastic, including size and buoyancy, which could in turn affect the vertical distribution of  
489 microplastics within the water column, transporting microplastics to the benthos (Kaiser, Kowalski and  
490 Waniek, 2017; Kooi *et al.*, 2017). This, in addition to the horizontal transport of microplastics via ocean  
491 currents and wind therefore means that microplastics have the capacity to transport microorganisms  
492 to new environments over vast distances, suggesting the potential for microplastics to act as a vector  
493 for the transfer of invasive pathogens to new environments.

494

495 High concentrations of microplastic debris in the North Pacific subtropical gyre have resulted in an  
496 increase in the pelagic insect *Halobates sericeus* and in *H. sericeus* egg densities (Goldstein, Rosenberg  
497 and Cheng, 2012). Jiang *et al.* (2018) profiled bacterial communities attached to microplastic samples  
498 taken from intertidal locations around the Yangtze estuary in China, and found a wide range of

499 bacterial taxa, including some that are associated with human and animal pathogens: *Vibrio* (0.4% of  
500 taxonomic abundance, found at Xiangshan bay); *Leptolyngbya* (1.6% abundance, found at Chongming  
501 island), and *Pseudomonas spp.* (<0.01% abundance, all plastics).

502

503 Harmful pathogens travelling large distances could have severe implications for food security. One  
504 potential example of this would be the colonisation of marine plastics by HAB (harmful algal bloom)  
505 species. When floating plastic debris collected along the North-west Mediterranean were analysed,  
506 several potentially harmful dinoflagellates were identified, including *Ostreopsis spp*, *Coolia spp* and  
507 *Alexandrium taylori* (Masó *et al.*, 2003), all of which can cause HABs. *Alexandrium spp.* can cause  
508 paralytic shellfish poisoning (PSP), which is hazardous to both marine organisms and humans.  
509 *Alexandrium catanella* has caused significant economic losses to the salmon industry in Chile, for  
510 example in 2009 when a large bloom was associated with a loss of over \$10 million to the Chilean  
511 Salmon industry (Mardones *et al.*, 2015). *Alexandrium taylori* has also been shown to produce  
512 paralytic shellfish toxins and has recently been identified for the first time in Malaysian waters (Lim *et*  
513 *al.*, 2005). Invasive HAB species, potentially transported by microplastics, could therefore be incredibly  
514 damaging to global fishery and aquaculture industries.

515

516 Marine plastic debris collected from multiple locations in the North Atlantic was analysed and bacterial  
517 assemblage sequenced to characterize the plastisphere community (Zettler, Mincer and Amaral-  
518 Zettler, 2013). In this diverse community, the bacteria genus *Vibrio* and dinoflagellate genus  
519 *Alexandrium* were identified. Both of these genii contain species that are pathogenic to both humans  
520 and animals. Several strains of *Vibrio spp.* including potentially pathogenic *Vibrio parahaemolyticus*  
521 were also detected on microplastics and in seawater from the North and Baltic sea by Kirstein *et al.*  
522 (2016). Microplastics samples from a transect taken along the Slovenian coast of the North Adriatic  
523 Sea were subjected to DNA extraction, amplification and phylogenetic analysis, and the bacterial  
524 pathogen *Aeromonas salmonicida* was identified on the particles (Viršek *et al.*, 2017). This species is  
525 pathogenic to several commercially important species, such as salmonids.

526

#### 527 **3.3.4 Chemical additives and adhered contaminants**

528 Microplastics contain chemicals added during plastic manufacture to enhance certain properties, and  
529 have also been shown to adsorb and concentrate contaminants from the environment such as PCBs,  
530 PAHs, and metals (Teuten *et al.*, 2007; Brennecke *et al.*, 2016). Many of these contaminants can be  
531 toxic to marine organisms. Several researchers have therefore investigated whether microplastics can

532 act as a vector for contaminant transfer to marine organisms, and whether this is a significant pathway  
533 compared to other methods of contaminant ingestion.

534

#### 535 3.3.4a Chemical additives

536 Chemical additives in plastics enhance the different properties that make plastics so useful; some act  
537 as fire retardants, while others may act as stabilisers, foaming agents or strength enhancers. When  
538 plastic pollution occurs, these additives slowly leach from plastics into their surrounding media, for  
539 example seawater. This has led to concerns that they may enter biological systems and affect the  
540 health of exposed organisms, however, there is also a growing set of evidence that the overall  
541 exposure of organisms to these chemicals from plastics is negligible compared to other sources.

542

543 The potential for leaching of nonylphenol (NP) and bisphenol A (BPA) in the GIT of *Arenicola marina*  
544 (lugworm) and *Gadus morhua* (Atlantic cod), and a comparison of exposure to these two substances  
545 by microplastics alone and total environmental exposure, was investigated utilising a biodynamic  
546 model by Koelmans, Besseling and Foekema (2014). They suggest that for cod, ingestion of  
547 microplastic is highly unlikely to lead to negative effects from NP and BPA and is negligible compared  
548 to uptake from water and prey. For lugworms, though ingestion of microplastic was hypothesised to  
549 be a substantial exposure pathway in certain conditions, the low concentrations of NP and BPA  
550 involved would not cause a risk to the lugworm.

551

#### 552 3.3.4b Adhered contaminants

553 In addition to leaching chemical additives, plastic particles can sorb contaminants from the  
554 environment, giving a possible route for the concentration of these chemicals, potentially increasing  
555 their toxicity if they are released into a marine organism. Teuten *et al.*, (2007) investigated the uptake  
556 and release of the hydrophobic organic contaminant phenanthrene by three virgin plastic polymers:  
557 polyethylene, polypropylene, and polyvinyl chloride. All three sorbed phenanthrene with varying  
558 efficiency, however all three plastics greatly exceeded the sorption of phenanthrene onto two natural  
559 sediments.

560

561 Ašmonaitė *et al.* (2018) investigated the effect of ingestion of large (100-400 µm) polystyrene  
562 microplastics (PS-MPs) on the rainbow trout (*Oncorhynchus mykiss*). Trout were exposed to virgin  
563 microplastics as well as microplastics exposed to either sewage effluent or environmental water in a  
564 harbour. All three sets of PS-MPs contained chemical contaminants including PAHs, plasticizers and  
565 surfactants, however, a wider variety of compounds were detected after exposure to sewage and

566 harbour water, confirming the ability for PS-MPs to sorb contaminants from the aquatic environment.  
567 Rainbow trout were experimentally exposed to these microplastics following a dietary-exposure  
568 protocol, however no significant changes in hepatic biomarker responses were observed, suggesting  
569 that PS-MPs did not induce adverse hepatic stress in rainbow trout; however, Ašmonaitė *et al.* (2018)  
570 theorize that this may be due to the size of the PS-MPs used, as oxidative stress effects have been  
571 observed for smaller polystyrene particles (Jeong *et al.*, 2016; Lei *et al.*, 2018). Ašmonaitė, Sundh, *et*  
572 *al.* (2018) also show that PS-MPs did not affect intestinal health in the same species.

573

574 A review and reinterpretation of the available literature by Koelmans *et al.* (2016) and a modelling  
575 study by Bakir *et al.* (2016), both investigating the relative importance of microplastics as a pathway  
576 for the transfer of adhered contaminants from microplastics to biota, suggest that this is not a  
577 significant route for exposure to adhered contaminants when compared to bioaccumulation from  
578 natural prey and water.

579

#### 580 3.3.4c Metals

581 Heavy metal pollution within the marine environment is increasingly becoming a serious threat to  
582 ecosystems (Naser, 2013) and may therefore become a risk to food security in the near future.  
583 Brennecke *et al.* (2016) examined the adsorption of two heavy metals, copper and zinc, leached from  
584 antifouling paint, to virgin polystyrene beads and aged polyvinylchloride fragments in seawater. Both  
585 heavy metals adsorbed onto the two microplastic types, with concentrations of Cu and Zn increasing  
586 significantly on PVC and PS over the 14-day experiment. Significantly greater adsorption of Cu onto  
587 PVC fragments was observed, with the authors theorizing this was due to the higher surface area and  
588 polarity of PVC.

589

590 The effect of exposure to microplastic (0.26 and 0.69 mg/L), mercury (0.010 and 0.016 mg/L) and  
591 mixtures of the two substances (same concentrations) on the gills and liver of juvenile European bass  
592 (*Dicentrarchus labrax*) over a 96-hour period showed that, while both alone caused oxidative stress in  
593 the gills and liver, the concentration of mercury in both gills and liver was significantly higher in the  
594 presence of microplastics than their absence (Barboza *et al.*, 2018b). This result is therefore indicative  
595 of a synergistic effect of microplastics on the accumulation of mercury within fish tissue. Heavy metals  
596 are proven environmental contaminants, and their interaction with microplastic debris therefore has  
597 potential to significantly alter the toxicity of microplastics within the marine environment.

598

599



600 **3.3.5 Transfer across biological membranes**

601 Microplastic ingestion may not be indicative of negative effects, as microplastics may be egested again  
602 quickly either by post-ingestion rejection or through faeces. However, if microplastics or nanoplastics  
603 are able to transfer into the tissues or circulatory system, by for example transfer across the gut lining  
604 or gill structures, this may lead to greater accumulation and negative effects as the organism may not  
605 be able to remove them. Transfer to tissues, organs and the circulatory system has been seen in  
606 laboratory studies in crabs (Farrell and Nelson, 2013; Watts *et al.*, 2014; Brennecke *et al.*, 2015),  
607 bivalves (Browne *et al.*, 2008; Von Moos, Burkhardt-Holm and Köhler, 2012; Al-Sid-Cheikh *et al.*, 2018)  
608 and fish (Avio, Gorbi and Regoli, 2015; Lu *et al.*, 2016).

609

610

611

612 Uptake of microplastics into the tissues of the blue mussel *Mytilus edulis* can cause changes on the  
613 cellular and tissue level (Von Moos, Burkhardt-Holm and Köhler, 2012). *M. edulis* were exposed to  
614 High-density polyethylene (HDPE) with irregularly shaped particles from >0-80 µm in size at a  
615 concentration of 2.5 g/L for up to 96 hours. Microplastic particles were found on the gills and in the  
616 digestive system, lysosomal system, connective tissue and digestive gland. Effects of microplastic  
617 exposure included granulocytoma formation after 3 hrs, and lysosomal membrane destabilization  
618 after 6 hrs; both effects are associated with the toxicological response of organisms to pollutants  
619 (Moore, 1985; Moore *et al.*, 2008).

620

621 Zebrafish *Danio rerio* exposed to polystyrene microplastic beads (5, and 20 µm) at 20 mg/L for up to  
622 7 days showed microplastic accumulation in the fish gills and gut (5 and 20 µm particles), and in the  
623 liver by 5 µm particles only (Lu *et al.*, 2016). Toxicity testing, exposing *D. rerio* to 5 and 70 µm particles  
624 at 20, 200 and 2000 µg/L for 3 weeks showed that at 2000 µg/L both particle sizes caused inflammation  
625 and lipid accumulation in the liver. Particle size did not cause any observable histopathological  
626 differences in fish tissues.

627

628 Smaller plastic particles are more likely to transfer across biological membranes than particles at the  
629 larger end of the micro-scale, for example through the villi or M-cells of the peyer's patches within the  
630 intestine (Galloway, 2015). However, biologically-facilitated fragmentation of microplastics to  
631 nanometre-sized fragments has been reported to occur through microplastic ingestion by Antarctic  
632 krill (*Euphausia superba*, Dawson *et al.*, 2018). Here, 31.5 µm polyethylene beads (average size, ±7.6  
633 standard deviation, S.D) were ingested by krill, and microplastic fragments identified in krill tissues

634 and faecal pellets were decreased by an average of 78% ( $7.1 \mu\text{m} \pm 6.2 \text{ S.D}$ ) and 81% ( $6.0 \mu\text{m} \pm 5.0 \text{ S.D}$ ).  
635 This is the first time that fragmentation of microplastics to nanoplastics has been reported in  
636 planktonic crustaceans, and could be indicative of a mechanism for microplastic translocation to  
637 tissues in crustaceans where initially they may have been too large.

638

639

## 640 **4. Discussion**

641

### 642 ***4.1 What does the data show?***

643 All of the commercially important organisms studied here, where data was available, were shown to  
644 contain microplastics. The population of animals shown to ingest microplastics varied widely by  
645 species, and when normalized for weight, the number of microplastics ingested per gram wet weight  
646 decreased with increasing trophic level. We conclude that commercially important organisms towards  
647 the base of the food chain (bivalves, crustaceans and small planktivorous fishes) are more likely to be  
648 contaminated with higher concentrations of microplastics, potentially posing a greater risk to their  
649 health and having implications for perceived or actual food safety.

650

651

652 The number of journal articles on the topic of microplastics has increased significantly over recent  
653 years: a search for 'microplastics' in Web of Science shows 473 papers published in 2018, up from 71  
654 published in 2014. However, there are still gaps in our knowledge, particularly pertaining to  
655 commercially important organisms. It is critically important that more targeted research is done to  
656 assess the risk of microplastics to commercially important seafood species; several species, such as  
657 Alaska pollock, Grass carp and Whiteleg shrimp have had no research published on their ingestion of  
658 microplastics within the natural environment. As similar species have shown microplastic ingestion  
659 we can surmise that they will most likely be ingesting plastics, but we have no idea of the scale of this  
660 or effects on these populations. As these three organisms had a combined production of 13.7 million  
661 tonnes of food in 2016, this is a huge gap in this research field and potentially an important risk to  
662 consider for worldwide food security.

663

664 The data presented in Fig. 2 and Table 5 suggests that microplastics do not biomagnify. Comparing  
665 microplastic concentrations within the GIT of different marine organisms to Hg concentrations within  
666 similar organisms (Fig. 2), normalizing by organism weight, shows contrasting trendlines; Hg presence  
667 in organism tissues (ppm) biomagnifies with increasing trophic level whereas the number of

668 microplastics  $\text{g}^{-1}$  w. w. decreases with increasing trophic level. Whilst the data presented here suggests  
669 that microplastics within marine organisms do not biomagnify, this may not be the case for  
670 nanoplastics. These particles are small enough to possibly pass through the gut lining and into the  
671 tissues of organisms (Al-Sid-Cheikh *et al.*, 2018), therefore they may be more likely to bioaccumulate  
672 in animal tissues and may potentially biomagnify through the food chain (although there is no data as  
673 of yet to support this hypothesis).

674

#### 675 **4.2 What factors influence microplastic consumption?**

676 Feeding strategy and environmental prevalence are primary drivers for microplastic consumption.  
677 Generally, lower trophic level organisms appear to ingest more microplastics due to feeding strategy,  
678 as observed by our biomagnification data (Fig. 2 and Table 5). However, there can be huge variations,  
679 for example although they occupy the same ecological niche, 76.6% Japanese anchovy were found  
680 with microplastics within their GIT (Tanaka and Takada, 2016), but only 0.9% Peruvian anchovy (Ory  
681 *et al.*, 2018). This is most likely due to the location where the fish were caught and the sample  
682 digestion methodology utilised. The Japanese anchovy were caught in Tokyo bay, which is in extremely  
683 close proximity to a very large level of anthropomorphic activity, with a drainage basin population of  
684 29 million people, whereas the Peruvian anchovy were caught in further offshore locations in  
685 proximity to smaller population centres, therefore less microplastic pollution may be expected.  
686 Tanaka and Takada (2016) also removed and digested the entire GIT, whereas Ory *et al.* (2018) instead  
687 removed and digested only the gut contents; such differences in methodology may lead to differing  
688 identification efficacies. These differences, in sampling site and methodology, may have resulted in  
689 the large difference in the number of anchovy caught containing microplastics, and care should always  
690 be taken when comparing ingestion studies to identify any sampling bias such as identified here.

691

692 Though trophic transfer does not appear to be an important factor in microplastic consumption, it is  
693 possible that organisms at aquaculture facilities may be exposed to dietary microplastic through  
694 contaminated fishmeal. In 2014, 15.8 million tonnes of fish were reduced to fishmeal (Green, 2016),  
695 for use as a feedstock in the agriculture sector. Miles and Chapman (2006) estimate that in 2010, 56%  
696 of fishmeal was used in the aquaculture sector, 20% in pig feed and 12% in chicken feed. This therefore  
697 represents a novel way for microplastics to be introduced into human food, with potential risks to  
698 many different agriculture industries. Fishmeal is advertised as a nutritious and protein-rich feedstock  
699 (Miles and Chapman, 2006), therefore microplastic contamination through the processing of  
700 contaminated organisms or contamination during fishmeal processing may affect this nutritional value  
701 and have knock-on effects on global agriculture.

702

### 703 **4.3 What are the issues with current studies?**

704 Problems with laboratory analysis of microplastics remain, with several papers likely underestimating  
705 the amount of microplastics found in organic material due to worries about contamination and the  
706 use of filters with pore sizes too large to catch smaller microplastics. Microplastic fibres are commonly  
707 removed from analysis due to concerns about contamination (Rochman *et al.*, 2015; Rummel *et al.*,  
708 2016; Ory *et al.*, 2018). Fibres are one of the most common types of microplastic debris worldwide  
709 (Lusher *et al.*, 2014; Barrows, Cathey and Petersen, 2018), therefore it is critical that research should  
710 utilise methodology to reduce contamination (laminar flow cabinets, non-synthetic laboratory  
711 consumables and clothing etc.), to allow for more robust and realistic analyses of environmental  
712 microplastic concentrations, as concentrations are very likely to be under-represented without the  
713 inclusion of microplastic fibres in results. Smaller microplastics are often missed from analysis due to  
714 equipment constraints, both in collection and analysis. Foekema *et al.*, (2013) and Rummel *et al.*  
715 (2016) only analysed particles larger than 0.2 and 0.5 mm respectively, due to the diameter of the  
716 sieve mesh used. Both Güven *et al.* (2017) and Foekema *et al.* (2013) investigated microplastic in the  
717 GIT of *Trachurus spp.*; Güven *et al.* filtered digested *Trachurus mediterraneus* stomach and intestine  
718 content through a 26 µm mesh, with the resulting percentage of *Trachurus* shown to ingest  
719 microplastics as 68% of the population; Foekema *et al.* filtered digested *Trachurus trachurus* samples  
720 through a 0.2 mm seive and found microplastics in 1% of the population. Güven *et al.* also included  
721 microplastic fibres in their results, while Foekema *et al.* did not. Mean microplastic size identified by  
722 Güven *et al.* was 656.18 µm ± 803.31 SD, median particle size observed by Foekema *et al.* was 800  
723 µm. Extrapolation of observed environmental concentrations of microplastics compared to their size  
724 shows that as mesh size or bead diameter decreases, the number of microplastics found per litre  
725 seawater increases by several orders of magnitude (Lenz, Enders and Nielsen, 2016). This shows a  
726 clear bias of microplastics identified due to methodology, and without standardization it is very  
727 difficult to accurately compare microplastic studies in a rigorous manner.

728

729 Methodological differences are also clear in the preparation of samples for microplastic analysis.  
730 When preparing fish digestive tracts for microplastic analysis, some researchers inspect the entire GIT,  
731 while others opt to inspect only the stomach contents. Both of these methods involve manually  
732 inspecting GIT contents for microplastics once scraped from their respective lining, while another  
733 method more commonly in use in newer studies is to digest the entire GIT, filtering this solution to  
734 remove most of the organic matter and make microplastics more visible and easier to quantify.  
735 Common solvents used to digest the organic material are H<sub>2</sub>O<sub>2</sub>, KOH, HNO<sub>3</sub> and HClO<sub>4</sub> (Foekema *et al.*,

736 2013; Li *et al.*, 2015, 2016; Van Cauwenberghe *et al.*, 2015; Davidson and Dudas, 2016; Jabeen *et al.*,  
737 2017; Phuong *et al.*, 2018; Qu *et al.*, 2018; Waite, Donnelly and Walters, 2018), with combinations of  
738 these solvents sometimes used to increase digestion efficacy (De Witte *et al.*, 2014; Devriese *et al.*,  
739 2015). Some of these treatments have been shown to have a destructive effect on microplastic  
740 particles (Cole *et al.*, 2014; Lusher *et al.*, 2017) therefore care should be taken to ensure microplastics  
741 are not damaged or eliminated due to the digestion protocol utilised. One option is to use digestive  
742 enzymes; for example Cole *et al.* (2014) and Courtene-Jones *et al.* (2017) have utilised enzymatic  
743 digestion with proteinase K and trypsin, respectively, with no observed impacts on microplastics.  
744 However, the methods utilized to effectively measure microplastics whilst avoiding microplastic  
745 alteration or destruction must be balanced against the cost, speed and effort required.

746

#### 747 **4.4 What are the risks of microplastics to fisheries and aquaculture?**

748 Measuring the cost of microplastic pollution to ecosystem services, such as food provisioning through  
749 fisheries and aquaculture, is very challenging, and research into this is still in its infancy. Measuring  
750 the economic cost of marine litter is complex due to the wide range of impacts on the environment,  
751 social and economic sectors (Newman *et al.*, 2015), and it can be expected to be even more  
752 challenging to look at the cost of only microplastics as a proportion of this. The close relationship  
753 between ecosystem services and the marine environment means that adverse environmental effects  
754 from microplastic pollution will have impacts on food provisioning, which could add risk to global food  
755 security. Research has been done to attempt to put a cost to large marine debris. A survey of Scottish  
756 fish vessels reported that 86% of vessels reported reduced catch and 95% reported snagging on their  
757 nets on seafloor debris, with an estimated cost of €11.7-13 million per year; the equivalent of 5% of  
758 the total revenue of affected fisheries (Mouat, Lozano and Bateson, 2010). Estimated values such as  
759 this are not available to look at the cost of microplastic pollution, however the risks of microplastics  
760 identified in this review may all add a cost to fisheries and aquaculture that we cannot currently  
761 quantify. Microplastics carrying pathogenic microbes or invasive species may decimate native  
762 populations of commercially important organisms such as shellfish and crustaceans. Increasing  
763 concentrations of microplastic within the marine environment may put a stress on the energetic  
764 burden of marine organisms if organisms have to spend more energy to consume nutritionally  
765 valuable food this will decrease the energy available for growth and reproduction, and could decrease  
766 mean population size and reproductive output. This would mean that commercially exploited  
767 organisms could take longer to reach a harvestable size, leading to decreased profits in the fisheries  
768 and aquaculture sector, and smaller organism size would lower the nutritional value of seafood.

769

770 Currently, there is no evidence that significant amounts of microplastics can translocate to the tissues  
771 of fish from e.g. the digestive tract or gills, and as most fish are consumed gutted or as processed  
772 pieces (e.g. fillets), there is little evidence that larger fish will transfer microplastics to humans through  
773 diet. However, in the case of smaller fish such as anchovies, as well as shellfish and edible seaweeds,  
774 where the whole organism is often consumed, there is a greater risk of humans consuming  
775 microplastics, with implications for food safety and food security. Studies have suggested that  
776 European consumers may consume 11,000 microplastics per year (Van Cauwenberghe and Janssen,  
777 2014) or 4620 microplastics per year (Catarino *et al.*, 2018) through seafood. Although it has been a  
778 concern that microplastics may leach additives or adsorbed chemical contaminants into humans upon  
779 ingestion, the estimated chemical exposure to humans of persistent organic pollutants and plastic  
780 additives following consumption of seafood is expected to be negligible, at <0.1% of total dietary  
781 exposure (FAO, 2017). Although risks from seafood ingestion are not currently clear, it is possible that  
782 studies such as these will affect the perception of consumers, leading to a change in consumer habits  
783 and diet, before robust studies can be performed to give a clear picture of the effects of plastic  
784 pollution (Koelmans *et al.*, 2017) on food safety and food security. The results of a survey by the  
785 German Environment Agency found that 62% of the population studied felt that they were strongly  
786 (39%) or moderately (23%) contaminated by plastic particles in food and drinking water (SAPEA, 2019);  
787 microplastics research that is reported whilst failing to address human health and food security  
788 concerns may heavily alter public perceptions in similar ways. This may cause a lowering of seafood  
789 value and reduced profits in the seafood and aquaculture sector, potentially impacting public health  
790 in areas which rely heavily on seafood diets. In addition to researching the prevalence and effect of  
791 microplastics that are ingested by organisms in the marine environment, significant numbers of  
792 microplastics may be added to seafood during processing stages and packaging; such concerns should  
793 be researched through analysing microplastic content throughout the production process, to  
794 eliminate any potential areas of contamination that may occur.

795

796 Microplastics are present in commonly consumed aquatic species sourced from both aquaculture and  
797 the marine environment. Processing steps may remove some microplastics, e.g. by removing the GIT  
798 of fish, or washing shellfish and molluscs, however microplastics have been identified in processed  
799 aquatic biota that is being sold for consumption (Karami *et al.*, 2017, 2018). The effect pathways of  
800 microplastics on the health of commercially important marine organisms, and possible risks to human  
801 health from consuming these organisms, must therefore be researched more thoroughly, to evaluate  
802 the potential effect of microplastic pollution to food security.

803

## 804 **5. Conclusion**

805

806 This review examined the presence of microplastics within commercially important marine organisms,  
807 and the risks they may have on organism health. All commercially important organisms analysed in  
808 this review were shown to contain microplastics. Investigation of microplastic concentrations at  
809 different trophic levels suggests that microplastics do not biomagnify, and organisms at lower trophic  
810 levels are at greater risk of microplastic contamination. While organisms higher up the food chain may  
811 not contain as many microplastics per gram body weight, risks are still present from contaminant  
812 transfer and chronic effects, potentially including increased feeding pressure as a result of the higher  
813 risk to lower trophic level organisms. This review highlights that some marine organisms that are  
814 important to global food security are omitted from current microplastics research, and that  
815 microplastics are a risk to the health of marine organisms worldwide. As fisheries and aquaculture are  
816 critical for global food security, this has implications for food security and food safety. Microplastics  
817 present an added risk to an already stressed environment, and further research on the effects of  
818 microplastic pollution is required to be able to perform comprehensive risk assessments on the effect  
819 of microplastics on food security.

820

## 821 **Acknowledgements**

822 This work was supported by the Natural Environment Research Council through the EnvEast Doctoral  
823 Training Partnership (grant number NE/L002582/1). MC and PL are funded by the Waitrose Plastic  
824 Plan Fund (Mussel Power).

825

## 826 **References**

827

- 828 Abbasi, S. *et al.* (2018) 'Microplastics in different tissues of fish and prawn from the Musa Estuary,  
829 Persian Gulf', *Chemosphere*. Pergamon, 205, pp. 80–87. doi: 10.1016/J.CHEMOSPHERE.2018.04.076.
- 830 Akhbarizadeh, R., Moore, F. and Keshavarzi, B. (2019) 'Investigating microplastics bioaccumulation  
831 and biomagnification in seafood from the Persian Gulf: a threat to human health?', *Food Additives &*  
832 *Contaminants: Part A*. Taylor & Francis, pp. 1–13. doi: 10.1080/19440049.2019.1649473.
- 833 Al-Sid-Cheikh, M. *et al.* (2018) 'Uptake, Whole-Body Distribution, and Depuration of Nanoplastics by  
834 the Scallop *Pecten maximus* at Environmentally Realistic Concentrations', *Environmental Science &*  
835 *Technology*. American Chemical Society, 52(24), pp. 14480–14486. doi: 10.1021/acs.est.8b05266.
- 836 Andrady, A. L. (1998) 'Biodegradation of Plastics: Monitoring what Happens.', in Pritchard, G. (ed.)  
837 *Plastics Additives*. Dordrecht: Springer Netherlands, pp. 32–40. doi: 10.1007/978-94-011-5862-6\_5.
- 838 Andrady, A. L. (2011) 'Microplastics in the marine environment', *Marine Pollution Bulletin*.

839 Pergamon, 62(8), pp. 1596–1605. doi: 10.1016/J.MARPOLBUL.2011.05.030.

840 Andradý, A. L. (2017) 'The plastic in microplastics: A review', *Marine Pollution Bulletin*. doi:  
841 10.1016/j.marpolbul.2017.01.082.

842 Ašmonaitė, G., Sundh, H., *et al.* (2018) 'Rainbow trout maintain intestinal transport and barrier  
843 functions following exposure to polystyrene microplastics', *Environmental Science & Technology*.  
844 American Chemical Society. doi: 10.1021/acs.est.8b04848.

845 Ašmonaitė, G., Larsson, K., *et al.* (2018) 'Size matters: ingestion of relatively large microplastics  
846 contaminated with environmental pollutants posed little risk for fish health and fillet quality',  
847 *Environmental Science & Technology*. American Chemical Society. doi: 10.1021/acs.est.8b04849.

848 Avio, C. G., Gorbi, S. and Regoli, F. (2015) 'Experimental development of a new protocol for  
849 extraction and characterization of microplastics in fish tissues: First observations in commercial  
850 species from Adriatic Sea', *Marine Environmental Research*. doi: 10.1016/j.marenvres.2015.06.014.

851 Bakir, A. *et al.* (2016) 'Relative importance of microplastics as a pathway for the transfer of  
852 hydrophobic organic chemicals to marine life', *Environmental Pollution*. Elsevier, 219, pp. 56–65. doi:  
853 10.1016/J.ENVPOL.2016.09.046.

854 Barboza, L. G. A., Dick Vethaak, A., *et al.* (2018a) 'Marine microplastic debris: An emerging issue for  
855 food security, food safety and human health', *Marine Pollution Bulletin*. Pergamon, pp. 336–348. doi:  
856 10.1016/j.marpolbul.2018.05.047.

857 Barboza, L. G. A., Vieira, L. R., *et al.* (2018b) 'Microplastics increase mercury bioconcentration in gills  
858 and bioaccumulation in the liver, and cause oxidative stress and damage in *Dicentrarchus labrax*  
859 juveniles', *Scientific Reports*. doi: 10.1038/s41598-018-34125-z.

860 Barnes, D. K. A. *et al.* (2009) 'Accumulation and fragmentation of plastic debris in global  
861 environments', *Philosophical Transactions of the Royal Society B: Biological Sciences*. doi:  
862 10.1098/rstb.2008.0205.

863 Barrows, A. P. W., Cathey, S. E. and Petersen, C. W. (2018) 'Marine environment microfiber  
864 contamination: Global patterns and the diversity of microparticle origins', *Environmental Pollution*.  
865 Elsevier, 237, pp. 275–284. doi: 10.1016/J.ENVPOL.2018.02.062.

866 Beaumont, N. J. *et al.* (2019) 'Global ecological, social and economic impacts of marine plastic',  
867 *Marine Pollution Bulletin*. Pergamon, 142, pp. 189–195. doi: 10.1016/J.MARPOLBUL.2019.03.022.

868 Bellas, J. *et al.* (2016) 'Ingestion of microplastics by demersal fish from the Spanish Atlantic and  
869 Mediterranean coasts', *Marine Pollution Bulletin*. Pergamon, 109(1), pp. 55–60. doi:  
870 10.1016/J.MARPOLBUL.2016.06.026.

871 Béné, C. *et al.* (2015) 'Feeding 9 billion by 2050 – Putting fish back on the menu', *Food Security*. doi:  
872 10.1007/s12571-015-0427-z.

873 Biginagwa, F. J. *et al.* (2016) 'First evidence of microplastics in the African Great Lakes: Recovery  
874 from Lake Victoria Nile perch and Nile tilapia', *Journal of Great Lakes Research*. Elsevier, 42(1), pp.  
875 146–149. doi: 10.1016/J.JGLR.2015.10.012.



876 Bordbar, L. *et al.* (2018) 'First evidence of ingested plastics by a high commercial shrimp species  
877 (Plesionika narval) in the eastern Mediterranean', *Marine Pollution Bulletin*. Pergamon, 136, pp.  
878 472–476. doi: 10.1016/J.MARPOLBUL.2018.09.030.

879 Borgå, K., Gabrielsen, G. . and Skaare, J. . (2001) 'Biomagnification of organochlorines along a  
880 Barents Sea food chain', *Environmental Pollution*. Elsevier, 113(2), pp. 187–198. doi: 10.1016/S0269-  
881 7491(00)00171-8.

882 Botterell, Z. L. R. *et al.* (2019) 'Bioavailability and effects of microplastics on marine zooplankton:  
883 A review', *Environmental Pollution*. Elsevier, 245, pp. 98–110. doi: 10.1016/J.ENVPOL.2018.10.065.

884 Boucher, J. and Friot, D. (2017) *Primary microplastics in the oceans: A global evaluation of sources*,  
885 *Primary microplastics in the oceans: A global evaluation of sources*. doi: 10.2305/iucn.ch.2017.01.en.

886 Bråte, I. L. N. *et al.* (2016) 'Plastic ingestion by Atlantic cod (*Gadus morhua*) from the Norwegian  
887 coast', *Marine Pollution Bulletin*. Pergamon, 112(1–2), pp. 105–110. doi:  
888 10.1016/J.MARPOLBUL.2016.08.034.

889 Bråte, I. L. N. *et al.* (2018) 'Mytilus spp. as sentinels for monitoring microplastic pollution in  
890 Norwegian coastal waters: A qualitative and quantitative study', *Environmental Pollution*. Elsevier,  
891 243, pp. 383–393. doi: 10.1016/J.ENVPOL.2018.08.077.

892 Brennecke, D. *et al.* (2015) 'Ingested microplastics (>100 µm) are translocated to organs of the  
893 tropical fiddler crab *Uca rapax*', *Marine Pollution Bulletin*. Pergamon, 96(1–2), pp. 491–495. doi:  
894 10.1016/J.MARPOLBUL.2015.05.001.

895 Brennecke, D. *et al.* (2016) 'Microplastics as vector for heavy metal contamination from the marine  
896 environment', *Estuarine, Coastal and Shelf Science*. doi: 10.1016/j.ecss.2015.12.003.

897 Browne, M. A. *et al.* (2008) 'Ingested microscopic plastic translocates to the circulatory system of the  
898 mussel, *Mytilus edulis* (L.)', *Environmental Science and Technology*. doi: 10.1021/es800249a.

899 Catarino, A. I. *et al.* (2018) 'Low levels of microplastics (MP) in wild mussels indicate that MP  
900 ingestion by humans is minimal compared to exposure via household fibres fallout during a meal',  
901 *Environmental Pollution*. Elsevier, 237, pp. 675–684. doi: 10.1016/J.ENVPOL.2018.02.069.

902 Van Cauwenberghe, L. *et al.* (2015) 'Microplastics are taken up by mussels (*Mytilus edulis*) and  
903 lugworms (*Arenicola marina*) living in natural habitats', *Environmental Pollution*. Elsevier, 199, pp.  
904 10–17. doi: 10.1016/J.ENVPOL.2015.01.008.

905 Van Cauwenberghe, L. and Janssen, C. R. (2014) 'Microplastics in bivalves cultured for human  
906 consumption', *Environmental Pollution*. Elsevier, 193, pp. 65–70. doi:  
907 10.1016/J.ENVPOL.2014.06.010.

908 Chakraborty, S. and Newton, A. C. (2011) 'Climate change, plant diseases and food security: An  
909 overview', *Plant Pathology*. doi: 10.1111/j.1365-3059.2010.02411.x.

910 Choy, C. A. and Drazen, J. C. (2013) 'Plastic for dinner? Observations of frequent debris ingestion by  
911 pelagic predatory fishes from the central North Pacific', *Marine Ecology Progress Series*, 485, pp.  
912 155–163. Available at: <https://www.int-res.com/abstracts/meps/v485/p155-163>.

913 Cole, M. *et al.* (2011) 'Microplastics as contaminants in the marine environment: A review', *Marine*  
914 *Pollution Bulletin*. Pergamon, 62(12), pp. 2588–2597. doi: 10.1016/J.MARPOLBUL.2011.09.025.

915 Cole, M. *et al.* (2014) 'Isolation of microplastics in biota-rich seawater samples and marine  
916 organisms', *Scientific Reports*. doi: 10.1038/srep04528.

917 Cole, M. *et al.* (2015) 'The impact of polystyrene microplastics on feeding, function and fecundity in  
918 the marine copepod *Calanus helgolandicus*', *Environmental Science and Technology*. doi:  
919 10.1021/es504525u.

920 Cole, M. *et al.* (2016) 'Microplastics Alter the Properties and Sinking Rates of Zooplankton Faecal  
921 Pellets', *Environmental Science and Technology*. doi: 10.1021/acs.est.5b05905.

922 Courtene-Jones, W. *et al.* (2017) 'Optimisation of enzymatic digestion and validation of specimen  
923 preservation methods for the analysis of ingested microplastics', *Analytical Methods*. doi:  
924 10.1039/c6ay02343f.

925 Critchell, K. and Hoogenboom, M. O. (2018) 'Effects of microplastic exposure on the body condition  
926 and behaviour of planktivorous reef fish (*Acanthochromis polyacanthus*)', *PLoS ONE*. doi:  
927 10.1371/journal.pone.0193308.

928 Davidson, K. and Dudas, S. E. (2016) 'Microplastic Ingestion by Wild and Cultured Manila Clams  
929 (*Venerupis philippinarum*) from Baynes Sound, British Columbia', *Archives of Environmental*  
930 *Contamination and Toxicology*, 71(2), pp. 147–156. doi: 10.1007/s00244-016-0286-4.

931 Dawson, A. L. *et al.* (2018) 'Turning microplastics into nanoplastics through digestive fragmentation  
932 by Antarctic krill', *Nature Communications*. doi: 10.1038/s41467-018-03465-9.

933 Defosse, J. M. and Hawkins, A. J. S. (1997) 'Selective feeding in shellfish: Size-dependent rejection of  
934 large particles within pseudofaeces from *Mytilus edulis*, *Ruditapes philippinarum* and *Tapes*  
935 *decussatus*', *Marine Biology*. doi: 10.1007/s002270050154.

936 Devriese, L. I. *et al.* (2015) 'Microplastic contamination in brown shrimp (*Crangon crangon*, Linnaeus  
937 1758) from coastal waters of the Southern North Sea and Channel area', *Marine Pollution Bulletin*.  
938 doi: 10.1016/j.marpolbul.2015.06.051.

939 De Witte, B. *et al.* (2014) 'Quality assessment of the blue mussel (*Mytilus edulis*): Comparison  
940 between commercial and wild types', *Marine Pollution Bulletin*. Pergamon, 85(1), pp. 146–155. doi:  
941 10.1016/J.MARPOLBUL.2014.06.006.

942 FAO *et al.* (2017) *The State of Food Security and Nutrition in the World 2017. Building Resilience for*  
943 *Peace and Food Security.*, Rome, FAO. doi: 10.1080/15226514.2012.751351.

944 FAO (2018) *The State of World Fisheries and Aquaculture Meeting the Sustainable Development*  
945 *Goals, Food and Agriculture Organization of the United Nations*. doi: 10.1364/OE.17.003331.

946 FAO (2017) *Microplastics in fisheries and aquaculture: status of knowledge on their occurrence and*  
947 *implications for aquatic organisms and food safety, FAO Fisheries and Aquaculture Technical Paper*.  
948 doi: 978-92-5-109882-0.

949 Farrell, P. and Nelson, K. (2013) 'Trophic level transfer of microplastic: *Mytilus edulis* (L.) to *Carcinus*

950 maenas (L.)', *Environmental Pollution*. Elsevier, 177, pp. 1–3. doi: 10.1016/J.ENVPOL.2013.01.046.

951 FDA (2017) *Mercury Levels in Commercial Fish and Shellfish (1990-2012)*. Available at:  
952 <https://www.fda.gov/food/metals/mercury-levels-commercial-fish-and-shellfish-1990-2012>  
953 (Accessed: 14 August 2019).

954 Foekema, E. M. *et al.* (2013) 'Plastic in north sea fish', *Environmental Science and Technology*. doi:  
955 10.1021/es400931b.

956 Frias, J. P. G. L. and Nash, R. (2019) 'Microplastics: Finding a consensus on the definition', *Marine*  
957 *Pollution Bulletin*. Pergamon, 138, pp. 145–147. doi: 10.1016/J.MARPOLBUL.2018.11.022.

958 Galloway, T. S. (2015) 'Micro- and nano-plastics and human health', in *Marine Anthropogenic Litter*.  
959 doi: 10.1007/978-3-319-16510-3\_13.

960 Galloway, T. S., Cole, M. and Lewis, C. (2017) 'Interactions of microplastic debris throughout the  
961 marine ecosystem', *Nature Ecology and Evolution*. doi: 10.1038/s41559-017-0116.

962 GESAMP (2016) *Sources, fate and effects of microplastics in the marine environment: part two of a*  
963 *global assessment*. Edited by C. M. Kershaw, P.J., Rochman. (IMO/FAO/UNESCO-  
964 IOC/UNIDO/WMO/IAEA/UN/ UNEP/UNDP Joint Group of Experts on the Scientific Aspects of Marine  
965 Environmental Protection). Rep. Stud. GESAMP No. 93, 220 p.

966 Geyer, R., Jambeck, J. R. and Law, K. L. (2017) 'Production, use, and fate of all plastics ever made',  
967 *Science Advances*. doi: 10.1126/sciadv.1700782.

968 Goldstein, M. C., Rosenberg, M. and Cheng, L. (2012) 'Increased oceanic microplastic debris  
969 enhances oviposition in an endemic pelagic insect', *Biology Letters*. doi: 10.1098/rsbl.2012.0298.

970 Green, K. (2016) 'Seafish Insight: The global picture – fishmeal production and trends', *FAO SOFIA*  
971 *Report*.

972 Gutow, L. *et al.* (2016) 'Experimental Evaluation of Seaweeds as a Vector for Microplastics into  
973 Marine Food Webs', *Environmental Science and Technology*. doi: 10.1021/acs.est.5b02431.

974 Güven, O. *et al.* (2017) 'Microplastic litter composition of the Turkish territorial waters of the  
975 Mediterranean Sea, and its occurrence in the gastrointestinal tract of fish', *Environmental Pollution*.  
976 Elsevier, 223, pp. 286–294. doi: 10.1016/J.ENVPOL.2017.01.025.

977 Hantoro, I. *et al.* (2019) 'Microplastics in coastal areas and seafood: implications for food safety',  
978 *Food Additives & Contaminants: Part A*. Taylor & Francis, 36(5), pp. 674–711. doi:  
979 10.1080/19440049.2019.1585581.

980 Hardesty, B. D. *et al.* (2017) 'Using Numerical Model Simulations to Improve the Understanding of  
981 Micro-plastic Distribution and Pathways in the Marine Environment ', *Frontiers in Marine Science* ,  
982 p. 30. Available at: <https://www.frontiersin.org/article/10.3389/fmars.2017.00030>.

983 Hartmann, N. B. *et al.* (2019) 'Are We Speaking the Same Language? Recommendations for a  
984 Definition and Categorization Framework for Plastic Debris', *Environmental Science and Technology*.  
985 doi: 10.1021/acs.est.8b05297.

986 Hermesen, E. *et al.* (2017) 'Detection of low numbers of microplastics in North Sea fish using strict  
987 quality assurance criteria', *Marine Pollution Bulletin*. Pergamon, 122(1–2), pp. 253–258. doi:  
988 10.1016/J.MARPOLBUL.2017.06.051.

989 Jabeen, K. *et al.* (2017) 'Microplastics and mesoplastics in fish from coastal and fresh waters of  
990 China', *Environmental Pollution*. Elsevier, 221, pp. 141–149. doi: 10.1016/J.ENVPOL.2016.11.055.

991 Jeong, C. B. *et al.* (2016) 'Microplastic Size-Dependent Toxicity, Oxidative Stress Induction, and p-JNK  
992 and p-p38 Activation in the Monogonont Rotifer (*Brachionus koreanus*)', *Environmental Science and  
993 Technology*. doi: 10.1021/acs.est.6b01441.

994 Jiang, P. *et al.* (2018) 'Microplastic-associated bacterial assemblages in the intertidal zone of the  
995 Yangtze Estuary', *Science of the Total Environment*. doi: 10.1016/j.scitotenv.2017.12.105.

996 Kaiser, D., Kowalski, N. and Waniek, J. J. (2017) 'Effects of biofouling on the sinking behavior of  
997 microplastics', *Environmental Research Letters*. doi: 10.1088/1748-9326/aa8e8b.

998 Karami, A. *et al.* (2017) 'Microplastics in eviscerated flesh and excised organs of dried fish', *Scientific  
999 Reports*. doi: 10.1002/advs.201700259.

1000 Karami, A. *et al.* (2018) 'Microplastic and mesoplastic contamination in canned sardines and sprats',  
1001 *Science of The Total Environment*. Elsevier, 612, pp. 1380–1386. doi:  
1002 10.1016/J.SCITOTENV.2017.09.005.

1003 Karbalaei, S. *et al.* (2019) 'Abundance and characteristics of microplastics in commercial marine fish  
1004 from Malaysia', *Marine Pollution Bulletin*. Pergamon, 148, pp. 5–15. doi:  
1005 10.1016/J.MARPOLBUL.2019.07.072.

1006 Kirstein, I. V. *et al.* (2016) 'Dangerous hitchhikers? Evidence for potentially pathogenic *Vibrio* spp. on  
1007 microplastic particles', *Marine Environmental Research*. Elsevier, 120, pp. 1–8. doi:  
1008 10.1016/J.MARENRES.2016.07.004.

1009 Koelmans, A. A. *et al.* (2016) 'Microplastic as a Vector for Chemicals in the Aquatic Environment:  
1010 Critical Review and Model-Supported Reinterpretation of Empirical Studies', *Environmental Science  
1011 & Technology*. American Chemical Society, 50(7), pp. 3315–3326. doi: 10.1021/acs.est.5b06069.

1012 Koelmans, A. A. *et al.* (2017) 'Risks of Plastic Debris: Unravelling Fact, Opinion, Perception, and  
1013 Belief', *Environmental Science and Technology*. doi: 10.1021/acs.est.7b02219.

1014 Koelmans, A. A., Besseling, E. and Foekema, E. M. (2014) 'Leaching of plastic additives to marine  
1015 organisms', *Environmental Pollution*. Elsevier, 187, pp. 49–54. doi: 10.1016/J.ENVPOL.2013.12.013.

1016 Kooi, M. *et al.* (2017) 'Ups and Downs in the Ocean: Effects of Biofouling on Vertical Transport of  
1017 Microplastics', *Environmental Science and Technology*. doi: 10.1021/acs.est.6b04702.

1018 Lavoie, R. A. *et al.* (2013) 'Biomagnification of Mercury in Aquatic Food Webs: A Worldwide Meta-  
1019 Analysis', *Environmental Science & Technology*. American Chemical Society, 47(23), pp. 13385–  
1020 13394. doi: 10.1021/es403103t.

1021 Law, K. L. and Thompson, R. C. (2014) 'Microplastics in the seas', *Science*. doi:  
1022 10.1126/science.1254065.

- 1023 Lei, L. *et al.* (2018) 'Polystyrene (nano)microplastics cause size-dependent neurotoxicity, oxidative  
1024 damage and other adverse effects in *Caenorhabditis elegans*', *Environmental Science: Nano*. doi:  
1025 10.1039/c8en00412a.
- 1026 Lenz, R., Enders, K. and Nielsen, T. G. (2016) 'Microplastic exposure studies should be  
1027 environmentally realistic', *Proceedings of the National Academy of Sciences*. doi:  
1028 10.1073/pnas.1606615113.
- 1029 Li, J. *et al.* (2015) 'Microplastics in commercial bivalves from China', *Environmental Pollution*. doi:  
1030 10.1016/j.envpol.2015.09.018.
- 1031 Li, J. *et al.* (2016) 'Microplastics in mussels along the coastal waters of China', *Environmental*  
1032 *Pollution*. Elsevier, 214, pp. 177–184. doi: 10.1016/J.ENVPOL.2016.04.012.
- 1033 Li, J. *et al.* (2019) 'Using mussel as a global bioindicator of coastal microplastic pollution',  
1034 *Environmental Pollution*. Elsevier, 244, pp. 522–533. doi: 10.1016/J.ENVPOL.2018.10.032.
- 1035 Liboiron, M. *et al.* (2016) 'Low plastic ingestion rate in Atlantic cod (*Gadus morhua*) from  
1036 Newfoundland destined for human consumption collected through citizen science methods', *Marine*  
1037 *Pollution Bulletin*. Pergamon, 113(1–2), pp. 428–437. doi: 10.1016/J.MARPOLBUL.2016.10.043.
- 1038 Lim, P. T. *et al.* (2005) 'First report of *Alexandrium taylori* and *Alexandrium peruvianum*  
1039 (Dinophyceae) in Malaysia waters', *Harmful Algae*. Elsevier, 4(2), pp. 391–400. doi:  
1040 10.1016/J.HAL.2004.07.001.
- 1041 Lu, Y. *et al.* (2016) 'Uptake and Accumulation of Polystyrene Microplastics in Zebrafish (*Danio rerio*)  
1042 and Toxic Effects in Liver', *Environmental Science and Technology*. doi: 10.1021/acs.est.6b00183.
- 1043 Lusher, A. L. *et al.* (2014) 'Microplastic pollution in the Northeast Atlantic Ocean: Validated and  
1044 opportunistic sampling', *Marine Pollution Bulletin*. doi: 10.1016/j.marpolbul.2014.08.023.
- 1045 Lusher, A. L. *et al.* (2017) 'Sampling, isolating and identifying microplastics ingested by fish and  
1046 invertebrates', *Analytical Methods*. doi: 10.1039/c6ay02415g.
- 1047 Lusher, A. L., McHugh, M. and Thompson, R. C. (2013) 'Occurrence of microplastics in the  
1048 gastrointestinal tract of pelagic and demersal fish from the English Channel', *Marine Pollution*  
1049 *Bulletin*. doi: 10.1016/j.marpolbul.2012.11.028.
- 1050 Mardones, J. I. *et al.* (2015) 'Fish gill damage by the dinoflagellate *Alexandrium catenella* from  
1051 Chilean fjords: Synergistic action of ROS and PUFA', *Harmful Algae*. Elsevier, 49, pp. 40–49. doi:  
1052 10.1016/J.HAL.2015.09.001.
- 1053 Markic, A. *et al.* (2018) 'Double trouble in the South Pacific subtropical gyre: Increased plastic  
1054 ingestion by fish in the oceanic accumulation zone', *Marine Pollution Bulletin*. Pergamon, 136, pp.  
1055 547–564. doi: 10.1016/J.MARPOLBUL.2018.09.031.
- 1056 Masó, M. *et al.* (2003) 'Drifting plastic debris as a potential vector for dispersing Harmful Algal Bloom  
1057 (HAB) species', *Scientia Marina*. doi: 10.3989/scimar.2003.67n1107.
- 1058 Miles, R. and Chapman, F. (2006) *The Benefits of Fish Meal in Aquaculture Diets*, FA122.

1059 Moore, M. N. (1985) 'Cellular responses to pollutants', *Marine Pollution Bulletin*. Pergamon, 16(4),  
1060 pp. 134–139. doi: 10.1016/0025-326X(85)90003-7.

1061 Moore, M. N. *et al.* (2008) 'Chapter Thirty-Three Lysosomes and Autophagy in Aquatic Animals',  
1062 *Methods in Enzymology*. Academic Press, 451, pp. 581–620. doi: 10.1016/S0076-6879(08)03233-3.

1063 Von Moos, N., Burkhardt-Holm, P. and Köhler, A. (2012) 'Uptake and effects of microplastics on cells  
1064 and tissue of the blue mussel *Mytilus edulis* L. after an experimental exposure', *Environmental  
1065 Science and Technology*. doi: 10.1021/es302332w.

1066 Mouat, J., Lozano, R. L. and Bateson, H. (2010) *Economic Impacts of Marine Litter, Kommunenes  
1067 Internasjonale Miljøorganisasjon (KIMO)*.

1068 Murphy, F. *et al.* (2017) 'The uptake of macroplastic & microplastic by demersal & pelagic fish in the  
1069 Northeast Atlantic around Scotland', *Marine Pollution Bulletin*. doi:  
1070 10.1016/j.marpolbul.2017.06.073.

1071 Murray, F. and Cowie, P. R. (2011) 'Plastic contamination in the decapod crustacean *Nephrops  
1072 norvegicus* (Linnaeus, 1758)', *Marine Pollution Bulletin*. doi: 10.1016/j.marpolbul.2011.03.032.

1073 Naser, H. A. (2013) 'Assessment and management of heavy metal pollution in the marine  
1074 environment of the Arabian Gulf: A review', *Marine Pollution Bulletin*. doi:  
1075 10.1016/j.marpolbul.2013.04.030.

1076 Nelms, S. E. *et al.* (2018) 'Investigating microplastic trophic transfer in marine top predators',  
1077 *Environmental Pollution*. Elsevier, 238, pp. 999–1007. doi: 10.1016/J.ENVPOL.2018.02.016.

1078 Neves, D. *et al.* (2015) 'Ingestion of microplastics by commercial fish off the Portuguese coast',  
1079 *Marine Pollution Bulletin*. Pergamon, 101(1), pp. 119–126. doi: 10.1016/j.marpolbul.2015.11.008.

1080 Newell, R. and Jordan, S. (1983) 'Preferential ingestion of organic material by the American oyster  
1081 *Crassostrea virginica*', *Marine Ecology Progress Series*. doi: 10.3354/meps013047.

1082 Newman, S. *et al.* (2015) 'The economics of marine litter', in *Marine Anthropogenic Litter*. doi:  
1083 10.1007/978-3-319-16510-3\_14.

1084 Oberbeckmann, S., Löder, M. G. J. and Labrenz, M. (2015) 'Marine microplastic-associated biofilms -  
1085 A review', *Environmental Chemistry*. doi: 10.1071/EN15069.

1086 Ogonowski, M. *et al.* (2017) 'Ingested microplastic is not correlated to HOC concentrations in Baltic  
1087 Sea herring'.

1088 Ory, N. *et al.* (2018) 'Low prevalence of microplastic contamination in planktivorous fish species from  
1089 the southeast Pacific Ocean', *Marine Pollution Bulletin*. doi: 10.1016/j.marpolbul.2017.12.016.

1090 Ory, N. C. *et al.* (2017) 'Amberstripe scad *Decapterus muroadsi* (Carangidae) fish ingest blue  
1091 microplastics resembling their copepod prey along the coast of Rapa Nui (Easter Island) in the South  
1092 Pacific subtropical gyre', *Science of the Total Environment*. doi: 10.1016/j.scitotenv.2017.01.175.

1093 Ory, N. C. *et al.* (2018) 'Capture, swallowing, and egestion of microplastics by a planktivorous  
1094 juvenile fish', *Environmental Pollution*. Elsevier, 240, pp. 566–573. doi:

1095 10.1016/J.ENVPOL.2018.04.093.

1096 Peeken, I. *et al.* (2018) 'Arctic sea ice is an important temporal sink and means of transport for  
1097 microplastic', *Nature Communications*, 9(1), p. 1505. doi: 10.1038/s41467-018-03825-5.

1098 Phuong, N. N. *et al.* (2018) 'Factors influencing the microplastic contamination of bivalves from the  
1099 French Atlantic coast: Location, season and/or mode of life?', *Marine Pollution Bulletin*. Pergamon,  
1100 129(2), pp. 664–674. doi: 10.1016/J.MARPOLBUL.2017.10.054.

1101 Plessi, M., Bertelli, D. and Monzani, A. (2001) 'Mercury and Selenium Content in Selected Seafood',  
1102 *Journal of Food Composition and Analysis*. Academic Press, 14(5), pp. 461–467. doi:  
1103 10.1006/JFCA.2001.1003.

1104 Qu, X. *et al.* (2018) 'Assessing the relationship between the abundance and properties of  
1105 microplastics in water and in mussels', *Science of The Total Environment*. Elsevier, 621, pp. 679–686.  
1106 doi: 10.1016/J.SCITOTENV.2017.11.284.

1107 Ribeiro, F. *et al.* (2017) 'Microplastics effects in *Scrobicularia plana*', *Marine Pollution Bulletin*.  
1108 Pergamon, 122(1–2), pp. 379–391. doi: 10.1016/J.MARPOLBUL.2017.06.078.

1109 Rochman, C. M. *et al.* (2015) 'Anthropogenic debris in seafood: Plastic debris and fibers from textiles  
1110 in fish and bivalves sold for human consumption', *Scientific Reports*. doi: 10.1038/srep14340.

1111 Rummel, C. D. *et al.* (2016) 'Plastic ingestion by pelagic and demersal fish from the North Sea and  
1112 Baltic Sea', *Marine Pollution Bulletin*. Pergamon, 102(1), pp. 134–141. doi:  
1113 10.1016/J.MARPOLBUL.2015.11.043.

1114 Scheld, A. M., Bilkovic, D. M. and Havens, K. J. (2016) 'The Dilemma of Derelict Gear', *Scientific*  
1115 *Reports*. doi: 10.1038/srep19671.

1116 Science Advice for Policy by European Academies (SAPEA) (2019) *A Scientific Perspective on*  
1117 *Microplastics in Nature and Society | SAPEA, Evidence Review Report*. doi: 10.26356/microplastics.

1118 Steer, M. *et al.* (2017) 'Microplastic ingestion in fish larvae in the western English Channel',  
1119 *Environmental Pollution*. doi: 10.1016/j.envpol.2017.03.062.

1120 Sundbæk, K. B. *et al.* (2018) 'Sorption of fluorescent polystyrene microplastic particles to edible  
1121 seaweed *Fucus vesiculosus*', *Journal of Applied Phycology*. doi: 10.1007/s10811-018-1472-8.

1122 Sussarellu, R. *et al.* (2016) 'Oyster reproduction is affected by exposure to polystyrene microplastics',  
1123 *Proceedings of the National Academy of Sciences*. doi: 10.1073/pnas.1519019113.

1124 Tanaka, K. and Takada, H. (2016) 'Microplastic fragments and microbeads in digestive tracts of  
1125 planktivorous fish from urban coastal waters', *Scientific Reports*. doi: 10.1038/srep34351.

1126 Teuten, E. L. *et al.* (2007) 'Potential for plastics to transport hydrophobic contaminants',  
1127 *Environmental Science and Technology*. doi: 10.1021/es071737s.

1128 Viršek, M. K. *et al.* (2017) 'Microplastics as a vector for the transport of the bacterial fish pathogen  
1129 species *Aeromonas salmonicida*', *Marine Pollution Bulletin*. Pergamon, 125(1–2), pp. 301–309. doi:  
1130 10.1016/J.MARPOLBUL.2017.08.024.

1131 Waite, H. R., Donnelly, M. J. and Walters, L. J. (2018) 'Quantity and types of microplastics in the  
1132 organic tissues of the eastern oyster *Crassostrea virginica* and Atlantic mud crab *Panopeus herbstii*  
1133 from a Florida estuary', *Marine Pollution Bulletin*. Pergamon, 129(1), pp. 179–185. doi:  
1134 10.1016/J.MARPOLBUL.2018.02.026.

1135 Ward, J. E. *et al.* (2019) 'Selective Ingestion and Egestion of Plastic Particles by the Blue Mussel  
1136 (*Mytilus edulis*) and Eastern Oyster (*Crassostrea virginica*): Implications for Using Bivalves as  
1137 Bioindicators of Microplastic Pollution', *Environmental Science & Technology*. American Chemical  
1138 Society, 53(15), pp. 8776–8784. doi: 10.1021/acs.est.9b02073.

1139 Watts, A. J. R. *et al.* (2014) 'Uptake and retention of microplastics by the shore crab *Carcinus*  
1140 *maenas*', *Environmental Science and Technology*. doi: 10.1021/es501090e.

1141 Watts, A. J. R. *et al.* (2015) 'Ingestion of Plastic Microfibers by the Crab *Carcinus maenas* and Its  
1142 Effect on Food Consumption and Energy Balance', *Environmental Science and Technology*. doi:  
1143 10.1021/acs.est.5b04026.

1144 Wegner, A. *et al.* (2012) 'Effects of nanopolystyrene on the feeding behavior of the blue mussel  
1145 (*Mytilus edulis* L.)', *Environmental Toxicology and Chemistry*. doi: 10.1002/etc.1984.

1146 Welden, N. A. C. and Cowie, P. R. (2016a) 'Environment and gut morphology influence microplastic  
1147 retention in langoustine, *Nephrops norvegicus*', *Environmental Pollution*. Elsevier, 214, pp. 859–865.  
1148 doi: 10.1016/J.ENVPOL.2016.03.067.

1149 Welden, N. A. C. and Cowie, P. R. (2016b) 'Environment and gut morphology influence microplastic  
1150 retention in langoustine, *Nephrops norvegicus*', *Environmental Pollution*. Elsevier, 214, pp. 859–865.  
1151 doi: 10.1016/J.ENVPOL.2016.03.067.

1152 Wójcik-Fudalewska, D., Normant-Saremba, M. and Anastácio, P. (2016) 'Occurrence of plastic debris  
1153 in the stomach of the invasive crab *Eriocheir sinensis*', *Marine Pollution Bulletin*. Pergamon, 113(1–  
1154 2), pp. 306–311. doi: 10.1016/J.MARPOLBUL.2016.09.059.

1155 Wollenberg, E. *et al.* (2016) 'Reducing risks to food security from climate change', *Global Food*  
1156 *Security*. doi: 10.1016/j.gfs.2016.06.002.

1157 Woods, M. N. *et al.* (2018) 'Microplastic fiber uptake, ingestion, and egestion rates in the blue  
1158 mussel (*Mytilus edulis*)', *Marine Pollution Bulletin*. Pergamon, 137, pp. 638–645. doi:  
1159 10.1016/J.MARPOLBUL.2018.10.061.

1160 Xu, X.-Y. *et al.* (2017) 'Microplastic ingestion reduces energy intake in the clam *Atactodea striata*',  
1161 *Marine Pollution Bulletin*. Pergamon, 124(2), pp. 798–802. doi: 10.1016/J.MARPOLBUL.2016.12.027.

1162 Yuan, W. *et al.* (2019) 'Microplastic abundance, distribution and composition in water, sediments,  
1163 and wild fish from Poyang Lake, China', *Ecotoxicology and Environmental Safety*. Academic Press,  
1164 170, pp. 180–187. doi: 10.1016/J.ECOENV.2018.11.126.

1165 Zettler, E. R., Mincer, T. J. and Amaral-Zettler, L. A. (2013) 'Life in the "plastisphere": Microbial  
1166 communities on plastic marine debris', *Environmental Science and Technology*. doi:  
1167 10.1021/es401288x.



