

2020-02

# Quantification and characterisation of microplastics ingested by selected juvenile fish species associated with mangroves in KwaZulu-Natal, South Africa

Naidoo, T

<http://hdl.handle.net/10026.1/17613>

---

10.1016/j.envpol.2019.113635

Environmental Pollution

Elsevier

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

1 Disclaimer

2 'This is a copy of the accepted paper as submitted for publication. Readers are advised to refer to the final version of the paper which can be  
3 found at

4 *Environmental Pollution*, 257, 113635. doi:[10.1016/j.envpol.2019.113635](https://doi.org/10.1016/j.envpol.2019.113635)

5

6 Quantification and characterisation of microplastics ingested by selected juvenile fish species associated with mangroves in KwaZulu-Natal, South  
7 Africa

8 Trishan Naidoo<sup>a,\*</sup>, Sershen<sup>b</sup>, Richard C. Thompson<sup>c</sup>, Anusha Rajkaran<sup>a</sup>

9 <sup>a</sup> Department for Biodiversity & Conservation Biology, University of the Western Cape, Private Bag X17, Bellville, 7535, South Africa

10 <sup>b</sup> School of Life Sciences, University of KwaZulu-Natal, Westville Campus, Private Bag X54001, Durban, 4000, South Africa

11 <sup>c</sup> Marine Biology and Ecology Research Centre, School of Biological and Marine Sciences, University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK

12

13 \* Corresponding author

14 Email address: trishan.naidoo2@gmail.com

15

16 Abstract

17 Though the number studies on microplastic ingestion by fish is growing, data on fish species characteristic of the South African coastline are  
18 scarce. This study quantified and characterised (physically and chemically) microplastics ingested by four species of juvenile fish (viz.  
19 *Oreochromis mossambicus* [Peters, 1852], *Terapon jarbua* [Forsskål, 1775], *Ambassis dussumieri* [Cuvier, 1828] and *Mugil* sp.), within four  
20 mangroves along the east coast of South Africa. Microplastics were isolated from whole fish using a proteinase K digestion method, and then  
21 quantified and characterised in terms of shape, chemical nature (plastic type), colour and length. Fibres (68%) and fragments (21%) were the  
22 dominant shapes found. Of the 174 fish sampled, 52% contained microplastic particles, with  $0.79 \pm 1.00$  particles per fish. The average number of  
23 particles per fish did not differ significantly across species within sites and across sites but was higher than in juvenile fish of other species  
24 sampled in oceanic habitats. The main plastic types collected using 10 µm filters and identified with Fourier Transform Infrared Spectroscopy  
25 (FTIR), were rayon (70.4%), polyester (10.4%), nylon (5.2%) and polyvinylchloride (3.0%). Particle length ranged from 0.1 - 4.8 mm, averaging  
26  $0.89 \pm 0.77$  mm, but irrespective of length, particles were mostly blue in colour. This study provides evidence that juvenile fish inhabiting  
27 mangroves are consuming significant quantities of microplastics. Importantly, it should be noted that rayon, though the most abundant plastic type

28 found, is a semi-synthetic, fibre made from regenerated cellulose that is commonly reported in studies of this nature. The habitats studied serve as  
29 nurseries for numerous fish species; however, more detailed studies are needed to assess whether microplastic ingestion could compromise the  
30 health of these fish or whether these effects are dependent on species, feeding habit and/or plastic type.

31 Capsule: South African juvenile fish associated with mangroves had  $0.79 \pm 1.00$  microplastic particles per fish. Rayon (70.4%), polyester (10.4%),  
32 nylon (5.2%) and polyvinylchloride (3.0%) were the main plastic types ingested.

33 Keywords: Estuary; ingestion; marine; plastic; rayon; debris

## 34 Introduction

35 Reports of microplastic ( $\leq 5$  mm in length) ingestion by aquatic organisms date back to the beginning of mainstream plastic production (Carpenter  
36 et al., 1972) and over the last decade have been recorded in a wide range of fauna Wright et al. (2013), including fish (Jovanović, 2017).  
37 Microplastic ingestion in fish in particular, has been reported for a wide variety of water bodies, including inland lakes (e.g. Biginagwa et al.,  
38 2016), ocean channels (Lusher et al., 2013), deep ocean (e.g. Anastasopoulou et al., 2013) and more recently estuarine habitats (Vendel et al.,  
39 2017). However, these reports represent a geographic bias in that they emanate largely from the developed world; data for African marine  
40 environments and species that typify the continent's coastal habitats are particularly scarce (but see Naidoo et al., 2016).

41 The interest in microplastic ingestion by fish is based on the negative health consequences, which have been reported from laboratory studies for a  
42 range of organisms and particle sizes. Whilst particles sizes used in experiments can differ greatly, from a lower limit of  $3 \mu\text{m}$  upwards, variable  
43 combinations of the following have been observed: decreased growth (Naidoo and Glassom, 2019), decreased feeding and/or weight loss  
44 (Besseling et al., 2013; de Sá et al., 2015), transfer to organs (Browne et al., 2008), inflammation (Wright et al., 2013), liver toxicity and pathology  
45 (Rochman et al., 2013b), endocrine disruption (Rochman et al., 2014) and decreased reproductive output (Sussarellu et al., 2016). Plastic particles  
46 can also act as vectors that transport persistent organic pollutants (Rios et al., 2007). This has been suggested to contribute to the negative health  
47 effects mentioned above (but see Bakir et al., 2016). Some authors do dispute their potential as the main vectors for these pollutants (e.g.  
48 Beckingham and Ghosh, 2017). However, in light of the possible negative health effects, microplastic ingestion by fish has the potential to affect  
49 fisheries by affecting fish growth (Markic and Nicol, 2014; Naidoo and Glassom, 2019). This could be especially problematic in developing  
50 countries that are heavily reliant on fish stocks (Lamberth and Turpie, 2003).

51 Whilst a number of studies have quantified and even characterised microplastic particles ingested by fish based on physical characteristics (e.g.  
52 Boerger et al., 2010; Davison and Asch, 2011; Kripa et al., 2014; Ferreira et al., 2016; Mizraji et al., 2017) reports that detail the chemical  
53 composition of particles ingested by fish are scarcer. The impact of these particles on organisms could be influenced by both the physical and the  
54 chemical nature of the plastic they are composed of; polyvinylchloride and polystyrene for example, can have carcinogenic properties, while other  
55 types may be less harmful (Rochman et al., 2013a). Fish feeding strategies may also influence the typology and abundance of microplastics

56 ingested by different species (Mizraji et al., 2017). Additionally, particle abundance and type can differ across habitats owing to density  
57 differences (Moore et al., 2005) and feeding strategies can differ across species (Ferreira et al., 2019) which suggests that different species could  
58 be exposed to different suites of plastics. It is therefore important to both quantify and characterise the different types of plastics ingested by fish in  
59 a wider range of natural and managed systems, and species. In this regard, only few studies have investigated plastic ingestion by fish in estuarine  
60 and specifically mangrove habitats within estuaries, compared with other marine environments (Vendel et al., 2017). Such information is  
61 important, since these systems are a crucial pathway through which microplastics reach the ocean (Possatto et al., 2015). These systems offer  
62 important ecosystem services by providing refugia to fish and thus food provision to humans (van Niekerk and Turpie, 2012). For example, the  
63 economic value associated with South African estuarine fisheries was estimated to be 1.2 billion rand per annum (van Niekerk and Turpie, 2012).  
64 These mangroves are also nursery sites for many species of juvenile fish, as they offer food and cover from predators (Laegdsgaard and Johnson,  
65 2001). There is a high probability that juvenile estuarine fish will interact with microplastics as these habitats are usually in close proximity to  
66 contamination sources and these particles are likely to be retained in this environment (Naidoo et al., 2015, 2016). Mangrove pneumatophores  
67 have also been suggested to increase the retention of plastics in estuarine systems (Martin et al., 2019). At the time of this study, there were no  
68 published reports on the abundance, type, length, colour and chemical composition of microplastics ingested by juvenile fish species associated  
69 with South African mangrove systems. This motivated the present study which quantified and characterised (physically and chemically)  
70 microplastics ingested by four species of juvenile fish (viz. Tilapia - *Oreochromis mossambicus* [Peters, 1852], Thornfish - *Terapon*  
71 *jarbua* [Forsskål, 1775], Glassfish - *Ambassis dussumieri* [Cuvier, 1828] and Mullet - *Mugil* sp.) in four mangroves in KwaZulu-Natal (KZN),  
72 South Africa. Many of the predominantly open estuaries on the east coast of South Africa support dense mangrove strands (Ward and Steinke,  
73 1982) and we compared three of them from an urban setting and one located in a semi-rural Ramsar site. The shape, chemical composition, colour  
74 and length of microplastics were recorded and compared among sites and species. Knowledge on systems and fish species can inform future  
75 pollution management strategies, particularly for mangroves which are subject to numerous anthropogenic pressures globally (Friess et al., 2019).  
76 Our focus on juvenile fish is based on the fact that many fish populations are rapidly depleting owing to overfishing, climate change, habitat  
77 destruction and other anthropogenic impacts; a healthy juvenile population is essential for sustaining fish stocks (Wallace et al., 1984; Harris and  
78 Cyrus, 1999).

## 79 Methods

### 80 Fish collection

81 In total, 174 juvenile fish, between the fry and fingerling stages of development, of four species (*O. mossambicus*, *T. jarbua*, *A. dussumieri* and  
82 *Mugil* sp.) were collected (if present) in the St. Lucia (28° 23'S, 32° 25'E), Umgeni (29° 48'S, 30° 02'E), Durban Harbour (29° 52'S, 31° 04'E) and  
83 Isipingo (30° 00'S, 30° 57'E) estuaries, in KZN, South Africa. Detailed information on these systems can be found in the National Biodiversity  
84 Assessment of these estuaries (see van Niekerk et al., 2019). St. Lucia has the largest mangrove area (288 ha), followed by Durban Harbour (13.4  
85 ha), Umgeni (27 ha) and the Isipingo (3.8 ha) (van Niekerk et al., 2019). While the St. Lucia and Isipingo estuarine systems are freshwater

86 dominated, the Durban Harbour and the Umgeni are more marine dominated (van Niekerk et al., 2019). Fish collection was done by actively  
87 pursuing fish, using dip nets on a single day at each estuary. Collections were done within January 2019, which corresponded to the wet season.  
88 Owing to species salinity preferences, *O. mossambicus* was present in the St. Lucia and Isipingo estuaries only, where salinity ranges from fresh to  
89 brackish conditions (Whitfield et al., 1981). *T. jarbua* and *A. dussumieri*, which are typical in marine conditions (Whitfield et al., 1981), occurred  
90 in the Umgeni and Durban Harbour estuaries only. Fish were placed in 99% anhydrous alcohol for euthanasia, a method approved by the Ethics  
91 Committee at the University of KwaZulu-Natal [AREC/011/016D]). Fish were transported to the laboratory in individual hermetically sealed  
92 plastic bags. Thereafter, the surface of each fish was dried with paper towel and its mass recorded to three decimal places. Fish were then placed  
93 individually in a new set of hermetically sealed polyethylene bags filled with commercial salt and stored at 4°C until further processing. Metal  
94 forceps (rinsed with deionized water before use) were used to handle the fish samples at all stages.

#### 95 Sample preparation

96 All glassware and handling equipment were washed and rinsed with deionized water before use. As suggested by Woodall et al. (2015), samples  
97 were filtered under vacuum in a clean room dedicated to microplastic research. The air ducts in this room are blocked off from the main air  
98 conditioning and white cotton lab coats are worn within. Glassware used for filtration were covered in aluminum foil before use and after the  
99 filtration of each sample. Petri dishes and metal forceps were pre-checked for plastic contamination under a dissecting microscope. Fish were  
100 removed from salt, rinsed and rehydrated in 18.2 MΩ Milli-Q water. Each fish was then examined under a dissecting microscope for surface  
101 contamination, possibly originating from salt during storage or prior handling. Any surface contamination found was removed and fish were  
102 transferred to 30 mL pre-cleaned glass bottles, which were then sealed with end caps to serve as digestion vials. All steps were carried out in a  
103 laminar flow hood to prevent contamination.

#### 104 Fish digestion and microplastic isolation

105 The direct inspection of fish guts for microplastics is challenging, particularly in juvenile fish, as the gut content and oil globules cloud  
106 observations. Therefore, many studies use nitric acid to digest fish tissue, aiding observation (Lusher et al., 2017). However, plastics such as nylon  
107 (polyamide), can be degraded by the acid, making this method more appropriate for large scale monitoring on a limited budget (Naidoo et al.,  
108 2017). Since we wanted to account for all plastics, digestions were carried out on whole fish using Proteinase K (3.0 - 15.0 unit/mg solid  
109 lyophilized powder, from *Engyodontium album*). The protease, filter papers and chemicals were all obtained from Sigma-Aldrich. The digestion  
110 method was adapted from Cole et al. (2014) and has been shown by Karlsson et al. (2017) to have a high recovery (97%) of plastics, with minimal  
111 alteration of their physical and chemical nature for identification.

112 One mg/mL of proteinase K in Tris EDTA Buffer (pH 8.0, prepared with Milli-Q water) was used for sample digestions. One mL of the solution  
113 was pipetted into each digestion vial and shaken for 1 min on a Hatti roto mixer to facilitate fish tissue digestion. All pipette tips and the tip of  
114 the pipette used for all digestion steps were pre-rinsed with deionised water before use. Three 1L replicates of Milli-Q water was filtered to serve

115 as controls and these revealed no contamination of fibres. Digestion vials were incubated in an oven at 39°C overnight. Upon recovery, samples  
116 were shaken once more and visually observed. Vials containing undigested fish tissue were placed back into the oven for further digestion. Fish  
117 bones, scales, otoliths and eye balls did not digest.

118 Once all digestible tissue was no longer visible, samples were filtered, under vacuum, through a 10 µm Whatman cyclopore polycarbonate filter  
119 (47 mm diameter). After the sample was poured into the sample holder, the sides of the digestion vial and the sample holder were rinsed three  
120 times, with 2 mL and 3 mL of Milli-Q water, respectively. The filter was then removed and placed within a closed Petri dish, which was then  
121 covered in foil and placed in an oven at 39°C overnight. The sample holder and filter holder were rinsed with Milli-Q water between samples.

## 122 Quantification and characterization of microplastics

123 After sample filtration, the filters were observed under a dissecting microscope (Microtec, HM-3), whilst still enclosed within Petri dishes.  
124 Particles that resembled microplastics were isolated and characterised, based on guidelines by Hidalgo-Ruz et al. (2012), placed on Whatman  
125 qualitative filters (No. 4) and enclosed in Petri dishes for imaging and plastic type diagnosis. Since 10 µm filters were used, this was the lower cut-  
126 off size for the diameter of fibres that would be retained. Exposure of filters to air was limited to < 1 min during particle transfer. To assess levels  
127 of potential airborne contamination during exposure of the filters to air two filters were left exposed on the lab bench within the room in which all  
128 sample viewing was conducted. Analysis of these exposed filters revealed the presence of a single clear fibre after 2 h. The transfer of particles  
129 from isolation filter to the one used for photography took <1 min, which validated the method used. It should be noted that visual observations of  
130 microplastic particles are subject to inaccuracies based on particle size and method(s) of observation (Song et al., 2015). For example, those  
131 authors reported that for sediment samples clear and white fragments can be disregarded as microplastics using microscopy for observation, yet  
132 spectral characteristics revealed otherwise. This under- representation of particles is probably widespread in the literature, since it is not always  
133 feasible to subject all particles to spectral analysis.

134 Imaging was done using a light microscope (Leica M205 C) at 5× magnification, with Petri dishes closed, and particle length was measuring using  
135 ImageJ. Of the 137 particles viewed, 91% were measured in terms of their longest length (mm). Thereafter samples were individually transferred  
136 to a Bruker Vertex 70 Fourier Transform Infrared Spectrometer (FTIR), equipped with a Hyperion 1000 Microscope attachment and a liquid  
137 nitrogen cooled mercury cadmium telluride detector (Bruker, Ettlingen, Germany). The instrument measured the absorbance of the sample using  
138 OPUS 7.5 software (Bruker, Ettlingen, Germany). Samples were prepared on a diamond compression cell and 30 background scans were run  
139 before analysing each sample. Sample spectra were based on 32 scans from 600 - 4000  $\lambda$ .  $\text{cm}^{-1}$ , at a resolution of 4  $\text{cm}^{-1}$  (Comnea-Stancu et al.,  
140 2017). Sample spectra were compared to reference spectra from a basic polymer library, a Bruker optics attenuated total reflection (ATR) polymer  
141 library and a synthetic fibres ATR library with 8, 234 and 337 entries, respectively. Sample evaluation was performed using a quality index  
142 adapted from Woodall et al. (2014). Polymer identification was based on the Euclidean distance between spectra, using the quick identity test  
143 option in the spectral software. Samples were compared to reference spectra and samples with a score of zero represented a perfect match, while

144 samples with a score of  $\leq 0.600$  were accepted as the reference polymer. Although the possibility of a cross-match between other organic matter  
145 and rayon does exist, as in the case of all other studies of this nature, care was taken to not include fibres that resembled natural materials by  
146 selecting fibres with no cellular structures, equal diameter and of a singular colour throughout (Reynolds and Ryan, 2018).

#### 147 Statistical analyses

148 Plastics found in fish were enumerated and thereafter characterised by shape, chemical nature (plastic type), colour and particle length (mm).  
149 Literature does point out the distinction between rayon and other microplastics since rayon is synthesized from naturally occurring polymers and  
150 hence considered semi-synthetic or regenerated fibre (Comnea-Stancu et al., 2017). However, rayon particles are frequently included in the  
151 category of microplastics for studies of this nature (e.g. Bessa et al., 2018; Halstead et al., 2018; Su et al., 2019b) given the fact that they are the  
152 product of a non-natural manufacturing process and are not a natural dietary component of aquatic organisms.

153 Given the uneven sample sizes for the different species and sites, nested ANOVA was used to test for any difference in mean particle abundance  
154 and mean particle length across species, within sites and across sites (data for all species within a site pooled). All differences were considered  
155 significant at the 0.05 level. The relationships among the total number of particles, particle length and fish mass were investigated using Spearman  
156 rank correlations. Particle length was averaged per fish if more than one particle was found.

#### 157 Results

158 Of the 174 fish analysed, 91 (52%) contained microplastics (Table 1). A total of 137 plastic particles were found, with an average number of  $0.787$   
159  $\pm 1.00$  particles per fish average (Table 1). The number of particles per fish did not differ significantly across sites ( $F = 2.58$ ,  $df = 3$ ,  $p = 0.06$ ) or  
160 among species, within sites ( $F = 1.41$ ,  $df = 2$ ,  $p = 0.25$ ). Nevertheless, data exhibited a few trends which are worth mentioning: (1) The maximum  
161 number of particles per fish was six, found in mullet collected from Durban Harbour, while tilapia from both St. Lucia and Isipingo and glassfish  
162 from Durban Harbour, contained a maximum of two particles per fish; (2) mullet from Umgeni exhibited the highest average number of particles;  
163 (3) the lowest average number and frequency of occurrence of ingested particles was observed in tilapia from the St. Lucia Estuary (Table 1).

164 The total number of particles per fish was not significantly correlated with fish mass ( $R = 0.106$ ,  $n = 174$   $p = 0.162$ ). Fish mass and particle length  
165 were also not significantly correlated ( $R = 0.086$ ,  $n = 86$ ,  $p = 0.433$ ). The average length of ingested particles was  $0.894 \pm 0.769$  mm and ranged  
166 from 0.1 - 4.8 mm and did not differ significantly across sites ( $F = 0.53$ ,  $df = 3$ ,  $p = 0.66$ ) or among species within sites ( $F = 1.46$ ,  $df = 2$ ,  $p = 0.24$ ).

167 The ingested particles fell into four shape categories and eleven plastic types (Figure 1). Despite this varied typology, the distribution of particles  
168 across these types was skewed. Fibres (68%) and fragments (21%) were the dominant shapes found, while rayon (70.4%), polyester (10.4%),  
169 nylon (5.2%) and polyvinylchloride (PVC, 3.0%) were the most abundant plastic types. Fibres composed of rayon dominated in all species and  
170 sites. Although the spectral sensitivities may vary in the four fish species sampled, the colour of particles are important since this may affect

171 particle selection by visually dependent feeders, especially if the microplastic resembles their prey items (e.g. Ory et al., 2017). The most common  
172 colours of microplastics found were blue, white, red, opaque and black (Figure 2).



173


174 Table 1. Characteristics of microplastics ingested by four fish species from each sampling site.

<b>Species</b>	<i>O. mossambicus</i>	<i>O. mossambicus</i>	<i>T. jarbua</i>	<i>Mugil sp.</i>	<i>Mugil sp.</i>	<i>A. dussumieri</i>	
<b>Common name</b>	<b>Tilapia</b>	<b>Tilapia</b>	<b>Thornfish</b>	<b>Mullet</b>	<b>Mullet</b>	<b>Glassfish</b>	<b>Total</b>
<b>Site</b>	<b>St. Lucia</b>	<b>Isipingo</b>	<b>Umgeni</b>	<b>Umgeni</b>	<b>Harbour</b>	<b>Harbour</b>	
Fish mass (g)*	0.13 ± 0.07	0.12 ± 0.18	0.07 ± 0.01	0.09 ± 0.05	0.41 ± 0.20	0.12 ± 0.09	0.16 ± 0.17
Fish with particles (number, %)	11 (38%)	13 (45%)	14 (48%)	17 (59%)	16 (55%)	20 (69%)	91 (52%)
No. of particles/fish*	0.41 ± 0.57	0.59 ± 0.73	0.66 ± 0.81	1.14 ± 1.25	1.00 ± 1.46	0.93 ± 0.75	0.79 ± 1.00
Min/Max no. of particles/fish	1/2	1/2	1/3	1/4	1/6	1/2	1/6
Particle length (mm) *	0.64 ± 0.46	0.92 ± 1.18	0.94 ± 0.81	0.88 ± 0.55	1.09 ± 0.77	0.76 ± 0.77	0.89 ± 0.77

175 All values are based on 29 individuals.


176 \*Values represent mean ± SD

177



178

179 Figure 1. The distribution (%) of microplastic particles ingested by juvenile fish across different (A) shape, (B) plastic type, (C) colour and (D)  
 180 size categories. The particles were extracted from four fish species sampled from four estuaries with mangroves in KwaZulu-Natal, South Africa.  
 181 Data for sites and species are pooled.



182

183 Figure 2. Microplastic particles found in juvenile fish of four fish species sampled from four mangroves in KwaZulu-Natal, South Africa. A:  
 184 particles of rayon (red and blue), B: blue fragment of PVC, C: opaque fragment of PVC from Durban Harbour mullet, D: polyester fibre, E: rayon  
 185 fibres and F: polyethylene fragment from a thornfish in Umgeni.

186 Discussion

187 *Comparison between sites and species*

188

189 The eastern seaboard of KZN, the location for all four study sites, is one of the most densely populated coastal regions in South Africa. Rapid  
190 urbanisation and a wide range of industrial and tourist activities along the coastline and upstream of the rivers that flow into many of its estuaries,  
191 introduce pollutants into many water bodies in the region (O'Donoghue and Marshall, 2003). Most recently, reports of microplastic ingestion in  
192 fish sampled in the region point towards the prevalence of plastic pollution in coastal systems in KZN (Naidoo et al., 2016). The findings of the  
193 present study support this assertion but also show the susceptibility of juvenile fish associated with KZN mangroves to microplastic ingestion.  
194 Furthermore, ingestion appears to be independent of species (at least for the four investigated here) and site; for instance, the quantity of  
195 microplastics ingested by fish in the most protected site (St. Lucia) was comparable with fish from Umgeni and Durban Harbour which are open to  
196 numerous anthropogenic activities. Whilst microplastic data were not assessed for this study, literature indicates that microplastics are present in  
197 the sediment and water fractions at the three urban sites investigated here (Naidoo et al., 2015). At the time of this study, environmental  
198 microplastic data were not available for St. Lucia. However, we believe that even mangrove systems that are sheltered from urbanisation can  
199 become contaminated by microplastic particles as a consequence of plastic inputs from runoff during rainfall events, domestic and industrial  
200 effluent that enter rivers that flow into estuaries and washing of clothes in rivers systems by local communities (Le, 2017). The plastic retention  
201 capability of mangrove forests is reported to be high (Martin et al., 2019) and thus particles entering this system are possibly trapped in, and could  
202 possibly be transferred to fish either directly by prey mis-identification and indirectly via their invertebrate prey (Farrell and Nelson, 2013).  
203 Coupled with this, in systems such as St. Lucia which has been closed for > 15 years (Tweddle et al., 2016), plastics could accumulate as a  
204 consequence of limited water exchange with the ocean (Claessens et al., 2011).

205 Previous studies suggest that there may be a difference in microplastic ingestion by intertidal fish of different feeding guilds (Mizraji et al., 2017).  
206 Literature suggests that whilst some species like mullet are generalistic feeding fish that are indiscriminant feeders, other species such as glassfish  
207 are selective feeders that feed largely on zooplankton (Dyer et al., 2015). Differences in plastic ingestion by species with different feeding  
208 strategies was also suggested to occur by Wright et al. (2013), but we found no significant evidence of this in the four species compared here.  
209 Similarly, Vendel et al. (2017) found that a fish's feeding guild did not determine the quantity of microplastics ingested when comparing 29  
210 species of estuarine fish. Halstead et al. (2018), also found no difference in the abundance of microplastics ingested by three species of benthic  
211 feeding fish in an urban estuary. It must be noted though that all the studies quoted above focused on adult and sub-adult fish as opposed to  
212 juvenile fish. Whilst all fish exhibit the tendency to feed near the sediment, certain species can adopt feeding strategies that could  
213 increase/decrease the number of microplastics ingested. Juvenile mullet, thornfish and tilapia most often feed near the sediment, on  
214 microphytobenthos or benthic invertebrates ([www.fishbase.org](http://www.fishbase.org)), in shallow water ecosystems, and this could explain the lack of significant

215 differences in microplastic ingestion levels between species within sites. Rayon particles, that are frequently reported in fish, are abundant in  
216 sediment and would this would explain the large portion of rayon particles in these species (Gago et al., 2018). Microplastics can also move  
217 through food chains (Setälä et al., 2014), and therefore glassfish that feed in the water column were also exposed to these particles.

218 Our study indicated that fish in this study area consumed similar levels of microplastics to other benthic feeding estuarine fish from urban settings.  
219 For example, we found that 52% of the 174 fish we investigated ingested microplastics, while Halstead et al. (2018), found that 43% of 93 fish  
220 from Sydney Harbour ingested microplastics. Estuarine and coastal fish seem to consume higher amounts of plastic particles than fish in oceanic  
221 habitats (see review by Jovanović, 2017), however it must also be noted that methodological differences among studies make comparisons  
222 difficult. As alluded to above, this is possibly because plastics can concentrate in estuaries, making their interaction with fish more frequent in  
223 these systems. For example Ramos et al. (2012) found that 13.4% of 425 fish had ingested blue nylon threads, in a tropical estuary where fishing is  
224 the main microplastic source, while Steer et al. (2017) found that only 2.9% of the 347 fish larvae they examined had ingested microplastics in the  
225 English channel, where fishing is common, but particles could possibly be wider spread. Some freshwater fish higher up rivers also seem to exhibit  
226 high levels of plastic ingestion. Karlsson et al. (2017), for example, found that 68% of 62 brown trout had consumed plastics. This suggests that  
227 fish feeding closer to terrestrial and freshwater plastic sources may be interacting more frequently with particles, prompting higher levels of  
228 ingestion.

#### 229 *Main plastic types and sources*

230 Most of the plastics found in fish were fibrous in shape. This is the case with many studies that have investigated microplastic ingestion in marine  
231 biota (as reviewed by Gago et al. (2018); and is especially so for estuarine fish (Ramos et al., 2012; Ferreira et al., 2016; Vendel et al., 2017;  
232 Halstead et al., 2018). The most common microfibrils reported in the environment and fish are acrylic, polypropylene, polyethylene, polyester and  
233 rayon (Gago et al., 2018; Halstead et al., 2018), which were all also encountered in this study. Rayon, also known as viscose in some locations,  
234 was by far the most abundant plastic ingested here. It is a semi-synthetic cellulosic fibre, manufactured from wood pulp and the derivatization of  
235 cellulose with carbon disulfide (Comnea-Stancu et al., 2017; Gago et al., 2018). Rayon is used in the manufacture of clothing and therefore fibres  
236 can be sourced from abrasion while washing (Comnea-Stancu et al., 2017).

237 Washing machine effluent can also contain polyester and acrylic fibres (Napper and Thompson, 2016), which were also found within fish in the  
238 current study. It has been estimated that > 1 900 fibres could be formed from machine washing a single garment (Browne et al., 2011). The type of  
239 material may also be important in creating fibres as it is estimated that 700 000 fibres could be released from a 6 kg load of acrylic material  
240 (Napper and Thompson, 2016). There have been reports of these fibres passing through sewage treatment plants into natural waterways (Browne  
241 et al., 2011), which is probably why they are so prevalent. Halstead et al. (2018), for example, estimated that a single mullet in an urban estuary  
242 has the potential to consume 11 000 microfibrils annually. Many of these fibres can ultimately end up in the deep sea (Woodall et al., 2014).

243 *Potential impacts of plastics and associated pollutants on fish health*

244 Although feeding performance was not tested here, fish can display decreased feeding performance when exposed to plastics < 90 µm (Miranda et  
245 al., 2019), however this is not true for all species (Jacob et al., 2019). Consumption of particles of the types found within the fish species studied  
246 here have been shown to have negative consequences. For example, glassfish fed a mixture of the plastics types found in this study, such as  
247 polyethylene, polystyrene and polyvinyl chloride, were also shown to have slower growth compared with controls during chronic exposure  
248 (Naidoo and Glassom, 2019). In other fish species, endocrine disruption was observed after chronic ingestion of both virgin and marine sourced  
249 polyethylene particles (Rochman et al., 2014). It must be noted however, that these were laboratory-based experiments in which fish were given a  
250 plastic feed daily, unlike the snap-shot we investigated here. The size of particles found in experiments mentioned above (~ 0.5 mm), and those  
251 found here (> 0.1 mm), were much larger than those shown to translocate into circulatory systems of fish (< 20 µm), by Su et al. (2019a).  
252 However, translocation of associated pollutants is possible (Bakir et al., 2014), which may have possible negative health impacts. For example,  
253 blue rayon fibres, are able to adsorb polycyclic aromatic hydrocarbons (PAHs) from estuaries, and have thus been used for monitoring  
254 environmental concentrations (Sakamoto and Hayatsu, 1990; Kummrow et al., 2006). Given that the bulk of the particles found here are rayon,  
255 this warrants further investigation into pollutants associated with these particles within mangroves which are widely reported to exhibit high levels  
256 of organic and inorganic pollutants.

257 *Particle length and colour*

258 The average length of ingested particles found here was  $0.894 \pm 0.769$  mm. This size is common for fibres and are within a size range that can  
259 easily be ingested by marine biota (Gago et al., 2018), including fish and their prey items (Ory et al., 2017). Fish mass did not correlate with  
260 particle length, which suggests that larger fish may not have consumed longer particles; i.e. these fish species appear to be indiscriminate in their  
261 selection of particles. As in other studies, example Vendel et al. (2017), the number of particles consumed also did not correlate with fish mass,  
262 negating the possibility that larger fish consume more particles than smaller fish.

263 Blue was the most common colour of fibres found in the mangrove associated fish examined here. Blue, black, white and red were also the most  
264 common colours of plastic found in other estuarine fish (Ramos et al., 2012; Ferreira et al., 2016). Blue fibres seem to be the most common colour  
265 of fibres in general, as confirmed by South African nearshore water (Nel and Froneman, 2015), global seawater and sediment microfibre  
266 assessments (Gago et al., 2018). Other common microplastic colours are transparent and black (Gago et al., 2018), which were also dominant here.  
267 Microplastic colour is important since it may influence uptake by visual predators. For example, Ory et al. (2017) found that small pelagic fish  
268 (*Decapterus muroadsi* [Temminck and Schlegel, 1844]), ingested mainly blue particles, possibly because this resembled the colour of the  
269 copepods they prey on. White and opaque colours, which were also found here, could also resemble copepods and other prey that that fish  
270 consume (Carpenter et al., 1972; Wright et al., 2013).

271 Concluding remarks and recommendations

272 The levels of microplastic ingestion by the juvenile fish studied here is important in the context of ingestion levels reported in other studies.  
273 Ingestion levels do not appear to be species or site dependent but high particle abundance and frequency of occurrence across all species may be  
274 reflective of the high levels of microplastics within mangrove systems. Rayon seems to be most prevalent in fish studied here which is in keeping  
275 with trends in the past and current literature. The main caveat of this study is that we could not obtain the same suite of species at all sites due to  
276 site and species peculiarities. Future studies should also consider comparing estuarine filter-feeding fish to benthic feeding fish. Fish do, however,  
277 change their feeding preferences during development (Ferreira et al., 2019) which should also be factored in when assessing the risks of  
278 microplastic ingestion in a species. In the context of assessing risk, the processing methods used in this study can serve as an effective approach  
279 for quantifying and characterising microplastic ingestion in fish. However, this should be coupled with techniques that identify the pollutants that  
280 adsorb onto microplastic particles for a more holistic assessment of how microplastics affect fish health and functioning. It must be acknowledged  
281 that the possibility of particles behind the gills contributing to abundance counts cannot be discounted entirely. Each fish was checked rigorously  
282 for surface contamination to minimize this possibility but we recommend that future studies consider removing the gills or head of specimens and  
283 analyzing these separately for microplastics.

284 Acknowledgements

285 The authors would like to acknowledge the Marine Biology and Ecology Research Centre (MBERC) team at the University of Plymouth,  
286 including Madeleine Steer for useful tips on proteinase digestion, Andy Atfield for organising lab equipment and Andrew Tonkin for assistance  
287 with the use of the  $\mu$ FTIR. Funding for this project was made possible by the National Research Foundation (NRF) grant number  
288 (SFP170605237015) and the Association of Common Wealth Universities, Blue Charter Fellowship, awarded to T. Naidoo. Mangrove research  
289 groups at the University of KwaZulu-Natal and the University of the Western Cape are funded by the NRF Marine and Coastal Research Grant  
290 (MCR180611344089).

291

292

293

294

295

296

297

298 References

- 299 Anastasopoulou, A., Mytilineou, C., Smith, C.J., Papadopoulou, K.N., 2013. Plastic debris ingested by deep-water fish of the Ionian Sea (Eastern  
300 Mediterranean). *Deep Sea Research Part I: Oceanographic Research Papers* 74, 11-13.
- 301 Bakir, A., O'Connor, I.A., Rowland, S.J., Hendriks, A.J., Thompson, R.C., 2016. Relative importance of microplastics as a pathway for the  
302 transfer of hydrophobic organic chemicals to marine life. *Environmental Pollution* 219, 56-65.
- 303 Bakir, A., Rowland, S.J., Thompson, R.C., 2014. Enhanced desorption of persistent organic pollutants from microplastics under simulated  
304 physiological conditions. *Environmental Pollution* 185, 16-23.
- 305 Beckingham, B., Ghosh, U., 2017. Differential bioavailability of polychlorinated biphenyls associated with environmental particles: Microplastic  
306 in comparison to wood, coal and biochar. *Environmental Pollution* 220, 150-158.
- 307 Bessa, F., Barría, P., Neto, J.M., Frias, J.P., Otero, V., Sobral, P., Marques, J., 2018. Occurrence of microplastics in commercial fish from a natural  
308 estuarine environment. *Marine Pollution Bulletin* 128, 575-584.
- 309 Besseling, E., Wegner, A., Foekema, E.M., Heuvel-Greve, M.J., Koelmans, A.A., 2013. Effects of microplastic on fitness and PCB  
310 bioaccumulation by the lugworm *Arenicola marina* (L.). *Environmental Science and Technology* 47, 593-600.
- 311 Biginagwa, F.J., Mayoma, B.S., Shashoua, Y., Syberg, K., Khan, F.R., 2016. First evidence of microplastics in the African Great Lakes: Recovery  
312 from Lake Victoria Nile perch and Nile tilapia. *Journal of Great Lakes Research* 42, 146-149.
- 313 Boerger, C.M., Lattin, G.L., Moore, S.L., Moore, C.J., 2010. Plastic ingestion by planktivorous fishes in the North Pacific Central Gyre. *Marine  
314 Pollution Bulletin* 60, 2275-2278.
- 315 Browne, M.A., Crump, P., Niven, S.J., Teuten, E., Tonkin, A., Galloway, T., Thompson, R., 2011. Accumulation of microplastic on shorelines  
316 worldwide: Sources and sinks. *Environmental Science and Technology* 45, 9175-9179.
- 317 Browne, M.A., Dissanayake, A., Galloway, T.S., Lowe, D.M., Thompson, R.C., 2008. Ingested microscopic plastic translocates to the circulatory  
318 system of the mussel, *Mytilus edulis* (L.). *Environmental Science and Technology* 42, 5026-5031.
- 319 Carpenter, E.J., Anderson, S.J., Harvey, G.R., Miklas, H.P., Peck, B.B., 1972. Polystyrene spherules in coastal waters. *Science* 178, 749-750.



- 320 Claessens, M., Meester, S.D., Landuyt, L.V., Clerck, K.D., Janssen, C.R., 2011. Occurrence and distribution of microplastics in marine sediments  
321 along the Belgian coast. *Marine Pollution Bulletin* 62, 2199-2204.
- 322 Cole, M., Webb, H., Lindeque, P.K., Fileman, E.S., Halsband, C., Galloway, T.S., 2014. Isolation of microplastics in biota-rich seawater samples  
323 and marine organisms. *Scientific Reports* 4, 1-8.
- 324 Comnea-Stancu, I.R., Wieland, K., Ramer, G., Schwaighofer, A., Lendl, B., 2017. On the identification of rayon/viscose as a major fraction of  
325 microplastics in the marine environment: Discrimination between natural and manmade cellulosic fibers using fourier transform infrared  
326 spectroscopy. *Applied Spectroscopy* 71, 939-950.
- 327 Davison, P., Asch, R.G., 2011. Plastic ingestion by mesopelagic fishes in the North Pacific Subtropical Gyre. *Marine Ecology Progress Series* 432,  
328 173-180.
- 329 de Sá, L.C., Luís, L.G., Guilhermino, L., 2015. Effects of microplastics on juveniles of the common goby (*Pomatoschistus microps*): Confusion  
330 with prey, reduction of the predatory performance and efficiency, and possible influence of developmental conditions. *Environmental Pollution*  
331 196, 359-362.
- 332 Dyer, D.C., Perissinotto, R., Carrasco, N.K., 2015. Temporal and spatial dietary dynamics of the longspine glassy (*Ambassis ambassis*) in the St  
333 Lucia estuarine system, iSimangaliso Wetland Park. *Water South Africa* 41, 91-104.
- 334 Farrell, P., Nelson, K., 2013. Trophic level transfer of microplastic: *Mytilus edulis* (L.) to *Carcinus maenas* (L.). *Environmental Pollution* 177, 1-3.
- 335 Ferreira, G.V., Barletta, M., Lima, A.R., 2019. Use of estuarine resources by top predator fishes. How do ecological patterns affect rates of  
336 contamination by microplastics? *Science of the Total Environment* 655, 292-304.
- 337 Ferreira, G.V., Barletta, M., Lima, A.R., Dantas, D.V., Justino, A.K., Costa, M.F., 2016. Plastic debris contamination in the life cycle of Acoupa  
338 weakfish (*Cynoscion acoupa*) in a tropical estuary. *ICES Journal of Marine Science* 73, 2695-2707.
- 339 Friess, D.A., Rogers, K., Lovelock, C.E., Krauss, K.W., Hamilton, S.E., Lee, S.Y., Lucas, R., Primavera, J., Rajkaran, A., Shi, S., 2019. The state  
340 of the world's mangrove forests: Past, present, and future. *Annual Review of Environment and Resources* 44, 1-27.
- 341 Gago, J., Carretero, O., Filgueiras, A., Viñas, L., 2018. Synthetic microfibers in the marine environment: A review on their occurrence in seawater  
342 and sediments. *Marine Pollution Bulletin* 127, 365-376.
- 343 Halstead, J.E., Smith, J.A., Carter, E.A., Lay, P.A., Johnston, E.L., 2018. Assessment tools for microplastics and natural fibres ingested by fish in  
344 an urbanised estuary. *Environmental Pollution* 234, 552-561.

- 345 Harris, S., Cyrus, D., 1999. Composition, abundance and seasonality of fish larvae in the mouth of Durban Harbour, KwaZulu-Natal, South Africa.  
346 South African Journal of Marine Science 21, 19-39.
- 347 Hidalgo-Ruz, V., Gutow, L., Thompson, R.C., Thiel, M., 2012. Microplastics in the marine environment: A review of the methods used for  
348 identification and quantification. Environmental Science and Technology 46, 3060–3075.
- 349 Jacob, H., Gilson, A., Lanctôt, C., Besson, M., Metian, M., Lecchini, D., 2019. No effect of polystyrene microplastics on foraging activity and  
350 survival in a post-larvae coral-reef fish, *Acanthurus triostegus*. Bulletin of Environmental Contamination and Toxicology 102, 457-461.
- 351 Jovanović, B., 2017. Ingestion of microplastics by fish and its potential consequences from a physical perspective. Integrated Environmental  
352 Assessment and Management 13, 510-515.
- 353 Karlsson, T.M., Vethaak, A.D., Almroth, B.C., Ariese, F., van Velzen, M., Hassellöv, M., Leslie, H.A., 2017. Screening for microplastics in  
354 sediment, water, marine invertebrates and fish: method development and microplastic accumulation. Marine Pollution Bulletin 122, 403-408.
- 355 Kripa, V., Nair, P.G., Dhanya, A., Pravitha, V., Abhilash, K., Mohammed, A.A., Vijayan, D., Vishnu, P., Mohan, G., Anilkumar, P., 2014.  
356 Microplastics in the gut of anchovies caught from the mud bank area of Alappuzha, Kerala. Marine Fisheries Information Service; Technical and  
357 Extension Series, 27-28.
- 358 Kummrow, F., Rech, C.M., Coimbra, C.A., Umbuzeiro, G.A., 2006. Blue rayon-anchored technique/*Salmonella microsoma* microsuspension  
359 assay as a tool to monitor for genotoxic polycyclic compounds in Santos Estuary. Mutation Research/Genetic Toxicology and Environmental  
360 Mutagenesis 609, 60-67.
- 361 Laegdsgaard, P., Johnson, C., 2001. Why do juvenile fish utilise mangrove habitats? Journal of Experimental Marine Biology and Ecology 257,  
362 229-253.
- 363 Lamberth, S., Turpie, J., 2003. The role of estuaries in South African fisheries: Economic importance and management implications. African  
364 Journal of Marine Science 25, 131-157.
- 365 Le, K., 2017. Microfiber Shedding: Hidden Environmental Impact. AATCC Review 17, 30-37.
- 366 Lusher, A., McHugh, M., Thompson, R., 2013. Occurrence of microplastics in the gastrointestinal tract of pelagic and demersal fish from the  
367 English Channel. Marine Pollution Bulletin 67, 94-99.
- 368 Lusher, A., Welden, N., Sobral, P., Cole, M., 2017. Sampling, isolating and identifying microplastics ingested by fish and invertebrates. Analytical  
369 Methods 9, 1346-1360.

- 370 Markic, A., Nicol, S., 2014. In a nutshell: Microplastics and fisheries. SPC Fisheries Newsletter 144, 27-29.
- 371 Martin, C., Almahasheer, H., Duarte, C.M., 2019. Mangrove forests as traps for marine litter. Environmental Pollution 247, 499-508.
- 372 Miranda, T., Vieira, L.R., Guilhermino, L., 2019. Neurotoxicity, behavior, and lethal effects of cadmium, microplastics, and their mixtures on  
373 *Pomatoschistus microps* juveniles from two wild populations exposed under laboratory conditions—implications to environmental and human risk  
374 assessment. International Journal of Environmental Research and Public Health 16, 2857.
- 375 Mizraji, R., Ahrendt, C., Perez-Venegas, D., Vargas, J., Pulgar, J., Aldana, M., Ojeda, F.P., Duarte, C., Galbán-Malagón, C., 2017. Is the feeding  
376 type related with the content of microplastics in intertidal fish gut? Marine Pollution Bulletin 116, 498-500.
- 377 Moore, C., Lattin, G., Zellers, A., 2005. Density of plastic particles found in zooplankton trawls from coastal waters of California to the North  
378 Pacific Central Gyre, The Plastic Debris Rivers to Sea Conference, Redondo Beach, California, USA.
- 379 Naidoo, T., Glassom, D., 2019. Decreased growth and survival in small juvenile fish, after chronic exposure to environmentally relevant  
380 concentrations of microplastic. Marine Pollution Bulletin 145, 254-259.
- 381 Naidoo, T., Glassom, D., Smit, A., J., 2015. Plastic pollution in five urban estuaries of KwaZulu-Natal, South Africa. Marine Pollution Bulletin  
382 101, 473-480.
- 383 Naidoo, T., Glassom, D., Smit, A., J., 2016. Plastic ingestion by estuarine mullet, *Mugil cephalus* (L., 1758), in an urban harbour, KwaZulu-Natal,  
384 South Africa. African Journal of Marine Science 38, 145-149.
- 385 Naidoo, T., Goordiyal, K., Glassom, D., 2017. Are nitric acid (HNO<sub>3</sub>) digestions efficient in isolating microplastics from juvenile fish? Water,  
386 Air, and Soil Pollution 228, 1-11.
- 387 Napper, I.E., Thompson, R.C., 2016. Release of synthetic microplastic plastic fibres from domestic washing machines: Effects of fabric type and  
388 washing conditions. Marine Pollution Bulletin 112, 39-45.
- 389 Nel, H., Froneman, P., 2015. A quantitative analysis of microplastic pollution along the south-eastern coastline of South Africa. Marine Pollution  
390 Bulletin 101, 274-279.
- 391 O'Donoghue, S., Marshall, D.J., 2003. Marine pollution research in South Africa: A status report: Review article. South African Journal of Science  
392 99, 349-356.

- 393 Ory, N.C., Sobral, P., Ferreira, J.L., Thiel, M., 2017. Amberstripe scad *Decapterus muroadsi* (Carangidae) fish ingest blue microplastics  
394 resembling their copepod prey along the coast of Rapa Nui (Easter Island) in the South Pacific Subtropical Gyre. *Science of the Total Environment*  
395 586, 430-437.
- 396 Possatto, F.E., Spach, H.L., Cattani, A.P., Lamour, M.R., Santos, L.O., Cordeiro, N.M.A., Broadhurst, M.K., 2015. Marine debris in a world  
397 heritage listed Brazilian estuary. *Marine Pollution Bulletin* 91, 548-553.
- 398 Ramos, J.A., Barletta, M., Costa, M.F., 2012. Ingestion of nylon threads by Gerreidae while using a tropical estuary as foraging grounds. *Aquatic*  
399 *Biology* 17, 29-34.
- 400 Reynolds, C., Ryan, P.G., 2018. Micro-plastic ingestion by waterbirds from contaminated wetlands in South Africa. *Marine Pollution Bulletin*  
401 126, 330-333.
- 402 Rios, L.M., Moore, C., Jones, P.R., 2007. Persistent organic pollutants carried by synthetic polymers in the ocean environment. *Marine Pollution*  
403 *Bulletin* 54, 1230-1237.
- 404 Rochman, C.M., Browne, M.A., Halpern, B.S., Hentschel, B.T., Hoh, E., Karapanagioti, H.K., Rios-Mendoza, L.M., Takada, H., Teh, S.,  
405 Thompson, R.C., 2013a. Policy: Classify plastic waste as hazardous. *Nature* 494, 169-171.
- 406 Rochman, C.M., Hoh, E., Kurobe, T., Teh, S., 2013b. Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. *Scientific*  
407 *Reports* 3, 1-7.
- 408 Rochman, C.M., Kurobe, T., Flores, I., Teh, S.J., 2014. Early warning signs of endocrine disruption in adult fish from the ingestion of  
409 polyethylene with and without sorbed chemical pollutants from the marine environment. *Science of the Total Environment* 493, 656-661.
- 410 Sakamoto, H., Hayatsu, H., 1990. A simple method for monitoring mutagenicity of river water. *Mutagens in Yodo River system, Kyoto-Osaka.*  
411 *Bulletin of Environmental Contamination and Toxicology* 44, 521-528.
- 412 Setälä, O., Fleming-Lehtinen, V., Lehtiniemi, M., 2014. Ingestion and transfer of microplastics in the planktonic food web. *Environmental*  
413 *Pollution* 185, 77-83.
- 414 Song, Y.K., Hong, S.H., Jang, M., Han, G.M., Rani, M., Lee, J., Shim, W.J., 2015. A comparison of microscopic and spectroscopic identification  
415 methods for analysis of microplastics in environmental samples. *Marine Pollution Bulletin* 93, 202-209.
- 416 Steer, M., Cole, M., Thompson, R.C., Lindeque, P.K., 2017. Microplastic ingestion in fish larvae in the western English Channel. *Environmental*  
417 *Pollution* 226, 250-259.

- 418 Su, L., Deng, H., Li, B., Chen, Q., Pettigrove, V., Wu, C., Shi, H., 2019a. The occurrence of microplastic in specific organs in commercially  
419 caught fishes from coast and estuary area of east China. *Journal of Hazardous Materials* 365, 716-724.
- 420 Su, L., Nan, B., Hassell, K.L., Craig, N.J., Pettigrove, V., 2019b. Microplastics biomonitoring in Australian urban wetlands using a common  
421 noxious fish (*Gambusia holbrooki*). *Chemosphere* 228, 65-74.
- 422 Sussarellu, R., Suquet, M., Thomas, Y., Lambert, C., Fabioux, C., Pernet, M.E.J., Le Goïc, N., Quillien, V., Mingant, C., Epelboin, Y., 2016.  
423 Oyster reproduction is affected by exposure to polystyrene microplastics. *Proceedings of the National Academy of Sciences* 113, 2430-2435.
- 424 Tweddle, G., Cyrus, D., Vivier, L., 2016. Penaeid and carid community changes in the St Lucia estuarine lake system, South Africa, under low  
425 water level, extended closed periods and marine reconnection conditions. *African Journal of Aquatic Science* 41, 251-266.
- 426 van Niekerk, L., Adams, J.B., Lamberth, S.J., MacKay, C.F., Taljaard, S., Turpie, J.K., S.P., W., Raimondo, D.C., (eds), 2019. South African  
427 National Biodiversity Assessment 2018: Technical Report. Volume 3: Estuarine Realm. CSIR report number CSIR/SPLA/EM/EXP/2019/0062/A.  
428 South African National Biodiversity Institute, Pretoria. Report Number: SANBI/NAT/NBA2018/2019/Vol3/A.
- 429 van Niekerk, L., Turpie, J., 2012. National Biodiversity Assessment 2011: Technical Report. Volume 3: Estuary Component. CSIR Report  
430 Number CSIR/NRE/ECOS/ER/2011/0045/B. Council for Scientific and Industrial Research, Stellenbosch. 1-202.
- 431 Vendel, A.L., Bessa, F., Alves, V., Amorim, A., Patrício, J., Palma, A., 2017. Widespread microplastic ingestion by fish assemblages in tropical  
432 estuaries subjected to anthropogenic pressures. *Marine Pollution Bulletin* 117, 448-455.
- 433 Wallace, J.H., Kok, H., Beckley, L., Bennett, B., Blaber, S., Whitfield, A., 1984. South African estuaries and their importance to fishes. *South  
434 African Journal of Science* 80, 203-207.
- 435 Ward, C., Steinke, T.D., 1982. A note on the distribution and approximate areas of mangroves in South Africa. *South African Journal of Botany* 1,  
436 51-53.
- 437 Whitfield, A., Blaber, S., Cyrus, D., 1981. Salinity ranges of some southern African fish species occurring in estuaries. *African Zoology* 16, 151-  
438 155.
- 439 Woodall, L.C., Gwinnett, C., Packer, M., Thompson, R.C., Robinson, L.F., Paterson, G.L., 2015. Using a forensic science approach to minimize  
440 environmental contamination and to identify microfibrils in marine sediments. *Marine Pollution Bulletin* 95, 40-46.
- 441 Woodall, L.C., Sanchez-Vidal, A., Canals, M., Paterson, G.L., Coppock, R., Sleight, V., Calafat, A., Rogers, A.D., Narayanaswamy, B.E.,  
442 Thompson, R.C., 2014. The deep sea is a major sink for microplastic debris. *Royal Society Open Science* 1, 1-8.

443 Wright, S.L., Thompson, R.C., Galloway, T.S., 2013. The physical impacts of microplastics on marine organisms: A review. *Environmental*  
444 *Pollution* 178, 483-492.

445

446