

2021-03

Chloroplast immunity illuminated

Littlejohn, George

<http://hdl.handle.net/10026.1/17850>

10.1111/nph.17076

New Phytologist

Wiley

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.



Tansley review

Chloroplast immunity illuminated

Author for correspondence:
Murray Grant
Email: m.grant@warwick.ac.uk

Received: 29 May 2020
Accepted: 12 September 2020

George R. Littlejohn^{1*} , Susan Breen^{2*} , Nicholas Smirnoff³  and Murray Grant² 

¹School of Biological and Marine Sciences, University of Plymouth, Plymouth PL4 8AA, UK; ²School of Life Sciences, University of Warwick, Coventry, CV4 7AL, UK; ³Biosciences, College of Life and Environmental Sciences, University of Exeter, Exeter, EX4 4QD, UK

Contents

Summary	3088	VI. Cellular reorganization during infection, stromules and perinuclear chloroplast movement	3099
I. Introduction	3088	VII. Functional significance of suppression nuclear encoded chloroplast genes (<i>NECGs</i>)	3101
II. Hormones and chloroplasts, a well-established link in plant immunity	3091	VIII. Concluding remarks	3103
III. cROS in immunity and insights from disruption of chloroplast components	3092	Acknowledgements	3103
IV. Immunity insights from perturbation of chloroplast metabolism and cROS production	3093	References	3103
V. Direct targeting of pathogen effectors to the chloroplast	3094		

Summary

New Phytologist (2021) **229**: 3088–3107
doi: 10.1111/nph.17076

Key words: chloroplast immunity, reactive oxygen species (ROS), PAMP-triggered immunity (PTI), effector-triggered immunity (ETI), photosystem, electron transport.

The chloroplast has recently emerged as pivotal to co-ordinating plant defence responses and as a target of plant pathogens. Beyond its central position in oxygenic photosynthesis and primary metabolism – key targets in the complex virulence strategies of diverse pathogens – the chloroplast integrates, decodes and responds to environmental signals. The capacity of chloroplasts to synthesize phytohormones and a diverse range of secondary metabolites, combined with retrograde and reactive oxygen signalling, provides exquisite flexibility to both perceive and respond to biotic stresses. These processes also represent a plethora of opportunities for pathogens to evolve strategies to directly or indirectly target ‘chloroplast immunity’. This review covers the contribution of the chloroplast to pathogen associated molecular pattern and effector triggered immunity as well as systemic acquired immunity. We address phytohormone modulation of immunity and surmise how chloroplast-derived reactive oxygen species underpin chloroplast immunity through indirect evidence inferred from genetic modification of core chloroplast components and direct pathogen targeting of the chloroplast. We assess the impact of transcriptional reprogramming of nuclear-encoded chloroplast genes during disease and defence and look at future research challenges.

II. Introduction

A plant’s initial response to a broad spectrum of different stresses, including pathogens, is through integrated signalling modules that recognize a common set of second messengers (calcium, reactive

oxygen species (ROS), nitric oxide (NO) and lipid molecules), often incorporating kinase-based signal transduction cascades. Understanding how cells specify the timing, amplitude and duration of signal outputs, and decode and integrate these signals locally and distally remains a key challenge in plant biology. What is

*These authors contributed equally to this work.

often neglected is that these signals are perceived not only at the cell surface and/or in the nucleus, but also by other organelles, which collectively contribute to orchestrating an effective response.

1. Plant immunity, more than membrane to nuclear signalling

Put simply, plant immunity comprises three core modules. Predominantly membrane-localized pattern recognition receptors (PRRs) perceive pathogen-associated molecular patterns (PAMPs) activating PAMP-triggered immunity (PTI). Pathogens deliver effectors (generally proteinaceous but also small molecules) directly or indirectly into the cell to collectively suppress PTI, often targeting the PRRs and their coreceptors. This effector triggered suppression (ETS) can be successfully overcome by intracellular plant disease resistance (R) proteins which activate effector-triggered immunity (ETI) effectively containing and eliminating the invading pathogen through a programmed cell death process known as the hypersensitive response (HR) (Jones & Dangl, 2006). However, there is a growing acceptance that PTI and ETI are not two distinct processes but are somewhat interdependent and contribute as a continuum to host immune responses (van der Burgh & Joosten, 2019). Superimposed on ETI is the initiation and establishment of broad-spectrum systemic immunity known as systemic acquired resistance (SAR) (Shine *et al.*, 2019). Because of the localization of PRRs and signalling components of ETI, most of the innate immunity research has been focused on the cell membrane, the nucleus and the role of MAPK (mitogen-activated protein kinase) signalling cascades in unravelling plant immunity mechanisms.

2. The chloroplast is a key hub in coordinating effective plant immune responses

Aside from oxygenic photosynthesis, chloroplasts act as both environmental signal integrators and metabolic hubs. Chloroplasts not only link to primary metabolism but synthesize phytohormones, fatty acids, amino acids and a plethora of other secondary metabolites. This therefore provides unprecedented flexibility in fine tuning complex signalling to specific environmental stresses, and the capacity to rapidly modulate and redeploy metabolic signalling. This review will focus on the role of the chloroplast in disease and defence and seek to provide the reader with an overview of current knowledge of chloroplast immunity. We will examine evidence of a pivotal role for chloroplasts both in orchestrating an effective immune response and as a pathogen target, to suppress immunity. Pathogens probably also reconfigure primary metabolism for nutrition, although experimental insight into this is limited. We will additionally touch on current concepts in retrograde signalling and draw parallels with other stress process that impact chloroplast homeostasis to explore commonalities in signalling responses.

3. Chloroplasts in plant immunity: an historical overview

In the past decade, the chloroplast has emerged as a central player in plant defence, initially in the context of its identification as a

genuine effector target but more recently in recognition of its contribution to defence. The importance of the chloroplast in immunity has been known for a long time. Kupeevicz (1947) first reported that viruses and other plant pathogens alter chlorophyll (Chl) accumulation during infection. By the 1990s, viral proteins, such as the coat protein of Tobacco Mosaic Virus (TMV), were identified within the chloroplast (Banerjee & Zaitlin, 1992). In comparison to virus research (which we only use here as exemplars), studies on how bacteria, fungal and oomycete pathogens target the chloroplast were limited until the emergence of 'effector biology' in the early 2000s.

4. The complexity of chloroplast immunity: where to begin?

We aim to leave readers with two key messages, the first being that chloroplast-derived reactive oxygen species (cROS) play a pivotal role in establishing effective plant immunity and, secondly, that pathogen effectors directly and indirectly target chloroplast processes to suppress immunity. Obviously, to effect these changes a plethora of processes are activated or suppressed. While recognising this is still an embryonic field, we will draw on relevant examples to provide insight into current state-of-knowledge of the complexity of chloroplast processes modified and known components targeted. We first briefly overview phytohormone modulation of immunity, and the contribution of the chloroplast to PTI, ETI and SAR in the context of cROS, and how effectors modulate/facilitate this.

We next document potential processes contributing to chloroplast immunity that have been revealed using genetic approaches, with a strong focus on the role of cROS. We then provide a comprehensive overview of proteinaceous pathogen effectors targeted to the chloroplast and their targets, if known. Finally, we examine transcriptional control of nuclear-encoded chloroplast genes in PTI and ETI and touch on the emerging role of subcellular reorganization. Chloroplast retrograde signalling has recently been comprehensively reviewed (Chan *et al.*, 2016; de Souza *et al.*, 2017), including possible roles for metabolites in immune signalling (Fernandez & Burch-Smith, 2019) and hence this is not addressed here, other than to highlight specific examples.

We have tried to illustrate a variety of key immune processes that impact the chloroplast throughout the review. A powerful technique to visualize the impact of pathogens on chloroplast physiology is through F_v/F_m measured by Chl fluorescence imaging (Baker, 2008). F_v/F_m provides sensitive quantitative temporal-spatial measurements of changes in the maximum (dark-adapted) quantum efficiency of photosystem II (PSII; a sensitive indicator of damage/downregulation of photosynthesis), while simultaneously enabling imaging of pathogen challenges in real time and, thus, is increasingly being used to monitor pathogen infection dynamics (de Torres Zabala *et al.*, 2015). Fig. 1 illustrates suppression of PTI by the virulent phyto-bacterial pathogen *Pseudomonas syringae* pv. *tomato* strain DC3000 (*Pst*) both visually (Fig. 1a) and quantitatively (Fig. 1b) and its relationship to *in planta* bacterial multiplication (Fig. 1c). F_v/F_m can also effectively capture changes in chloroplast physiology caused by fungal challenges (Fig. 1d).




Fig. 1 Photosystem II quantum efficiency (F_v/F_m) captures early chloroplast changes in response to virulent and avirulent pathogens. (a,b) Challenge with the virulent apoplasmic bacterial phytopathogen *P. syringae* pv. *tomato* DC3000 (*Pst*) but not mock ($MgCl_2$) or the disarmed *hrpA* mutant results in reduced F_v/F_m 7–8 hpi (h post-infection) as illustrated visually (a) or quantitatively during disease establishment (b). (c) *Pst* multiplication significantly increase above initial inoculation levels at 8 h post-infiltration, coincident with reduction in F_v/F_m . Error bars, \pm SD. (d) Spray infection with spores of the virulent rice pathogen *Magnaporthe oryzae* Guy11 similarly induces localized decreases in F_v/F_m . (e) Challenge with the vascular pathogen *Xanthomonas campestris* pv. *campestris* (*Xcc*) or *P. syringae* pv. *maculicola* suppresses F_v/F_m during infection, the extent of which is directly correlated with virulence of the strains. Pretreatment with ABA, which is rapidly induced *de novo* following virulent bacterial infections, dramatically enhances suppression of F_v/F_m in both *Xcc* and *Pst*. (f) ETI induced either by RPM1 or RPS4 following challenge with *Pst* carrying the respective avirulence genes, *AvrRpm1* or *AvrRps4*, causes a rapid suppression of F_v/F_m , the timing of which is unique to the specific R protein and correlates with speed of HR development. Kindly provided by: (a–c, f) M. Grant & S. Breen; (d) G. Littlejohn; (e) de Torres *et al.* (2015: Fig. S4).

III. Hormones and chloroplasts, a well-established link in plant immunity

Being the main site of phytohormone precursor synthesis, the chloroplast is central to integrating signals from PTI and ETI and an obvious target for effector modulation. Hormonal crosstalk in plant–microbe interactions is now well established (Robert-Seilaniantz *et al.*, 2011; Burger & Chory, 2019). Thus, here we only briefly overview the core roles of, or selected new insights into, the three key immunity modulating hormones salicylic acid (SA), jasmonic acid (JA) and abscisic acid (ABA) to provide context for further reference in later sections.

1. Salicylic acid

Salicylic acid (SA) is the archetypal defence hormone effective against biotrophic and hemibiotrophic pathogens (Ding & Ding, 2020). Rapid SA biosynthesis in response to pathogens occurs through formation of isochorismate by the chloroplast-localized isochorismate synthase (ICS). The decades long challenge to understand how isochorismate was converted to SA was recently resolved. EDS5 (ENHANCED DISEASE SUSCEPTIBILITY 5) exports isochorismate to the cytosol where the amidotransferase PBS3 (*avrPphB* SUSCEPTIBLE3), originally identified in a genetic screen for loss of RPS5-specified resistance (Warren *et al.*, 1999), catalyses its conjugation to glutamate, forming isochorismate-9-glutamate, which spontaneously decomposes into SA (Rekhter *et al.*, 2019; Torrens-Spence *et al.*, 2019). SA may directly interfere with pathogen virulence strategies through its interaction with nonexpressor of pathogenesis-related gene (NPR) SA receptors. Increases in SA inhibit the transcriptional corepressors NPR3 and NPR4, but activate the transcriptional coactivator NPR1, collectively inducing SA responsive defence genes, including key regulators of plant immunity (Ding *et al.*, 2018). SA can also function indirectly by inhibiting ROS scavenging enzymes such as catalase and ascorbate peroxidases (Durner & Klessig, 1995; Zhang *et al.*, 2016). More recently, SA was proposed as a retrograde signal generated by impaired PSII proteostasis (Duan *et al.*, 2019), although whether biotic stress leads to sufficient accumulation of photodamaged proteins to instigate SA retrograde signalling remains to be demonstrated.

2. Jasmonates

Classically, jasmonates are associated with core biotrophic pathogen virulence strategies to suppress SA signalling. JA also acts synergistically with ethylene in defence against necrotrophic pathogens and ABA during herbivory (Robert-Seilaniantz *et al.*, 2011; Zhang *et al.*, 2017; Yang *et al.*, 2019). Linolenic and linoleic acid, derived from chloroplast galactolipids, provide the 18-carbon fatty acid substrate which is oxidized at the C-13 position by chloroplast lipoxygenase then cyclized to 12-oxo-phytodienoic acid (OPDA) via the consecutive activities of allene oxide synthase and allene oxide cyclase. OPDA is exported to the peroxisome where it is converted, *via* a series of beta oxidation steps, to JA which is conjugated to isoleucine to form bioactive JA-Ile. JA may undergo alternative modifications, although biological understanding of

their significance is currently limited (Wasternack & Hause, 2013). With a predominant focus on jasmonate antagonism of biotrophic defences, it is often overlooked that jasmonates are also produced *de novo* during ETI (Andersson *et al.*, 2006; Zoeller *et al.*, 2012) and have been implicated in both SAR (Truman *et al.*, 2007) and induced systemic resistance (ISR) (van Wees *et al.*, 2000).

3. The role of ABA in repressing chloroplast immunity

While early studies revealed that ABA treatment suppressed resistance to biotrophic and hemibiotrophic bacterial, fungal and oomycete pathogens (Henfling *et al.*, 1980; Mohr & Cahill, 2003), it was not until subsequent, transcriptomic and genetic studies with ABA biosynthetic and signalling mutants demonstrating that pathogens hijack host ABA signalling to promote virulence that ABA became universally recognized as a key player in suppression of biotrophic immunity. *De novo* ABA synthesis induced by virulent *Pst* is remarkably rapid, occurring within 6 h of challenge, significantly preceding bacterial multiplication (de Torres-Zabala *et al.*, 2007, 2009). Pathogen-induced ABA requires transcriptional upregulation of genes encoding the chloroplast-localized 9-*cis*-epoxycarotenoid dioxygenase (NCED) and cytosolic abscisic aldehyde oxidase (AAO) – key enzymes in the final steps of ABA biosynthesis (Truman *et al.*, 2006; de Torres-Zabala *et al.*, 2007, 2009; Peng *et al.*, 2019). Concomitantly, transcripts encoding protein phosphatase 2Cs (PP2C), negative regulators of ABA signalling, are suppressed (Truman *et al.*, 2006) (see Fig. 4c later for a summary).

Carotenoid intermediates provide the precursors for ABA biosynthesis. Zeaxanthin, derived from β -carotene – whose oxidation products are themselves potential chloroplast signalling molecules (reviewed by Havaux, 2014) – is converted to violaxanthin, and then via *trans*-neoxanthin into 9'-*cis*-neoxanthin and 9'-*cis*-violaxanthin. These substrates are converted by NCED to the 15-carbon xanthoxin which is transported to the cytosol where it is converted into abscisic aldehyde and finally to ABA via AAO (Seo & Koshiba, 2002). *De novo* ABA induced by *Pst* is proposed to suppress PTI-induced cROS (de Torres Zabala *et al.*, 2015) as well as antagonizing later SA signalling (de Torres Zabala *et al.*, 2009). ABA biosynthetic mutants (*aa3*) are more resistant to *Pst* and other biotrophic pathogens. Notably, pretreatment with ABA abolished PTI, enhanced the decrease of F_v/F_m (Fig. 1e) and, analogous to ABA suppression of ROS in imbibed seeds (Ye *et al.*, 2012), induced ROS (de Torres Zabala *et al.*, 2015). As ABA can repress transcription of many plastid genes through PP2C-dependent activation of nuclear genes (Yamburenko *et al.*, 2015), the recent demonstration that *Xanthomonas* effectors of both rice and *Arabidopsis* pathogens promote virulence by suppressing transcripts encoding chloroplast-localized PP2Cs (Akimoto-Tomiyama *et al.*, 2018) reinforces the complex role of ABA in effector modulation of chloroplast immunity.

IV. cROS in immunity and insights from disruption of chloroplast components

Specificity in ROS signalling is achieved via the spatiotemporal control of production and scavenging at different organellar and

subcellular locations. During plant defence, recognition of PAMPs by PRRs activates plasma membrane-localized NADPH oxidase (Zhou *et al.*, 2019) and apoplastic type III peroxidases (Daudi *et al.*, 2012) generating, within minutes, a rapid burst of H₂O₂ comprising synthesis of short-lived superoxide and its more stable dismutation product hydrogen peroxide in the apoplast (Smirnov & Arnaud, 2019). Hydrogen peroxide can enter the cytosol *via* plasma membrane aquaporins (Rodrigues *et al.*, 2017). ROS are also produced in organelles by oxygen reduction during electron transport and by oxidase enzymes in peroxisomes (Asada, 2006; Mullineaux *et al.*, 2018; Waszczak *et al.*, 2018; Smirnov & Arnaud, 2019). The prominent routes for cROS generation are oxygen photoreduction at PSI (Mehler reaction) and possibly via the PSII electron acceptor plastoquinone (Dietz *et al.*, 2016; Vetoshkina *et al.*, 2017). Singlet oxygen (¹O₂), a highly reactive species, is formed in PSII by transfer of excitation energy from triplet-state Chl (Mullineaux *et al.*, 2018a; Dogra *et al.*, 2019) and is the major ROS involved in ETI-induced lipid peroxidation (Zoeller *et al.*, 2012).

1. PTI and cROS

As part of PTI, chloroplasts of *Arabidopsis* leaves challenged with virulent *Pst* generate reactive species (as determined by 2′7′-dichlorodihydrofluorescein diacetate (H₂DCFDA) oxidation) which are suppressed within 4 h by *Pst* effectors (de Torres Zabala *et al.*, 2015). DCMU (3-(3,4-dichlorophenyl)-1,1-dimethylurea), which blocks photosynthetic electron transport between PSII and plastoquinone (Metz *et al.*, 1986), also blocks 2′7′-dichlorodihydrofluorescein oxidation, indicating that this burst is probably generated by oxygen photoreduction producing superoxide/H₂O₂ downstream of PSII (Mubarakshina *et al.*, 2010; Exposito-Rodriguez *et al.*, 2017). Interestingly, *Pst* ROS suppression coincides with a decrease in F_v/F_m (Fig. 1a,b) and photosynthesis (de Torres Zabala *et al.*, 2015) and an increase in bacterial growth (Fig. 1c) indicating that effectors (some of which are targeted to the chloroplast) interfere with critical photosynthetic components that have yet to be identified. Notably, ABA mimics DCMU application, suggesting that *Pst*-induced *de novo* ABA biosynthesis may play a key role in suppressing ROS production. Indeed, pretreatment of leaves with ABA strongly enhances the *Pst*-induced decrease in F_v/F_m and this is common to other, less virulent pathogens such as *Xanthomonas campestris* pv. *campestris* (Fig. 1e). At the same time, *Pst* (and other pathogens) suppress the expression of a large set of nuclear-encoded chloroplast genes including photosynthesis-related and antioxidant enzyme transcripts (Bilgin *et al.*, 2010; de Torres Zabala *et al.*, 2015; Su *et al.*, 2018). The signalling mechanism driving PTI-generated cROS is unclear but may involve calcium and/or retrograde signalling as discussed below.

2. cROS and ETI

The interaction between high light, phytochrome and pathogen responses has been documented (Bechtold *et al.*, 2005; Ballare, 2014). Light is required for, or enhances, ETI-triggered HR

(Torres *et al.*, 2006; Nomura *et al.*, 2012). These observations suggest the interaction of cROS with photosynthesis, SA production (Chaouch *et al.*, 2010, 2012) and additionally NO (Zaninotto *et al.*, 2006; Yun *et al.*, 2011; Yun *et al.*, 2016). The development of an HR is rapid and effectively contains the pathogen. The HR is widely thought to be triggered by ¹O₂ generation, which leads to lipid peroxidation (Havaux, 2014). Pioneering analytical studies of the temporal accumulation of oxidation products derived from unsaturated fatty acids during the HR strongly support a ¹O₂ burst. Notably, the HR leads to an early and massive accumulation of both enzymatic and nonenzymatic chloroplast galactolipid-derived oxylipins (Andersson *et al.*, 2006; Zoeller *et al.*, 2012) with huge increases in JA measured within 5 h of infection with *PstavrRpm1* (Zoeller *et al.*, 2012). This timing is consistent with the earlier biophoton production following *PstavrRpm1* challenge (Bennett *et al.*, 2005), which is indicative of lipid oxidation (Havaux *et al.*, 2006). For example, HR in *Arabidopsis* inoculated with *PstavrRpm1* is enhanced by increased light intensity and associated with disruption of the PSII light harvesting complex, decreased F_v/F_m (Fig. 1f) and accumulation of the Chl catabolite pheophorbide, a potent photosensitizer that generates ¹O₂ (Mur *et al.*, 2010). Indeed, F_v/F_m provides a powerful readout to accurately capture and quantify the timing of specific R protein activation, before visible symptoms, as illustrated for the RPM1–AvrRpm1 and RPS4–AvrRPS4 interactions (Fig. 1f; for a recent review see also Perez-Bueno *et al.*, 2019).

Further evidence that chloroplast-sourced ROS are involved in ETI and mediated by MAPK pathways are provided by studies in *Nicotiana benthamiana* (Liu *et al.*, 2007) and *Arabidopsis* (Su *et al.*, 2018). *PstavrRpt2* (like *PstavrRpm1*, Fig. 1f) causes a much larger and earlier decrease in PSII quantum efficiency than *Pst* (Fig. 1a) (de Torres Zabala *et al.*, 2015), and a more prolonged activation of MAPKs (Su *et al.*, 2018). This response is mirrored by conditional induction of MAPKs (MPK3/6) leading to cell death. Both *PstavrRpt2* and MAPK activation increase cROS within 6 h in a light-dependent manner (consistent with biophoton generation at *c.* 7 h post-inoculation (hpi); Bennett *et al.*, 2005), accompanied by visible disruption of PSII. Comparison of apparent chloroplast-sourced ROS between PTI and ETI in this system indicates that chloroplast-targeted effectors decrease photosynthesis and suppress cROS production (de Torres Zabala *et al.*, 2015) whereas ETI involves a more aggressive effect on photosynthesis, as illustrated by rapid decreases in F_v/F_m and an increase in ROS (Su *et al.*, 2018), consistent with extensive chloroplast galactolipid oxidation recorded during early ETI (Andersson *et al.*, 2006; Zoeller *et al.*, 2012).

3. A role for cROS in systemic immunity?

The chloroplast is becoming increasingly linked to effective SAR, a process conferring broad-spectrum and lasting immunity to pathogens of diverse lifestyles (Fernandez & Burch-Smith, 2019). Classic SAR is established following successful ETI leading to the HR. Chloroplast lipids and cROS appear to be central to generation of SAR inducing signal(s) following ETI-activated HR (Wendehenne *et al.*, 2014; Shine *et al.*, 2019), supported by the SAR-deficient phenotypes of fatty acid desaturase (*sfd2*;

SUPPRESSOR OF FATTY ACID DESATURASE DEFICIENCY2) mutants. The chloroplast galactolipid mutants *mgd1* and *dgd1* (*monogalactosyl synthase 1*, *digalactosyl synthase 1*), responsible for monogalactosyldiacylglycerol (MGDG) and digalactosyldiacylglycerol (DGDG) synthesis respectively, function nonredundantly in initial SAR signal perception (Gao *et al.*, 2014; Shah *et al.*, 2014). As noted above, ETI-generated ROS modify fatty acids on chloroplast galactolipids leading to rapid JA accumulation (Andersson *et al.*, 2006; Zoeller *et al.*, 2012). Although JA is classically associated with suppression of SA signalling in biotrophic interactions, jasmonates have been implicated in both SAR (Truman *et al.*, 2007; Liu *et al.*, 2016) and ISR (van Wees *et al.*, 2000). While somewhat speculative, interesting parallels are emerging between the role of ROS in signalling pathways that regulate SAR and systemic acquired acclimation (SAA) and wounding, where a wave of ROS signalling appears to be a common early mediator of systemic signalling responses although on different time scales (Zandalinas *et al.*, 2019).

V. Immunity insights from perturbation of chloroplast metabolism and cROS production

The following sections describe components of the photosynthetic electron transport system and the main sources of ROS occurring in the chloroplast. The reader is referred to the schematic in Fig. 2 for context. The removal of H₂O₂ in chloroplasts is carried out by a diverse set of enzymes providing robustness to PTI. These include ascorbate peroxidases localized to the stroma or attached to the thylakoid membrane along with glutathione and associated enzymes to regenerate oxidized ascorbate: glutathione peroxidase-like and peroxiredoxins (Smirnoff & Arnaud, 2019) (Fig. 2). Oxidized peroxiredoxins are regenerated by thioredoxin with involvement of NADPH-dependent thioredoxin reductase C (NTRC) (Perez-Ruiz *et al.*, 2017). While mutants of individual peroxiredoxins (Prx) in *Arabidopsis* (there are four chloroplast Prx isoforms in *Arabidopsis*; Tripathi *et al.*, 2009), have normal *Pst* responses, an NTRC mutant (*ntrc*) shows increased cell death and increased peroxide production as determined by 3,3'-diaminobenzidine staining (Ishiga *et al.*, 2011) but no difference in *Pst* growth compared to wild-type Col-0. Interestingly, the authors also showed that NTRC-silenced tomato plants showed accelerated necrotic cell death and enhanced symptom development in response to the necrotrophic soil pathogen *Sclerotinia sclerotiorum*. A similar response was elicited by nonhost *P. syringae*, although pathovars varied in specific responses (Ishiga *et al.*, 2016). Notably, these symptoms were absent in plants inoculated with a coronatine (COR)-deficient *Pst* strain, implicating a role for COR in cROS-induced disease-associated necrosis (Ishiga *et al.*, 2016). Antisense knockdown of two chloroplast GPX-like enzymes in *Arabidopsis* increases H₂O₂ and high light-induced SA. These plants had elevated PTI to *Pst* and *P. syringae* pv. *maculicola* (*Psm*), and more extensive HR following *PstavrRpm1*-initiated ETI (Chang *et al.*, 2009). Manipulation of chloroplast APX also impacts pathogen response. Rice lines overexpressing thylakoid membrane-bound APX exhibited increased initial tolerance to rice bacterial blight conferred by *Xanthomonas oryzae*

pv. *oryzae*, whereas RNAi lines were more susceptible, and this was correlated with H₂O₂ levels, presumably chloroplast-derived (Jiang *et al.*, 2016). In *Arabidopsis*, conditional silencing of thylakoid-bound APX showed that accumulation of chloroplastic H₂O₂ triggered retrograde signalling leading to induction of nuclear-encoded pathogen defence genes in the absence of any pathogen challenge (Maruta *et al.*, 2012). While not confined to chloroplasts, the concentration of the antioxidants ascorbate and glutathione, which are involved in H₂O₂ removal and redox regulation, influence pathogen responses. Ascorbate-deficient mutants have increased H₂O₂, PR levels, camalexin and SA accumulation and have increased basal resistance to *Pst* and the oomycete *Hyaloperonospora* (Barth *et al.*, 2004; Pavet *et al.*, 2005; Colville & Smirnoff, 2008; Mukherjee *et al.*, 2010). Consistent with these observations, glutathione-deficient mutants have decreased resistance to *PstavrRpm1* (Ball *et al.*, 2004; Parisy *et al.*, 2007).

Expressing the cyanobacterial electron transport protein flavodoxin in tobacco chloroplasts improves robustness of photosynthesis to various stresses including methyl viologen (MV; a redox cycling compound that generates superoxide at PSI) and high light. This appears to be associated with decreased cROS production (Tognetti *et al.*, 2006) and altered pathogen responses (Zurbriggen *et al.*, 2009; Rossi *et al.*, 2017). The reason that flavodoxin, which has a flavin cofactor, improves stress resistance and decreases cROS production is not immediately apparent. It functionally replaces the plant PSI electron acceptor ferredoxin (Tognetti *et al.*, 2006) which has a 2Fe–2S reaction centre. One possibility is that electron transport through flavodoxin decreases oxygen photoreduction at PSI (the Mehler reaction) (Fig. 2). Alternatively, because Fe–S proteins are a target for superoxide and H₂O₂, which can demetallate them (Imlay, 2013), chloroplastic ferredoxin may be sensitive to inactivation by ROS. Indeed, superoxide inactivates spinach ferredoxin (Fisher *et al.*, 2016), consistent with the marked increase in resistance to MV (Tognetti *et al.*, 2006). This may account for the significant reduction in localized cell death induced by the nonhost pathogen *Xanthomonas campestris* pv. *vesicatoria* (*Xcv*) in flavodoxin-expressing tobacco leaves, which was associated with decreased cROS production (Zurbriggen *et al.*, 2009). Similarly, infection of flavodoxin-overexpressing tobacco with the necrotrophic fungus *Botrytis cinerea* significantly restricted hyphal growth, lesion development, *Pathogenesis Related* (*PR*) gene expression and phytoalexin accumulation (Rossi *et al.*, 2017). Expression of flavodoxin in *Arabidopsis* chloroplasts decreases ROS production and disassembly of PSII in response to *PstavrRpt2*, attenuating ETI (Su *et al.*, 2018). These studies highlight a central role for cROS in effective PTI and ETI. A mutant in the main chloroplast ferredoxin (*fd2*; Fig. 2) exhibiting altered pathogen responses provides additional evidence linking electron transport from PSI with PTI (Wang *et al.*, 2018). *fd2* was more susceptible to *Pst*, possibly as a direct result of the elevated JA observed. By contrast, ETI elicited by *AvrRpt2* was stronger, with twice as much H₂O₂ generation. This result is part of a growing body of evidence for possible photosystem-specific roles for ROS generation during ETI and PTI, with ROS generated by ETI being primarily derived from PSII whereas PTI may generally require electron transport to PSI, which is compromised in *fd2* plants. This is also consistent




Fig. 2 An outline of the photosynthetic electron transport system showing the main sources of reactive oxygen species (ROS; singlet oxygen ($^1\text{O}_2$), superoxide (O_2^-) and hydrogen peroxide (H_2O_2)) at photosystems I and II. Proteins that are validated effector targets are shown in red. APX, ascorbate peroxidase; 2-Cys Prx, 2-cysteine peroxidase; Fd, ferredoxin; FNR, ferredoxin-NADPH reductase; GPX, glutathione peroxidase-like; NTRC, NADPH-dependent thioredoxin reductase; PQ, plastoquinone; PC, plastocyanin; SOD, superoxide dismutase; Trx, thioredoxin.

with PSI being the source of H_2O_2 for PTI (de Torres Zabala *et al.*, 2015).

In conclusion, various lines of evidence show that cROS is induced by PTI and ETI, and we speculate that H_2O_2 derived from PSI may be the primary ROS underpinning PTI whereas ETI elicits rapid accumulation of $^1\text{O}_2$ (Fig. 2). The intensity and duration of the response, and hence the eventual pathogenic outcome is dictated by a complex interaction, its outcome being dictated by the specific pathogen virulence strategy and host resistance protein complement. Higher H_2O_2 levels can improve basal immunity, but effectors collaborate to directly or indirectly repress cROS production, probably by inhibiting electron transport to PSI. By contrast, and somewhat counterintuitively, ETI appears to elicit an extensive disruption of photosynthesis, including breakdown of PSII leading to greater ROS production and the HR. This is likely to be driven by $^1\text{O}_2$. However, at this point a fuller understanding of these mechanisms is limited by the poor specificity of the ROS assays (Smirnov & Arnaud, 2019). New genetically encoded reporters (Nietzel *et al.*, 2019) and nanosensors (Lew *et al.*, 2020) offer better specificity and temporal spatial resolution to better dissect these processes.

VI. Direct targeting of pathogen effectors to the chloroplast

The previous sections show that pathogen effectors modulate chloroplast function, either directly or indirectly, which implies

that effectors may themselves localize to the chloroplast and directly interact with chloroplast-located targets. Here we summarize direct and indirect experimental evidence for effector localization to the chloroplast.

1. Bacterial effectors

Evidence for physical targeting of chloroplasts by bacterial effectors did not emerge until the mid-2000s (Jelenska *et al.*, 2007; Lee *et al.*, 2008) yet remarkably and more than 10 of *Pseudomonas syringae*'s core effector repertoire of 30–40 proteins have been predicted or experimentally shown to localize to the chloroplast (Table 1). More recently, a number of effector proteins from *Ralstonia solanocera* have been shown to localize to the chloroplast (Table 1), although knowledge of their host targets is limited (Jelenska *et al.*, 2007; Lee *et al.*, 2008; Rodriguez-Herva *et al.*, 2012).

Pseudomonas syringae The recently described *P. syringae* pan genome (Laflamme *et al.*, 2020) has provided a rich resource to further expand our knowledge of effectors targeted to the chloroplast.

HopI1 One of the first bacterial effectors found to target the chloroplast was HopI1 from *Psm*. HopI1 has a redundant chloroplast-targeting sequence and contains P/Q-rich repeats (to facilitate protein folding) and a J domain, through which it directly

Table 1 Pathogen effectors predicted to be chloroplast localised.

Effector	Origin species	Pathovar	Localisation	Experimental method for localisation	Chloroplastic Interacting partner	Interacting partner identification	ETI/ETS	Reference
Bacteria								
HopI1	<i>Pseudomonas syringae</i>	<i>maculicola</i>	Chloroplast	Transgenic Arabidopsis, chloroplast fract, c onfocal in transient Nb	HSP70	Yeast complementation	ETS	Jelenska et al. (2007)
AvrRps4	<i>P. syringae</i>	<i>pisi</i>	Chloroplast, Nucleus	GFP fusion			Extachloro-plastic ETI	Li et al. (2014)
HopK1	<i>P. syringae</i>	<i>tomato</i>	Cytoplasm	transient in Nb			ETS	Li et al. (2014)
HopO1-2	<i>P. syringae</i>	<i>tomato</i>	Chloroplast, Nucleus	GFP fusion				
	<i>P. syringae</i>	<i>tomato</i>	Cytoplasm	transient in Nb				
	<i>P. syringae</i>	<i>tomato</i>	Chloroplast	Import assay (isolated pea chloroplasts)			ETS	de Torres Zabala et al. (2015)
HopR1	<i>P. syringae</i>	<i>tomato</i>	Chloroplast	Import assay (isolated pea chloroplasts)	PTF1, CBSX2	Y2H	ETI -transient expression in Nb	Mukhtar et al. (2011), de Torres Zabala et al. (2015)
HopN1	<i>P. syringae</i>	<i>tomato</i>	Chloroplast	GFP fusion	PsbQ	<i>in vitro</i> pull-down	ETS	Rodriguez-Herva et al. (2012)
HopBB1	<i>P. syringae</i>	<i>tomato</i>	Chloroplast	transient in Nb	PTF1	Y2H		Mukhtar et al. (2011)
HopM1	<i>P. syringae</i>	<i>actinidiae</i>	Chloroplast	Predicted – <i>in silico</i> YFP fusion				Choi et al. (2017)
HopU1	<i>P. syringae</i>	<i>tomato</i>	Chloroplast	transient in Nb	RBP31 plus two additional chloroplast RNA-binding proteins	Y2H		Mukhtar et al. (2011)
HopZ1	<i>P. syringae</i>	<i>tomato</i>	Chloroplast		AT4G39050, AT3G11590, AT3G07780	Y2H		Mukhtar et al. (2011)
RipAL	<i>Ralstonia solanacearum</i>		Chloroplast	GFP fusion				Nakano & Mukaihara (2018)
RipAD	<i>R. solanacearum</i>		Chloroplast	transient in Nb				Jeon et al. (2020)
RipG3, RipG7	<i>R. solanacearum</i>		Chloroplast	YFP fusion				
Fungi								
ToxA	<i>Pyrenophora tritici-repentis</i>		Chloroplast	transient in Nb	Nbcab13, NbrbcX, NbrbcS	Y2H		Dahal et al. (2018)
CTP1, CTP2, MLP124111	<i>Melampsora larici-populina</i>		Chloroplast	GFP fusion	ToxABP1	Y2H	ETS	Manning et al. (2007), Sperschneider et al. (2017)
CTP3	<i>M. lini</i>		Chloroplast	transient in Nb	Thf1			Petre et al. (2015), Petre et al. (2016)
			Chloroplast	GFP fusion				Petre et al. (2015), Petre et al. (2016)

Table 1 (Continued)

Effector	Origin species	Pathovar	Localisation	Experimental method for localisation	Chloroplastic interacting partner	Interacting partner identification	ETI/ETS	Reference
PST03196, PST18220	<i>Puccinia striiformis</i> f. sp. <i>tritici</i>		Chloroplast	GFP fusion transient in Nb				Petre <i>et al.</i> (2016a,b)
PGTG_00164, PGTG_06076	<i>Puccinia graminis</i> f. sp. <i>tritici</i>		Chloroplast	GFP fusion transient in Nb				Sperschneider <i>et al.</i> (2017) Tang <i>et al.</i> (2020)
SsITL	<i>Sclerotinia sclerotiorum</i>		Chloroplast		CAS (calcium-sensing)			
Oomycete PhRXLR-C20 PhRXLR-C27*	<i>Plasmopara halstedii</i>		Chloroplast	YFP fusion transient in Nb				Pecrix <i>et al.</i> (2019)
PVXLR54, PVXLR61, PVXLR86, PVXLR161	<i>Plasmopara viticola</i>		Chloroplast, nucleus, mitochondria	Experimental				Liu <i>et al.</i> (2018)

*Chloroplast associated; CBS, CBS Domain containing protein; HSP70 Heat shock 70 protein; Nb, *Nicotiana benthamiana*; PTF1, PHYTOCHROME AND FLOWERING TIME; Nbrbc, ribulose biphosphate carboxylase small chain from *N. benthamiana*; Nbcab13, chlorophyll a-b binding protein 13 from *N. benthamiana*; Thf1, Thylakoid formation 1.

binds to and affects the activity and/or specificity of chloroplast-associated cytosolic Heat shock protein 70 (Hsp70). While *in planta* interaction with the Hsp70 chloroplast isomer has yet to be demonstrated, HopI1 induces altered thylakoid structure and reduced SA accumulation (Jelenska *et al.*, 2007), although how HopI1 enters the chloroplast remains to be determined.

HopN1 HopN1 suppresses ROS accumulation, callose deposition and HR cell death (López-Solanilla *et al.*, 2004; Rodriguez-Herva *et al.*, 2012), these activities being dependent on its cysteine protease activity. HopN1 localizes to the thylakoid membrane, interacting with and degrading PsbQ from PSII, reducing oxygen production, electron transport and attenuating cROS. Collectively, these studies have shown that PsbQ quantitatively contributed to both PTI and nonhost HR.

AvrRps4/HopK1 AvrRps4 is more commonly associated with triggering ETI when recognized by RPS4 in *A. thaliana*. However, AvrRps4 localizes to both the nucleus and the chloroplast and has high N-terminal sequence homology to another effector protein, HopK1. Both AvrRps4 and HopK1 target the chloroplast via a cleavable transit peptide (Li *et al.*, 2014) and their chloroplast localization is required to suppress the classical PTI responses, ROS production and callose deposition, and to enhance bacterial growth. The generation of combinations of chimeric effectors between C- and N-terminal domains of AvrRps4 and HopK1 demonstrated that AvrRps4 contributes to bacterial virulence in *Pst* lacking HopK1, although the chloroplast targets of these effectors remain to be determined. HopK1^N-AvrRps4^C but not AvrRps4^N-HopK1^C chimeras induced a strong HR delivered through *Pseudomonas fluorescens* (Halane *et al.*, 2018). However, AvrRps4^N not only directly interacted with EDS1 but also contributed to bacterial virulence in *Pst* lacking HopK1, establishing AvrRps4 as an evolved bipartite effector with dual nuclear and chloroplast functions (Halane *et al.*, 2018).

HopM1 *Pst* HopM1 localizes to the *trans*-Golgi network where it interacts with the ADP-ribosylation factor guanine nucleotide exchange factor, AtMIN7, to suppress vesicle-trafficking (Nomura *et al.*, 2011). However, HopM1 from *P. syringae* pv. *actinidiae*, with 67% amino acid identity to *Pst* HopM1, localizes to the chloroplast (Choi *et al.*, 2017), suggesting an intriguing evolution of alternative functions for these proteins.

Of the remaining *P. syringae* effectors that are known to localize to the chloroplast, the predicted ADP-ribosyl transferase HopO1-2 and HopR1 translocate into isolated chloroplasts (de Torres Zabala *et al.*, 2015) although further functional insight is lacking. HopR1 and HopO1-2 were amongst a number of effectors identified to interact with predicted chloroplast-localized proteins in yeast two-hybrid screens, including HopU1, HopZ1, HopW1 and HopBB1 (Lee *et al.*, 2008; Mukhtar *et al.*, 2011).

Other bacterial effectors Evidence for effectors targeting to the chloroplast is emerging from other bacterial pathogens. The chloroplastic phospholipase A1 RipAL (*Ralstonia*-injected proteins) from *Ralstonia solanacearum* (Nakano & Mukaihara, 2018)

shares homology with DEFECTIVE IN ANTHET DEHISCENCE1 (Ishiguro *et al.*, 2001), which catalyses the release of linoleic acid, a critical precursor of JA biosynthesis, from chloroplast membranes. RipAL localizes to the chloroplast and wild type, but not a lipase active site mutant, suppressed PTI in *N. benthamiana* via enhanced JA signalling and JA/JA-isoleucine content, with a concomitant decrease in SA and associated SA-signalling genes (Nakano & Mukaihara, 2018). The F-box domain RipG effector family comprises seven members, of which RipG3 and RipG7 interact with chloroplast proteins – possible targets for ubiquitination and proteasomal degradation (Dahal *et al.*, 2018). RipAD is also localized to chloroplasts, although its host target(s) remain unknown (Jeon *et al.*, 2020). Notably, both RipAL and RipAD interfere with flg22-triggered ROS production presumably from the chloroplast (Nakano & Mukaihara, 2018; Jeon *et al.*, 2020).

2. Effectors from filamentous pathogens

Chloroplast-localized effector proteins from fungi and oomycetes are now being identified, indicating that filamentous pathogens have also evolved to target the chloroplast (Table 1).

Rusts Transient expression in *N. benthamiana* has localized eight effector proteins from rusts (Table 1) (Petre *et al.*, 2015; Petre *et al.*, 2016; Sperschneider *et al.*, 2017). Notably, the program LOCALISER has proved useful for *in silico* prediction of chloroplast and other cellular effector addresses (Sperschneider *et al.*, 2017), identifying a

further two chloroplast-localized effectors from the biotrophic rust *Puccinia graminis* f. sp. *tritici*, PGTG_00164 and PGTG_06076, which were experimentally validated.

Given the dearth of experimentally validated chloroplast-localized effector proteins from other fungi, this may reflect a rust virulence strategy or lack of experimental endeavour.

Oomycete Oomycete effectors are largely of the ‘RXLR’ class. RXLRs are defined by a secretion signal peptide followed by a conserved N-terminal domain comprising the RXLR (Arg–Xaa–Leu–Arg) consensus sequence, where X is any amino acid that shares a conserved structural fold (Win *et al.*, 2012). A high-throughput screen of 83 candidate RXLR effectors of the obligate biotrophic oomycete *Plasmopara viticola* (Liu *et al.*, 2018) identified four effectors localized to the chloroplast (Table 1). Only one contained a cleavable N-terminal transit peptide and was specifically targeted to the chloroplast, PvRXLR86, whereas the others had multiple organellular addresses (Liu *et al.*, 2018). PvRXLR61 and PvRXLR161 localized to the chloroplast and nuclei whereas PvRXLR54 additionally targeted the mitochondria (Liu *et al.*, 2018). A chloroplast-localized effector was also identified from the related sunflower powdery mildew, *Plasmopara halstedii*. PhRXLR-C20, expressed during pathogen colonization, was observed in the chloroplast and stromules (Pecrix *et al.*, 2019). Notably, PhRXLR-C27 targeted plastid-associated membranes (Pecrix *et al.*, 2019). The host targets of these two effectors remain unknown.




Fig. 3 Convergent targeting of Thylakoid formation 1 (Thf1), a negative regulator of cell death, by diverse pathogens. Thf1 plays an important role in photosystem II (PSII) – light harvesting complex II dynamics and is targeted by necrotrophs, biotrophs and viruses. (a) The effector protein ToxA found in a variety of necrotrophic wheat fungal pathogens, *Parastagonospora nodorum* (*Pn*), *Pyrenophora tritici-repentis* (*Ptr*) and *Bipolaris sorokiniana* (*Bs*), targets the wheat Thf1 orthologue, ToxA Binding Protein 1 (ToxABP), inducing necrosis via ROS accumulation through reduction in PSI and PSII protein complex abundance. The wheat sensitivity protein, Tsn1, is required for ToxA-dependent necrosis and may monitor binding of ToxA to ToxABP1. (b) The hemibiotrophic bacterium *Pseudomonas syringae* pv. *tomato* (*Pst*) delivers effectors (yellow circles) which appear to disrupt Thf1 function, again leading to enhanced lesion formation, although it remains to be determined whether this is by direct or indirect interaction. (c) *The Tobamovirus* (TBV) N' virus resistance protein, belongs to the conserved Solanaceae I2 class of CC-NBS-LRR resistance protein. that also confers resistance to *Phytophthora* and *Fusarium* sp. TBV's CC domain physically targets and destabilizes TBV-coat protein in a light-dependent manner to enhance resistance. Based on analogy to the cyanobacterium *Synechocystis* Thf1 orthologue, Psb29, Thf1 destabilization affects accumulation of the FtsH ATP-dependent zinc metalloproteases, FTSH2 and FTSH5 (also known as VAR2 and VAR1 respectively), which are involved in the selective degradation of PSII subunits, such as D1 during PS repair. This would lead to PSII disassembly and increased ROS production.




Fig. 4 Modulation of calcium and reactive oxygen species (ROS) during suppression of PAMP-triggered immunity (PTI). (a) PTI initiates stromal Ca^{2+} spikes via mitogen-activated protein kinase (MAPK) activation of the chloroplast-localized CAS (calcium sensing protein), and these changes in $[\text{Ca}^{2+}]_{\text{chl}}$ are necessary for callose deposition and stomatal closure. *cas* mutants are compromised in resistance to both virulent and avirulent *Pseudomonas syringae* pv. *tomato* DC3000 (*Pst*). (b) The integrin-like effector SsITL (blue) from the soil fungal pathogen *Sclerotinia sclerotiorum* directly targets CAS to suppress immunity. Virulent *Pst* attenuates cROS by rapid induction of *de novo* ABA synthesis. *Pst* effector delivery rapidly induces expression of *NCED3* and *AAO3*, encoding key enzymes in ABA biosynthesis while concomitantly suppressing expression of the *PP2Cs*, encoding negative regulators of ABA signalling. ABA application suppresses PTI-induced chloroplastic ROS generation and enhances effector suppression of F_v/F_m . RipAL and RipAD, *Ralstonia solanacearum* effectors, also target the chloroplast and suppress cROS, although the mechanism remains to be determined. CAS, Ca^{2+} -sensing protein; NCED, 9-*cis*-epoxycarotenoid dioxygenase; AAO3, abscisic aldehyde oxidase 3; PP2C, protein phosphatase 2C; Rip, *Ralstonia*-injected protein; PQ, plastocyanin; SsITL, *Sclerotinia sclerotiorum* integrin-like protein; PS, photosystem; PS II, photosystem II; PS I, photosystem I (see Fig. 2 for more details).

Necrotrophic fungal effectors ToxA, a 178 amino acid secreted necrotrophic effector protein was first isolated from the fungus *Pyrenophora tritici-repentis* (Sarma *et al.*, 2005) and was more recently identified in *Parastagonospora nodorum* and *Bipolaris sorokiniana* (McDonald *et al.*, 2017). ToxA targets the chloroplast ToxA Binding Protein 1 (ToxABP1), inducing ROS accumulation through decrease in PSI and PSII protein complex abundance (Manning *et al.*, 2007; Faris *et al.*, 2010). The sensitivity in wheat to ToxA is governed by the *Tsn1* locus, encoding classical nucleotide binding, leucine rich repeat disease resistance proteins, suggesting these may monitor ToxA activity. The severity of necrosis can be restricted by preventing ROS accumulation or silencing ToxABP1 (Manning *et al.*, 2007). The *A. thaliana* homologue of the wheat ToxABP1, known as Thylakoid formation 1 (Thf1), is also targeted by multiple pathogens (see below), suggesting convergent evolution of effector targets. The *S. sclerotiorum* effector SsITL has recently been shown to localize to the chloroplast and interact with the chloroplast-localized calcium-sensing receptor (CAS, see below) (Tang *et al.*, 2020). The interaction of SsITL with CAS interferes with the SA signalling pathway to reduce SA accumulation during early infection while overexpression of CAS increased resistance to *S. sclerotiorum* (Tang *et al.*, 2020).

3. Convergent targeting of Thf1, a negative regulator of cell death, by diverse pathogens

Aside from being a target of ToxA, chloroplast-localized Thf1 is involved in a range of host–microbe interactions (necrotrophic,

biotrophic, viral), mediating both PTI and ETI (Fig. 3). Thf1 is an orthologue of ToxABP1 which binds ToxA (see above, Fig. 3a) and plays a central role in controlling PSII–light-harvesting complex II (LHCII) dynamics during dark-induced senescence and light acclimation (Huang *et al.*, 2013). It has also been linked to DC3000 virulence and virus infection (Fig. 3b,c). Both virus-induced gene-silenced *SLALC*, the tomato Thf1 orthologue, and *Arabidopsis thf1* mutants exhibited accelerated lesion formation upon DC3000 challenge, and *SLALC1* chloroplast localization was affected by coronatine (Wangdi *et al.*, 2010). Interestingly, Thf1 was additionally identified as an interactor with the CC domain of the Solanaceae I2-like class of CC-NLRs (Ori *et al.*, 1997), which provide immunity against a range of pathogens including *Fusarium oxysporum* f. sp. *lycopersici* (Hamel *et al.*, 2016), *Phytophthora infestans* in potato (Huang *et al.*, 2005) and *Tobamovirus* coat protein in pepper (Tomita *et al.*, 2011). Using *N'*, an I2 CC-NLR which recognizes *Tobamovirus* coat protein (Hamel *et al.*, 2016) demonstrated that Thf1 functions as a negative regulator of cell death, and activation of *N'* results in the destabilization of Thf1 in a light-dependent manner (Fig. 3c). Notably, like the TMV N protein interaction with chloroplast-localized NRIP protein (see below (Caplan *et al.*, 2008)), the *N'*–Thf1 interaction appears to take place in the cytosol. Possible insight into how Thf1 destabilization impacts chloroplast immunity is provided by the demonstration that a cyanobacterial Thf1 homologue Psb29 is required for the accumulation of the FtsH ATP-dependent zinc metalloproteases, which function in selective degradation of PSII subunits during repair (Beckova *et al.*, 2017). Normally, inactivation of PSII is restored through a repair cycle replacing damaged




Fig. 5 Physical responses of chloroplasts to pathogen infection. Stromule formation is common to both ETI and PTI, possibly providing a conduit of physical retrograde communication. (a) Confocal micrographs of tobacco N protein TMV p50-mediated ETI with chloroplasts visualized in N-containing NRIP1-Cerulean plants. Upper left: stromules wrapped around nuclei (maximum-intensity projection of a z stack). Upper right: direct connection to the nucleus of clusters of stromule tips (single z stack plane). Lower panels: nuclei with a mixture of tip or surrounding stromule connections (transparent projections of z stacks). Bars, 10 μm (from Caplan *et al.*, 2015). (b) Confocal images of reactive oxygen species (visualized by 2',7'-dichlorodihydrofluorescein diacetate [H_2DCFDA] staining) in nucleus and chloroplasts of leaf cells challenged with the nonvirulent *Pseudomonas syringae* pv. *tomato* (*Pst*) *hrpA* mutant eliciting PAMP-triggered immunity (PTI; left panel, bar 20 μm) or virulent *Pst* capturing effector-triggered susceptibility (ETS; right panel, bar 10 μm) visualized *c.* 5 h post-inoculation. White arrows denote chloroplasts sitting on the nucleus – both organelles show strong H_2DCFDA staining. Yellow arrow represents an H_2DCFDA F-stained chloroplast whose stromule is associated with the nucleus. Red fluorescence corresponds to Chl and green channel to the H_2DCFDA signal. Bars, 10 μm . (c) Compared with mock challenge (left panel) chloroplast aggregation is seen during *Pst* ETS in *A. thaliana* (18 h post-infection (hpi)). Red fluorescence signal is derived from Chl and green fluorescence from *Pst* labelled with GFP (adapted from Hutt *et al.*, 2014).

protein subunits, mainly the D1 reaction centre subunit, with functional copies. Damaged D1 repair is usually mediated through proteolysis by members of the Arabidopsis FTSH family. Thf1 is required for normal accumulation of FTSH2 and FTSH5 (also known as *VAR2* and *VARI* respectively; Wu *et al.*, 2013). Thus N' destabilization of Thf1 would diminish FTSH2/5 levels, impacting PSII repair, and lead to the production of ROS and presumably HR cell death (Fig. 3c).

4. Getting the message across: is calcium signalling involved?

There are common and distinct roles for ROS and calcium signalling in activating and uncoupling chloroplast immunity. Calcium signalling, like ROS signalling, is probably via a propagative wave, initiated at the plasma membrane upon PRR activation and transmitting to the chloroplast and nucleus, although current knowledge of this remains sparse. Twenty years ago, rapid transient cytosolic calcium ($\text{Ca}^{2+}_{\text{cyt}}$) increases in response to PTI (*Pst*, *PsthrpA* and *Pst avrRpm1* challenges) were recorded using the calcium-sensitive reporter aequorin (Grant *et al.*, 2000). *PstavrRpm1* (ETI) elicited an additional slow, sustained increase in $\text{Ca}^{2+}_{\text{cyt}}$, yet it is still unclear whether this is a signal perceived by other organelles, or indicative of loss of Ca^{2+} homeostasis coincident with HR development.

A role for calcium in establishment of chloroplast immunity is evidenced from studies on the thylakoid-membrane-localized Ca^{2+} -sensing protein (CAS), which generates stromal Ca^{2+} spikes via Ca^{2+} release from thylakoid membranes (Fig. 4a). The *cas-1* mutant was strongly compromised in resistance to virulent and avirulent *Pst* (Fig. 4b). Additionally, classical PTI responses such as callose deposition and stomatal closure were attenuated in *cas-1*. Biochemical characterization of CAS-silenced *N. benthamiana* plants positioned CAS downstream of activated MAPK signalling cascades and upstream of ROS signalling (Nomura *et al.*, 2012). Recently, the *S. sclerotiorum* integrin-like effector SsITL was shown to directly target CAS to suppress immunity (Fig. 4b) (Tang *et al.*, 2020). SsITL-expressing transgenic plants were more susceptible and CAS overexpression enhanced resistance to *S. sclerotiorum*, consistent with the previously reported role of SsITL in suppression of JA/ethylene signalling (Zhu *et al.*, 2013). Thus, stromal calcium signalling appears important in mediating broad-spectrum immunity.

VII. Cellular reorganization during infection, stromules and perinuclear chloroplast movement

Subcellular reorganization is well documented during plant–pathogen interactions. In addition to the generation of specialized interfaces between plant cells and invading pathogens (e.g. the extrahaustorial membrane (EHM) and biotrophic interfacial complex (BIC)), cellular components are recruited to sites of infection, often mediated by actin microfilaments or microtubules, as recently reviewed (Park *et al.*, 2018b; Boevink *et al.*, 2020). Chloroplasts move around the cell on actin microfilaments, but




Fig. 6 Are photosystems I and II specifically targeted in disease and defence? (a) Rapid PAMP triggered immunity (PTI) responses upon PAMP recognition activate MAPK signalling and transcription, with strong suppression of *NECGs* (*Nuclear Encoded Chloroplast Genes*) relative to mock treatment (c. 30–40% of all differentially expressed genes) within 2 h of challenge. Early chloroplast responses include calcium spiking induced by CAS and increased H_2O_2 generation at photosystem I (PSI) which may act as a retrograde signal. (b) Effector-triggered suppression of PTI (ETS) can occur by interfering with the pattern recognition receptor, at the coreceptor complex, through modulation of MAPK signalling, via transcriptional reprogramming or directly within the chloroplast, but probably a combination thereof. Early ETS responses include attenuation of MAPK signalling and reprogramming of *NECG* expression. In the chloroplast, calcium spiking and reactive oxygen species (ROS) generation are suppressed by either interference with photosystems themselves or electron transfer between PSII and PSI. Most direct effectors targets remain unidentified but virulence strategies of diverse pathogens include direct or indirect targeting of Thylakoid formation 1 (see Fig. 3). (c) During effector triggered immunity (ETI) many of these virulence processes are overridden, with effector recognition inducing a stronger and sustained activation of MAPK signalling, an increase in $[Ca^{2+}]_{cyt}$ and rapid collapse of the quantum efficiency of PSII (F_v/F_m), which is associated with an increase in 1O_2 generation at PSII. This results in lipid peroxidation and appears to be the catalyst for generation of local and systemic signalling molecules. Hop, Hrp-dependent *outer* protein; *NECG*, nuclear encoded chloroplast genes; PRR, pattern recognition receptor; BAK1, BRASSINOSTEROID INSENSITIVE 1-associated receptor kinase 1 – an exemplar PRR coreceptor; CAS, calcium sensing protein; PQ, plastocyanin. SAR, systemic acquired resistance; ISR, induced systemic resistance.

there is compelling evidence which shows that stromule formation (finger-like tubular stroma-filled chloroplast extensions) is mainly microtubule-dependent (Caplan *et al.*, 2015; Erickson *et al.*, 2018). However, treatment with the microtubule-depolymerizing agent oryzalin indicates additional microtubule-independent stromule formation with each type characterized also by its speed of movement (Erickson *et al.*, 2018).

1. Stromules and perinuclear chloroplast movement – ROS as a retrograde immune signal?

Systematic studies of chloroplasts during pathogen challenge are limited. Pioneering work on the TMV N resistance protein/TMV

p50 effector demonstrated a cytosolic interaction of N with chloroplastic localized N Receptor Interacting Protein 1 (NRIP1) (Caplan *et al.*, 2008). N-mediated ROS-induced stromules in a CHloroplast Unusual Positioning 1 (CHUP1)-dependent manner (Fig. 5a) (Caplan *et al.*, 2015). While stromules can be induced *in vitro*, indicating this is a chloroplast autonomous response, actin microfilament remodelling to facilitate perinuclear chloroplast movement appears to be an active ETI strategy to establish a conduit for possible retrograde ROS (or metabolite) signals (Fig. 5a) (Caplan *et al.*, 2015; Kumar *et al.*, 2018; Park *et al.*, 2018a; Park *et al.*, 2018b; Fernandez & Burch-Smith, 2019).

Stromules were observed following flg22 treatment, but not 20 hpi with *Pst*hrc (Caplan *et al.*, 2015). This apparent anomaly

may represent a timing issue as cROS is produced during early PTI (4–5 hpi with *PsthrpA*; de Torres Zabala *et al.*, 2015 (Fig. 5b)). Strikingly, PTI-induced cROS, as determined by H₂DCFDA staining, was only detected in perinuclear chloroplasts or those with stromules that appear to physically contact the nucleus following *PsthrpA* challenge. Interestingly, chloroplasts staining for ROS were significantly smaller than the others (Fig. 5b), suggesting heterogeneity in chloroplasts as reported for high light responses (Exposito-Rodriguez *et al.*, 2017), possibly a direct consequence of stromule formation. Additionally, there is evidence for chloroplast aggregation late in successful infections, as illustrated in Fig. 5(c) and described by Hutt *et al.* (2017). As many of these studies use different cell types (epidermal vs mesophyll), the importance of cell type on chloroplast function and chloroplast heterogeneity in specific pathogen immune responses requires further investigation. Chloroplasts also appear to be recruited to the EHM in *P. infestans* infections of *N. benthamiana*, where the anchoring of chloroplasts to the EHM is also CHUP1-mediated (Toufexi *et al.*, 2019). Silencing of *CHUP1* reduced chloroplast recruitment to the EHM, reduced stromule formation and led to higher levels of *P. infestans* hyphal growth, reinforcing the importance of CHUP1 and highlighting a role for chloroplast dynamics in establishment of plant immunity.

Thus, organization of chloroplasts during infection is typified by perinuclear chloroplast localization and the CHUP1-dependent extension of stromules toward the nucleus, each of which provide a physical basis for retrograde signalling (Erickson *et al.*, 2018; Mullineaux *et al.*, 2020). Indeed, perinuclear positioning of chloroplasts in immunity appears generic, being reported during viral infections (Fig. 5a) and in both avirulent (*Pst*), virulent (*PsthopQ1-1*) and *Agrobacterium tumefaciens* challenges of *N. benthamiana*, transient expression of effectors or viral proteins such as p50, or following exogenous application of ROS (Erickson *et al.*, 2014; Caplan *et al.*, 2015; Ding *et al.*, 2019). Pathogen effects on stromule formation and chloroplast–nuclear association is remarkably similar to cROS-mediated high light responses (Exposito-Rodriguez *et al.*, 2017) and oxidative stress imposed by silencing of *NTRC* (Brunkard *et al.*, 2015).

Recent evidence for effector suppression of stromules comes from studies with the *Xcv* E3 ubiquitin ligase effector XopL. Overexpression of XopL but not an XopL E3 ubiquitin ligase mutant in *N. benthamiana* abolished stromule formation in lower epidermal cells induced by *A. tumefaciens* (Erickson *et al.*, 2014; Erickson *et al.*, 2018). By contrast, XopQ, known to elicit ETI in *N. benthamiana*, increased stromule formation by over 50%. Notably, perinuclear chloroplast localization was still observed with XopL overexpression, implying nuclear recruitment of chloroplasts and formation of stromules to be independent mechanisms in immunity.

VIII. Functional significance of suppression nuclear-encoded chloroplast genes (*NECGs*)

While suppression of *NECGs* has been reported previously (e.g. Bilgin *et al.*, 2010), a detailed time course comparing *Pst* with its type III secretion-deficient *hrpA* mutant revealed that wholesale

suppression of *NECGs* was a PTI response, with c. 35% of all differentially suppressed genes within 3 hpi representing *NECGs* (de Torres Zabala *et al.*, 2015; Lewis *et al.*, 2015). This appears to indicate an active defence response to prioritize defence at the expense of growth. Notably, neither *hrpA* (nor *flg22*) challenge markedly affected Chl fluorescence parameters (de Torres Zabala *et al.*, 2015), yet within 3 hpi, *Pst* effectors differentially regulate a subset of *hrpA*-suppressed *NECGs* (Fig. 6). These transcriptional changes occur in parallel to suppression of cROS and before measurable differences in F_v/F_m or decrease in photosynthesis rate (de Torres Zabala *et al.*, 2015).

A meta-analysis of rice transcriptomic datasets also reported extensive downregulation of *NECGs* under both biotic and abiotic stress (Cohen & Leach, 2019). Considering the 11 diverse datasets and disparate temporal sampling, a core set of 85 photosynthesis-related genes were identified as suppressed across eight experiments. Thus, rapid transcriptional suppression of *NECGs* is a core response to retrograde stress signals, possibly representing a universal strategy to maximize resource allocation to defence by short-term attenuation of photosynthetic capacity, but possibly collaterally decreasing the capacity to repair effector targets.

Increasing evidence suggests that MAPKs mediate the transcriptional reprogramming of *NECGs*. MAPKs are rapidly activated following PAMP recognition and the subsequent apoplastic ROS burst (Meng & Zhang, 2013) (Fig. 6a). MAPKs can be induced by ROS but can themselves modulate ROS production. A body of evidence is emerging that the MPK3/MPK6 pathway also orchestrates ETI responses downstream of R protein activation that contribute to elevated cROS. Conditional activation of tobacco MPK3/6 orthologues SIPK/Ntf4/WIPK led to rapid, light-dependent suppression of CO₂ fixation, resulting in excess excitation energy, the generation of cROS and HR-like cell death (Liu *et al.*, 2007). ETI induced by constitutively active *Nicotiana tabacum* MAPK kinase 2 (NtMEK2_{DD}) led to sustained activation of MPK3/MPK6 in *Arabidopsis* (Fig. 6c). Similarly, conditional induction of *AvrRpt2* activated MPK3/MPK6, resulting in a rapid inhibition of PSII and accumulation of singlet oxygen and H₂O₂ in chloroplasts (Su *et al.*, 2018). How MAPKs impose specificity in modulating chloroplast immunity and transcriptional regulation of *NECGs* requires further investigation. It has recently been shown that fluctuating light activates local and systemic transcriptional reprogramming, including overrepresentation of genes involved in photoprotection, photosynthesis and photorespiration (Kumar *et al.*, 2018; Schneider *et al.*, 2019). Whether MPK6 integrates the retrograde signals to drive this adaptive, photoprotective response remains to be determined.

1. Emerging examples of indirect transcriptional modulation of ETI- and PTI-mediated chloroplast immunity

Here we review two examples of chloroplast immunity impacted by differential gene regulation. The first involves ETI activation of the *P. infestans* R protein Rpi-vnt1.1 by its effector AVRvnt1, which depends on light-driven alternative promoter selection. Light is required for expression of full-length tomato and potato *glycerate 3-kinase* (*GLYK*) transcripts encoding a chloroplast transit sequence.




Fig. 7 Differential regulation of nuclear encoded chloroplast genes impact immunity. (a) A chloroplast targeting sequence, encoded by light-driven alternative promoter selection of the solanaceous glycerate 3-kinase (GLYK) transcript, provides a cytosolic target for the *Phytophthora infestans* effector AVRvnt1, which is recognized by the Resistance to *P. infestans* protein Rpi-vnt1.1. AVRvnt1 binds to GLYK's chloroplast targeting sequence (CTP), intercepting its trafficking to chloroplast where it plays a role in basal and systemic immunity. Depletion of GLYK is indirectly sensed by Rpi-vnt1.1 activating effector triggered immunity (ETI). (b) Infection by the rice blast fungus *Magnaporthe oryzae* activates PAMP triggered immunity (PTI) in the rice variety *japonica* (but not *indica*). This is due to a light-dependent expression polymorphism in the promoter of *Light Harvesting Complex of Photosystem II 5* (*LHCBS5*). *LHCBS5* is cytoplasmically phosphorylated at Thr24 of its chloroplast transit sequence, leading to accumulation of both *LHCBS5* and superoxide in the chloroplast and enhanced basal immunity. Phosphorylated *LHCBS5* is predicted to form a trimeric complex, disrupting its binding to *PsbS*, which leads to reduced electron transfer, increased cROS and enhanced basal resistance. (c) *WKS1* (Wheat Kinase START 1), an atypical resistance protein encoded at the YR36 (Yellow Rust resistance) locus, confers partial race-nonspecific resistance to *P. striiformis* f. sp. *tritici*, is N-terminally processed and localizes to the chloroplast where it phosphorylates the thylakoid-associated ascorbate peroxidase (tAPX). *WKS1* also binds and phosphorylates the PSII component *PsbO*, decreasing its ability to bind to the supercomplex and resulting in a gradual accumulation of reactive oxygen species (ROS). SAR, systemic acquired resistance; HR, hypersensitive response.

AVRvnt1 binds to this chloroplast-targeting sequence and activates resistance (independent of GLYK kinase activity), impairing accumulation of GLYK in both total and chloroplast fractions of potato (Gao *et al.*, 2020). This is somewhat analogous to the TMV N–NRIP1 interaction described above (Caplan *et al.*, 2008), but in this case AVRvnt1 intercepts GLYK's trafficking to the chloroplast, the depletion of which (probably via proteasomal degradation) is indirectly sensed by Rpi-vnt1.1 activating ETI (Fig. 7a).

The second example requires both differential expression of the rice *Light Harvesting Complex of Photosystem II 5* (*LHCBS5*) and its light-dependent phosphorylation. During infection by the rice blast fungus *Magnaporthe oryzae*, *japonica* but not *indica* rice varieties show elevated PTI due to a simple nucleotide polymorphism in the *japonica LHCBS5* promoter leading to increased expression of *LHCBS5* (Liu *et al.*, 2019). Cytosolic phosphorylation

of *LHCBS5*'s chloroplast transit sequence on Thr24 leads to accumulation of both *LHCBS5* and superoxide in the chloroplast and enhanced basal immunity. Interestingly, *LHCBS5* was not phosphorylated during ETI. *LHCBS5* overexpression lines were more resistant, and RNAi knockdown lines were more susceptible, to *M. oryzae*. *LHCBS5* binds *PsbS*, a thylakoid sensor that is involved in nonphotochemical quenching (NPQ). As phosphorylated *LHCBS5* accumulating in the chloroplast can form a trimeric complex, the authors predicted that during *M. oryzae japonica* infection *PsbS* binding is disrupted, resulting in decreased electron transfer, increased cROS and enhanced basal resistance (Fig. 7b) (Liu *et al.*, 2019). This is supported by studies on *Arabidopsis* and rice plants deficient in *PsbS*, which have higher levels of cROS with rice mutants showing enhanced resistant to *M. oryzae* (Zulfugarov *et al.*, 2014). This may help mechanistically explain the results of

Gohre *et al.* (2012) where the observed flg22-induced decrease in PsbS abundance may be associated with increased cROS (Gohre *et al.*, 2012).

2. Direct targeting of 'resistance' proteins to the chloroplast

Given effector localization to the chloroplast, it is not unreasonable to propose classical R proteins to be associated with the chloroplast to monitor activity, and R proteins have been experimentally predicted to be chloroplast-associated (<http://suba.live/>). Indeed, the atypical chloroplast-localized Wheat Kinase START 1 (WKS1) confers partial race-nonspecific resistance to *P. striiformis* f. sp. *tritici* and is encoded at the *YR36* (*Yellow Rust resistance*) locus. N-terminally processed WKS1 localizes to the chloroplast, binding to and phosphorylating both thylakoid-associated ascorbate peroxidase (tAPX) potentially restricting cROS detoxification (Gou *et al.*, 2015) and PSII component PsbO, decreasing its ability to bind to the supercomplex (Wang *et al.*, 2019) (Fig. 7c). Both *psbo-A1* mutant and RNAi lines exhibited induction of chlorosis and reduced *P. striiformis* f. sp. *tritici* growth (Wang *et al.*, 2019). The authors concluded that WKS1 initially triggers chlorosis by phosphorylating PsbO, and the gradual accumulation of ROS (exacerbated by the phosphorylation of tAPX) induces cell death. It is currently unknown whether WKS1 is a target for *P. striiformis* f. sp. *tritici* effectors. Thus, the PSII supercomplex is emerging as a common effector target, as further evidenced by HopN1 targeting of PsbQ (Rodriguez-Herva *et al.*, 2012).

IX. Concluding remarks

Chloroplasts are a central hub in plant metabolism, enabling them to act as environmental sensors and communicate *via* a diversity of retrograde signals to the nucleus. It is now clear that chloroplasts play an essential role in plant immunity, and effectors from diverse pathogens have evolved to directly or indirectly target chloroplast function. At the biochemical level, the underlying mechanisms are complex, involving chloroplast-sourced oxylipins, hormones, hydrogen peroxide and singlet oxygen. An emerging theme is that PTI is associated with simultaneous repression of *NECGs* and induction of cROS, predominantly generated at PSI. Effector-mediated suppression includes modulating *NECGs*, and manipulating hormonal balance and various strategies to attenuate cROS *via* disassembly of the photosystems, although a detailed understanding of this remains elusive. Recent evidence suggests that plant resistance proteins can monitor perturbations to chloroplast homeostasis or recognize chloroplast-targeted effectors to activate ETI. Although further evidence is needed, it appears that in contrast to PTI, ETI drives $^1\text{O}_2$ generation *via* PSII disassembly, the resultant lipid oxidation products contributing to HR. Impairment of photosystem function is potentiated by chloroplast-targeted effectors, some of which have been shown to interact with components of the photosystems likely to affect their function and stability. Furthermore, pathogen infection elicits chloroplast repositioning and formation of stromules that might facilitate retrograde signalling. Not considered in this review, but equally important, are the interacting roles of NO and interorganelle

interactions with mitochondria (which have well-known roles in cell death) and peroxisomes.

Further challenges in this relatively embryonic field are multiple. We need to better understand the role of the multiple retrograde chloroplast to nucleus signalling pathways, in addition to ROS, which have been proposed to influence light response, and how these might interact with pathogens (Vogel *et al.*, 2014). Furthermore, in some cases ROS production could be a side reaction associated with other changes that comprise the actual signalling mechanism. This would not be easy to resolve but should be considered when assessing results. The challenge of understanding the relationship between the production of ROS by organelles and from the initial apoplastic PAMP-induced oxidative burst will require the use of probes with high spatial and chemical specificity. It will also require understanding the chloroplast targets manipulated by pathogens to suppress immunity.

Identifying the chloroplast targets of effectors and characterizing their interaction will not only provide important insight into how pathogens have evolved to target chloroplast immunity, but may potentially identify new herbicide leads. Aside from this, other particularly fundamental questions remain. Does the increasingly observed heterogeneity in size and positioning of chloroplasts in the cell reflect different metabolism and signalling roles in response to pathogens? How are PTI-induced cROS generated and how many chloroplasts need to respond to confer effective immunity? How many effectors, both in number and in diversity, need to target a (specific) chloroplast to suppress cROS? That being the case, do R proteins effectively guard chloroplasts?

Answers to these and other questions will not only contribute to fundamental understanding of chloroplast biology, but place the chloroplast at the forefront of endeavours to develop crops with improved pathogen resistance.

Acknowledgements

We thank Trupti Gaikwad (Warwick) and Naofel Aljafer and Dan Horton (Plymouth) for providing fluorescence images of *Arabidopsis* and *M. oryzae* infections of barley respectively. Marian Littlejohn assisted in drafting cartoons. Work related to this review is supported by ERDF Agri-tech Grant 05R16P00366 (GRL), and BBSRC/UKRI grants BB/P002560/1 (MG), BB/I020004/1 (NS), BB/P002366/1 (NS) and BB/N001311/1 (NS).

ORCID

Susan Breen  <https://orcid.org/0000-0001-5232-9973>
Murray Grant  <https://orcid.org/0000-0003-3357-0264>
George R. Littlejohn  <https://orcid.org/0000-0002-8768-2598>
Nicholas Smirnoff  <https://orcid.org/0000-0001-5630-5602>

References

Akimoto-Tomiya C, Tanabe S, Kajiwara H, Minami E, Ochiai H. 2018. Loss of chloroplast-localized protein phosphatase 2Cs in *Arabidopsis thaliana* leads to enhancement of plant immunity and resistance to *Xanthomonas campestris* pv. *campestris* infection. *Molecular Plant Pathology* 19: 1184–1195.

- Gerwick WH, Gobel C, Feussner I, Ellerstrom M. 2006. Oxylipin profiling of the hypersensitive response in *Arabidopsis thaliana*. Formation of a novel oxo-phytyldienoic acid-containing galactolipid, Arabidopside E. *Journal of Biological Chemistry* 281: 31528–31537.
- Asada K. 2006. Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiology* 141: 391–396.
- Baker NR. 2008. Chlorophyll fluorescence: a probe of photosynthesis *in vivo*. *Annual Review of Plant Biology* 59: 89–113.
- Ball L, Accotto GP, Bechtold U, Creissen G, Funck D, Jimenez A, Kular B, Leyland N, Mejia-Carranza J, Reynolds H *et al.* 2004. Evidence for a direct link between glutathione biosynthesis and stress defense gene expression in *Arabidopsis*. *Plant Cell* 16: 2448–2462.
- Ballare CL. 2014. Light regulation of plant defense. *Annual Review of Plant Biology* 65: 335–363.
- Banerjee N, Zaitlin M. 1992. Import of tobacco mosaic virus coat protein into intact chloroplasts *in vitro*. *MPMI* 5: 466–471.
- Barth C, Moeder W, Klessig DF, Conklin PL. 2004. The timing of senescence and response to pathogens is altered in the ascorbate-deficient *Arabidopsis* mutant *vitamin C-1*. *Plant Physiology* 134: 1784–1792.
- Bechtold U, Karpinski S, Mullineaux PM. 2005. The influence of the light environment and photosynthesis on oxidative signalling responses in plant-biotrophic pathogen interactions. *Plant, Cell & Environment* 28: 1046–1055.
- Beckova M, Gardian Z, Yu J, Konik P, Nixon PJ, Komenda J. 2017. Association of Psb28 and Psb27 proteins with PSII-PSI supercomplexes upon exposure of *Synechocystis* sp. PCC 6803 to high light. *Molecular Plant* 10: 62–72.
- Bennett M, Mehta M, Grant M. 2005. Biophoton imaging: a nondestructive method for assaying R gene responses. *Molecular Plant–Microbe Interactions* 18: 95–102.
- Bilgin DD, Zavala JA, Zhu J, Clough SJ, Ort DR, DeLucia EH. 2010. Biotic stress globally downregulates photosynthesis genes. *Plant, Cell & Environment* 33: 1597–1613.
- Boevink PC, Birch PR, Turnbull D, Whisson SC. 2020. Devastating intimacy: the cell biology of plant-*Phytophthora* interactions. *New Phytologist* 228: 445–458.
- Brunkard JO, Runkel AM, Zambryski PC. 2015. Chloroplasts extend stromules independently and in response to internal redox signals. *Proceedings of the National Academy of Sciences, USA* 112: 10044–10049.
- Burger M, Chory J. 2019. Stressed out about hormones: how plants orchestrate immunity. *Cell Host & Microbe* 26: 163–172.
- van der Burgh AM, Joosten M. 2019. Plant immunity: thinking outside and inside the box. *Trends in Plant Science* 24: 587–601.
- Caplan JL, Kumar AS, Park E, Padmanabhan MS, Hoban K, Modla S, Czymmek K, Dinesh-Kumar SP. 2015. Chloroplast stromules function during innate immunity. *Developmental Cell* 34: 45–57.
- Caplan JL, Mamillapalli P, Burch-Smith TM, Czymmek K, Dinesh-Kumar SP. 2008. Chloroplast protein NRIP1 mediates innate immune receptor recognition of a viral effector. *Cell* 132: 449–462.
- Chan KX, Phua SY, Crisp P, McQuinn R, Pogson BJ. 2016. Learning the languages of the chloroplast: retrograde signaling and beyond. *Annual Review of Plant Biology* 67: 25–53.
- Chang CC, Slesak I, Jorda L, Sotnikov A, Melzer M, Miszalski Z, Mullineaux PM, Parker JE, Karpinska B, Karpinski S. 2009. *Arabidopsis* chloroplast glutathione peroxidases play a role in cross talk between photooxidative stress and immune responses. *Plant Physiology* 150: 670–683.
- Chaouch S, Queval G, Noctor G. 2012. AtRbohF is a crucial modulator of defence-associated metabolism and a key actor in the interplay between intracellular oxidative stress and pathogenesis responses in *Arabidopsis*. *The Plant Journal* 69: 613–627.
- Chaouch S, Queval G, Vanderauwera S, Mhamdi A, Vandorpe M, Langlois-Meurinne M, Van Breusegem F, Saundrenan P, Noctor G. 2010. Peroxisomal hydrogen peroxide is coupled to biotic defense responses by ISOCHORISMATE SYNTHASE1 in a daylength-related manner. *Plant Physiology* 153: 1692–1705.
- Choi S, Jayaraman J, Segonzac C, Park HJ, Park H, Han SW, Sohn KH. 2017. *Pseudomonas syringae* pv. *actinidiae* type III effectors localized at multiple cellular compartments activate or suppress innate immune responses in *Nicotiana benthamiana*. *Frontiers in Plant Science* 8: 2157.
- Cohen SP, Leach JE. 2019. Abiotic and biotic stresses induce a core transcriptome response in rice. *Scientific Reports* 9: 6273.
- Colville L, Smirnoff N. 2008. Antioxidant status, peroxidase activity, and PR protein transcript levels in ascorbate-deficient *Arabidopsis thaliana vtc* mutants. *Journal of Experimental Botany* 59: 3857–3868.
- Dahal A, Chen L, Kiba A, Hikichi Y, Ohnishi K. 2018. Chloroplastic proteins are targets for the RipG effectors of *Ralstonia solanacearum*. *International Journal of Environmental Science and Technology* 5: 147–156.
- Daudi A, Cheng Z, O'Brien JA, Mammarella N, Khan S, Ausubel FM, Bolwell GP. 2012. The apoplastic oxidative burst peroxidase in *Arabidopsis* is a major component of pattern-triggered immunity. *Plant Cell* 24: 275–287.
- Dietz KJ, Turkan I, Krieger-Liszskay A. 2016. Redox- and reactive oxygen species-dependent signaling into and out of the photosynthesizing chloroplast. *Plant Physiology* 171: 1541–1550.
- Ding P, Ding Y. 2020. Stories of salicylic acid: a plant defense hormone. *Trends in Plant Science* 25: 549–565.
- Ding X, Jimenez-Gongora T, Krenz B, Lozano-Duran R. 2019. Chloroplast clustering around the nucleus is a general response to pathogen perception in *Nicotiana benthamiana*. *Molecular Plant Pathology* 20: 1298–1306.
- Ding Y, Sun T, Ao K, Peng Y, Zhang Y, Li X, Zhang Y. 2018. Opposite roles of salicylic acid receptors NPR1 and NPR3/NPR4 in transcriptional regulation of plant immunity. *Cell* 173: 1454–1467.
- Dogra V, Li M, Singh S, Li M, Kim C. 2019. Oxidative post-translational modification of EXECUTER1 is required for singlet oxygen sensing in plastids. *Nature Communications* 10: 2834.
- Duan J, Lee KP, Dogra V, Zhang S, Liu K, Caceres-Moreno C, Lv S, Xing W, Kato Y, Sakamoto W *et al.* 2019. Impaired PSII proteostasis promotes retrograde signaling via salicylic acid. *Plant Physiology* 180: 2182–2197.
- Durner J, Klessig DF. 1995. Inhibition of ascorbate peroxidase by salicylic acid and 2,6-dichloroisonicotinic acid, two inducers of plant defense responses. *Proceedings of the National Academy of Sciences, USA* 92: 11312–11316.
- Erickson JL, Adlung N, Lampe C, Bonas U, Schattat MH. 2018. The Xanthomonas effector XopL uncovers the role of microtubules in stromule extension and dynamics in *Nicotiana benthamiana*. *The Plant Journal* 93: 856–870.
- Erickson JL, Ziegler J, Guevara D, Abel S, Klosgen RB, Mathur J, Rothstein SJ, Schattat MH. 2014. *Agrobacterium*-derived cytokinin influences plastid morphology and starch accumulation in *Nicotiana benthamiana* during transient assays. *BMC Plant Biology* 14: 127.
- Exposito-Rodríguez M, Laissue PP, Yvon-Durocher G, Smirnoff N, Mullineaux PM. 2017. Photosynthesis-dependent H₂O₂ transfer from chloroplasts to nuclei provides a high-light signalling mechanism. *Nature Communications* 8: 49.
- Faris JD, Zhang Z, Lu H, Lu S, Reddy L, Cloutier S, Fellers JP, Meinhardt SW, Rasmussen JB, Xu SS *et al.* 2010. A unique wheat disease resistance-like gene governs effector-triggered susceptibility to necrotrophic pathogens. *Proceedings of the National Academy of Sciences, USA* 107: 13544–13549.
- Fernandez JC, Burch-Smith TM. 2019. Chloroplasts as mediators of plant biotic interactions over short and long distances. *Current Opinion in Plant Biology* 50: 148–155.
- Fisher B, Yarmolinsky D, Abdel-Ghany S, Pilon M, Pilon-Smits EA, Sagi M, Van Hoewyk D. 2016. Superoxide generated from the glutathione-mediated reduction of selenite damages the iron-sulfur cluster of chloroplastic ferredoxin. *Plant Physiology and Biochemistry* 106: 228–235.
- Gao C, Xu H, Huang J, Sun B, Zhang F, Savage Z, Duggan C, Yan T, Wu CH, Wang Y *et al.* 2020. Pathogen manipulation of chloroplast function triggers a light-dependent immune recognition. *Proceedings of the National Academy of Sciences, USA* 117: 9613–9620.
- Gao QM, Yu K, Xia Y, Shine MB, Wang C, Navarre D, Kachroo A, Kachroo P. 2014. Mono- and digalactosyldiacylglycerol lipids function nonredundantly to regulate systemic acquired resistance in plants. *Cell Reports* 9: 1681–1691.
- Gohre V, Jones AM, Sklenar J, Robatzek S, Weber AP. 2012. Molecular crosstalk between PAMP-triggered immunity and photosynthesis. *Molecular Plant–Microbe Interactions* 25: 1083–1092.
- Gou JY, Li K, Wu K, Wang X, Lin H, Cantu D, Uauy C, Dobon-Alonso A, Midorikawa T, Inoue K *et al.* 2015. Wheat stripe rust resistance protein WKS1 reduces the ability of the thylakoid-associated ascorbate peroxidase to detoxify reactive oxygen species. *Plant Cell* 27: 1755–1770.

- Grant M, Brown I, Adams S, Knight M, Ainslie A, Mansfield J. 2000. The RPM1 plant disease resistance gene facilitates a rapid and sustained increase in cytosolic calcium that is necessary for the oxidative burst and hypersensitive cell death. *The Plant Journal* 23: 441–450.
- Halane MK, Kim SH, Spears BJ, Garner CM, Rogan CJ, Okafor EC, Su J, Bhattacharjee S, Gassmann W. 2018. The bacterial type III-secreted protein AvrRps4 is a bipartite effector. *PLoS Pathogens* 14: e1006984.
- Hamel LP, Sekine KT, Wallon T, Sugiwaka Y, Kobayashi K, Moffett P. 2016. The chloroplastic protein THF1 interacts with the coiled-coil domain of the disease resistance protein N' and regulates light-dependent cell death. *Plant Physiology* 171: 658–674.
- Havaux M. 2014. Carotenoid oxidation products as stress signals in plants. *The Plant Journal* 79: 597–606.
- Havaux M, Triantaphyllides C, Genty B. 2006. Autoluminescence imaging: a non-invasive tool for mapping oxidative stress. *Trends in Plant Science* 11: 480–484.
- Henfling J, Bostock R. 1980. Effect of abscisic-acid on rishitin and lubimin accumulation and resistance to *Phytophthora infestans* and *Cladosporium cucumerinum* in potato-tuber tissue-slices. *Phytopathology* 70: 1074–1078.
- Huang S, van der Vossen EA, Kuang H, Vleeshouwers VG, Zhang N, Borm TJ, van Eck HJ, Baker B, Jacobsen E, Visser RG. 2005. Comparative genomics enabled the isolation of the R3a late blight resistance gene in potato. *The Plant Journal* 42: 251–261.
- Huang W, Chen Q, Zhu Y, Hu F, Zhang L, Ma Z, He Z, Huang J. 2013. Arabidopsis thylakoid formation 1 is a critical regulator for dynamics of PSII-LHCII complexes in leaf senescence and excess light. *Molecular Plant* 6: 1673–1691.
- Hutt H, Everson R, Love J, Littlejohn GR. 2014. How clumpy is my image? Scoring in crowdsourced annotation tasks. *Soft Computing* 19: 1541–1552.
- Imlay JA. 2013. The molecular mechanisms and physiological consequences of oxidative stress: lessons from a model bacterium. *Nature Reviews Microbiology* 11: 443–454.
- Ishiga Y, Ishiga T, Ikeda Y, Matsuura T, Mysore KS. 2016. NADPH-dependent thioredoxin reductase C plays a role in nonhost disease resistance against *Pseudomonas syringae* pathogens by regulating chloroplast-generated reactive oxygen species. *PeerJ* 4: e1938.
- Ishiga Y, Ishiga T, Wangdi T, Mysore KS, Uppalapati SR. 2011. NTRC and chloroplast-generated reactive oxygen species regulate *Pseudomonas syringae* pv. *tomato* disease development in tomato and Arabidopsis. *Molecular Plant–Microbe Interactions* 25: 294–306.
- Ishiguro S, Kawai-Oda A, Ueda J, Nishida I, Okada K. 2001. The DEFECTIVE IN ANOTHER DEHISCENCE1 gene encodes a novel phospholipase a1 catalyzing the initial step of jasmonic acid biosynthesis, which synchronizes pollen maturation, anther dehiscence, and flower opening in Arabidopsis. *Plant Cell* 13: 2191–2209.
- Jelenska J, Yao N, Vinatzer BA, Wright CM, Brodsky JL, Greenberg JT. 2007. A J domain virulence effector of *Pseudomonas syringae* remodels host chloroplasts and suppresses defenses. *Current Biology* 17: 499–508.
- Jeon H, Kim W, Kim B, Lee S, Jayaraman J, Jung G, Choi S, Sohn KH, Segonzac C. 2020. *Ralstonia solanacearum* type III effectors with predicted nuclear localization signal localize to various cell compartments and modulate immune responses in *Nicotiana* spp. *Plant Pathology Journal* 36: 43–53.
- Jiang G, Yin D, Zhao J, Chen H, Guo L, Zhu L, Zhai W. 2016. The rice thylakoid membrane-bound ascorbate peroxidase OsAPX8 functions in tolerance to bacterial blight. *Scientific Reports* 6: 26104.
- Jones JD, Dangl JL. 2006. The plant immune system. *Nature* 444: 323–329.
- Kumar AS, Park E, Nedo A, Alqarni A, Ren L, Hoban K, Modla S, McDonald JH, Kambhmettu C, Dinesh-Kumar SP *et al.* 2018. Stromule extension along microtubules coordinated with actin-mediated anchoring guides perinuclear chloroplast movement during innate immunity. *Elife* 7: e23625.
- Kupcevicz VF. 1947. *The physiology of the diseased plant in relation to the general questions of parasitism*. Moscow, Russia: USSR Academy of Sciences.
- Lafamme B, Dillon MM, Martel A, Almeida RND, Desveaux D, Guttman DS. 2020. The pan-genome effector-triggered immunity landscape of a host–pathogen interaction. *Science* 367: 763–768.
- Lee MW, Jelenska J, Greenberg JT. 2008. Arabidopsis proteins important for modulating defense responses to *Pseudomonas syringae* that secrete HopW1-1. *The Plant Journal* 54: 452–465.
- Lew TTS, Koman VB, Silmore KS, Seo JS, Gordiichuk P, Kwak SY, Park M, Ang MC, Khong DT, Lee MA *et al.* 2020. Real-time detection of wound-induced H₂O₂ signalling waves in plants with optical nanosensors. *Nature Plants* 6: 404–415.
- Lewis LR, Polanski Z, de Torres-Zabala M, Jayaraman S. 2015. Transcriptional dynamics driving basal defense and pathogen effector mediated immunosuppression in Arabidopsis leaves following infection with *Pseudomonas syringae* pv. *tomato* DC3000. *Plant Cell* 27: 3038–3064.
- Li G, Froehlich JE, Elowsky C, Msanne J, Ostosh AC, Zhang C, Awada T, Alfano JR. 2014. Distinct *Pseudomonas* type-III effectors use a cleavable transit peptide to target chloroplasts. *The Plant Journal* 77: 310–321.
- Liu L, Sonbol FM, Huot B, Gu Y, Withers J, Mwimba M, Yao J, He SY, Dong X. 2016. Salicylic acid receptors activate jasmonic acid signalling through a non-canonical pathway to promote effector-triggered immunity. *Nature Communications* 7: 13099.
- Liu M, Zhang S, Hu J, Sun W, Padilla J, He Y, Li Y, Yin Z. 2019. Phosphorylation-guarded light-harvesting complex II contributes to broad-spectrum blast resistance in rice. *Proceedings of the National Academy of Sciences, USA* 116.
- Liu Y, Lan X, Song S, Yin L, Dry IB, Qu J, Xiang J, Lu J. 2018. In planta functional analysis and subcellular localization of the oomycete pathogen *Plasmopara viticola* candidate RXLR effector repertoire. *Frontiers in Plant Science* 9: 286.
- Liu Y, Ren D, Pike S, Pallardy S, Gassmann W, Zhang S. 2007. Chloroplast-generated reactive oxygen species are involved in hypersensitive response-like cell death mediated by a mitogen-activated protein kinase cascade. *The Plant Journal* 51: 941–954.
- López-Solanilla E, Bronstein PA, Schneider AR, Collmer A. 2004. HopPtoN is a *Pseudomonas syringae* Hrp (type III secretion system) cystein protease effector that suppresses pathogen-induced necrosis associated with both compatible and incompatible plant interactions. *Molecular Microbiology* 54: 353–365.
- Manning VA, Hardison LK, Ciuffetti LM. 2007. Ptr ToxA interacts with a chloroplast-localized protein. *Molecular Plant–Microbe Interactions* 20: 168–177.
- Maruta T, Noshi M, Tanouchi A, Tamoi M, Yabuta Y, Yoshimura K, Ishikawa T, Shigeoka S. 2012. H₂O₂-triggered retrograde signaling from chloroplasts to nucleus plays specific role in response to stress. *Journal of Biological Chemistry* 287: 11717–11729.
- McDonald MC, Ahren D, Simpfendorfer S, Milgate A, Solomon PS. 2017. The discovery of the virulence gene *ToxA* in the wheat and barley pathogen *Bipolaris sorokiniana*. *Molecular Plant Pathology* 19: 432–439.
- Meng X, Zhang S. 2013. MAPK cascades in plant disease resistance signaling. *Annual Review of Phytopathology* 51: 245–266.
- Metz JG, Pakrasi HB, Seibert M, Arntzer CJ. 1986. Evidence for a dual function of the herbicide-binding D1 protein in photosystem II. *FEBS Letters* 205: 269–274.
- Mohr PG, Cahill DM. 2003. Abscisic acid influences the susceptibility of *Arabidopsis thaliana* to *Pseudomonas syringae* pv. *tomato* and *Peronospora parasitica*. *Functional Plant Biology* 30: 461–469.
- Mubarakshina MM, Ivanov BN, Naydov IA, Hillier W, Badger MR, Krieger-Liszkay A. 2010. Production and diffusion of chloroplastic H₂O₂ and its implication to signalling. *Journal of Experimental Botany* 61: 3577–3587.
- Mukherjee M, Larrimore KE, Ahmed NJ, Bedick TS, Barghouthi NT, Traw MB, Barth C. 2010. Ascorbic acid deficiency in Arabidopsis induces constitutive priming that is dependent on hydrogen peroxide, salicylic acid, and the *NPR1* gene. *Molecular Plant–Microbe Interactions* 23: 340–351.
- Mukhtar MS, Carvunis AR, Dreze M, Epple P, Steinbrenner J, Moore J, Tasan M, Galli M, Hao T, Nishimura MT *et al.* 2011. Independently evolved virulence effectors converge onto hubs in a plant immune system network. *Science* 333: 596–601.
- Mullineaux PM, Exposito-Rodriguez M, Laissue PP, Smirnov N. 2018. ROS-dependent signalling pathways in plants and algae exposed to high light: comparisons with other eukaryotes. *Free Radical Biology and Medicine* 122: 52–64.
- Mullineaux PM, Exposito-Rodriguez M, Laissue PP, Smirnov N, Park E. 2020. Spatial chloroplast-to-nucleus signalling involving plastid-nuclear complexes and stromules. *Philosophical Transactions of the Royal Society of London. Series B, Biological sciences* 375: 20190405.
- Mur LA, Aubry S, Mondhe M, Kingston-Smith A, Gallagher J, Timms-Taravella E, James C, Papp I, Hortensteiner S, Thomas H *et al.* 2010. Accumulation of

- chlorophyll catabolites photosensitizes the hypersensitive response elicited by *Pseudomonas syringae* in Arabidopsis. *New Phytologist* 188: 161–174.
- Nakano M, Mukaiyama T. 2018. *Ralstonia solanacearum* type III effector RipAL targets chloroplasts and induces jasmonic acid production to suppress salicylic acid-mediated defense responses in plants. *Plant Cell Physiology* 59: 2576–2589.
- Nietzel T, Elsasser M, Ruberti C, Steinbeck J, Ugalde JM, Fuchs P, Wagner S, Ostermann L, Moseler A, Lemke P *et al.* 2019. The fluorescent protein sensor roGFP2-Orp1 monitors *in vivo* H₂O₂ and thiol redox integration and elucidates intracellular H₂O₂ dynamics during elicitor-induced oxidative burst in Arabidopsis. *New Phytologist* 221: 1649–1664.
- Nomura H, Komori T, Uemura S, Kanda Y, Shimotani K, Nakai K, Furuichi T, Takebayashi K, Sugimoto T, Sano S *et al.* 2012. Chloroplast-mediated activation of plant immune signalling in Arabidopsis. *Nature Communications* 3: doi: doi.org/10.1038/ncomms1926.
- Nomura K, Mecey C, Lee YN, Imboden LA, Chang JH, He SY. 2011. Effector-triggered immunity blocks pathogen degradation of an immunity-associated vesicle traffic regulator in Arabidopsis. *Proceedings of the National Academy of Sciences, USA* 108: 10774–10779.
- Ori N, Eshed Y, Paran I, Presting G, Aviv D, Tanksley S, Zamir D, Fluhr R. 1997. The I2C family from the wilt disease resistance locus *I2* belongs to the nucleotide binding, leucine-rich repeat superfamily of plant resistance genes. *Plant Cell* 9: 521–532.
- Parisy V, Poinssot B, Owsianowski L, Buchala A, Glazebrook J, Mauch F. 2007. Identification of PAD2 as a gamma-glutamylcysteine synthetase highlights the importance of glutathione in disease resistance of Arabidopsis. *The Plant Journal* 49: 159–172.
- Park E, Caplan JL, Dinesh-Kumar SP. 2018a. Dynamic coordination of plastid morphological change by cytoskeleton for chloroplast-nucleus communication during plant immune responses. *Plant Signaling & Behavior* 13: e1500064.
- Park E, Nedo A, Caplan JL, Dinesh-Kumar SP. 2018b. Plant-microbe interactions: organelles and the cytoskeleton in action. *New Phytologist* 217: 1012–1028.
- Pavet V, Olmos E, Kiddle G, Mowla S, Kumar S, Antoniw J, Alvarez ME, Foyer CH. 2005. Ascorbic acid deficiency activates cell death and disease resistance responses in Arabidopsis. *Plant Physiology* 139: 1291–1303.
- Pecrix Y, Buendia L, Penouilh-Suzette C, Maréchaux M, Legrand L, Bouchez O, Rengel D, Gouzy J, Cottret L, Vear F *et al.* 2019. Sunflower resistance to multiple downy mildew pathotypes revealed by recognition of conserved effectors of the oomycete *Plasmopara halstedii*. *The Plant Journal* 97: 730–748.
- Peng Z, Hu Y, Zhang J, Hugueta-Tapia JC, Block AK, Park S, Sapkota S, Liu Z, Liu S, White FF. 2019. *Xanthomonas translucens* commandeers the host rate-limiting step in ABA biosynthesis for disease susceptibility. *Proceedings of the National Academy of Sciences, USA* 116: 20938–20946.
- Perez-Bueno ML, Pineda M, Baron M. 2019. Phenotyping plant responses to biotic stress by chlorophyll fluorescence imaging. *Frontiers in Plant Science* 10: 1135.
- Perez-Ruiz JM, Naranjo B, Ojeda V, Guinea M, Cejudo FJ. 2017. NTRC-dependent redox balance of 2-Cys peroxiredoxins is needed for optimal function of the photosynthetic apparatus. *Proceedings of the National Academy of Sciences, USA* 114: 12069–12074.
- Petre B, Saunders DGO, Sklenar J, Lorrain C, Krasileva KV, Win J, Duplessis S, Kamoun S. 2016. Heterologous expression screens in *Nicotiana benthamiana* identify a candidate effector of the wheat yellow rust pathogen that associates with processing bodies. *PLoS ONE* 11: e0149035.
- Petre B, Saunders DGO, Sklenar J, Lorrain C, Win J, Duplessis S, Kamoun S. 2015. Candidate effector proteins of the rust pathogen *Melampsora larici-populina* target diverse plant cell compartments. *Molecular Plant-Microbe Interactions* 28: 689–700.
- Rekhter D, Ludke D, Ding Y, Feussner K, Zienkiewicz K, Lipka V, Wiermer M, Zhang Y, Feussner I. 2019. Isochorismate-derived biosynthesis of the plant stress hormone salicylic acid. *Science* 365: 498–502.
- Robert-Seilaniantz A, Grant M, Jones JD. 2011. Hormone crosstalk in plant disease and defense: more than just jasmonate-salicylate antagonism. *Annual review of Phytopathology* 49: 317–343.
- Rodrigues O, Reshetnyak G, Grondin A, Saijo Y, Leonhardt N, Maurel C, Verdoucq L. 2017. Aquaporins facilitate hydrogen peroxide entry into guard cells to mediate ABA- and pathogen-triggered stomatal closure. *Proceedings of the National Academy of Sciences, USA* 114: 9200–9205.
- Rodriguez-Herva JJ, Gonzalez-Melendi P, Cuartas-Lanza R, Antunez-Lamas M, Rio-Alvarez I, Li Z, Lopez-Torrejon G, Diaz I, Del Pozo JC, Chakravarthy S *et al.* 2012. A bacterial cysteine protease effector protein interferes with photosynthesis to suppress plant innate immune responses. *Cellular Microbiology* 14: 669–681.
- Rossi FR, Krapp AR, Bisaro F, Maiale SJ, Pieckenstein FL, Carrillo N. 2017. Reactive oxygen species generated in chloroplasts contribute to tobacco leaf infection by the necrotrophic fungus *Botrytis cinerea*. *The Plant Journal* 92: 761–773.
- Sarma GN, Manning VA, Ciuffetti LM, Karplus PA. 2005. Structure of Ptr ToxA: an RGD-containing host-selective toxin from *Pyrenophora tritici-repentis*. *Plant Cell* 17: 3190–3202.
- Schneider T, Bolger A, Zeier J, Preiskowski S, Benes V, Trenkamp S, Usadel B, Farre EM, Matsubara S. 2019. Fluctuating light interacts with time of day and leaf development stage to reprogram gene expression. *Plant Physiology* 179: 1632–1657.
- Seo M, Koshiba T. 2002. Complex regulation of ABA biosynthesis in plants. *Trends in Plant Science* 7: 41–48.
- Shah J, Chaturvedi R, Chowdhury Z, Venables B, Petros RA. 2014. Signaling by small metabolites in systemic acquired resistance. *The Plant Journal* 79: 645–658.
- Shine MB, Xiao X, Kachroo P, Kachroo A. 2019. Signaling mechanisms underlying systemic acquired resistance to microbial pathogens. *Plant Science* 279: 81–86.
- Smirnov N, Arnaud D. 2019. Hydrogen peroxide metabolism and functions in plants. *New Phytologist* 221: 1197–1214.
- de Souza A, Wang JZ, Dehesh K. 2017. Retrograde signals: integrators of interorganellar communication and orchestrators of plant development. *Annual Review of Plant Biology* 68: 85–108.
- Sperschneider J, Catanzariti A-M, DeBoer K, Petre B, Gardiner DM, Singh KB, Dodds PN, Taylor JM. 2017. LOCALIZER: subcellular localization prediction of both plant and effector proteins in the plant cell. *Scientific Reports* 7: 44598.
- Su J, Yang L, Zhu Q, Wu H, He Y, Liu Y, Xu J, Jiang D, Zhang S. 2018. Active photosynthetic inhibition mediated by MPK3/MPK6 is critical to effector-triggered immunity. *PLoS Biology* 16: e2004122.
- Tang L, Yang G, Ma M, Liu X, Li B, Xie J, Fu Y, Chen T, Yu Y, Chen W *et al.* 2020. An effector of a necrotrophic fungal pathogen targets the calcium-sensing receptor in chloroplasts to inhibit host resistance. *Molecular Plant-Microbe Interactions* 21: 686–701.
- Tognetti VB, Palatnik JF, Fillat MF, Melzer M, Hajirezaei MR, Valle EM, Carrillo N. Functional replacement of ferredoxin by a cyanobacterial flavodoxin in tobacco confers broad-range stress tolerance. *Plant Cell* 18: 2035–2050.
- Tomita R, Sekine KT, Mizumoto H, Sakamoto M, Murai J, Kiba A, Hikichi Y, Suzuki K, Kobayashi K. 2011. Genetic basis for the hierarchical interaction between *Tobamovirus* spp. and L resistance gene alleles from different pepper species. *Molecular Plant-Microbe Interactions* 24: 108–117.
- Torrens-Spence MP, Bobokalonova A, Carballo V, Glinkerman CM, Pluskal T, Shen A, Weng JK. 2019. PBS3 and EPS1 complete salicylic acid biosynthesis from isochorismate in Arabidopsis. *Molecular Plant* 12: 1577–1586.
- Torres MA, Jones JD, Dangl JL. 2006. Reactive oxygen species signaling in response to pathogens. *Plant Physiology* 141: 373–378.
- de Torres Zabala M, Bennett MH, Truman WH, Grant MR. 2009. Antagonism between salicylic and abscisic acid reflects early host-pathogen conflict and moulds plant defence responses. *The Plant Journal* 59: 375–386.
- de Torres Zabala M, Littlejohn G, Jayaraman S, Studholme D, Bailey T, Lawson T, Tillich M, Licht D, Bolter B, Delfino L *et al.* 2015. Chloroplasts play a central role in plant defence and are targeted by pathogen effectors. *Nature Plants* 1: 15074.
- de Torres-Zabala M, Truman W, Bennett MH, Lafforgue G, Mansfield JW, Rodriguez Egea P, Bogre L, Grant M. 2007. *Pseudomonas syringae* pv. *tomato* hijacks the Arabidopsis abscisic acid signalling pathway to cause disease. *EMBO Journal* 26: 1434–1443.
- Toufexi A, Duggan C, Pandey P, Savage Z, Segretin ME, Yuen LH, Gaboriau DCA, Leary AY, Khandare V, Ward AD *et al.* 2019. Chloroplasts navigate towards the pathogen interface to counteract infection by the Irish potato famine pathogen. *BioRxiv*. doi:10.1101/516443.
- Tripathi BN, Bhatt I, Dietz KJ. 2009. Peroxiredoxins: a less studied component of hydrogen peroxide detoxification in photosynthetic organisms. *Protoplasma* 235: 3–15.

- Truman W, Bennett MH, Kubigsteltig I, Turnbull C, Grant M. 2007. Arabidopsis systemic immunity uses conserved defense signaling pathways and is mediated by jasmonates. *Proceedings of the National Academy of Sciences, USA* **104**: 1075–1080.
- Truman W, Zabala MdT, Grant M. 2006. Type III effectors orchestrate a complex interplay between transcriptional networks to modify basal defense responses during pathogenesis and resistance. *The Plant Journal* **46**: 14–33.
- Vetoshkina DV, Ivanov BN, Khorobrykh SA, Proskuryakov II, Borisova-Mubarakshina MM. 2017. Involvement of the chloroplast plastoquinone pool in the Mehler reaction. *Physiologia Plantarum* **161**: 45–55.
- Vogel MO, Moore M, König K, Pecher P, Alsharafa K, Lee J, Dietz KJ. 2014. Fast retrograde signaling in response to high light involves metabolite export, MITOGEN-ACTIVATED PROTEIN KINASE6, and AP2/ERF transcription factors in Arabidopsis. *Plant Cell* **26**: 1151–1165.
- Wang M, Rui L, Yan H, Shi H, Zhao W, Lin JE, Zhang K, Blakeslee JJ, Mackey D, Tang D *et al.* 2018. The major leaf ferredoxin Fd2 regulates plant innate immunity in Arabidopsis. *Molecular Plant Pathology* **19**: 1377–1390.
- Wang S, Li QP, Wang J, Yan Y, Zhang GL, Zhang H, Wu J, Chen F, Wang X, Kang Z *et al.* 2019. YR36/WKS1-mediated phosphorylation of PsbO, an extrinsic member of photosystem II, inhibits photosynthesis and confers stripe rust resistance in wheat. *Molecular Plant* **12**: 1639–1650.
- Wangdi T, Uppalapati SR, Nagaraj S, Ryu CM, Bender CL, Mysore KS. 2010. A virus-induced gene silencing screen identifies a role for Thylakoid Formation1 in *Pseudomonas syringae* pv *tomato* symptom development in tomato and Arabidopsis. *Plant Physiology* **152**: 281–292.
- Warren RF, Merritt PM, Holub E, Innes RW. 1999. Identification of three putative signal transduction genes involved in *R* gene-specified disease resistance in Arabidopsis. *Genetics* **152**: 401–412.
- Wasternack C, Hause B. 2013. Jasmonates: biosynthesis, perception, signal transduction and action in plant stress response, growth and development. An update to the 2007 review in *Annals of Botany*. *Annals of Botany* **111**: 1021–1058.
- Waszczak C, Carmody M, Kangasjarvi J. 2018. Reactive oxygen species in plant signaling. *Annual Review of Plant Biology* **69**: 209–236.
- van Wees SC, de Swart EA, van Pelt JA, van Loon LC, Pieterse CM. 2000. Enhancement of induced disease resistance by simultaneous activation of salicylate- and jasmonate-dependent defense pathways in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* **97**: 8711–8716.
- Wendehenne D, Gao QM, Kachroo A, Kachroo P. 2014. Free radical-mediated systemic immunity in plants. *Current Opinion in Plant Biology* **20**: 127–134.
- Win J, Krasileva KV, Kamoun S, Shirasu K, Staskawicz BJ, Banfield MJ. 2012. Sequence divergent RXLR effectors share a structural fold conserved across plant pathogenic oomycete species. *PLoS Pathogens* **8**: e1002400.
- Wu W, Zhu Y, Ma Z, Sun Y, Quan Q, Li P, Hu P, Shi T, Lo C, Chu IK *et al.* 2013. Proteomic evidence for genetic epistasis: ClpR4 mutations switch leaf variegation to virescence in Arabidopsis. *The Plant Journal* **76**: 943–956.
- Yamburenko MV, Zubo YO, Borner T. 2015. Abscisic acid affects transcription of chloroplast genes via protein phosphatase 2C-dependent activation of nuclear genes: repression by guanosine-3'-5'-bisdiphosphate and activation by sigma factor 5. *The Plant Journal* **82**: 1030–1041.
- Yang J, Duan G, Li C, Liu L, Han G, Zhang Y, Wang C. 2019. The crosstalks between jasmonic acid and other plant hormone signaling highlight the involvement of jasmonic acid as a core component in plant response to biotic and abiotic stresses. *Frontiers in Plant Science* **10**: 1349.
- Ye N, Zhu G, Liu Y, Zhang A, Li Y, Liu R, Shi L, Jia L, Zhang J. 2012. Ascorbic acid and reactive oxygen species are involved in the inhibition of seed germination by abscisic acid in rice seeds. *Journal of Experimental Botany* **63**: 1809–1822.
- Yun BW, Feechan A, Yin M, Saidi NB, Le Bihan T, Yu M, Moore JW, Kang JG, Kwon E, Spoel SH *et al.* 2011. S-nitrosylation of NADPH oxidase regulates cell death in plant immunity. *Nature* **478**: 264–268.
- Yun BW, Skelly MJ, Yin M, Yu M, Mun BG, Lee SU, Hussain A, Spoel SH, Loake GJ. 2016. Nitric oxide and S-nitrosoglutathione function additively during plant immunity. *New Phytologist* **211**: 516–526.
- Zandalinas SI, Sengupta S, Burks D, Azad RK, Mittler R. 2019. Identification and characterization of a core set of ROS wave-associated transcripts involved in the systemic acquired acclimation response of Arabidopsis to excess light. *The Plant Journal* **98**: 126–141.
- Zaninotto F, La Camera S, Polverari A, Delledonne M. 2006. Cross talk between reactive nitrogen and oxygen species during the hypersensitive disease resistance response. *Plant Physiology* **141**: 379–383.
- Zhang L, Zhang F, Melotto M, Yao J, He SY. 2017. Jasmonate signaling and manipulation by pathogens and insects. *Journal of Experimental Botany* **68**: 1371–1385.
- Zhang ZS, Xu YY, Xie ZW, Li XY, He ZH, Peng XX. 2016. Association-dissociation of glycolate oxidase with catalase in rice: A potential switch to modulate intracellular H₂O₂ levels. *Molecular Plant* **9**: 737–748.
- Zhou Z, Zhao Y, Bi G, Liang X, Zhou JM. 2019. Early signalling mechanisms underlying receptor kinase-mediated immunity in plants. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* **374**: 20180310.
- Zhu W, Wei W, Fu Y, Cheng J, Xie J, Li G, Yi X, Kang Z, Dickman MB, Jiang D. 2013. A secretory protein of necrotrophic fungus *Sclerotinia sclerotiorum* that suppresses host resistance. *PLoS ONE* **8**: e53901.
- Zoeller M, Stingl N, Kruschke M, Fekete A, Waller F, Berger S, Mueller MJ. 2012. Lipid profiling of the Arabidopsis hypersensitive response reveals specific lipid peroxidation and fragmentation processes: biogenesis of pimelic and azelaic acid. *Plant Physiology* **160**: 365–378.
- Zulfugarov IS, Tovuu A, Eu Y-J, Dogsom B, Poudyal RS, Nath K, Hall M, Banerjee M, Yoon UC, Moon Y-H *et al.* 2014. Production of superoxide from Photosystem II in a rice (*Oryza sativa* L.) mutant lacking PsbS. *BMC Plant Biology* **14**: 242.
- Zurbriggen MD, Carrillo N, Tognetti VB, Melzer M, Peisker M, Hause B, Hajirezaei MR. 2009. Chloroplast-generated reactive oxygen species play a major role in localized cell death during the non-host interaction between tobacco and *Xanthomonas campestris* pv. *vesicatoria*. *The Plant Journal* **60**: 962–973.