

2021-10-20

# The Effect of Pre-operative Exercise Intervention on Patient Outcomes Following Bariatric Surgery: a Systematic Review and Meta-analysis

Durey, BJ

<http://hdl.handle.net/10026.1/18175>

---

10.1007/s11695-021-05743-w

Obesity Surgery

Springer Science and Business Media LLC

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

**The effect of pre-operative exercise intervention on patient outcomes following bariatric surgery: a systematic review and meta-analysis**

Shortened title: Exercise pre bariatric surgery.

Belinda J Durey<sup>1,3\*</sup>, Dominic Fritche<sup>2</sup>, Daniel S Martin<sup>4,5,6</sup>, Lawrence MJ Best<sup>3,7</sup>

1. Alliance for Research in Exercise, Nutrition and Activity (ARENA), School of Health Sciences, University of South Australia, GPO Box 2471, Adelaide, SA, 5001, Australia.
2. University College London, Medical School, WC1E 6BT, UK
3. UCL Division of Surgery and Interventional Science, University College London, London, WC1E 6BT, UK
4. UCL Division of Surgery and Interventional Science, Royal Free Hospital, Pond St, London, NW3 2PS, UK
5. Intensive Care Unit, University Hospitals Plymouth, Derriford Road, Plymouth, Devon, PL6 8DH, UK
6. Peninsula Medical School, University of Plymouth, John Bull Building, Tamar Science Park, Plymouth, PL6 8BU, UK
7. Northern Care Alliance NHS Group, Mayo Building, Salford Royal, Stott Lane, Salford, M68HD

Correspondence to:

Belinda J Durey

Allied Health & Human Performance, University of South Australia

Alliance for Research in Exercise, Nutrition and Activity (ARENA), School of Health Sciences, University of South Australia, GPO Box 2471, Adelaide, SA, 5001, Australia.

E-mail: Belinda.norton@unisa.edu.au

**ABSTRACT**

This systematic review aimed to assess the effect of pre-operative exercise intervention on short- and long-term health and clinical outcomes for adult patients undergoing bariatric surgery (BS). We searched MEDLINE, EMBASE, Cochrane Central Register of Controlled

Trials (CENTRAL), SPORTDiscus, and reference lists of relevant papers, through March 2021. Five randomized controlled trials were included (n = 199 patients). Modest increases in cardiorespiratory fitness ( $VO_{2max}$ ) were found at both pre-operative ( $0.73 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P \leq 0.001$ ) and maximum follow-up time points ( $0.98 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P \leq 0.04$ ). There was no significant effect of exercise intervention on %total weight loss (%TWL). Pre-operative exercise can induce significant short- and long-term improvements in fitness in individuals with obesity. There is insufficient evidence to determine whether pre-operative training impacts other post-operative clinical outcomes.

### **(3) Key Points**

- In obese patients, pre-BS exercise can induce short- and long-term fitness improvements
- There is insufficient evidence on pre-BS training and post-operative clinical outcomes
- More exercise interventions vs clinical outcomes are required in bariatric patients

## INTRODUCTION

Obesity is a significant public health problem globally. The World Health Organization estimated that in 2016 more than 1.9 billion adults were overweight. Of these, over 650 million were obese (1). Class 2 obesity, defined as having a BMI  $\geq 35$ , has a significantly negative impact on quality of life (2), physical health (3) and life expectancy (4). Despite the costs and invasiveness, bariatric surgery (BS) is recognised as the most successful long-term treatment for weight loss (5). The documented benefits of surgery include an improved metabolic profile and, in the majority of those with type 2 diabetes, remission of the disease. However, the long term efficacy of BS is less promising, with diabetes recurring in between 21-43% of these patients within nine years (6). In addition, most BS patients experience an initial loss of lean muscle mass (7) which negatively impacts metabolism. Overall, the long term 'failure rate' in achieving optimal BMI is approximately 20% for those with Class 2 obesity and 35% for Class 3 (8).

There is growing evidence that modifiable health behaviours may play a significant role in patients' health outcomes and weight loss following BS (9-11). Among individuals who have attempted weight loss, the vast majority of patients who keep the weight off report high levels of physical activity (PA) (12, 13). Candidates for BS typically exhibit low levels of PA, both before and after surgery (14, 15), and have reduced functional capacity (16). Compromised cardiometabolic health, reduced fitness, and excess pre-operative body weight are individually acknowledged as predictors of post-operative mortality in a range of populations (14, 16-20). Despite this knowledge, there are currently no evidence-based exercise guidelines for BS candidates, but instead broad PA recommendations (21). Specific pre-operative exercise intervention has been proposed as one strategy to improve fitness, body composition, and other markers of health in patients undergoing BS (21).

The pre-operative waiting period may represent an ideal opportunity to facilitate healthy lifestyle change, as patients are engaged in the process of preparing for surgery and are not yet facing post-operative recovery challenges. Substantial evidence supports pre-operative exercise interventions in other types of major surgery, such as cardiothoracic surgery, orthopaedic and abdominal cancer (22-24), where it has been demonstrated that a fitter surgical population will have lower incidence of post-operative morbidity and mortality. There remain limited studies, especially reviews, which have examined this association in BS patients.

The benefits of exercise for those with obesity include improved cardiovascular disease risk factors including cardiorespiratory fitness ( $VO_{2max}$ ) (25) and reduced risk of mortality (26). In the pre-operative setting, the improvement in  $VO_{2max}$  is particularly important because of its inverse association with higher risk of surgical complications and mortality (27).

Although a positive relationship between exercise and body composition has been established, few studies support the efficacy of exercise alone to achieve weight loss. However, even when exercise does not result in weight loss, it has been shown to confer significant health benefits to people who are overweight or obese (25, 26). In addition, studies do support the efficacy of exercise to prevent weight gain (28). A review published in 2018 synthesized findings from 15 studies involving post-operative exercise training programs. The authors concluded that exercise interventions performed after BS were effective in optimising weight and specifically fat mass loss, which resulted in improved physical fitness (29). It is important to acknowledge that previous PA interventions in obese

populations have described exercise barriers that are unique to this cohort, in addition to those experienced by the general population (30, 31).

There have so far been no systematic reviews to investigate the effects of pre-operative exercise intervention on these variables among BS candidates. Therefore, we aimed to summarize the evidence from randomized controlled trials (RCTs) comparing the effects of pre-operative exercise-based interventions on fitness and clinical outcomes for BS patients. The hypothesis being tested in this review is that patients with obesity who undertake preoperative exercise training will have improved fitness and %TWL, fewer complications, and reduced hospital length of stay.

## **MATERIALS AND METHODS**

### **Literature search strategy**

The PICO (Population, Intervention, Comparison, and Outcome) framework was used to design the search. A comprehensive list of search terms was established after consultation with a specialist librarian, and included: ‘bariatric surgery’, ‘prehabilitation’, ‘exercise therapy’, ‘exercise’, ‘physical activity’, ‘fitness’, ‘physical fitness’, ‘crossover procedure’, ‘double-blind procedure’, ‘randomised controlled trial’, ‘single-blind procedure’, ‘factorial’, ‘crossover’, ‘RCT’.

MeSH terms/keywords and free text were used together with the terms (AND/OR/NOT).

Language or time restrictions were not used.

A literature search was conducted to identify all published and unpublished randomised controlled trials. The literature search identified potential studies in all languages. The

following electronic databases were searched: MEDLINE, Embase, Cochrane Central Register of Controlled Clinical Trials, SPORTDiscus, Web of Science and Scopus. The main database search was performed on 1<sup>st</sup> May 2020. This search was updated on 2<sup>nd</sup> March 2021.

Two independent reviewers (BD and DF) appraised all the potential studies identified by the search. Relevant full texts were retrieved, and the reviewers identified studies for inclusion, then recorded reasons for exclusion of ineligible studies. Disagreements were resolved by consensus. Duplicate references were identified and excluded, and multiple reports derived from the same study were collated.

Reference lists of all primary studies and review articles were checked for additional references. Further, authors of identified trials were contacted and asked to identify other published and unpublished studies.

### **Inclusion and Exclusion Criteria**

Eligible studies included RCTs of pre-operative exercise intervention in a study population of adult BS patients. Studies included were reported as full text, those published as abstract only, and unpublished data. We specifically excluded studies with interventions that were limited only to a specific muscle group, for example inspiratory muscle training, because we viewed these as fundamentally different from whole-body PA trials. We also excluded any studies without a control group. Our protocol was prospectively registered with PROSPERO International Prospective Register of Systematic Reviews (PROSPERO 2020 CRD42020173918) available

from: [https://www.crd.york.ac.uk/prospere/display\\_record.php?ID=CRD42020173918](https://www.crd.york.ac.uk/prospere/display_record.php?ID=CRD42020173918)

We included adults (aged 18 years and above) with obesity ( $\text{BMI} \geq 30 \text{ kg}\cdot\text{m}^{-2}$ ) included in exercise intervention RCTs whilst awaiting BS.

Included trials compared exercise intervention with usual care in the pre-operative setting for BS. Exercise is defined as any planned and structured activity carried out to sustain or improve health and fitness. The pre-operative period is specified as between initial primary care referral, to the date of surgery.

Studies that used mixed-modal pre-operative intervention, for example, exercise plus other lifestyle modification deviating from standard care, were excluded. This was to ensure any measured effect could be attributed to exercise intervention alone. Interventions included in this category involved nutrition, or cognitive behaviour therapy.

A set of agreed standardised core outcomes has been developed for bariatric and metabolic surgery using formal consensus methods involving patients and health professionals (32). These nine items, as well as two additional outcomes (serious adverse events and fitness), were included in this review.

The primary outcomes assessed were 1) all-cause mortality in the short-term (30-days) and/or longer-term (maximal follow-up), 2) post-operative short-term morbidity (measured via post-operative morbidity survey [POMS]), 3) overall quality of life (measured via QoL scale / questionnaire), and 4) serious adverse events (short-term and longer-term, respectively, measured using Clavien-Dindo)



The secondary outcomes assessed were treatment associated costs, length of hospital stay, number of days of lost work (maximal follow-up), changes in fitness (pre-operative and maximal follow-up), re-operation / re-intervention and its classification of severity, change in weight (%TWL) diabetes status, technical complications of the specific operation, and micronutrient status.

### **Data extraction**

Two review authors (BD and DF) independently extracted study characteristics and outcome data from included studies using a pre-piloted extraction form. Data collected included the following study characteristics: year and country of publication, number of patients in each treatment arm, number of post-randomisation exclusions, details of each intervention, and baseline patient characteristics. If outcomes were reported multiple times (for example, change in weight was reported at 6 weeks and 3 months) the later time point (i.e., 3 months) was chosen for data extraction.

Where an identified study was published as abstract only, we contacted investigators to verify key study characteristics and obtain missing numerical outcome data. If we were unable to obtain the information from study investigators, we imputed the standard deviation from standard error, interquartile range, or P values according to the Cochrane Handbook for Systematic Reviews of Interventions (33).

### **Assessment of risk of bias in included studies**

The quality of reporting each trial was evaluated by two independent reviewers (BD and DF) using the revised Cochrane risk of bias 2 (RoB2) tool (34). The results of the RoB2 quality assessment (whether a study was of ‘low risk’, ‘there are some concerns’, or ‘high risk’ of

bias) were used to judge the quality of evidence using GRADE criteria (35). Disagreements between the reviewers were resolved by consensus.

### **Statistical analysis**

Data of included studies were pooled using relevant effect measures. Dichotomous data (all-cause mortality, proportion of people with, and number of, serious adverse events, re-operation/re-intervention) were analysed as risk ratios with 95% confidence interval.

Continuous data were analysed as mean difference with 95% confidence interval, when the outcome was reported or converted to the same units in all the trials. Conversion equations were used for fitness data where it was not presented as  $\text{VO}_2$  (36).

Meta-analysis was undertaken only when considered meaningful, for instance, if the treatments, participants, and the underlying clinical questions were similar. Meta-analysis was performed using RevMan 5.4 software (37).

### **Subgroup analysis**

The following subgroup analyses were planned:

1. Different BMI categories (Obesity Class 1, 2 and 3)
2. Trials at low risk of bias compared to trials at high risk of bias
3. Definition used by authors for serious adverse events and any adverse events (International Conference on Harmonisation Expert Working Group vs other methods)
4. Any differences in methods of bariatric surgery
5. Based on the period of follow-up (short term vs medium or long term)
6. Based on the length of time prior to surgery (more or less than a year)

7. Based on different delivery method of exercise (supervised vs unsupervised)

All primary outcomes were to be used in the subgroup analyses.

### **Assessment of heterogeneity**

Effect sizes were presented as weighted mean differences with 95% confidence intervals.

Heterogeneity was assessed on the basis of study design, intervention treatments, stratification, subject type, and statistical methods.

## **RESULTS**

Identification of studies is described in the PRISMA flow diagram (Figure 1). 2,423 articles were found by the systematic search. Following removal of duplicates, 2099 documents were considered, with full-text screening required for 34.

Articles were excluded if the intervention, sample, study design, or outcome did not fulfil the criteria listed in the PICO. The full texts of the remaining 34 studies were reviewed. Five studies satisfied our selection criteria and were included in the systematic review (38-42), with a combined total of 199 participants. Within the studies the mean age ranged from 38 to 47 years and mean baseline BMI ranged from 45 to 47 kg•m<sup>-2</sup>. Post-intervention follow-up ranged from pre-operatively (2 weeks) (41) to 12 months post-operatively (38).

### **Included studies**

Among the included trials, no outcome data were identified on mortality, morbidity, quality of life, serious adverse events (long-term), treatment associated costs, days of lost work, re-operation, diabetes status, technical complications of the specific operation or micronutrient status. Four studies assessed fitness outcomes in a total of 163 obese adults. Three studies

examined body weight outcomes in 142 obese adults. One study assessed serious adverse events and length of hospital stay in 22 participants. Three of the studies were conducted in Canada (38, 41, 42) and 2 were conducted in the USA (39, 40). A summary of the included studies is shown in Table 1.

*Table 1: Characteristics of included studies*

Study	Intervention	Participants	Outcomes
<b>Baillot et al 2018</b>	12-week pre-operative PA intervention.	N=25 (20 females) adult participants (mean age 43 years) with obesity (mean BMI 46 kg•m <sup>-2</sup> )	Cardiorespiratory fitness via six-minute walk test (6MWT), measured at baseline, pre-operatively (12 weeks) and 12-months post-operatively.
<b>Bond et al 2017</b>	6-week pre-operative PA intervention.	N= 36 (31 females) adult participants (mean age 47 years) with obesity (mean BMI 45.8 kg•m <sup>-2</sup> )	Weight, measured at baseline, pre- operatively (6 weeks) and post-operatively (6-months).
<b>Creel et al 2016</b>	6-week pre-operative PA intervention.	N= 36 (31 females) adult participants (mean age 47 years) with obesity (mean BMI 45.8 kg•m <sup>-2</sup> )	Weight, measured at baseline, pre- operatively (6 weeks) and post-operatively (6-months).
<b>Kwok et al 2016</b>	12-week pre-operative PA intervention.	N=32 (25 females) adult participants	Cardiorespiratory fitness (6MWT)

		(mean age 47 years) with obesity (mean BMI 45.7 kg•m <sup>-2</sup> )	measured at baseline and pre-operatively (12 weeks).
<b>Li et al 2013</b>	8-week pre-operative PA trial.	N=22 (15 females) adult participants (mean age 38 years) with obesity (mean BMI 47 kg•m <sup>-2</sup> ) awaiting gastric bypass surgery in Canada.	%TWL and cardiorespiratory fitness (VO <sub>2max</sub> ) were measured at baseline, before surgery (8 weeks), and 10-weeks after surgery.

Intervention characteristics are outlined in Table 2. There was a behavioural component to the interventions in all studies, including exercise counselling, reviewing self-monitoring records and progress towards goals, problem-solving barriers, teaching behaviour change strategies, developing action plans to achieve goals, and interdisciplinary lifestyle management. Durations of interventions ranged from 2 to 12 weeks, and exercise modalities varied from self-determined walking to individually prescribed high-intensity interval training (HIIT).

*Table 2: Intervention Characteristics*

	<b>Baillet et al</b>	<b>Bond et al</b>	<b>Creel et al</b>	<b>Kwok et al</b>	<b>Li et al</b>
<b>Duration:</b>	12 weeks	6 weeks	2 weeks	12 weeks	8 weeks
<b>Supervision:</b>	Partial	Unsupervised	Unsupervised	Full	Partial

<b>Modality:</b>	Aerobic, strength	Walking	Walking	Aerobic, strength	Aerobic, strength, HIIT
<b>Intensity:</b>  <b>Low (L),</b>  <b>Moderate (M),</b>  <b>Vigorous (V)</b>	M - V:  55 – 85%  heart rate  reserve  (HRR)	M:  Habitual  exercise plus  additional  5,000  steps/day OR  30 min/day	M:  10,000  steps/day	M - V	M - V:  45 – 100%  HRR
<b>Frequency/week:</b>	3 days	7 days	7 days	3 days	5 days
<b>Time/week:</b>	240 min	210 min	n/a	180-225 min	120-210 min

### **Risk of bias in included studies**

The results of the risk of bias assessment are reported in Figure 2. Three studies were reported to have high overall risk of bias, and 2 had some concerns. Subjects and assessors were not blinded in any study, however, the difficulty of blinding in exercise interventions is acknowledged. More than 15% of patients randomised were excluded in 3 of 5 studies (38-40) due to surgery not going ahead as planned. All 5 studies performed intention-to-treat analyses so these patients were included in analysis.

### **Effects of interventions**

It was not possible to evaluate all-cause mortality, post-operative short-term morbidity, quality of life, number of days of lost work, re-operation / re-intervention and its classification of severity, diabetes status, technical complications of the specific operation, micronutrient status or cost-effectiveness using meta-analyses, as no studies assessed these outcomes.

The risk ratio for serious adverse events following BS (short-term) showed no difference between intervention and control groups, however this was only based on one study (42) (Figure 3). Only one study reported length of hospital stay (Li 2013) (Figure 4). They reported no difference between intervention (mean 2.6 days) and control groups (mean 2.5 days). Confidence intervals were not estimable. Estimates of the between-study variances are presented in Figures 5, 6 and 7.

Meta-analyses were conducted on the changes in fitness and %TWL. Results indicated that there were increases in  $VO_{2max}$  ( $mL \cdot kg^{-1} \cdot min^{-1}$ ) at both pre-operative (mean difference (MD) 0.73,  $P \leq 0.001$ ) and maximum follow-up time points (MD 0.98,  $P \leq 0.04$ ), relative to baseline, as illustrated in Figures 5 and 6. Figure 7 illustrates no significant effect of exercise intervention on %TWL (MD 0.94, 95% CI -1.61 - 3.48).

### **Heterogeneity**

There was no significant heterogeneity for both short and longer-term fitness change analysis. The corresponding  $I^2$  values were 62% and 0%, respectively. There was significant heterogeneity in weight loss analysis ( $I^2 = 70\%$ ).

### **DISCUSSION**

This systematic review found 5 RCTs that investigated the effects of pre-operative exercise intervention in bariatric patients. The pre-operative exercise intervention had a small but significant and lasting effect on cardiorespiratory fitness (0.98  $mL \cdot kg^{-1} \cdot min^{-1}$  increase in  $VO_{2max}$ ). However, the clinical significance of the increase in fitness could not be determined since few relevant clinical measures were reported in the included studies.

Only 1 of the 5 studies (42) measured serious adverse events. Of the 22 patients in this study, one control group participant experienced 1 serious adverse event: anastomotic leakage following surgery. One study (42) measured length of hospital stay, which showed no difference between groups following an 8 week pre-operative exercise intervention. However, within other clinical populations (43, 44) positive benefits of pre-operative exercise interventions on clinical outcomes, such as length of hospital stay and serious adverse events, have been demonstrated.

The modest cardiorespiratory improvements identified both pre-and post-operatively in the current review ( $0.73 - 0.98 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ) equate to approximately 1/3 MET. Three studies (38, 41, 42) measured pre-operative changes in fitness with interventions ranging from 8-12 weeks, and 120-240 minutes per week of moderate – vigorous intensity. Three studies measured post-operative changes in fitness, where maximum follow-up was assessed between 10 weeks and 12-months post-operatively (38, 40, 42) with interventions ranging from 2-12 weeks. Across all studies only one (41) measured adherence (60% within the intervention group) and compliance (75% attendance at planned sessions).

Cardiorespiratory fitness, or the efficiency of utilising oxygen, has been shown to be a powerful predictor of all-cause mortality and life expectancy in both the general population (45) and in clinical populations (46-48). Franklin et al (47) identified that for every 1 MET increase ( $3.5 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ) there is an associated improvement in all-cause mortality from 22-30%. However, there is a lack of studies that objectively measure aerobic capacity and clinical outcomes following surgery. It is therefore impossible to accurately quantify the clinical benefits following physiological adaptation.



It should be noted that we found substantial evidence of heterogeneity for short-term cardiorespiratory fitness changes based on the  $I^2$  value of 62%. Therefore, these results should be interpreted cautiously.

Body weight changes were reported in 3 (39, 40, 42) of the 5 studies at maximum follow-up. A meta-analysis showed no significant effect of pre-operative exercise intervention on %TWL. The remaining two studies only reported BMI. Collectively the mean was not different between the intervention and control groups, averaging 28.9 and 28.0 kg•m<sup>-2</sup>, respectively. However, measurement of %TWL does not differentiate between fat mass and fat-free mass, therefore no conclusions could be drawn about the contribution of each to overall weight change. Since lean tissue loss following BS is common, exercise should be prioritised to attenuate these negative impacts on metabolism (7). There was substantial evidence of heterogeneity for %TWL based on the  $I^2$  value of 70%. Therefore, these results should be interpreted cautiously.

## CONCLUSION

This review and meta-analysis have shown modest improvements in short and long-term cardiorespiratory fitness as a result of pre-operative exercise interventions. However, given the small number of studies and sample sizes, together with inconsistent exercise prescription and outcomes measures, it was not possible to draw conclusions about clinical outcomes. To help overcome this shortcoming, and in line with Coulman and colleagues (32), we recommend adherence to the Core Outcome Set (COS) for future research in this patient cohort. Further, high quality RCTs, adhering to the COS are required in order to definitively

investigate whether pre-operative exercise interventions are beneficial for patients undergoing bariatric surgery.

This review supports the rationale that pre-operative exercise should be initiated prior to surgery for those awaiting bariatric surgery, for the purpose of improving patients' fitness – a key aspect of health.

We conclude that there is a need for further studies investigating the association between pre-operative exercise training and clinical outcomes in bariatric patients. The 5 studies included in this review showed a wide variation of interventions, and few common outcomes.

### **Figures:**

Fig.1 PRISMA flow diagram.

Fig. 2 Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

Fig. 3 Risk ratio for serious adverse events (number of events) following bariatric surgery using a random-effects model.

Fig. 4 Length of hospital stay following bariatric surgery in a random-effects model.

Fig. 5 Meta-analysis of pre-operative change in  $VO_{2max}$  ( $mL \cdot kg^{-1} \cdot min^{-1}$ ) in a random-effects model

Fig. 6 Meta-analysis of change in  $VO_{2max}$  ( $mL \cdot kg^{-1} \cdot min^{-1}$ ) at maximum follow-up post bariatric surgery in a random-effects model.

Fig. 7 Meta-analysis of %total weight loss following bariatric surgery in a random-effects model.

**Conflict of Interest:** Belinda Durey, Dominic Fritche, Daniel Martin and Lawrence Best declare that they have no conflict of interest.

**Statement of informed consent:** As this is a review, informed consent does not apply.

**Ethical approval:** All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. For this type of study formal consent is not required.

## **References to included studies**

### **Baillot 2018**

Baillot A, Mampuya WM, Dionne IJ, Comeau E, Meziat-Burdin A, Langlois MF. Impacts of Supervised Exercise Training in Addition to Interdisciplinary Lifestyle Management in Subjects Awaiting Bariatric Surgery: a Randomized Controlled Study. *Obesity Surgery* 2016;26(11):2602-10.

Baillot A, Mampuya WM, Dionne IJ, Comeau EM, Meziat-Burdin A, Langlois M. Adding supervised group exercise training to interdisciplinary lifestyle management in subjects awaiting bariatric surgery: A randomized controlled study. *Obesity Surgery* 2014;24(8):1357.

\* Baillot A, Vallee CA, Mampuya WM, Dionne IJ, Comeau E, Meziat-Burdin A, et al. Effects of a Pre-surgery Supervised Exercise Training 1 Year After Bariatric Surgery: a Randomized Controlled Study. *Obesity Surgery* 2018;28(4):955-62.

**Bond 2017**

Bond DS, Thomas JG, King WC, Vithiananthan S, Trautvetter J, Unick JL, et al. Exercise improves quality of life in bariatric surgery candidates: Results from the Bari-Active trial. *Obesity* 2015;23(3):536-42.

\* Bond DS, Thomas JG, Vithiananthan S, Unick J, Webster J, Roye GD, et al. Intervention-related increases in pre-operative physical activity are maintained 6-months after Bariatric surgery: Results from the bari-active trial. *International Journal of Obesity* 2017;41(3):467-70.

Bond DS, Thomas JG, Vithiananthan S, Webster J, Unick J, Ryder BA, et al. Changes in enjoyment, self-efficacy, and motivation during a randomized trial to promote habitual physical activity adoption in bariatric surgery patients. *Surgery for Obesity and Related Diseases* 2016;12(5):1072-9. [Other: 1550-7289]

Bond DS, Vithiananthan S, Thomas JG, Trautvetter J, Unick JL, Jakicic JM, et al. Bari-Active: a randomized controlled trial of a pre-operative intervention to increase physical activity in bariatric surgery patients. *Surgery for Obesity and Related Diseases* 2015;11(1):169-80. [Other: 1550-7289]

**Creel 2016**

Creel DB, Schuh LM, Reed CA, Gomez AR, Hurst LA, Stote J, et al. A randomized trial comparing two interventions to increase physical activity among patients undergoing bariatric surgery. *Obesity* 2016;24(8):1660-8.

### **Kwok 2016**

Kwok K, Hardy K, Bouchard D, Vergis A. The impact of a pre-operative exercise program on patients awaiting bariatric surgery. *Surgery for Obesity and Related Diseases* 2016;12 (7 Supplement 1):S201-2.

### **Li 2013**

Li C, Zavorsky GS, Kim DJ, Christou NV, Feldman LS, Carli F. Effects of a bariatric pre-operative exercise program: A pilot randomized study. *Surgical Endoscopy and Other Interventional Techniques* 2013;27:S431.

## **REFERENCES**

1. WHO. Obesity and Overweight Geneva: World Health Organization 2018; Available from: <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight>.
2. Major P, Matlok M, Pedziwiatr M, Migaczewski M, Budzynski P, Stanek M, et al. Quality of Life After Bariatric Surgery. *Obes Surg* 2015 Sep;25(9):1703-10.
3. Must A, Spadano J, Coakley EH, Field AE, Colditz G, Dietz WH. The disease burden associated with overweight and obesity. *JAMA* 1999 Oct 27;282(16):1523-9.
4. Fontaine KR, Redden DT, Wang C, Westfall AO, Allison DB. Years of Life Lost Due to Obesity. *JAMA* 2003;289(2):187-93.

5. Colquitt JL, Pickett K, Loveman E, Frampton GK. Surgery for weight loss in adults. Cochrane Database of Systematic Reviews2014(8).
6. Yu J, Zhou X, Li L, Li S, Tan J, Li Y, et al. The Long-Term Effects of Bariatric Surgery for Type 2 Diabetes: Systematic Review and Meta-analysis of Randomized and Non-randomized Evidence. Obesity Surgery2015 2015/01/01;25(1):143-58.
7. Alba DL, Wu L, Cawthon PM, Mulligan K, Lang T, Patel S, et al. Changes in Lean Mass, Absolute and Relative Muscle Strength, and Physical Performance After Gastric Bypass Surgery. J Clin Endocrinol Metab2019 Mar 1;104(3):711-20.
8. Christou NV, Look D, Maclean LD. Weight gain after short- and long-limb gastric bypass in patients followed for longer than 10 years. Ann Surg2006 Nov;244(5):734-40.
9. Egberts K, Brown WA, Brennan L, O'Brien PE. Does Exercise Improve Weight Loss after Bariatric Surgery? A Systematic Review. Obesity Surgery2012 Feb;22(2):335-41.
10. Jacobi D, Ciangura C, Couet C, Oppert J-M. Physical activity and weight loss following bariatric surgery. Obesity Reviews2011;12(5):366-77.
11. Shah M, Snell PG, Rao S, Adams-Huet B, Quittner C, Livingston EH, et al. High-volume exercise program in obese bariatric surgery patients: A randomized, controlled trial. Obesity2011 September;19(9):1826-34.
12. Thomas JG, Bond DS, Phelan S, Hill JO, Wing RR. Weight-loss maintenance for 10 years in the National Weight Control Registry. Am J Prev Med2014 Jan;46(1):17-23.
13. Karfopoulou E, Mouliou K, Koutras Y, Yannakoulia M. Behaviours associated with weight loss maintenance and regaining in a Mediterranean population sample. A qualitative study. Clin Obes2013 Oct;3(5):141-9.
14. Gilbertson NM, Paisley AS, Kranz S, Weltman A, Kirby JL, Hallowell PT, et al. Bariatric Surgery Resistance: Using Preoperative Lifestyle Medicine and/or Pharmacology for Metabolic Responsiveness. Obes Surg2017 Dec;27(12):3281-91.
15. King WC, Belle SH, Eid GM, Dakin GF, Inabnet WB, Mitchell JE, et al. Physical activity levels of patients undergoing bariatric surgery in the Longitudinal Assessment of Bariatric Surgery study. Surgery for Obesity & Related Diseases2008;4(6):721-8.

16. McCullough PA, Gallagher MJ, Dejong AT, Sandberg KR, Trivax JE, Alexander D, et al. Cardiorespiratory fitness and short-term complications after bariatric surgery. *Chest* 2006 Aug;130(2):517-25.
17. Hennis PJ, Meale PM, Hurst RA, O'Doherty AF, Otto J, Kuper M, et al. Cardiopulmonary exercise testing predicts postoperative outcome in patients undergoing gastric bypass surgery. *Br J Anaesth* 2012 Oct;109(4):566-71.
18. Pico-Sirvent I, Aracil-Marco A, Pastor D, Moya-Ramon M. Effects of a Combined High-Intensity Interval Training and Resistance Training Program in Patients Awaiting Bariatric Surgery: A Pilot Study. *Sports (Basel)* 2019 Mar 25;7(3).
19. O'Doherty AF, West M, Jack S, Grocott MP. Preoperative aerobic exercise training in elective intra-cavity surgery: a systematic review. *Br J Anaesth* 2013 May;110(5):679-89.
20. Duceppe E, Parlow J, MacDonald P, Lyons K, McMullen M, Srinathan S, et al. Canadian Cardiovascular Society Guidelines on Perioperative Cardiac Risk Assessment and Management for Patients Who Undergo Noncardiac Surgery. *Can J Cardiol* 2017 Jan;33(1):17-32.
21. King WC, Bond DS. The importance of preoperative and postoperative physical activity counseling in bariatric surgery. *Exerc Sport Sci Rev* 2013 Jan;41(1):26-35.
22. Hoogeboom TJ, Oosting E, Vriesevink JE, Veenhof C, Siemonsma PC, de Bie RA, et al. Therapeutic validity and effectiveness of preoperative exercise on functional recovery after joint replacement: a systematic review and meta-analysis. *PLoS One* 2012;7(5):e38031.
23. Hulzebos EH, Smit Y, Helders PP, van Meeteren NL. Preoperative physical therapy for elective cardiac surgery patients. *Cochrane Database Syst Rev* 2012 Nov 14;11:CD010118.
24. Hijazi Y, Gondal U, Aziz O. A systematic review of prehabilitation programs in abdominal cancer surgery. *Int J Surg* 2017 Mar;39:156-62.
25. Shaw KA, Gennat HC, O'Rourke P, Del Mar C. Exercise for overweight or obesity. *Cochrane Database of Systematic Reviews* 2006(4).
26. Ross R, Blair S, de Lannoy L, Despres JP, Lavie CJ. Changing the endpoints for determining effective obesity management. *Prog Cardiovasc Dis* 2015 Jan-Feb;57(4):330-6.

27. Moran J, Wilson F, Guinan E, McCormick P, Hussey J, Moriarty J. Role of cardiopulmonary exercise testing as a risk-assessment method in patients undergoing intra-abdominal surgery: a systematic review. *Br J Anaesth* 2016 Feb;116(2):177-91.
28. Fonseca-Junior SJ, Sá CG, Rodrigues PA, Oliveira AJ, Fernandes-Filho J. Physical exercise and morbid obesity: a systematic review. *Arq Bras Cir Dig* 2013;26 Suppl 1:67-73.
29. Bellicha A, Ciangura C, Poitou C, Portero P, Oppert JM. Effectiveness of exercise training after bariatric surgery-a systematic literature review and meta-analysis. *Obesity Reviews* 2018 Nov;19(11):1544-56.
30. Zabatiero J, Hill K, Gucciardi DF, Hamdorf JM, Taylor SF, Hagger MS, et al. Beliefs, Barriers and Facilitators to Physical Activity in Bariatric Surgery Candidates. *Obesity Surgery* 2016 2016/05/01;26(5):1097-109.
31. Pfeil M, Pulford A, Mahon D, Ferguson Y, Lewis MP. The Patient Journey to Gastric Band Surgery: A Qualitative Exploration. *Bariatric Surg Pract Patient Care* 2013;8(2):69-76.
32. Coulman KD, Hopkins J, Brookes ST, Chalmers K, Main B, Owen-Smith A, et al. A Core Outcome Set for the Benefits and Adverse Events of Bariatric and Metabolic Surgery: The BARIACT Project. *PLoS Med* 2016 Nov;13(11):e1002187.
33. Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA. *Cochrane Handbook for Systematic Reviews of Interventions*: Cochrane; 2020. Available from: [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook).
34. Sterne JAC SJ, Page MJ, Elbers RG, Blencowe NS, Boutron I, Cates CJ, Cheng H-Y, Corbett MS, Eldridge SM, Hernán MA, Hopewell S, Hróbjartsson A, Junqueira DR, Jüni P, Kirkham JJ, Lasserson T, Li T, McAleenan A, Reeves BC, Shepperd S, Shrier I, Stewart LA, Tilling K, White IR, Whiting PF, Higgins JPT. . RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;366(l4898).
35. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336(7650):924-6.
36. Ross RM, Murthy JN, Wollak ID, Jackson AS. The six minute walk test accurately estimates mean peak oxygen uptake. *BMC Pulmonary Medicine* 2010 2010/05/26;10(1):31.
37. Centre TNC. *Review Manager (RevMan)*. 5.3 ed. Copenhagen: The Cochrane Collaboration; 2014.



38. Baillot A, Vallee CA, Mampuya WM, Dionne IJ, Comeau E, Meziat-Burdin A, et al. Effects of a Pre-surgery Supervised Exercise Training 1 Year After Bariatric Surgery: a Randomized Controlled Study. *Obesity Surgery* 2018 01 Apr;28(4):955-62.
39. Bond DS, Thomas JG, Vithiananthan S, Unick J, Webster J, Roye GD, et al. Intervention-related increases in preoperative physical activity are maintained 6-months after Bariatric surgery: Results from the bari-active trial. *International Journal of Obesity* 2017 01 Mar;41(3):467-70.
40. Creel DB, Schuh LM, Reed CA, Gomez AR, Hurst LA, Stote J, et al. A randomized trial comparing two interventions to increase physical activity among patients undergoing bariatric surgery. *Obesity* 2016 01 Aug;24(8):1660-8.
41. Kwok K, Hardy K, Bouchard D, Vergis A. The impact of a pre-operative exercise program on patients awaiting bariatric surgery. *Surgery for Obesity and Related Diseases*. [Conference Abstract]. 2016 August;12 (7 Supplement 1):S201-S2.
42. Li C, Zavorsky GS, Kim DJ, Christou NV, Feldman LS, Carli F. Effects of a bariatric preoperative exercise program: A pilot randomized study. *Surgical Endoscopy and Other Interventional Techniques*. [Conference Abstract]. 2013 April;27:S431.
43. Valkenet K, van de Port IG, Dronkers JJ, de Vries WR, Lindeman E, Backx FJ. The effects of preoperative exercise therapy on postoperative outcome: a systematic review. *Clin Rehabil* 2011 Feb;25(2):99-111.
44. Cavalheri V, Granger C. Preoperative exercise training for patients with non-small cell lung cancer. *Cochrane Database Syst Rev* 2017 Jun 7;6(6):Cd012020.
45. Lee DC, Artero EG, Sui X, Blair SN. Mortality trends in the general population: the importance of cardiorespiratory fitness. *J Psychopharmacol* 2010 Nov;24(4 Suppl):27-35.
46. Ross R, Blair SN, Arena R, Church TS, Despres JP, Franklin BA, et al. Importance of Assessing Cardiorespiratory Fitness in Clinical Practice: A Case for Fitness as a Clinical Vital Sign: A Scientific Statement From the American Heart Association. *Circulation* 2016 Dec 13;134(24):e653-e99.
47. Franklin BA, Lavie CJ, Squires RW, Milani RV. Exercise-based cardiac rehabilitation and improvements in cardiorespiratory fitness: implications regarding patient benefit. *Mayo Clin Proc* 2013 May;88(5):431-7.

48. Swank AM, Horton J, Fleg JL, Fonarow GC, Keteyian S, Goldberg L, et al. Modest increase in peak VO<sub>2</sub> is related to better clinical outcomes in chronic heart failure patients: results from heart failure and a controlled trial to investigate outcomes of exercise training. *Circ Heart Fail* 2012 Sep 1;5(5):579-85.