

2022-03-14

The Role of NEDD4 E3 UbiquitinProtein Ligases in Parkinson's Disease

Conway, James

<http://hdl.handle.net/10026.1/18993>

10.3390/genes13030513



Genes

MDPI AG

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Review

The Role of NEDD4 E3 Ubiquitin–Protein Ligases in Parkinson’s Disease

James A. Conway , Grant Kinsman  and Edgar R. Kramer * 

Faculty of Health, Plymouth Institute of Health and Care Research, Peninsula Medical School, University of Plymouth, Plymouth PL6 8BU, UK; james.conway@plymouth.ac.uk (J.A.C.); grant.kinsman@postgrad.plymouth.ac.uk (G.K.)

* Correspondence: edgar.kramer@plymouth.ac.uk; Tel.: +44-(0)-1752-583204

Abstract: Parkinson’s disease (PD) is a debilitating neurodegenerative disease that causes a great clinical burden. However, its exact molecular pathologies are not fully understood. Whilst there are a number of avenues for research into slowing, halting, or reversing PD, one central idea is to enhance the clearance of the proposed aetiological protein, oligomeric α -synuclein. Oligomeric α -synuclein is the main constituent protein in Lewy bodies and neurites and is considered neurotoxic. Multiple E3 ubiquitin-protein ligases, including the NEDD4 (neural precursor cell expressed developmentally downregulated protein 4) family, parkin, SIAH (mammalian homologues of *Drosophila* seven in absentia), CHIP (carboxy-terminus of Hsc70 interacting protein), and SCF^{FBXL5} SCF ubiquitin ligase assembled by the S-phase kinase-associated protein (SKP1), cullin-1 (Cul1), a zinc-binding RING finger protein, and the F-box domain/Leucine-rich repeat protein 5-containing protein FBXL5), have been shown to be able to ubiquitinate α -synuclein, influencing its subsequent degradation via the proteasome or lysosome. Here, we explore the link between NEDD4 ligases and PD, which is not only via α -synuclein but further strengthened by several additional substrates and interaction partners. Some members of the NEDD4 family of ligases are thought to crosstalk even with PD-related genes and proteins found to be mutated in familial forms of PD. Mutations in NEDD4 family genes have not been observed in PD patients, most likely because of their essential survival function during development. Following further in vivo studies, it has been thought that NEDD4 ligases may be viable therapeutic targets in PD. NEDD4 family members could clear toxic proteins, enhancing cell survival and slowing disease progression, or might diminish beneficial proteins, reducing cell survival and accelerating disease progression. Here, we review studies to date on the expression and function of NEDD4 ubiquitin ligases in the brain and their possible impact on PD pathology.

Keywords: NEDD4; E3 ubiquitin–protein ligase; α -synuclein; Parkinson’s disease; dopaminergic neurons



Citation: Conway, J.A.; Kinsman, G.; Kramer, E.R. The Role of NEDD4 E3 Ubiquitin–Protein Ligases in Parkinson’s Disease. *Genes* **2022**, *13*, 513. <https://doi.org/10.3390/genes13030513>

Academic Editor: Allison D. Ebert

Received: 1 February 2022

Accepted: 3 March 2022

Published: 14 March 2022

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Parkinson’s disease (PD) is characterised by the loss of midbrain dopaminergic neurons in the substantia nigra, which is frequently accompanied by an accumulation of α -synuclein in β -sheet filaments in these neurons (so-called Lewy bodies) and neurites [1]. This aggregation process is thought to underlie the disease’s toxicity, with intermediate α -synuclein oligomers being the toxic agent [1]. The accumulation of misfolded α -synuclein in PD is considered to be due to increased expression [2,3] or reduced degradation via the ubiquitin proteasome, the lysosome, and the autophagy system [4,5].

α -synuclein is known to undergo mono- and polyubiquitination; the former modification is normally involved in regulating protein trafficking, and the latter is considered a prerequisite for degradation. However, for small proteins such as α -synuclein, monoubiquitination seems to be sufficient for proteasomal degradation [4–7] (Figure 1). For α -synuclein, it has been proposed that the non-ubiquitinated protein might be slowly degraded by autophagy, the monoubiquitinated protein might be degraded by the proteasome, and the

polyubiquitinated protein may be degraded by the proteasome and lysosome [4,8]. α -synuclein can be ubiquitinated on nine different lysine residues, lysines 6, 10, 12, 21, 23, 32, 34, 43 and 96, with different preferences in monomeric, oligomeric, and aggregated α -synuclein, as N-terminal monoubiquitination stimulates aggregation and proteasomal degradation [4,5].




Figure 1. NEDD4-mediated ubiquitination of protein substrates. Attachment of ubiquitin (Ub) to proteins is usually catalysed by an enzymatic cascade of a ubiquitin-activating enzyme E1, a ubiquitin-binding/conjugating enzyme E2, and a ubiquitin–protein ligase enzyme E3 that catalyses the transfer of the C-terminal carboxyl group of ubiquitin to the lysine (K) ϵ -amino group of the specific substrate. The process of ubiquitination can occur on transmembrane proteins (e.g., RET, ion channels) and on intracellular proteins (e.g., α -synuclein). The fate of the protein is dependent upon the number of ubiquitin moieties attached to each other on a substrate and which amino acid in ubiquitin the chain is extended: one of the seven lysines (K6, K11, K27, K29, K33, K48, K63) or, through its N-terminal, methionine (M1). Monoubiquitination and multimono-ubiquitination of a transmembrane protein generally result in its transport, internalisation, and recycling. Linear and branched polyubiquitination with K48-linked chains results in proteasomal degradation of the substrate, and that with K63 extension regulates protein–protein interactions, protein activity, DNA

repair, autophagy, immunity, inflammation, and protein trafficking to the lysosome [9]. The primary role(s) of each of the eight distinct polyubiquitin chains formed at one of the seven lysine residues or the primary methionine are indicated (orange box) [4,5,8]. ER = endoplasmic reticulum; ERAD = Endoplasmic-reticulum-associated protein degradation; TCR = T-cell receptor; TLR2/4 = Toll-like receptor 2 and 4.

The attachment of ubiquitin to proteins (“ubiquitination”) is usually catalysed by an enzymatic cascade of a ubiquitin-activating enzyme E1 (only two in the human genome), a ubiquitin-binding/conjugating enzyme E2 (around 35 in the human genome), and a ubiquitin–protein ligase enzyme E3 (around 600 in the human genome) that catalyses the transfer of the C-terminal carboxyl group of ubiquitin to the lysine (K) ϵ -amino group of the specific substrate. E3 ligases have at least two domains: a region to interact with an E2 enzyme and a region to recognise the specific substrate proteins. Based on the E2 interaction domain, E3 enzymes can be grouped into two families, HECT (homologous to human papillomavirus oncogene E6-associated protein carboxy-terminus) domain E3s and the more frequent single- and multisubunit RING (really interesting new gene; two zinc ions in a cross-braced arrangement of eight cysteines and histidines) and RING-like (U-box found in the polyubiquitin chain elongation protein E4 *saccharomyces cerevisiae* Ufd2 protein with noncovalent interactions of core amino acids forming a RING-like tertiary structure without zinc and plant homeodomain/proline-hydroxylase-domain/leukaemia-associated protein (PHD/LAP) with zinc) domain E3s [10].

Several different E3 ubiquitin–protein ligases have been described to be able to ubiquitinate different forms of α -synuclein [5], but it remains a matter of debate which E3 ubiquitin–protein ligases might be crucial for α -synuclein degradation, how mutations and misfolding of α -synuclein reduce its recognition by E3 enzymes as a substrate, and whether the E3 enzyme activity itself might be altered in PD. Besides the possible redundancy of different E3 ligases, it seems a common theme that E3 ligases ubiquitinate several different substrates. Therefore, accumulated α -synuclein might block E3 ubiquitin ligase activity and lead to the accumulation of other substrates, which may subsequently contribute to the disease aetiology.

After summarising the different E3 ubiquitin–protein ligases, which have been suggested to use α -synuclein as a substrate, we focus on one group of them—the NEDD4 family, which has many additional PD-linked substrates—and support the idea that NEDD4 family members can be considered as therapeutic targets to treat PD.

2. E3 Ligases Ubiquitinating α -Synuclein

Interestingly, members of the single-subunit (parkin, SIAH, CHIP) and multiple-subunit ($\text{SCF}^{\text{F}^{\text{XBL5}}}$) RING domain E3 ligase family, as well as the HECT domain family of E3 ligases (NEDD4 family), have been found capable of ubiquitinating α -synuclein.

Parkin was the first E3 ubiquitin–protein ligase described to ubiquitinate α -synuclein *in vitro* and required the presence of the E2 ubiquitin-conjugating enzyme UbcH7 [11]. However, parkin was only able to ubiquitinate a post-translationally modified form of α -synuclein, a specific 22-kilodalton O-glycosylated form of α -synuclein that could also be detected in PD and dementia with Lewy body patients [11]. Parkin was found to be mutated in some familial cases of PD [12], and all parkin mutations seemed to block ubiquitination activity [13]. Parkin is usually autoinhibited, requires self-ubiquitination for its activation and has been shown to label proteins for degradation by the proteasome or lysosome [14]. Parkin has also been shown to ubiquitinate the α -synuclein-interacting protein synphilin-1, which is a presynaptic protein localised to synaptic vesicles, like α -synuclein, and is a constituent of Lewy bodies, like α -synuclein and parkin [15,16]. Recently, α -synuclein was shown to lead to S-nitrosylation, autoubiquitination, and degradation of parkin [17]. However, the relevance of this crosstalk between parkin and α -synuclein to the development and progression of PD remains uncertain.

Next, the two members of the SIAH (mammalian homologues of *Drosophila* seven in absentia) family of E3 ligases, SIAH-1 and SIAH-2 [18], were reported to polyubiquitinate,

with the E2 enzyme UbcH5 synphilin-1 promoting their degradation through the ubiquitin–proteasome pathway. α -synuclein was only mono- or diubiquitinated by SIAH-2 and was reported not to be degraded by one laboratory [19] and subsequently degraded via the proteasome pathway by another laboratory [8]. USP9X was shown to be able to remove α -synuclein monoubiquitination generated by SIAH-2 and thereby prevent α -synuclein protein degradation [8]. In addition, SIAH1 was shown to mono- and diubiquitinate α -synuclein on lysines 10, 12, 21, 23, 34, 43, and 96 together with the E2 enzyme UbcH8, which did not affect the degradation of α -synuclein but increased its insolubility, aggregation, and cellular toxicity [20,21]. Interestingly, only the autosomal dominant mutation A30P of α -synuclein in familial PD (and not A53T) abolished SIAH-1 mediated ubiquitination [20]. As for parkin, the in vivo relevance of SIAH-dependent ubiquitination of α -synuclein remains to be shown.

The E3 ligase CHIP (carboxy-terminus of Hsc70 interacting protein) is a multidomain chaperone with a tetratricopeptide/Heat shock protein 70 binding domain and a U-box/ubiquitin ligase domain [5]. Interestingly, it was shown that the tetratricopeptide repeat domain of CHIP is critical for proteasomal degradation of α -synuclein, whereas the U-box domain of CHIP is sufficient to direct α -synuclein toward the lysosomal degradation pathway [22]. Subsequently, it was suggested that CHIP selectively reduced α -synuclein oligomerisation and toxicity in a tetratricopeptide domain-dependent, U-box-independent manner by specifically degrading toxic α -synuclein oligomers [23]. The ubiquitination of oligomeric α -synuclein by CHIP and UbcH5b can be negatively regulated by the Hsp70-mediated association with the co-chaperone BCL-2-associated athanogene 5 (BAG5) with CHIP [24]. The proof that this is critical for PD pathology still needs to be provided.

More recently, an SCF ubiquitin ligase assembled by the S-phase kinase-associated protein (SKP1), cullin-1 (Cul1), a zinc-binding RING finger protein, and the F-box domain/Leucine-rich repeat protein 5-containing protein FBXL5 (SCF^{FBXL5}) was shown to target exogenous α -synuclein and inhibit aggregation in vitro and in vivo in mice [25]. This observation is interesting in regard to α -synuclein seeding and spreading along the gut–brain axis and inside the brain but awaits independent confirmation.

Finally, from the nine human NEDD4 (neural precursor cell expressed developmentally downregulated protein 4) family members, which are NEDD4-1/NEDD4, NEDD4-2/NEDD4L (NEDD4-like), ITCH/AIP4 (itchy/atrophin-1 interacting protein 4), SMURF1 (SMAD-specific E3 ubiquitin–protein ligase 1), SMURF2, WWP1 (WW domain-containing E3 ubiquitin–protein ligase 1), WWP2/AIP2, NEDL1 (NEDD4-like ubiquitin–protein ligase 1), and NEDL2, at least five have been characterised to ubiquitinate α -synuclein and promote its degradation [26,27]. NEDD4-1 together with UbcH5 and UbcH7 used mainly ubiquitin K63 but also K29 and K33 to polyubiquitinate α -synuclein and enhance its lysosomal degradation [28,29] (see Figure 1). Other WW domain/HECT-domain E3s, NEDD4-2, SMURF1, and SMURF2, were reportedly unable to ubiquitinate α -synuclein to the same extent as NEDD4-1 [28]. As detailed below, NEDD4 ligases have three or four tryptophan-rich (WW) domains that mediate protein–protein interactions with an xPxY (PY) motif (often PPxY or LPSY with x being any amino acid) motif on substrates and adaptors. α -synuclein does not contain a PY sequence and instead has proline-rich regions near its C-terminus [27]. It has been proposed that these stretches may mediate recognition of α -synuclein by NEDD4 ligases [28,29]. Recognition of α -synuclein by NEDD4 family enzymes is thought to happen not only through conventional binding to the WW domains of NEDD4 but through the C2 and HECT domains of NEDD4 [27]. It has previously been demonstrated that NEDD4-1 recognises the C-terminus of α -synuclein and subsequently also ubiquitinates α -synuclein on K21 and K96 [28]. Interestingly, overexpression of NEDD4-1 in *Drosophila* rescued the rough eye phenotype induced by α -synuclein overexpression, and in rats, adeno-associated virus (AAV)-mediated NEDD4-1 expression rescued the loss of midbrain dopaminergic neurons induced by AAV-mediated overexpression of human A53T α -synuclein [30] (see also Tables 1–3).

Table 1. Substrates, adaptors, and modifiers/regulators of NEDD4-1.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
ABCB1 (ATP-binding cassette sub-family B member 1/P-Glycoprotein)	✓			PDY	Polyubiquitination	An ATP Binding Cassette transporter that exports β -amyloid from Blood-brain barrier endothelial cells. Potential for intervention in Alzheimer's disease	In vitro: Sf21 cells	[31]
α -synuclein	✓			PDNEAYEMP, PLY, PPLP, PPSP, PFY	Monoubiquitination and Polyubiquitination (K63-linked)	Lysosomal degradation. Potential protection mechanism against Parkinson's Disease pathogenesis	In vitro: SH-SY5Y cells In vivo: <i>Drosophila</i> and Rat	[28–30]
Beclin 1	✓			LPLY	Beclin 1: polyubiquitination (K63 and K11-linked)	Subunit of the class III phosphatidylinositol 3-kinase complex. Autophagy-related protein. Proteasomal degradation. Inhibited autophagy and cell survival	In vitro: HeLa cells	[32]
β 2-AR (β 2-Arrestin Receptor)	✓				Ubiquitination	Involved in internalised receptor degradation and lysosomal trafficking. Degraded via the lysosome	In vitro: HEK293 cells	[33]
β -arrestin 1		✓				Adaptor protein for NHE1 ubiquitination	In vitro: HEK293 cells, mouse embryonic fibroblasts	[34]
β -arrestin 2		✓	✓			Adaptor protein for β 2-adrenergic receptor ubiquitination	In vitro: HEK293 cells	[33]
Caspase-1, -3, -4, -6, -7, -11	✓		✓		Truncation	NEDD4-1 can be cleaved by caspases and can K48-polyubiquitinate caspase 11	In vitro: Jurkat cells A549	[14,35–37]
Cbl-b (Casitas B-lineage Lymphoma b)	✓		✓		Polyubiquitination	Impedes NEDD4-1 interaction with PTEN and also polyubiquitinates Cbl-b for degradation	In vivo: <i>Cbl-b</i> ^{C373A} and <i>NEDD4</i> ^{Gt(IRES βgeo)} 249Lex mice	[38,39]
CNrasGEF (Cyclic Nucleotide rat sarcoma virus Guanine nucleotide Exchange Factor)	✓			PPGY, PPDY	Polyubiquitination	RAS guanine nucleotide exchange factors that are degraded via the proteasome. NEDD4-1 overexpression promoted migration and invasion of glioma cells	In vitro: HEK293T and Glioma cells	[40,41]
Connexin43	✓			PPGY	Ubiquitination	Gap junction protein. Proteasome and lysosome degradation	In vitro: WB-F344 rat liver epithelial cells	[42]

Table 1. Cont.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
c-Src (Proto-oncogene tyrosine-protein kinase Src)			✓			Tyrosine kinase that activates NEDD4-1 through phosphorylation of its HECT and C2 domains. Phosphorylation inhibits auto-regulation	In vitro: HeLa, HEK293T and Platinum E cells	[43]
Δ Np63 transcriptional target	✓	✓		PPPY	Ubiquitination and polyubiquitination	A homologue of p53 tumour suppressor. Protein destabilisation of ΔNp63α and degradation. Downregulates NEDD4-1 leading to the suppression of nuclear PTEN in basal layer keratinocytes	In vitro: HEK293-EBNA, HaCaT, A431 and H1299 cells In vivo: Zebrafish embryos	[44,45]
EPS15 (Epithelial growth factor receptor substrate 15)	✓				Monoubiquitination and polyubiquitination	An endocytic protein that is targeted for degradation by NEDD4-1	In vitro: HeLa and B82L cells	[46]
FGFR1 (Fibroblast Growth Factor Receptor 1)	✓	✓		VLLVRP SRLSSSG	Ubiquitination	FGFR1 is a tyrosine kinase involved in cell proliferation and differentiation during development. Inhibited neural stem cell differentiation. Activates c-Src that subsequently activates NEDD4-1	In vitro: Human embryonic stem cells In vivo: Zebrafish embryos	[47,48]
GAG (group specific antigen) protein	✓			PPPY, PTAP	Monoubiquitination	From the HTLV-1 (human T-lymphotropic virus type 1) involved in hijacking multivesicular body (MVB) pathway proteins required for viral budding	In vitro: HEK293T cells	[49]
γ2-adaptin	✓	✓		PPAY	Monoubiquitination and multi-polyubiquitination	A member of the clathrin adaptor protein family. Forms a complex with NEDD4-1 and is involved in endosomal/multivesicular body (MVB) pathway and the assembly and release of the HBV	In vitro: HuH-7 and HeLa cells	[50,51]
HBV X protein (Hepatitis B virus)	✓				Ubiquitination	A multifunctional regulator that is encoded by the HBV genome. It is degraded via the K48 proteasomal pathway	In vitro: HEK293T, HBV-related HCC cell lines - HepG2.215, HepG3B, SNU182, SNU387, PLC/PRF/5, and MHCC97H	[52]

Table 1. Cont.

Name	Substrate Adaptor Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
HER3 (human epidermal growth factor receptor 3)	✓	PPRY	Polyubiquitination	A member of the EGRF family. Degraded via the proteasome. Inhibited cancer cell proliferation and tumour growth. NEDD4-1 knockdown induces apoptosis in DU145 cells	In vivo: Chinese Hamster ovary cells In vitro: MDA-MB-453, MCF-7, and DU145 cells	[53,54]
HGS (Hepatocyte growth factor- regulated tyrosine kinase substrate)	✓	PPEY	Ubiquitination	Binding partner for NEDD4-1 involved in EGFR lysosome degradation	In vitro: Chinese hamster ovary and HeLa cells	[55]
IFITM3 (Interferon (IFN)-induced transmem- brane protein 3)	✓	PPNY	Polyubiquitination	A cell-intrinsic factor that limits influenza virus and other viral infections such as SARS	In vitro: HEK293T, A549, NCI-H358, NCI-H2009 cells and MEFs	[56]
IGF1R (Insulin-like growth factor 1 receptor)	✓		Ubiquitination	A tyrosine kinase receptor. Expression can be downregulated by NEDD4-1 through the indirect effect on the oxidation of very-low-density lipoproteins. Ubiquitination and degradation require a C1060 site	In vitro: Hepatocytes from Landes goose embryos In vivo: Intracerebral haemorrhage mice, Sprague Dawley and Tg2576 mice	[57]
IGPR-1 (Immunoglob- ulin and Proline-rich receptor-1, also known as TMIGD2/ CD28H)	✓	PPR	Polyubiquitination	A cell adhesion molecule involved in, for example, autophagy, angiogenesis and cell adhesion. Is degraded via the lysosomal pathway	In vitro: HEK293 cells	[58]
IRS-2 (insulin receptor substrate)	✓		Monoubiquitination	NEDD4-1 recruits IRS-2 to the membrane to enhance IGF signalling	In vivo: Zebrafish embryos In vitro: HEK293 cells	[59]
ISG15 (Interferon- stimulated gene 15)	✓		ISGylation	Can attach to NEDD4-1 inhibiting its ubiquitination properties	In vitro: HEK293, HeLa cells and MEFs	[60–62]
KLF8 (Krueppel- like factor 8)	✓		Ubiquitination	The function of this transcription factor is regulated by NEDD4-1	In vitro: HEK293 cells	[63,64]

Table 1. Cont.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
LATS1 (large tumour suppressor kinase 1)	✓				Ubiquitination	A serine/threonine kinase involved in the suppression of tumours	In vitro: HEK293 cells	[65]
LC3 (Microtubule-associated protein 1A/1B-light chain 3)		✓		WEII, WVVL, WFFL, WDKL		An autophagy-related protein. LC3 binds to NEDD4-1, but is not a ubiquitination substrate of NEDD4-1	In vitro: HEK293 cells	[66]
LDLRAD4 (Low density lipoprotein receptor class A domain containing 4)	✓				Ubiquitination	Degraded via the lysosome and is a negative regulator of TGF- β signalling	In vivo: Female BALB/c nude mice In vitro: L02 and HepG2 cells	[67]
LMP2A (Latent membrane protein 2A)	✓			PPPPY	Ubiquitination	A latent Epstein–Barr virus (EBV) infection protein involved in B cell signal transduction	In vitro: BJAB, Ramos, Raji, Jurkat, HPB.ALL and M12 cells	[68]
MDM2 (Mouse double minute 2 homolog)	✓				Polyubiquitination (K63-linked)	Is an E3 ubiquitin ligase involved in negative regulation of p53. MDM2 is stabilised via NEDD4-1 interacting with its RING domain. NEDD4-1 overexpression reduces p53 levels	In vivo: NEDD4-1 KO mouse embryonic fibroblasts	[69]
MEKK5 (mitogen-activated protein kinase 5) (Apoptosis Signal-regulating Kinase 1 (ASK1))		✓				A serine/threonine kinase that regulates NEDD4-1 cell migration signalling in lung cancer	In vitro: HEK293T, NCI-H1650, and A549 cells	[58]
N4BP (NEDD4-binding protein)	✓			PPLP, PPEY, PPPY	Monoubiquitination and Polyubiquitination	N4BP is degraded via the proteasome. NEDD4-1 regulates N4BP1 at promyelocytic leukaemia nuclear bodies	In vitro: HEK293 cells and MEFs	[70,71]
NAB (N-aryl benzimidazole)		✓?				NAB2 reduces the ratio of K63-linked ubiquitin chains on A53T α -synuclein by an unknown mechanism. NAB2 binds NEDD4-1 but does not alter conformation or enzymatic activity.	In vitro: SH-SH5Y cells	[72,73]

Table 1. Cont.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
NEDD4-1 (Neural precursor cell-expressed developmentally-down-regulated protein 4-1)	✓	✓			Autoubiquitination	K29-linked autoubiquitination. C2 and HECT domains bind resulting in autoubiquitination	In vitro: HEK293T, HeLa, THP-1 and A549 cells	[74,75]
NHE1 (Sodium-Hydrogen antiporter 1)	✓				Multi-monoubiquitination and polyubiquitination	The Na(+)/H(+) exchanger 1 is ubiquitinated for degradation by NEDD4-1 but requires β -arrestin 1	In vitro: HEK293 cells	[34]
N-Myc (N-myc proto-oncogene protein/ basic helix-loop-helix protein 37)	✓				Polyubiquitination	An oncoprotein that is degraded via the proteasome. NEDD4-1 suppresses neuroblastoma and pancreatic cancer cell proliferation	In vitro: Neuroblastoma BE(2)-C, CHP134, pancreatic cancer MiaPaca-2 and HEK293 cells	[76]
Notch	✓			PPSY	Polyubiquitination	A plasma membrane receptor that is ubiquitinated for degradation via the proteasome	In vivo: <i>Drosophila</i> and Conditional NEDD4-1 overexpression in Wistar Rat	[77–79]
Rap2a (RAS-related protein 2a)	✓				Monoubiquitination of K63	Rap2a Is a member of the RAS-related protein family. NEDD4-1 Inhibits GTP-Rap2a activity subsequently promoting the migration and invasiveness of glioma cells	In vitro: Human glioma cell lines U251 and U87	[80]
RAS (Rat sarcoma virus)	✓			PPGY, PPDY	Polyubiquitination and monoubiquitination	Small guanosine triphosphatases involved in a multitude of different cellular processes by acting as a molecular switch. RAS is regulated via NEDD4-1ubiquitination sending it for degradation to the lysosome. This regulation suppressed tumorigenesis	In vitro: HEK293T, HeLa, NIH 3T3, MEF and HepG2 cells	[40,81]

Table 1. Cont.

Name	Substrate Adaptor Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
RET (Rearranged during transfection)	✓		Polyubiquitination	A receptor tyrosine kinase. The short form (Ret9) becomes localised and internalised into the endosomal network through clathrin-coated pits following NEDD4-1 ubiquitination. This causes inhibition of Ret9-mediated neurotrophic signalling at the cell surface and promotion of post-internalisation signalling. This mechanism could potentially impact neurotrophic signalling of dopaminergic neurons and play a role in Parkinson's disease	In vitro: HEK293 and SH-SY5Y cells	[82]
RNAPII (Ribonucleic acid Polymerase II)	✓		Monoubiquitination and polyubiquitination	A multiprotein involved in the transcription of DNA into mRNA that is degraded via the proteasome after being ubiquitinated by NEDD4-1. This ubiquitination is dependent on NEDD4-1 interacting with the ElonginA/B/C-Cullin 5 complex	In vitro: HEK293, MRC5 and <i>S. cerevisiae</i>	[83,84]
RTP801/REDD1	✓		Polyubiquitination	A pro-apoptotic protein that is targeted for degradation by NEDD4-1 via K63 ubiquitin linkages. NEDD4-1 loss may elevate RTP801 proteins leading to an increase in neuronal death in Parkinson's disease	In vivo: <i>NEDD4-1^{f/f}</i> , <i>Emx1</i> Cre mice In vitro: PC12, HEK293 cells and rat primary cortical neurons	[85]
SAG (S-Arrestin)	✓		Polyubiquitination	An anti-apoptotic cellular survival protein that is degraded by the proteasome. NEDD4-1 reduction of SAG resulted in etoposide-induced apoptosis in cancer cells. SAG does not bind to WW domains as it lacks PY motifs but interacts with NEDD4-1 via its RING domain	In vitro: HEK293T	[86]
SCAMP3 (Secretory Carrier Membrane Protein 3)	✓	PPAY, PSAP, PTEP	Multi-monoubiquitination	Integral membrane proteins involved in the cell surface recycling system. SCAMP3 is a NEDD4-1 substrate that is involved in the degradation of EGFR via the lysosome	In vitro: HeLa and HEK293T cells	[87]

Table 1. Cont.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
Spy1A	✓				Polyubiquitination	A cyclin-like protein that is needed for a cell to progress through the G ₁ /S phase. Spy1A is required for p53-mediated tumour suppression. Spy1A is degraded in a cell cycle-dependent manner during mitosis via the ubiquitin-proteasome system	In vitro: Human mammary breast cancer, MCF7, and HEK293cells	[88,89]
SQSTM1 (p62)	✓				Polyubiquitination (K63-type)	An autophagy-related protein. NEDD4-1 ubiquitinates its PB1 domain. Lack of NEDD4-1 leads to accumulation of aberrant enlarged inclusion bodies	In vitro: HEK293T, HEK293A and A549 cells	[66,90]
VEGF-R2 (vascular endothelial growth factor receptor-2)	✓				Monoubiquitination	This receptor is degraded by NEDD4-1 but Grb10 regulates this process by interacting with NEDD4-1	In vitro: HEK-293 EBNA cells	[91]

?, experimental uncertainty. Further investigation required to confirm result. Abbreviations: PDY, Proline Aspartate Tyrosine; PDNEAYEMP, Proline Aspartate Asparagine Glutamate Alanine Tyrosine, Glutamate, Methionine Proline; PLY, Proline Leucine Tyrosine; PPLP, Proline Proline Leucine Proline; PPSP, Proline Proline Serine Proline; PFY, Proline Phenylalanine Tyrosine; LPLY, Leucine Proline Leucine Tyrosine; PPGY, Proline Proline Glycine Tyrosine; PPDY, Proline Proline Aspartate Tyrosine; PPPY, Proline Proline Proline Tyrosine; VLLVRPSRLSSSG, Valine Leucine Leucine Valine Arginine Proline Serine Arginine Leucine Serine Serine Serine Glycine; PTAP, Proline Threonine Alanine Proline; PPAY, Proline Proline Alanine Tyrosine; PPRY, Proline Proline Aspartate Tyrosine; PPEY, Proline Proline Glutamate Tyrosine; PPNY, Proline Proline Asparagine Tyrosine; PPR, polyproline rich; WEIL, Tryptophan Glutamate Isoleucine Isoleucine; WVVL, Tryptophan Valine Valine Leucine; WFFL, Tryptophan Phenylalanine; WDKL, Tryptophan Aspartate Lysine Leucine; PPPPY, Proline Proline Proline Proline Tyrosine; PPSY, Proline Proline Serine Tyrosine; PSAP, Proline Serine Alanine Proline; PTEP, Proline Threonine Glutamate Proline.

Table 2. Substrates, adaptors, and modifiers/regulators of NEDD4-2.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
14-3-3	✓	✓	✓			14-3-3 is an inhibitory binding partner for NEDD4-2 through a PI3-kinase/SGK1-dependent manner. This interaction is dependent on the phosphorylation of key residues Ser ³⁴² and Ser ⁴⁴⁸ on NEDD4-2	In vitro: Kidney tubule epithelial mpkCCDC14, HECT293 and <i>E. coli</i> BL21(DE3) cells	[92,93]

Table 2. Cont.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
α -Arrestins	✓	✓		PPLP, PPEY, PPLY, PPSY, PPNY, PPPY	Ubiquitination; adaptor/regulator	α -Arrestins are scaffolding molecules involved in regulating receptor trafficking and cell signalling. These can be both substrates and adaptors for NEDD4-2. α -arrestins are implicated in the regulation of DMT1. It is activated through polymerisation or membrane tethering and is ubiquitinated by NEDD4-2	In vitro: CHO, Caco-2, HepG2 and HEK293T cells In vivo: <i>Arrdc1^{tm1(KOMP)Vlcr}</i> (VG17312) and <i>Arrdc4^{tm1(KOMP)Vlcr}</i> (VG18749) embryonic stem cells	[27,94,95]
AMPK (AMP-activated protein kinase)		✓				AMPK is a metabolic sensor that inhibits ENaC. It activates NEDD4-2 by phosphorylation, promotes ENaC-NEDD4-2 interaction and subsequent ENaC degradation. AMPK also signals in the mTOR pathway where it plays a role in cell death.	In vitro: HEK293 cells In vivo: <i>Xenopus</i> oocytes	[96]
ATA2 (Analogue Terminal Adaptor II)	✓				Polyubiquitination	NEDD4-2 regulates amino acid transporter ATA2 activity on the cell surface by proteasomal degradation	In vitro: 3T3-L1 adipocytes Ex vivo: <i>Xenopus</i> oocytes	[97]
CFTR (Cystic fibrosis transmembrane conductance regulator)	✓				Ubiquitination	CFTR is downregulated by NEDD4-2 possibly via the proteasome and lysosome degradation (contested by [98]). Ref. [99], however, show that CFTR is degraded via NEDD4-2 but has to interact with 14-3-3 ϵ and be activated by SGK1 phosphorylation	In vitro: CFPAC-1 (Δ F508) and CFBE41o- Δ F cells	[98–100]
CHT1 (high-affinity choline transporter 1)	✓				Ubiquitination	NEDD4-2-mediated ubiquitination regulates cell surface expression of CHT1 thereby impeding choline uptake and HC-3 binding	In vitro: HEK293 Cells	[101]
CLC-5 (Chloride Voltage-Gated Channel 5)	✓			PPLPPY	Ubiquitination	Voltage-gated channels that function as dimers. NEDD4-2 decreases cell surface expression of CLC-5 through ubiquitination	Ex vivo: <i>Xenopus</i> oocytes In vivo: NEDD4-2 null mice	[102–104]
CLC-K (Chloride channel protein CLC-Ka) /barttin	✓			PPYVRL (located on barttin)	Ubiquitination	A chloride channel that requires barttin to be functional. Downregulation of CLC-Ka/barttin comes as a result of NEDD4-2-mediated ubiquitination	Ex vivo: <i>Xenopus</i> oocytes	[105,106]
CRTC3 (CREB Regulated Transcription Coactivator 3)	✓			PPPY	Polyubiquitination	NEDD4-2 is responsible for the downregulation of CRTC3 in a proteasome-dependant manner in response to prolonged cAMP signalling	In vitro: HEK293T cells	[107]

Table 2. Cont.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
DAT (Dopamine Transporter)	✓				Polyubiquitination	A transporter for dopamine. PKC-dependent DAT ubiquitination by NEDD4-2 requires its WW3 and WW4 domains. May have implications in Parkinson's disease	In vitro: HEK 293 and PAE cells	[108]
DMT1 (Divalent metal transporter 1)	✓				Polyubiquitination	A metal transporter that is ubiquitinated by NEDD4-2 but requires the adaptor protein Ndfip1	In vitro: SH-SY5Y and HEK293T cells	[109]
DVL2 (Dishevelled 2)	✓			PPPY	Polyubiquitination	NEDD4-2 negatively regulates Wnt signalling by targeting dishevelled for proteasomal degradation. Wnt5a induces JNK-mediated phosphorylation of NEDD4-2, which in turn promotes Dvl2 degradation	In vitro: HEK293T and HeLa cells	[110,111]
EAAT1/2 (Excitatory amino acid transporter 1 and 2. Also known as Glutamate transporter 1 and 2)	✓			PPPD	Ubiquitination	Regulation of EAAT1/2 through NEDD4-2 depends on SGK kinases. NEDD4-2 knockdown with shRNA decreases GLT-1 ubiquitination, promoting glutamate uptake and increases GLT-1 expression. This may play a role in glutamatergic signalling in dementia	Ex vivo: <i>Xenopus</i> oocytes In vitro: MPP ⁺ treated astrocytes	[112–114]
GluA1 (Glutamate receptor 1)	✓			PKY	Ubiquitination	NEDD4-2 ubiquitinates GluA1 at lysine-868 and mediates its surface expression. This may play a role in glutamatergic signalling in dementia	In vitro: HEK293 cells In vivo: <i>NEDD4-2^{andi}</i> mice	[115]
hERG(1) (human ether-à-gogo-related gene (1))	✓			PPAY	Monoubiquitination and polyubiquitination	The human ether-a-go-go-related gene protein (hERG) is a voltage-gated cardiac potassium channel. Caveolin-3 (Cav3), hERG, and NEDD4-2 form a complex. hERG expression in the plasma membrane is regulated by Cav3 through NEDD4-2 ubiquitination	In vitro: HEK293 cells (Patch clamp) In vivo: Guinea pig	[38,116]
IKKβ (inhibitor of nuclear factor kappa-B kinase subunit β)		✓	✓			IKKβ activates NEDD4-2 via phosphorylation that results in the regulation of ENaC	In vitro: HEK-293, HEK-293T and mpkCCD _{c14} cells. Ex vivo: <i>Xenopus</i> oocytes (TEV)	[117]
JNK1 (Janus Kinase 1)		✓	✓			JNK1 activates NEDD4-2 via phosphorylation that results in the regulation of ENaC	In vitro: HEK293 cells and mpkCCD _{c14} cells. Ex vivo: <i>Xenopus</i> oocytes (TEV)	[118]

Table 2. Cont.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
KCNQ (Voltage-Gated Potassium Channel Subunits) 1, 2/3, 3/5	✓			PPDPPY	Polyubiquitination	Amplitude of K ⁺ currents mediated by KCNQ2/3 and KCNQ3/5 were reduced by NEDD4-2. NEDD4-2 is activated by AMPK leading to reduced KCNQ1 expression	In vitro: HEK293 cells Ex vivo: <i>Xenopus</i> oocytes	[116,119–121]
Na _v s (Voltage-gated sodium channels) 1.2, 1.3, 1.5, 1.6, 1.7, 1.8	✓			PPSY, LPSY, PGSP	Ubiquitination	Are vital in creating and propagating action potentials and reside in the membrane. NEDD4-2 interaction inhibits activity of multiple Na _v s, including the cardiac (Na _v 1.5) and neuronal Na _v s (Na _v 1.2, Na _v 1.7, and Na _v 1.8)	In vitro: HEK293 cells In vivo: SNS- <i>NEDD4-2</i> ^{−/−} mice, Pulldown of mouse brain lysates Ex vivo: <i>Xenopus</i> oocytes	[122–124]
NCC (Sodium Chloride symporter)	✓				Ubiquitination	NCC ubiquitination at the cell surface is achieved by NEDD4-2 and its deficiency upregulate NCC. NEDD4-2 may require another protein to achieve this	In vitro: HEK293 and mDCT ₁₅ cells Ex vivo: <i>Xenopus</i> Oocyte In vivo: <i>Pax8-rtTA</i> and <i>TRE-LC1</i> transgenic mice	[125,126]
NEDD4-2 (Neural precursor cell-expressed developmentally-downregulated gene/protein 4-2)	✓	✓		LPPY	Inhibitory self-ubiquitination of NEDD4-2	Promotes NEDD4-2 stabilisation through auto-ubiquitination involving its own PY motif located on its HECT domain. This interaction may be between an active and non-active form	In vitro: HEK293 cells Ex vivo: <i>Xenopus</i> oocytes	[104,127, 128]
NEDD8 (Neural precursor cell-expressed developmentally-downregulated gene/protein 8)		✓			Neddylation	Neddylation is a process whereby the ubiquitin-like protein, Nedd8, is conjugated to NEDD4-2 resulting in its degradation	In vitro: Mouse M1 kidney and mouse NCTC1469 liver cells In vivo: <i>CYP4F2</i> transgenic mice	[129]
NHE3 (Sodium–hydrogen antiporter 3)	✓			PPNY	Ubiquitination	An Na ⁺ /H ⁺ exchanger that is expressed within the kidney where it is involved in blood pressure regulation through NaCl and HCO ₃ [−] absorption. NHE is ubiquitinated by NEDD4-2. Disruption of NEDD4-2 interaction elevates human NHE3 expression and activity	In vitro: HEK 293 and PS120 cells	[130]

Table 2. Cont.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
NKCC1/2 (Na-K-Cl co-transporter 1/2)	✓					An Na ⁺ /K ⁺ /2Cl [−] co-transporter where NEDD4-2 is involved in its downregulation. NEDD4-2 indirectly suppresses NKCC1 expression	In vitro: HEK293T In vivo: Tam-induced <i>NEDD4-2^{fl/fl}</i> ; Vil-Cre ^{E^{ERT2}} mice	[36]
OAT (Organic ion transporters) 1/3	✓				Polyubiquitination and multiubiquitination	NEDD4-2 regulates cell surface OAT1/3 expression and its transport activities	In vitro: COS-7 and HEK293T cells	[131,132]
Occludin	✓			PPPY	Polyubiquitination	An integral membrane protein that NEDD4-2 ubiquitinates. NEDD4-2 overexpression reduced occluding at tight junctions	In vitro: HEK293 and mplCCD _{c14} cells	[133]
PKA (Protein Kinase A)			✓			Inhibitory phosphorylation of NEDD4-2. cAMP regulates ENaC through phosphorylation & inhibition of NEDD4-2	In vitro: COS-7, FRT epithelial cells	[134]
SCAMP3 (Secretory Carrier Membrane Protein 3)	✓			PPAY	Multi-monoubiquitination	SCAMP3 ubiquitination is involved in the degradation of EGFR via the lysosome	In vitro HeLa and HEK293T cells	[87]
SGK1 (Serum/Glucocorticoid Regulated Kinase 1)		✓	✓	PPFY		Regulates the activity of several ion transport proteins. Inhibitory Phosphorylation of NEDD4-2 causes its interaction with 14-3-3 and subsequent degradation	In vitro: Kidney tubule epithelial and COS7 cells	[92,135, 136]
SGLT1 (sodium-glucose linked transporter 1)	✓				Ubiquitination	NEDD4-2 ubiquitinates and downregulates SGLT1	Ex vivo: <i>Xenopus</i> oocytes	[137]
Smad (Mothers against decapentaplegic homolog) 2, 3, 4, 7	✓			PPPY	Polyubiquitination	NEDD4-2, interacts with Smads, inducing their polyubiquitination and degradation. This is not the case for Smad3	In vitro: COS7 (Smad2) and HEK293T (Smad 2,3,4&7) cells	[136,138–141]
SP-C (Surfactant Protein C)	✓			PPDY	Monoubiquitination or biubiquitination at K6	NEDD4-2-mediated ubiquitination regulates luminal relocation of SP-C, leading to its processing and secretion	In vitro: HEK293 cells	[142]
TGF-βR1 (Transforming Growth Factor-β Receptor 1)	✓					Plays a role in epithelial-mesenchymal transition via phosphorylation of small mothers against decapentaplegic (SMAD). NEDD4-2 suppresses its signalling	In vivo: Sprague Dawley rats. In vitro: NRK-52E cells	[143]

Table 2. Cont.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
TrkA (Tropomyosin-related kinase Trk A)	✓			PPVY, PPSY, PPIY	Multi-monoubiquitination	Tropomyosin-related kinase (Trk) A is a receptor specifically for nerve growth factor and is downregulated by NEDD4-2	In vitro: PC12-615 cells and primary cortical neurons	[144–146]
Tweety	✓			PPTY	Ubiquitination	A family of chloride ion channels. NEDD4-2-mediated ubiquitination of TTYH2 regulates both cell surface and total levels of Tweety proteins	In vitro: HEK293 cells	[147]
ULK1 (Unc-51 like autophagy activating kinase 1)	✓				Polyubiquitination	A serine–threonine kinase involved in autophagy. NEDD4-2 ubiquitinates ULK1 and targets it for proteasomal degradation	In vitro: HeLa cells	[148]
Usp2-45 (Ubiquitin-specific protease 2-45)		✓				Adaptor of NEDD4-2 for ENaC Ubiquitination	In vitro: HEK293 cells	[149]
WNK1 (With No Lysine Kinase)	✓			PPQY, PFY	Ubiquitination	Serine-threonine kinases that regulate potassium, sodium, and blood-pressure homeostasis. Hormonal (Aldosterone, insulin and vasopressin) regulation of NEDD4-2 and WNK to regulate NCC (thiazide-sensitive NaCl cotransporter)	In vitro: mpkCCD _{c14} and HEK294T Cells In vivo: NEDD4-2 ^{fl/fl} Pax8-rtTATRE-LC1 (Renal-specific NEDD4-2 KO) mice	[150]

Abbreviations: PPLP (Proline Proline Leucine Proline), PPEY (Proline Proline Glutamate Tyrosine), PPLY (Proline Proline Leucine Tyrosine), PPSY (Proline Proline Serine Tyrosine), PPNY (Proline Proline Asparagine Tyrosine), PPPY (Proline Proline Proline Tyrosine), PPLPPY (Proline Proline Leucine Proline Proline Tyrosine), PPYVRL (Proline Proline Tyrosine Valine Arginine Leucine), PPPD (Proline Proline Proline Aspartate), PKY (Proline Lysine Tyrosine), PPAY (Proline Proline Alanine Tyrosine), PPDPY (Proline Proline Aspartate Proline Proline Tyrosine), LPTY (Leucine Proline Threonine Tyrosine), LPSY (Leucine Proline Serine Tyrosine), PGSP (Proline Glycine Serine Proline), LPPY (Leucine Proline Proline Tyrosine), PPFY (Proline Proline Phenylalanine Tyrosine) (Proline Proline Asparagine Tyrosine), PPDY (Proline Proline Aspartate Tyrosine), PPVY (Proline Proline Valine Tyrosine), PPIY (Proline Proline Isoleucine Tyrosine), PPQY (Proline Proline Glutamine Tyrosine), PFY.

Table 3. Substrates, adaptors, and modifiers/regulators of both NEDD4-1 and NEDD4-2.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
ACK-1 (activated Cdc42-associated kinase 1) (NEDD4-1)	✓			PPAY	Monoubiquitination, Polyubiquitination	ACK-1 is a cytoplasmic tyrosine kinase and is a NEDD4-1 and NEDD4-2 substrate. Its degradation through the proteasome results in downregulation of ACK-1. These authors suggest only NEDD4-1 to ubiquitinate ACK-1	In vitro: COS7, HEK293T, HeLa, T47D, and A549 cells	[151]

Table 3. Cont.

Name	Substrate	Adaptor Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
ACK-1 (activated Cdc42- associated kinase 1) (NEDD4-2)	✓		PPAY	Polyubiquitination	Its degradation through the proteasome results in downregulation of ACK-1 although this is contested [151]	In vitro: HeLa and COS7 cells	[152]
α -synuclein filaments (NEDD4-1)	✓		PDNE AYEMP, PLY, PPLP PPSP, PFY	Monoubiquitination and Polyubiquitination (K63-linked)	Degraded via the lysosome. Potential protection mechanism against Parkinson's Disease pathogenesis	In vitro: SH-SY5Y In vivo: Drosophila and Rat	[28–30]
α -synuclein filaments (NEDD4-2)	✓			Polyubiquitination		In vitro: SH-SY5Y cells	[29]
AKT (Protein Kinase B) (NEDD4-1)	✓		LPEY, LPFY	Polyubiquitination (K63 and K48-linked) and multi- monoubiquitination	A critical effector kinase that regulates numerous cellular processes such as cell growth, death, differentiation, and migration. NEDD4-1 regulates nuclear trafficking of the activated form of AKT via the proteasome & enhances bortezomib sensitivity	In vitro: MCF-7, HeLa, and NEDD4 ^{+/+} , and NEDD4 ^{-/-} cells and MEFs	[153,154]
AKT (Protein Kinase B) (NEDD4-2)		✓			Inhibitory phosphorylation of NEDD4-2. Inhibits NEDD4-2, increasing ENaC expression and Na ⁺ absorption	In vitro: FRT cells	[155]
AMPA (NEDD4-1)	✓			Ubiquitination	An ionotropic glutamate receptor. Ubiquitination leads to AMPAR internalisation and subsequent degradation	In vivo: Rat dissociated hippocampal or cortical neurons In vitro: HEK293 cells	[156–158]
AMPA (NEDD4-2)	✓			Ubiquitination	Ubiquitinates the GRIA1 subunit of AMPA thereby mediating neuronal excitation	In vivo: Sprague Dawley (SD) rats, Nedd4-2 ^{and} and GluA1 knockout mice In vitro: HEK293 cells	[115,159, 160]
AQP2 (Aquaporin 2) (NEDD4-1)	✓			Polyubiquitination	Ubiquitinates and degrades AQP2 but requires NDFIP1 and NDFIP2 adaptors	In vitro: HEK293 and mpkCCD cells	[161]
AQP2 (Aquaporin 2) (NEDD4-2)	✓			Polyubiquitination	Ubiquitinates and degrades AQP2 but requires NDFIP1 and NDFIP2 adaptors	In vitro: HEK293 and mpkCCD cells	[161]

Table 3. Cont.

Name	Substrate	Adaptor Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
Ca ²⁺ (Calcium ions) (NEDD4-1)		✓			Binds to the C2 domain of NEDD4-1 leading to the activation of its ligase activity and inhibition of auto-regulation. PIP2/IP3 ratio dictates its function in either the proximity of the membrane (PIP2) or cytoplasm (IP3)	Used biophysical techniques to complement the literature (NMR Spectroscopy)	[128]
DLG3 Discs Large MAGUK Scaffold Protein 3) (NEDD4-1 and NEDD4-2)	✓		PPGY, PPDY	Monoubiquitination	NEDD4-1 interaction results in Dlg3 monoubiquitination, apical membrane recruitment, and tight junction consolidation	In vitro: MDCK cells In vivo: Co-IP of mouse brain lysates	[162]
DVL2 (NEDD4-1 and NEDD4-2)	✓			Ubiquitination	A protein involved in the Wnt pathway. Reported to be ubiquitinated for degradation via NEDD4-1 and NEDD4-2	In vivo: NEDD4-1 ^{fl/fl} and NEDD4-2 ^{fl/fl} mice In vitro: HEK293T, DLD1 and HCT116 cells	[163]
ENaC (Epithelial Sodium Channel) (NEDD4-1)	✓		PPNY, PPRY, PPAY	Ubiquitination of α and γ subunits	The epithelial Na ⁺ channel is downregulated through proteasome degradation. Lack of binding motif in Liddle's syndrome causes hyperactivity	In vitro: rat foetal distal lung epithelial and <i>Xenopus</i> -derived A6 cells Ex vivo: <i>Xenopus</i> oocyte	[164,165]
ENaC (Epithelial Sodium Channel) (NEDD4-2)	✓		PPAY, PPNY, PPRY, PPKY	Monoubiquitination	NEDD4-2 catalyses ubiquitination and reduces expression of ENaC at the cell surface through lysosomal degradation. NEDD4-2 also targets Na ⁺ /Cl [−] cotransporter (NCC) in the kidney	In vitro: HEK293T cells	[116,126, 166,167]
Grb10 (Growth Factor Receptor Bound Protein 10) (NEDD4-1)		✓			An adaptor for NEDD4-1 to ubiquitinate and degrade IGF-IR. Required for indirect ubiquitination and degradation of VEGFR2	In vitro: p6, p6/Grb10, HEK-293 EBNA and HUVEC cells	[42,168–170]
Grb10 (Growth Factor Receptor Bound Protein 10) (NEDD4-2)		✓ ✓	PQTPF		Associates with NEDD4-2 leading to its regulation and inhibition of ubiquitination and degradation of K _v 1.3 channels	In vitro: HEK293 cells	[171]

Table 3. Cont.

Name	Substrate	Adaptor Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
LGR5 (Leucine-rich repeat containing G protein-coupled receptor 5) (NEDD4-1 and NEDD4-2)	✓			Ubiquitination	A receptor for R-spondin and is a protein involved in the Wnt pathway. It is apparently degraded via the lysosome and proteasome involving NEDD4-1 and NEDD4-2	In vivo: <i>NEDD4-1^{fl/fl}</i> and <i>NEDD4-2^{fl/fl}</i> mice In vitro: HEK293T, DLD1 and HCT116 cells	[163]
MTMR4 (NEDD4-1)	✓		PPLY	Ubiquitination	Myotubularin-related protein 4, an inositol phosphatase that regulates endosomal signalling	In vitro: HEK293 or HeLa cells	[172,173]
MTMR4 (NEDD4-2)		✓	PPLY		MTMR4 reduces NEDD4-2-mediated proteasome degradation of wild type and mutant KCNQ1 and hERG channels via dephosphorylation	In vitro: Induced pluripotent stem cardiomyocytes from human dermal fibroblasts	[174]
Nav (Voltage-gated Sodium channels)1.2 and 1.7 (NEDD4-1)	✓		PPSY (Proline Proline Serine Tyrosine)	Ubiquitination	Regulates sodium channels through lysosomal degradation	Ex vivo: <i>Xenopus Oocytes</i>	[175]
Navs (Voltage-gated Sodium channels) 1.2, 1.3,1.5, 1.6, 1.7, 1.8 (NEDD4-2)	✓		PPSY, LPSY	Ubiquitination	NEDD4-2 ubiquitination inhibits activity of multiple Navs, including the cardiac (Nav1.5) and neuronal Navs (Nav1.2, Nav1.7, and Nav1.8)	In vitro: HEK-293 cells. In vivo: SNS- <i>NEDD4-2^{-/-}</i> mice, Pulldown of mouse brain lysates Ex vivo: <i>Xenopus oocytes</i>	[42,116, 122–124]
Ndfip1/2 (NEDD4 family-interacting proteins) (NEDD4-1)		✓ ✓			NEDD4 family-interacting protein that activates the HECT domain of NEDD4-1. Ndfip1 binds substrates such as PTEN to act as an adaptor for NEDD4-1 ligase activity	In vitro: HEK-293T, SH-SY5Y, PC3 cells and primary MEFs In vivo: <i>C57BL/6J</i> , Emx1-Cre (<i>C57BL/6</i>) and <i>Nestin-Cre</i> mice (<i>B6.Cg-Tg(Nes-cre)1Kln/J</i>)	[176]

Table 3. Cont.

Name	Substrate	Adaptor	Modifier/Regulator	Binding Motifs	Modification	Function	Experimental Model Used	References
Ndfip1/2 (NEDD4 family-interacting proteins) (NEDD4-2)		✓	✓	PPPY, PPSY, LPTY, PSY, PTY		Ndfip activates HECT domain of NEDD4-2	Ex vivo: <i>Xenopus</i> oocytes	[42,161, 176]
OAT1 (organic anion transporter 1) (NEDD4-1)	✓				Ubiquitination	Organic anion transporter (OAT). NEDD4-1 is an important regulator for hOAT1 ubiquitination, expression, and function via its WW2 and WW3 domains	In vitro: COS-7 and HEK293T cells	[131]
OAT1/3 (Organic ion transporters 1/3) (NEDD4-2)	✓				Ubiquitination	NEDD4-2 ubiquitination regulates cell surface OAT1/3 expression with their WW3 and WW4 domains	In vitro: COS-7 and HEK293T cells	[131,132]
PTEN (Phosphatase and tensin homolog) (NEDD4-1& NEDD4-2)	✓?	✓		PRR	Multi-monoubiquitination Polyubiquitination (Ubiquitination independent of NEDD4-1)	PTEN is a tumour suppressor. Nuclear importation occurs via monoubiquitination NEDD4-1 single and NEDD4-1 & NEDD4-2 double knock out mice do not exhibit stability, subcellular activity or localisation differences of PTEN	In vitro: PC3 and HEK293T cells In vivo: <i>Xenopus</i> tectum In vivo: Murine	[111,176–182]

?, experimental uncertainty. Further investigation required to confirm result. Abbreviations: PPAY (Proline Proline Alanine Tyrosine), PDNEAYEMP (Proline Aspartate Asparagine Glutamate Alanine Tyrosine Glutamate Methionine Proline), PLY (Proline Leucine Tyrosine), PPLP (Proline Proline Leucine Proline), PPSP (Proline Proline Serine Proline), LPEY (Leucine Proline Glutamate Tyrosine), LPFY (Leucine Proline Phenylalanine Tyrosine), PPGY (Proline Proline Glycine Tyrosine), PPDY (Proline Proline Aspartate Tyrosine), PPNY (Proline Proline Asparagine Tyrosine), PPRY (Proline Proline Arginine Tyrosine), PPAY (Proline Proline Alanine Tyrosine), PPKY (Proline Proline Lysine Tyrosine), PQTPF (Proline Glutamine Threonine Proline Phenylalanine), PPLY (Proline Proline Leucine Tyrosine), PPSY (Proline Proline Serine Tyrosine), LPSY (Leucine Proline Serine Tyrosine), LPTY (Leucine Proline Threonine Tyrosine), PTY (Proline Threonine Tyrosine), PRR (Proline rich region).

More recently, in vitro-generated, β -sheet-containing α -synuclein filaments were found to be a better substrate for ubiquitination than monomeric α -synuclein, and wild-type α -synuclein was observed to be a better substrate than the mutated human A53T α -synuclein when testing the NEDD4 family members NEDD4-1, NEDD4-2, ITCH, SMURF2, and WWP2 [27]. Fibrils of α -synuclein enter the cytosol through a dynamin-dependent mechanism or by penetrating the plasma membrane directly [29]. NEDD4-1 in the cytosol binds the c-terminus of cytosolic α -synuclein through its WW, C2, and HECT domains and preferentially ligates a lysine⁶³-linked polyubiquitin chain to the protein. This ubiquitination facilitates the targeting of α -synuclein to endosomes. The ESCRT (endosomal sorting complex required for transport) complex then recognises the ubiquitinated α -synuclein and subsequently transports it to the late endosome via invagination of the endosomal membrane. This may then promote lysosomal degradation of the α -synuclein. The A53T mutation is located close to a region that is thought to form the core of the β -sheet-rich region [183]. It has been suggested that the A53T mutation may reduce the surface hydrophobicity of the β -sheet structure, in turn hindering binding of the ligase [27]. It may be that patients with the A53T mutation develop early-onset PD because mutant A53T α -synuclein forms filaments more rapidly than the wild-type protein [184]. In addition, it is plausible that lack of recognition, ubiquitination, and degradation may contribute to the

accumulation and spread of A53T α -synuclein, which the wildtype protein is not subject to (see also Tables 1–3).

Interestingly, the Lindquist group developed a phenotypic model of α -synuclein toxicity in yeast. They discovered a small molecule, N-arylbenzimidazole (NAB), that was able to alleviate many major phenotypic markers of α -synuclein toxicity [73,185]. Counter genetic screening showed that NAB activity was dependent on the yeast NEDD4 orthologue Rsp5. A further investigation in mammalian cell models indicated that NAB activity was conserved through evolution and was dependent on NEDD4-1 in these cells. An NAB derivative, NAB2, was found through structure–activity relationship optimisation of the NAB scaffold. NAB2 exhibited improved activity over that of NAB [73,185]. Although NEDD4-1 has a potential role in the cellular response to α -synuclein toxicity, it is considered a noncanonical drug target, since it lacks a discrete active site and has a relatively complex mechanism of activation involving multiple additional enzymes. Despite its complex requirements for activation, NEDD4-1 is thought to be the only member of this signalling pathway that directly interacts with substrates. This allows for great specificity for the manipulation of ubiquitination by drugs. A recent study showed that NAB2 engages with NEDD4-1 at its N-terminus [72]. Treatment with NAB2 significantly increases co-localisation of NEDD4-1 with the early endosome marker Rab5a. This may complement data that have shown that NEDD4-1 traffics α -synuclein to the endosome via K63-linked ubiquitination [28]; however, this should be explored further. α -synuclein toxicity in SH-SY5Y cells was also analysed, and it was found that the trafficking from ER (endoplasmic reticulum) to Golgi regulator (TFG), which is known to regulate ER to Golgi trafficking (a process disrupted in PD), is also an interacting protein of NEDD4-1 [72]. In short, studies into NAB/NAB2–NEDD4-1– α -synuclein interaction show promise for reducing α -synuclein load and toxicity, providing some hope for translation into PD patients in the future (see also Tables 1–3).

This summary of different α -synuclein ubiquitinating E3 ubiquitin–protein ligases suggests a possible redundancy in E3 ubiquitin–protein ligases—though they might have different preferences for modified or aggregated forms of α -synuclein—and highlights the importance of validating the significance of these data in vivo under physiological and pathophysiological conditions. Interestingly, so far, in none of the single knockout mice for these E3 ubiquitin–protein ligases has a clear PD phenotype such as α -synuclein accumulation or midbrain dopaminergic cell death, been described, neither in mice deficient for parkin [186], SIAH1a [187], SIAH1b [188], SIAH2 [189], CHIP [190], FBXL5 [191], ITCH [192], SMURF1 [193,194], SMURF2 [194], WWP1 [195], WWP2 [196], NEDD4-1 [197], nor NEDD4-2 [198]. In SIAH1b-, NEDD4-1-, and NEDD4-2-deficient mice, embryonic lethality may have interfered with a careful analysis of the adult dopaminergic system, and conditional knockout approaches for these genes might allow for the investigation of PD-related phenotypes in the near future [197–199]. These E3 ubiquitin–protein ligases have been shown to ubiquitinate and thereby regulate not only α -synuclein but other important proteins in the midbrain dopaminergic system, which might also contribute to PD pathology. So far, no mutations or single nucleotide polymorphisms (SNPs) in *Siah1b*, *NEDD4-1*, or *NEDD4-2* have been associated with PD, but these ligases are likely to be important players in the protein network altered in PD [200–202]. For SIAH1 function in PD, we refer the reader to the published literature [18,49,52]. Here, we now focus on the two NEDD4 family members NEDD4-1 and NEDD4-2 and discuss their expression, structure, regulation, substrates, and function in the midbrain dopaminergic system, as well as their links to the pathology and treatment of PD.

3. NEDD4-1 and NEDD4-2 Expression

The human *NEDD4-1* gene is located on chromosome 15q21.3 and is comprised of 33 exons transcribed in three mRNAs of 6.4, 7.8, and 9.5 kbp in size. It encodes the NEDD4-1 protein, which has a molecular weight of around 120 kDa [203]. In mice, the *NEDD4-1* gene is located on chromosome 9, and the protein has a similar molecular weight [204].

The *NEDD4-2* (*NEDD4L*) gene is located on chromosome 18q12.31 in humans, with 40 exons, and might result in at least five different transcripts, which appear tissue dependent [205,206]. Variability in these transcripts exists in the N-termini, with varying WW domains and sgk1 phosphorylation sites [206,207]. *NEDD4-2* has been detected to be marginally smaller than *NEDD4-1*, with *NEDD4-2*-specific antibodies detected in two bands on a Western blot. In most tissues, these bands lie at the ~110–115 kDa mark, with one varying in size depending on the tissue type it is expressed in [207,208]. In mice, the *NEDD4-2* gene is also localised on chromosome 18. The human *NEDD4-2* gene is around 78% homologue to the human *NEDD4-1* gene, and the proteins *NEDD4-1* and *NEDD4-2* share 63% sequence identity. *NEDD4-1* gene homologues can be found in all eukaryotic organisms, although *NEDD4-2* is found only in vertebrates. It is therefore thought that *NEDD4-2* arose much later in evolution by gene duplication [42,209].

The two *NEDD4-1* and *NEDD4-2* cDNAs are highly expressed in the developing embryonic and postnatal mouse brain and are subsequently downregulated in the adult brain [210].

NEDD4-1 is also ubiquitously expressed in humans, in the endocrine tissue, lung, proximal digestive tract, gastrointestinal tract, liver, gallbladder, pancreas, kidney, urinary bladder, gonads, muscle, skin, bone marrow, and lymphoid tissue [211]. *NEDD4-1* protein was detected in the dopaminergic system in neuromelanin-positive neurons and in reactive glia cells in the substantia nigra and locus coeruleus of Parkinson's disease and in Lewy body dementia patient brains containing Lewy bodies. It was also detected in lower amounts in neuromelanin-positive neurons in human control brains [28,85]. *NEDD4-1* mRNA was shown to be increased in brain regions with Lewy body pathology [212]. This suggests an important role for *NEDD4-1* in disease. It may be a possibility that *NEDD4-1* accumulation in Lewy body-containing neurons occurs as a result of neuronal damage. However, a more likely explanation is that *NEDD4-1* regulation is representative of a neuroprotective response that leads to a reduction in α -synuclein accumulation. *NEDD4-1* mRNA and protein were also detected in the brain stem of mice [197], but so far, no detailed cellular expression study of *NEDD4-1* has been conducted in the dopaminergic system of mice. In other parts of the central nervous system *NEDD4-1* has been described to be expressed in oligodendrocytes [110]. In cultured cells, *NEDD4-1* is predominantly expressed in the cytosol, near the nucleus, and can be found in neurites after neuronal differentiation. It can, however, be recruited with E2 enzymes to the nucleus [203]. *NEDD4-1* has also been shown to be active at the cell membrane and exosomes [213]. *NEDD4-2* is broadly expressed in humans, in the endocrine tissue, heart, lung, gastrointestinal tract, liver, gallbladder, pancreas, kidney, urinary bladder, gonads, muscle, bone marrow, and lymphoid tissue [211,214]. In the dopaminergic system, *NEDD4-2* protein expression was confirmed in the substantia nigra of mice [114] but not in human brain sections. The *NEDD4-2* protein was found mainly in the cytosol of substantia nigra neurons and astrocytes [114].

The precise cell type-specific expression of *NEDD4-1* and *NEDD4-2* in the midbrain dopaminergic system requires further analysis. However, their expression in midbrain dopaminergic neurons not only during development but during adulthood and ageing, as well as in PD brain samples, supports the idea of an important role here for protein homeostasis.

4. *NEDD4-1* and *NEDD4-2* Structure und Posttranslational Modifications

NEDD4-1 and *NEDD4-2* proteins have modular structures, which are conserved throughout the family (see Figure 2). These modules consist of a C2 calcium-dependent phospholipid-binding domain at the N-terminus (mediating plasma membrane localisation), which can be involved in targeting substrates and adaptors; three or four WW domains, which mediate protein–protein interactions with an xPxY (variable amino acid–proline–variable amino acid–tyrosine) motif on substrates and adaptors; and a catalytic HECT domain at the C-terminus (catalytic cysteine Cys⁸⁶⁷ in *NEDD4-1* and Cys⁹⁴² in

NEDD4-2), which forms a thioester bond with activated ubiquitin that has been transferred from an E2 conjugase before transferring ubiquitin moieties to specific substrates [26,42] (see Figure 2). The WW3 and WW4 domains seem to bind to the PY motif in the substrates, with WW3 generally exhibiting higher substrate affinity than WW4 [42]. Recognition of substrates by NEDD4 ligases involves not only the classical E3 ligase binding of PY motifs to WW domains but the C2 and HECT domains of the ligase [27]. The C2 domain can be cleaved off by caspases during apoptosis, allowing fast degradation of the leftover WW and HECT domains [35]. Details on substrates, adaptors, modifiers, and regulators of NEDD4-1 (see Table 1), NEDD4-2 (see Table 2), and both E3 ubiquitin ligases (see Table 3) are summarised in the Tables 1–3.




Figure 2. Schematic structural representation of NEDD4-1 and NEDD4-2 proteins in humans, mice, and fruit flies. The NEDD4 family of ligases is defined by its modular structure, a lipid-binding/ Ca^{2+} (C2) domain at the N-terminus, a number of WW domains in the middle section, and a HECT ubiquitin ligase at the C-terminus, the latter of which is required for its E3 ubiquitin ligase function. NEDD4's WW domains can interact with PY (proline-tyrosine) motifs to recruit them for ubiquitination. This includes NEDD4's own PY motifs located on the C terminus. Alternative splicing in mice has led some NEDD4-2 variants to lack a C2 domain, although in neurons, NEDD4-2 predominantly contains a C2 domain. WW domains regulate substrate recruitment for ubiquitination and may be expanded in higher-order organisms [111,215]. Common NEDD4 phosphorylation sites are indicated in red.

Human NEDD4-1 and NEDD4-2 were shown to work together with the same E2 ligases: Ubc4, UbcH5A, UbcH5B, UbcH5C, UbcH6, and UbcH7 [203,216]. NEDD4-1 can mono- and polyubiquitinate substrates with K48 and K63, but is also involved in K6 and K27 linkage [111]. NEDD4-2 can also mono- and polyubiquitinate its substrate [214].

The activity of NEDD4-1 and NEDD4-2 is normally blocked via autoinhibition, which also stabilises NEDD4 proteins by preventing autoubiquitination and subsequent proteasomal degradation [127,217]. To form this inhibitory protein conformation, the C2 domain can bind the HECT domain in NEDD4-1 [218], or two conserved tryptophans in the WW domain can bind the PY substrate recognition motif in the HECT domain in NEDD4-2 [127] (see also Tables 1 and 2).

The autoinhibitory conformation can be disrupted, and the substrate specificity altered, by posttranslational modifications including phosphorylation, ubiquitination, neddylation, and SUMOylation, as well as by the presence of calcium binding to the C2 domain of NEDD4-1 [219], autoubiquitination of NEDD4-2 [127], or binding of adaptor and scaffold proteins such as 14-3-3, Numb, or NEDD4 family-interacting proteins (NDFIP1 and NDFIP2) [93,220,221]. In general, substrate-binding seems able to change the conformation of NEDD4 proteins, allowing self-ubiquitination and subsequent degradation, which results in downregulation of NEDD4 once it has ubiquitinated its targets [127].

Autoubiquitination of NEDD4 proteins can be triggered by different interaction partners and can lead to activation, inactivation, or different substrate specificity. The low-density lipoprotein receptor class A domain containing 3 (LRAD3), a member of the low-density lipoprotein (LDL) receptor family, has been found to bind NEDD4 proteins, leading to NEDD4 autoubiquitination and subsequent degradation [222]. Self-catalysed monoubiquitination of NEDD4-1 can enhance substrate recruitment, as shown for the clathrin-coated pit adaptor protein EPS15 (epidermal growth factor receptor substrate 15), which is monoubiquitinated by NEDD4-1 as well as parkin [217,223]. Monoubiquitination of EPS15 leads to an intramolecular binding of ubiquitin to the ubiquitin interaction motive of EPS15 and prevents, for example, EPS15-dependent recruitment of monoubiquitinated EGFR to clathrin-coated pits for internalisation and deactivation [223]. However, ubiquitination of a conserved lysine residue in the HECT domain α 1-helix of one NEDD4-1 protein was also suggested to expose the α 1-helix to bind to the HECT ubiquitin-binding patch of another NEDD4-1 protein, allowing NEDD4-1 to form an inactive trimer [224]. The binding of the adaptor protein Numb to NEDD4-1 has also been shown to stimulate NEDD4-1-mediated ubiquitination of the tumour suppressor PTEN (phosphatase and tensin homologue) and its subsequent degradation [225].

NEDD4-2 is a target for NEDDylation, which is a similar posttranslational modification process to ubiquitination that conjugates a ubiquitin-like protein, NEDD8 (neural precursor cell-expressed developmentally downregulated gene 8), to a substrate with the help of E1, E2, and E3 enzymes (mainly cullin-RING ubiquitin ligases) [129]. Neddylation of NEDD4-2 increases its ubiquitination activity regarding the sodium-coupled bicarbonate cotransporter 1 (NBCe1), which is essential for acid–base homeostasis in the kidney, and leads to proteasomal degradation of NBCe1 and its translocation from the cell membrane into the cytosol [226].

Preliminary data have suggested that NEDD4-1 can be SUMOylated, which is, again, a similar posttranslational modification process to ubiquitination that links SUMO to a substrate in the presence of E1, E2 (Ubc9), and E3 (Smt3p) enzymes [62]. Sumoylation of NEDD4-1 seems to occur not on the consensus site (K357) but on an unknown site; it decreases NEDD4-1 autoubiquitination activity [62].

Alterations in the phosphorylation of NEDD4-1 and NEDD4-2 have been widely observed to regulate their ubiquitination activity and alter the binding of adaptor or scaffold proteins. Fibroblast growth factor receptor 1 (FGFR1) is a substrate of NEDD4-1 ubiquitination that triggers c-Src kinase-dependent phosphorylation of NEDD4-1 at Tyr⁴³ in the C2 domain and Tyr⁵⁸⁵ in the HECT domain, supporting activation [43]. For NEDD4-2, G-protein-coupled receptor (GPCR) protease-activated receptor-1 (PAR1) stimulates c-Src-mediated Tyr⁴⁸⁵ phosphorylation within the 2,3-linker peptide between WW domains 2 and 3 and leads to NEDD4-2 activation [227]. Phosphorylation of *Xenopus* NEDD4-2 on Ser³³⁸ or Ser⁴⁴⁴ by the serine/threonine kinase serum- and glucocorticoid-induced kinase 1 (SGK1) was shown to lead to a reduction in its affinity for the natural NEDD4 substrate epithelia Na⁺

channel (ENaC), which regulates whole-body Na^+ balance and blood pressure [228,229]. Human NEDD4-2 phosphorylation by aldosterone-induced SGK1 on Ser³⁴² and Ser⁴⁴⁸ (and Thr³⁶⁷) was shown to facilitate 14-3-3 protein binding to NEDD4-2, leading to inhibition of the interaction between NEDD4-2's HECT and WW domains, stabilisation of ENaC in the kidney, and enhanced ubiquitination of the AMPA (α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid) receptor subunit GluA1 (glutamate ionotropic receptor AMPA type subunit 1) in the brain. For *Xenopus* NEDD4-2, it was suggested that the 14-3-3 dimer binds first on NEDD4-2 P-Ser⁴⁴⁴, the high-affinity (major) site, and subsequently on one of the lower-affinity (minor) sites, P-Ser³³⁸ or P-Thr³⁶³ [230]. For human NEDD4-2 it has been shown that the 14-3-3 dimer simultaneously anchors on two of the three phosphorylation sites, P-Ser³⁴², P-Thr³⁶⁷ and P-Ser⁴⁴⁸, of NEDD4-2, with P-Ser⁴⁴⁸ being the key residue [93]. SGK1 also leads to phosphorylation of human NEDD4-2 Ser⁴⁶⁸ and an increase in ENaC protein [92,231]. Interestingly, SGK1 has also been suggested to be a NEDD4-2 substrate, leading to its own degradation and generating a negative feedback loop [232]. The same three SGK1 phosphorylation sites, Ser³⁴², Ser⁴⁴⁸ and Thr²⁶⁷, of NEDD4-2 are also used by vasopressin-induced cyclic AMP-dependent protein kinase A (PKA). Furthermore, insulin activates SGK1 and Akt (protein kinase B) and leads to Ser³⁴² and Ser⁴²⁸ phosphorylation of human NEDD4-2, upregulating ENaC on the membrane [155]. IkappaB kinase β (IKK β) has been found not only to bind to ENaC and enhance its activity but to phosphorylate *Xenopus* NEDD4-2 on Ser⁴⁴⁴, preventing NEDD4-2-dependent ENaC ubiquitination [117]. Interestingly, 14-3-3 η has also been shown to bind and inhibit the ubiquitination activity of wildtype parkin but not of parkin with R42P, K161N, and T240R mutations associated with autosomal recessive juvenile parkinsonism [233]. The parkin/14-3-3 inhibitory complex could be prevented by wildtype α -synuclein but not by A30P and A53T mutations, causing PD [233]. These data define the chaperone-like protein 14-3-3 as an important inhibitor of E3 ligases associated with PD.

The regulatory mechanisms occurring in the midbrain dopaminergic system to change the activity or substrate specificity of NEDD4-1 and NEDD4-2 have so far not been investigated but are likely to utilise at least some of the aforementioned posttranslational modifications and interaction partners. With further information about important *in vivo* substrates and functions of NEDD4-1 and NEDD4-2 in dopaminergic neurons, it will be a revealing task to study the detailed regulatory mechanisms.

5. NEDD4-1 and NEDD4-2 Substrates, Adaptors, Regulators, and Function

Upon the discovery of NEDD4-2, it was proposed that NEDD4-1 and NEDD4-2 may have redundant functions with shared interaction partners and substrates, however there appears to be adaptors, substrates, and functions specific or unique to NEDD4-1 and NEDD4-2. Below, we list some possible shared and unique NEDD4-1 and NEDD4-2 substrates, adaptors, and regulators.

The different phenotypes of NEDD4-1 and NEDD4-2 knockout (KO) mice suggest that their main substrates are distinct, and the redundancy might be limited to a few substrates and functions [42]. The predominant phenotype of NEDD4-1 KO mice is embryonic lethality at midgestation, with pronounced heart defects (double-outlet right ventricle and atrioventricular cushion defects) and vasculature abnormalities leading to growth retardation (with a body weight less than 40% of that of wild-type littermates) [46,117,118]. In contrast, NEDD4-2 KO mice show perinatal lethality, with increased ENaC levels that seem to cause premature foetal lung fluid clearance, resulting in a failure to inflate the lungs [198]. Only a few of these mice survived up to 22 days [198]. This phenotype was also confirmed in lung-specific NEDD4-2 deficient mice [234]. When crossing floxed NEDD4-2 mice with EIIa-Cre mice (B6.FVB-Tg(EIIa-cre)C5379Lmgd/J) [235] expressing Cre in a mosaic pattern in the embryo before implantation in the uterine wall, the NEDD4-2 KO mice might not be a complete null for NEDD4-2 [236]. These mice were viable but showed defects in the respiratory, renal, cardiac, neural, and immune systems and high blood pressure, indicating that NEDD4-2 is a key regulator of Na^+ homeostasis and that ENaC is

one of its most important physiological substrates [236]. This suggests that even ENaC can be a substrate of NEDD4-1 and NEDD4-2 [42]. The NEDD4-1 and NEDD4-2 KO mouse data suggested that in vivo NEDD4-2 is most likely the more important E3 ubiquitin ligase for ENaC. Interestingly, the G protein-coupled receptor kinase 2 (Grk2) can phosphorylate ENaC on Ser⁶³³ in the C-terminus of the β -subunit, which increases ENaC activity and prevents ENaC ubiquitination by NEDD4 ligases and subsequent degradation [237]. GRK2 and other GRK family members have also been described to phosphorylate α -synuclein on Serine¹²⁹, which is common in PD patients, however it has not been reported whether this phosphorylation negatively influences ubiquitination by NEDD4 ligases [238].

This suggests that NEDD4-1 and NEDD4-2 may have common but also unique functions, and this might depend on the specific tissue investigated. Therefore, it seems important to investigate how far NEDD4-1 and NEDD4-2 have redundant and unique functions in vivo during the development and maintenance of the midbrain dopaminergic system as well as in pathophysiological conditions leading to PD.

Most NEDD4-1 and NEDD4-2 substrates and adaptors have so far been investigated only in vitro and await in vivo confirmation. PTEN (phosphatase and tension homologue) is a good example to illustrate the importance of verifying possible NEDD4 substrates in vivo under physiological conditions in an organism such as the mouse. Cell culture experiments and human cancer tissue suggested that PTEN might be a NEDD4-1 and NEDD4-2 substrate [177–180,239]), but analysis of NEDD4-1 and NEDD4-2 knockout mice has shown that PTEN stability, subcellular localisation, and activity are not altered in the absence of NEDD4-1 and/or NEDD4-2 [181,182]. Furthermore, more recent cell culture experiments have not supported PTEN as a NEDD4 substrate [225]. Therefore, more research would be required to finally resolve this controversy between in vitro and in vivo data.

The adaptor proteins NDFIP1 (NEDD4 family-interacting protein 1 or NEDD4 WW domain-binding protein 5 (N4WBP5)) and NDFIP2 (NEDD4 family-interacting protein 2 or NEDD4 WW domain-binding protein 5A (N4WBP5A)) are small, endosomal, PY-motif-containing membrane proteins that can both function as adaptors for NEDD4-1, NEDD4-2, ITCH/AIP4, WWP1, and WWP2, facilitating their binding to proteins that lack PY motifs, preventing autoinhibition of the ligase, and possibly serving as ubiquitination substrates. Overexpression of NDFIP1 is able to recruit NEDD4-1, NEDD4-2, and ITCH to neuronal exosomes, which are normally free of these E3 ligases, for secretion [109,203]. The positive effect of NDFIP1/NEDD4-1 in improving neuronal survival during brain injury suggests that perhaps exosomal NEDD4-1 might enhance transport and degradation of unwanted proteins [203]. As microglial exosomes facilitate the transmission of α -synuclein in PD [240], it would be of interest to study the role of NEDD4 ligases in this process. NDFIP1 and NDFIP2 are physically and functionally associated with multiple components of the epidermal growth factor (EGF) signalling cascade, and their levels modulate the relative output of different signalling pathways. They associate with the EGF receptor and the phosphatase and tension homologue (PTEN) and control the ubiquitination and abundance of PTEN, cellular Casitas B-lineage Lymphoma E3 ligase (c-CBL), and cellular Sarcoma family kinases (c-Src). NDFIP2, but not NDFIP1, also binds to and is phosphorylated by two c-Src kinases (Src and Lyn) and can act as a scaffold for Src phosphorylation of NDFIP1 and potentially other substrates. Depletion of NDFIP1 inhibits serine/threonine kinase Akt (protein kinase B, PKB) activation in EGF-stimulated HeLa cells, stimulates activation of cellular transcription factor c-Jun-N-terminal Kinase (Jnk), and enhances cell multiplication. Interestingly, increased iron is often found in the substantia nigra of PD patients and has been associated with increased NDFIP1 levels [239]. It would be of interest to examine whether iron misregulation may serve to be protective to nigral dopaminergic neurons by upregulating NDFIP1 and facilitating NEDD4-1-mediated ubiquitination of α -synuclein.

Adaptor proteins such as NDFIP1 and NDFIP2 seem to use different members of the NEDD4 family in vivo. NDFIP1-deficient mice showed a reduced life expectancy, with severe inflammation of the skin and lung, enhanced T-cell activation, proliferation and

differentiation to T helper 2 cells, and a prolonged JunB half-life such as that in Itchy mutant mice lacking functional ITCH protein [192,241]. NDFIP2-deficient mice showed no overt immunopathology, but NDFIP2 deficiency seemed to enhance the NDFIP1 knockout phenotype, leading to further accumulation of effector CD4⁺ T cells and an increase in JAK (Janus kinase) protein, which might be explained by reduced Itch or NEDD4-2 activation [242]. Further research has to be done to confirm the use of the adaptor proteins NDFIP1 and NDFIP2 by NEDD4-1 and NEDD4-2.

NEDD4-1 activation has also been shown to be important for autophagy and mitophagy [66,243]. LC3 (MAP1LC3, microtubule-associated protein 1 light chain 3) is essential in autophagy by functioning in elongation of the phagophore double-layer membrane and in the recruitment of proteins for autophagic processes. LC3 activates and recruits NEDD4-1 to the phagophore assembly site (PAS) by binding the conserved WXXL LC3-binding motive between the C2 and the WW2 domains. LC3-I is activated to LC3-II by cleavage and conjugation to phosphatidylethanolamine (PE) and is recruited to autophagosomes by binding LIR (LC-3 interacting region domain-containing protein). Subsequently, NEDD4-1 ubiquitinates the LC3-interacting protein p62 (sequestosome-1, SQSTM1) and beclin-1 (BECN1), which seems required to recruit downstream effectors for autophagosome formation [32,66]. More recently, NEDD4-1 lysine²⁹-linked autoubiquitination on lysine¹²⁷⁹ was shown to recruit USP13 (ubiquitin-specific protease 13) to form a deubiquitination complex, which stabilised VPS34 to promote autophagy by removing the lysine⁴⁸-linked polyubiquitin chains from VPS34 at lysine⁴¹⁹ [74]. Surprisingly, in mice, endoplasmic reticulum (ER) stress and activation of the unfolded protein response (UPR) increased autophagy and NEDD4-2 expression in the liver, but not NEDD4-1 expression. In addition, in cell culture, high amounts of NEDD4-2 correlated with increased autophagy, while low amounts of NEDD4-2 correlated with reduced autophagy [244]. In PD, reduced autophagy is a common phenotype that can be triggered by α -synuclein accumulation and might be enhanced by NEDD4-1 and NEDD4-2 [245].

Interestingly, NEDD4 ubiquitination activity is required for the release of some retroviruses but might be inhibited as a cellular defence mechanism. The Gag protein of human oncoretrovirus HTLV-1 (human T-lymphotropic virus type 1) has a tandem PPPY/PTAP motif and needs to be ubiquitinated by the E3 ligase NEDD4-1 at the plasma membrane. It also requires Tsg101 (tumour susceptibility gene 101) recruitment at the ESCRT (endosomal sorting complexes required for transport) pathway in late endosomes/multivesicular bodies for driving virus budding [49]. Despite the HIV (human immunodeficiency virus) Gag protein lacking a PY motif, it also uses NEDD4-1 and NEDD4-2 for its ubiquitination, to stimulate budding; Nedd4-2, with the adaptor protein AMOT-1 (angiomin-1 protein); and NEDD4-1, by binding and ubiquitinating adaptor protein ALIX (apoptosis-linked gene 2 (ALG-2)-interacting protein X, programmed cell death 6-interacting protein) [246]. However, NEDD4 family members might be inhibited in cells after viral or bacterial infection by binding with upregulated interferon-induced ubiquitin-like protein ISG15 (interferon-stimulated gene 15). The binding of ISG15 to NEDD4-1, NEDL1, NEDL2, or WWP2 can block their interaction with ubiquitin-E2 enzymes and interfere with the ubiquitination of retroviral group-specific antigen precursors and matrix proteins, such as VP40 of Ebola with a PPxY motif, which is essential for the release/budding of Ebola, vesicular stomatitis, and rabies virus particles [247,248]. DNA virus proteins such as the latent membrane protein 2A (LMP2A) of Epstein–Barr Virus (EBV, human herpesvirus 4) also interact via their PPPPY motif with the NEDD4 family members NEDD4-1, ITCH, and WWP2 [68]. This interaction leads to the ubiquitination of LMP2A and LMP2A-associated proteins such as the protein tyrosine kinases Lyn (Lck/Yes novel tyrosine kinase of the Src kinase family) and Syk (spleen tyrosine kinase), which might be important for EBV latency and the regulation of B-cell signal transduction [68]. Taken together, this suggests an important role of NEDD4 family members in vesicular transport and retrovirus propagation being regulated by the immune system, which have been found to be altered also in neurodegenerative diseases such as PD [249].

In the context of PD, most research in the past focused on NEDD4-1 after NEDD4-1 was found to ubiquitinate α -synuclein as detailed above [28,29]. An interesting NEDD4-1 substrate in this regard is the receptor tyrosine (and serine) kinase RET (abbreviation for rearranged during transfection), the canonical receptor for the TGF- β (transforming growth factor- β)-related neurotrophic factor family member GDNF (glial cell line-derived neurotrophic factor), which is currently in clinical trials in PD patients [245,250–252] (see Figure 1). The RET receptor is important for the maintenance, protection, and regeneration of midbrain dopaminergic neurons [253–255]. In cell culture experiments, the turnover of the long splice isoform of RET, RET51, is mediated after activation and autophosphorylation by binding of the adaptor protein GRB2 (growth factor receptor-bound protein 2) to RET tyrosine¹⁰⁹⁶ and subsequent recruitment of the E3 ubiquitin–protein ligase c-CBL [82]. However, the short RET isoform, RET9, binds to the adaptor proteins GRB10 or SHANK2 (SH3 And Multiple Ankyrin Repeat Domains 2). This depends on its phosphorylated tyrosine¹⁰⁶² and the C-terminal PDZ-binding motif (PDZ is an initialism combining the first letters of the first three proteins discovered to share the domain—postsynaptic density protein (PSD95), Drosophila disc large tumour suppressor (Dlg1), and zonula occludens-1 protein (zo-1)) and recruits NEDD4-1 [82]. RET polyubiquitination triggers receptor internalisation from clathrin-coated pits at the cell membrane into endosomal compartments for receptor recycling to the cell surface or lysosomal degradation [256]. RET51 shows more K63 ubiquitin linkage—in contrast to RET9—and can be sorted to RAB11-positive recycling endosomes for signalling, intracellular trafficking, and return to the cell surface or targeted for lysosomal degradation [257,258]. RET9 ubiquitination chains are more K48 linked, targeting the protein more for proteasomal degradation [259]. Determining whether NEDD4-1-dependent ubiquitination of RET9 also occurs in dopaminergic neurons in vivo and influences survival and physiology requires further investigation.

Other receptor tyrosine kinases have also been suggested to be substrates of NEDD4-1, such as the fibroblast growth factor receptor 1 (FGFR1) [43] and the epidermal growth factor receptor (EGFR) members ErbB1 (erythroblastic leukaemia viral oncogene homologous-B2 receptor tyrosine kinase 1) [225] and HER3/ErbB3 (human epidermal growth factor receptor 3) [53]. NEDD4-1 also mediates the adaptor protein β -arrestin2's agonist-dependent ubiquitination and lysosomal degradation of the β 2-adrenergic receptor (β 2AR) [33]. IGF1R (insulin-like growth factor I receptor) can bind the adaptor protein GRB10, and this was suggested to mark IGF1R for NEDD4-1-dependent polyubiquitination and degradation [168,260,261] or protect IGF1R from NEDD4-1 ubiquitination [262]. Further work is needed to understand this controversial NEDD4-1 and IGF1R crosstalk. VEGFR-2 (vascular endothelial growth factors receptor 2), but not VEGFR-1, has also been suggested to be protected from NEDD4-1-induced degradation by binding GRB10, although VEGFR-2 might not be a direct NEDD4-1 ubiquitination substrate [91]. VEGF stimulation of VEGFR-2 increases GRB10 expression and c-Src-dependent tyrosine phosphorylation of GRB10, which subsequently increases VEGFR-2 protein levels [263]. Interestingly, in NEDD4-1 knockout mice, GRB10 protein levels were increased and IGF1 and insulin signalling were reduced, while *GRB10* gene deletion rescued the NEDD4-1 knockout lethality, suggesting a negative regulatory function of GRB10 for IGF1 and insulin signalling [262]. These data suggest that NEDD4-1 might not only directly targets receptor tyrosine kinases as substrates but also indirectly regulate receptor tyrosine kinase signalling by targeting associated adaptor proteins. For example, NEDD4-1 monoubiquitinates insulin receptor substrate (IRS)-2, which promotes its binding to the clathrin-coated pit adaptor protein epsin-1 and the recruitment of IGF1R, which phosphorylates IRS-2, stimulating downstream signalling [59]. Another NEDD4-1 monoubiquitination substrate seems to be the adaptor protein HGS (hepatocyte growth factor-related kinase substrate, HRS), which leads to intramolecular binding of ubiquitin to the HGS ubiquitin-interaction domain (UIM), leading in turn to reduced endocytosis of EGFR [55]. The secretory carrier membrane protein-3 (SCAMP3) has a PY motif and seems to be multimonomubiquitinated by NEDD4-1, which allows HGS interaction and prevents EGFR degradation [87]. The endosomal sorting

complexes (ESCRT-0, -I, -II, -III, VPS4-VTA1 (vacuolar protein sorting protein 4 and vesicle trafficking protein 1), and ALIX (apoptosis-linked gene 2-interacting protein X) homodimer) are peripheral membrane protein complexes required together for degradation of damaged or unwanted plasma membrane and cytosolic proteins, lysosome and multivesicular body (MVB) biogenesis, autophagy, and viral budding [264,265]. The ESCRT-0 complex sorts ubiquitinated membrane proteins into MVB. It consists of HGS and STAM (signal transduction adaptor molecules STAM1 and STAM2) proteins and can be associated with the ubiquitin-binding domain-containing protein EPS15B (epidermal growth factor receptor pathway substrate 15B) to mediate EGFR degradation [265,266]. EPS15 is associated with clathrin-coated pit adaptor protein 2 (AP2) and plays a role in EGFR internalisation [266]. The NEDD4 family members NEDD4-1 and ITCH/AIP4 have been found associated with ESCRT complexes, which seems important not only for viral GAG protein ubiquitination and budding but for degradation of a membrane-associated pool of Lys⁶³ polyubiquitinated α -synuclein [28,265]. ESCRT proteins (VPS4 from ESCRT-0, charge multivesicular body protein 2B (CHMP2B) from ESCRT-III) have been found to be important for lysosomal targeting of α -synuclein and are also localised to Lewy bodies [29,267–269].

Both NEDD4-1 and NEDD4-2 can bind the PY motif containing non-receptor tyrosine and serine/threonine kinase ACK (activated Cdc42-associated tyrosine kinase, TNK2). NEDD4-1 leads to polyubiquitination and subsequent lysosomal degradation of ACK along with EGFR in response to EGF stimulation, while the data for NEDD4-2 in regard to ACK ubiquitination remain inconsistent [151,152]. Interestingly, FGFR3 activation leads to NEDD4-1 phosphorylation, which subsequently targets the transmembrane protein programmed death-ligand 1 (PD-L1, cluster of differentiation 274 (CD274), B7 homologue 1 (B7-H1)), which is involved in immune system suppression for Lys⁴⁸-linked polyubiquitination and degradation [270]. Furthermore, activation of the small GTPase RAS (rat sarcoma virus protein) is reduced by NEDD4-1 ubiquitinating the PY motif containing the RAS activator CNrasGEF (cyclic nucleotide RAS guanine-nucleotide exchange factor) for proteasomal degradation [40].

A receptor tyrosine kinase that is suggested to be a NEDD4-2 substrate is the NGF (nerve growth factor) receptor TRKA (tropomyosin receptor kinase A; NTRK1, neurotrophic receptor tyrosine kinase 1) with a PPXY motif, which is, in its activated/phosphorylated state, marked for degradation by NEDD4-2-dependent ubiquitination and is more abundant in the dorsal root ganglia of NEDD4-2 deficient mice [145,271,272]. However, the closely related BDNF (brain-derived neurotrophic receptor) receptor TRKB lacks a PPXY motif and seems not to be a NEDD4-2 substrate [145]. TRKA is normally not expressed in midbrain dopaminergic neurons, but the ectopic expression of TRKA in vivo in mice combined with NGF treatment protected dopaminergic neurons from 6-OHDA-induced cell death [273]. TRKB is found in midbrain dopaminergic neurons but is not essential for development and maintenance [253]. However, TRKB seems to protect cells from MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine/1-methyl-4-phenylpyridinium) toxicity in mice [274].

NEDD4-2 and other NEDD4 family members have also been found to regulate the signalling of the transforming growth factor beta (TGF- β) family of ligands (33 human genes encoding, for example, TGF- β 1, 2, and 3; activins; inhibins; and bone morphogenetic protein (BMPs)). TGF- β family members activate the TGF- β receptor by inducing the heterotetramerisation of the two single transmembrane serine–threonine (and tyrosine) kinase receptors, TGF- β receptors type I and type II [275,276]. TGF- β receptor signalling leads to transcriptional regulation of target genes through the canonical SMAD signalling pathway (the SMAD abbreviation refers to the homologies to the *Caenorhabditis elegans* SMA (“small” worm phenotype) and the MAD family (“Mothers Against Decapentaplegic”) of genes in *Drosophila*) protein involving signalling pathway and the non-canonical (SMAD independent, involving tyrosine-autophosphorylation) [275,276]. TGF- β receptor type I has also been described as a NEDD4-2, SMURF1, SMURF2, and WWP1 substrate in conjunction with the inhibitory SMAD7 protein as an adaptor. It is marked for degradation by ubiquitination. The TGF- β I and II receptors signal together through the receptor-regulated SMADs

(R-SMADs 1, 2, 3, 5, and 8/9), which can partner with SMAD4 (Co-SMAD) or be suppressed by the inhibitory SMADs (I-SMAD 6 and 7) or the SMAD corepressor SnoN (Ski-related novel protein N) to downregulate TGF- β receptor signalling [140,276]. Interestingly NEDD4-2 was able to bind to all SMAD proteins with a PY motif (1,2,3,5,6,7), but not to those without one (4 and 8), and it induced ubiquitination and subsequent degradation of SMAD2 but not of SMAD3 [140]. Although NEDD4-2 and SMURF2 can both bind to SnoN via SMAD2, 3, or 4, only SMURF2 can ubiquitinate SnoN and lead to its degradation [140]. The TGF- β I and II receptors are expressed in dopaminergic neurons, and TGF- β II-deficient mice showed reduced TGF- β I receptor levels, reduced dendritic growth and spine formation, a decreased range of excitatory-to-inhibitory synapses, a reduced excitation/inhibitory ratio (ratio of evoked miniature excitatory postsynaptic currents (mEPSC) to miniature inhibitory postsynaptic currents ratio (MIPSC)), hyperactivity, and a reversal-learning defect but no change in dopaminergic cell counts [277].

As for α -synuclein, several different ubiquitin–protein ligases have been proposed to influence internalisation, signalling and degradation of most receptor and cytosolic tyrosine and serine–threonine kinases. Therefore, very careful *in vivo* analysis is needed to understand the roles of NEDD4-1 and NEDD4-2 in regulating a specific kinase, considering in detail the context, such as the organism, developmental stage and age, tissue, and environmental and physiological challenges.

Concerning PD and the function of the dopaminergic system, it is also important to mention that NEDD4-2 can ubiquitinate neurotransmitter transporters. The dopamine transporter (DAT) is required for the reuptake of dopamine into dopaminergic neurons and has been suggested to be ubiquitinated by NEDD4-2, but not NEDD4-1, leading to endocytosis by binding to epsin and Eps15 on clathrin-coated pits [108,278]. NEDD4-2 seems to cooperate with the E2 enzymes UBE2D and UBE2L3 to conjugate, in a PKC-dependent reaction, primarily lysine⁶³-linked ubiquitin chains onto DAT [108]. Reduced DAT levels increase the amount of extracellular dopamine and prolong the stimulation of pre- and postsynaptic dopamine receptors, leading in mice to locomotion defects [279]. Another NEDD4-2 substrate that is ubiquitinated in an PKC-dependent manner and reduced in neurodegenerative diseases is the glutamate transporter-1 (GLT-1), which is subsequently internalised and degraded [280]. In MPP⁺-treated astrocytes and MPTP-treated mice, NEDD4-2 mediated the ubiquitination of GLT-1 [114]. Conversely, NEDD4-2 knockdown increased glutamate transporter protein levels. In MPTP-treated mice, NEDD4-2 knockdown ameliorated movement disorders, increased tyrosine hydroxylase expression in the midbrain, and attenuated astrogliosis and reactive microgliosis associated with glutamate excitotoxicity [114]. To support the idea that NEDD4-2 might be a therapeutic target for the treatment of PD, it would be of interest to confirm these results in conditional NEDD4-2 KO mice treated with MPTP or a more physiological challenge such as mild overexpressed α -synuclein [245].

Both NEDD4-1 and NEDD4-2 seem to ubiquitinate, in a PKC-dependent reaction (phosphorylation of NEDD4-2 threonine¹⁹⁷, serine²²¹, serine³⁵⁴, and serine⁴²⁰), the human organic anion transporter 1 (hOAT1) in kidney proximal tubule cells, which is important for the release of anti-HIV drugs, anti-tumour drugs, antibiotics, and anti-inflammatory drugs. Ubiquitination of hOAT1 leads to its reduced activity, internalisation and degradation [131,281]. NEDD4-2 has been suggested to also ubiquitinate hOAT3 [282]. NEDD4-1 seems capable of ubiquitinating the ATP binding cassette transporter B1 (ABCB1), which can export the neurotoxic peptide β -amyloid from endothelial cells in the blood–brain barrier to protect the brain [31]. In Alzheimer's patients, ABCB1 protein levels were reduced, while NEDD4-1 protein levels were increased, suggesting NEDD4-1 as a therapeutic target for the treatment of Alzheimer's disease.

Another interesting NEDD4-1 substrate is the proapoptotic protein RTP801 (regulated in development and DNA damage responses, Redd1; DNA-damage-inducible transcript 4, DDIT4; or dexamethasone-induced gene 2 encoded protein, Dig2), a mTOR suppressor that has previously been shown to cause neuronal death in both cellular and animal

models of PD [85]. RTP801 has been found upregulated in toxin-induced PD animal models such as 6-hydroxydopamine (6-OHDA), MPTP/MPP+ and rotenone, as well as in the substantia nigra dopaminergic neurons of PD patients, while NEDD4-1 seems to be downregulated [283]. An *in vitro* study showed that RTP801 can be subjected to lysosomal degradation and is conjugated with K63-linked polyubiquitin chains by NEDD4-1 [85]. It has been proposed that RTP801 is stress-induced upregulated at the early stage of PD to maintain cellular function, but sustained elevation and mTOR and AKT inhibition might lead to dopaminergic cell death [284]. However, it has also been suggested that NEDD4-1 acts as a downstream target of the PI3K/PTEN–mTORC1 signalling pathway to promote neurite growth and regeneration [182].

As ion channel dysfunction become increasingly intertwined with PD pathology, it is worth investigating the capacity of NEDD4 ligases to interact with them, particularly *in vivo*. Dopaminergic neurons are characterised electrophysiologically by their spontaneous discharge from pacemaker activity to burst-firing [285–290]. Tonic action potential firing which contributes to the pacemaker activity of dopaminergic neurons is controlled by ion channels [291]. Ion channels can play critical roles in the neuronal excitability, cell volume and the regulation of neurotransmitter release.

Voltage-gated sodium channels (Na_vs) reside in the cell membrane are essential for the creation and propagation of action potentials. Studies examining the interaction of NEDD4-2 with Na_vs have been carried out *in vivo*, *in vitro* and *ex-vivo* and demonstrated that Na_vs 1.2, 1.3, 1.5, 1.6, 1.7, 1.8 are substrates of NEDD4-2 mediated through interaction with PPSY, PLSY and PGSP motifs [122–124]. In a patch-clamp experiment, Na_v1.2 was shown to be downregulated when NEDD4-2 was expressed in HEK293 cells [122]. Na_v1.6 is essential for neuronal excitability and a number of motor disorders are associated with mutations in Na_v1.6. Homozygous-null mutations in Na_v1.6 lead to juvenile mortality in mice between P19 and P21 [292]. Phosphorylated Pro-Gly-Ser⁵⁵³-Pro motif on Na_v1.6 is a putative binding site for NEDD4 ubiquitin ligases. One study hypothesised that NEDD4-2 contributes to the ubiquitination and subsequent internalisation of Na_v1.6. In cultured cells, NEDD4-2 was indeed shown to interact with Na_v1.6 through a C-terminal Pro-Ser-Tyr¹⁹⁴⁵ motif, causing a reduction in Na_v1.6 current density. This regulation appears to require both the Pro-Gly-Ser-Pro motif in L1 and the Pro-Ser-Tyr motif in the C terminus of Na_v1.6. When NEDD4-2 binding to the Pro-Ser-Tyr motif was prevented, a stress-mediated increase in Na_v1.6 current density was observed. Phosphorylation of the Pro-Gly-Ser-Pro motif in the L1 of Na_v1.6 seems necessary for stress-induced current modulation. Positive or negative regulation appears to depend on the availability of the PRO-Ser-Tyr motif in the C-terminus to bind NEDD4-2 [124].

A study examined cognitive impairments in a rat model of PD [293]. In these 6-OHDA lesioned rats, Na_v1.1 was substantially elevated in reactive hippocampal astrocytes 28 days after lesioning, which reduced after 49 days. No changes were observed in Na_v1.6 levels at 28 days, but was elevated in hippocampal neurons at a later time-point of 49 days post-lesion. The predominantly embryonically expressed Na_v1.3 appeared to be re-expressed in hippocampal CA neurons at 49 days post-lesion. In this study 6-OHDA lesioned rats were treated with the Na_v blocker, phenytoin. These rats exhibited improved spatial learning and memory in the Morris water maze compared to lesioned rats not given the phenytoin [293].

Na_vs, in particular Na_v1.6, appear to play an important role in neuronal physiology and may play a role in the genesis of cognitive deficits in PD. It would be of interest to examine the *in vivo* role of Na_vs in dopaminergic neurons specifically and how this might be altered in the presence or absence of NEDD4 ligases. It would also be of interest to investigate how these channels behave in α -synuclein-mediated models of PD and how this could impact the fate of dopaminergic neurons and their interaction with proximal glia.

In addition to voltage-gated sodium channels, NEDD4 ligases have also been demonstrated to interact with potassium ion channels. In the brain KCNQ2 and KCNQ4 potassium ion channels are mostly limited to the substantia nigra and ventral tegmental area of the midbrain [294–297]. In the striatum, dopaminergic nerve termini express KCNQ2 and

KCNQ3 [297]. The muscarine-sensitive K(+) current (M-current) stabilises neuronal resting potential, therefore limiting neuronal excitability. M-current is mediated through heteromeric ion channels comprised of KCNQ3 subunits which associate with either KCNQ2 or KCNQ5 subunits. In a study examining the regulation of KCNQ2/3/5 channels it was revealed that NEDD4-2 but not NEDD4-1 could reduce K(+) currents mediated by KCNQ2/3 and KCNQ3/5 in a *Xenopus* oocyte expression system. Through deletion experiments it was shown that the KCNQ3 subunit is required for NEDD4-2 to regulate the heteromeric channels. Co-immunoprecipitation and Glutathione S-transferase fusion pulldown experiments demonstrated that NEDD4-2 and KCNQ2/3 interact directly. NEDD4-2 was also able to ubiquitinate KCNQ2/3 in transfected HEK293T cells [119]. Other Potassium channels in this family have also been shown to interact with NEDD4 ligases (particularly NEDD4-2) [116,120,121], however these channels have not currently been implicated in PD or dopaminergic system pathology. It appears however that in the nervous system NEDD4-2 is potentially an important M-current activity regulator.

6. Conclusions and Future Directions

The collected data support the notion that further research is required to clarify the unique and common features of NEDD4-1 and NEDD4-2 especially in the midbrain dopaminergic system affected in PD. The large number of possible mechanisms for regulating NEDD4 ubiquitination activity, substrate specificity, and protein interactions make an *in silico* prediction of possible outcomes extremely difficult. The regulation, function, and substrate specificity of NEDD4-1 and NEDD4-2 need to be studied *in vivo* in a tissue- and cell-type-specific fashion before strategies can be designed to propose them as therapeutic targets for neurodegenerative diseases such as PD. The possible substrates suggest that NEDD4-1 and/or NEDD4-2 could be beneficial or harmful in the disease context. Currently, it is not clear if NEDD4-1 and/or NEDD4-2 protein levels should be increased or decreased to improve the conditions in dopaminergic neurons under pathophysiological conditions such as PD. Research on NEDD4 proteins in neurodegenerative diseases remains an exciting field in which many surprising findings can still be expected. Unbiased approaches should therefore be applied to remain open to all possible outcomes. In the recent years, the toolkit for studying NEDD4-1 and NEDD4-2 has dramatically improved, with the availability of conditional animal models for NEDD4-1 and NEDD4-2, good antibodies, and specific knockdown possibilities, which will facilitate further investigations.

Author Contributions: Conceptualization, E.R.K.; writing—original draft preparation, review and editing, J.A.C., G.K. and E.R.K.; visualization, J.A.C.; supervision, E.R.K.; funding acquisition, J.A.C. and E.R.K. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by BRACE, Bristol, UK (E.R.K.), Alzheimer’s Research UK (J.A.C., E.R.K.) and the Northcott Devon Medical Foundation (J.A.C., E.R.K.).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Goedert, M.; Spillantini, M.G.; Del Tredici, K.; Braak, H. 100 years of Lewy pathology. *Nat. Rev. Neurol.* **2013**, *9*, 13–24. [[CrossRef](#)] [[PubMed](#)]
- Singleton, A.; Gwinn-Hardy, K. Parkinson’s disease and dementia with Lewy bodies: A difference in dose? *Lancet* **2004**, *364*, 1105–1107. [[CrossRef](#)]
- Lassot, I.; Mora, S.; Lesage, S.; Zieba, B.A.; Coque, E.; Condroyer, C.; Bossowski, J.P.; Mojsa, B.; Marelli, C.; Soulet, C.; et al. The E3 Ubiquitin Ligases TRIM17 and TRIM41 Modulate α -Synuclein Expression by Regulating ZSCAN21. *Cell Rep.* **2018**, *25*, 2484–2496.e2489. [[CrossRef](#)] [[PubMed](#)]

4. Abeywardana, T.; Lin, Y.H.; Rott, R.; Engelender, S.; Pratt, M.R. Site-specific differences in proteasome-dependent degradation of monoubiquitinated α -synuclein. *Chem. Biol.* **2013**, *20*, 1207–1213. [\[CrossRef\]](#)
5. Zhang, J.; Li, X.; Li, J.-D. The Roles of Post-translational Modifications on α -Synuclein in the Pathogenesis of Parkinson's Diseases. *Front. Neurosci.* **2019**, *13*, 381. [\[CrossRef\]](#)
6. Hasegawa, M.; Fujiwara, H.; Nonaka, T.; Wakabayashi, K.; Takahashi, H.; Lee, V.M.; Trojanowski, J.Q.; Mann, D.; Iwatsubo, T. Phosphorylated α -synuclein is ubiquitinated in α -synucleinopathy lesions. *J. Biol. Chem.* **2002**, *277*, 49071–49076. [\[CrossRef\]](#)
7. Tofaris, G.K.; Razzaq, A.; Ghetti, B.; Lilley, K.S.; Spillantini, M.G. Ubiquitination of α -synuclein in Lewy bodies is a pathological event not associated with impairment of proteasome function. *J. Biol. Chem.* **2003**, *278*, 44405–44411. [\[CrossRef\]](#)
8. Rott, R.; Szargel, R.; Haskin, J.; Bandopadhyay, R.; Lees, A.J.; Shani, V.; Engelender, S. α -Synuclein fate is determined by USP9X-regulated monoubiquitination. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 18666–18671. [\[CrossRef\]](#)
9. Deng, L.; Meng, T.; Chen, L.; Wei, W.; Wang, P. The role of ubiquitination in tumorigenesis and targeted drug discovery. *Signal Transduct. Target. Ther.* **2020**, *5*, 11. [\[CrossRef\]](#)
10. Metzger, M.B.; Hristova, V.A.; Weissman, A.M. HECT and RING finger families of E3 ubiquitin ligases at a glance. *J. Cell Sci.* **2012**, *125*, 531. [\[CrossRef\]](#)
11. Shimura, H.; Schlossmacher, M.G.; Hattori, N.; Frosch, M.P.; Trockenbacher, A.; Schneider, R.; Mizuno, Y.; Kosik, K.S.; Selkoe, D.J. Ubiquitination of a new form of α -synuclein by parkin from human brain: Implications for Parkinson's disease. *Science* **2001**, *293*, 263–269. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Kitada, T.; Asakawa, S.; Hattori, N.; Matsumine, H.; Yamamura, Y.; Minoshima, S.; Yokochi, M.; Mizuno, Y.; Shimizu, N. Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* **1998**, *392*, 605–608. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Dawson, T.M.; Dawson, V.L. The role of parkin in familial and sporadic Parkinson's disease. *Mov. Disord.* **2010**, *25*, S32–S39. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Liu, X.; Liao, X.; Rao, X.; Wang, B.; Zhang, J.; Xu, G.; Jiang, X.; Qin, X.; Chen, C.; Zou, Z. The lysosomal membrane protein LAMP-2 is dispensable for PINK1/Parkin-mediated mitophagy. *FEBS Lett.* **2020**, *594*, 823–840. [\[CrossRef\]](#) [\[PubMed\]](#)
15. Chung, K.K.K.; Zhang, Y.; Lim, K.L.; Tanaka, Y.; Huang, H.; Gao, J.; Ross, C.A.; Dawson, V.L.; Dawson, T.M. Parkin ubiquitinates the α -synuclein-interacting protein, synphilin-1: Implications for Lewy-body formation in Parkinson disease. *Nat. Med.* **2001**, *7*, 1144–1150. [\[CrossRef\]](#)
16. Ribeiro, C.S.; Carneiro, K.; Ross, C.A.; Menezes, J.R.L.; Engelender, S. Synphilin-1 Is Developmentally Localized to Synaptic Terminals, and Its Association with Synaptic Vesicles Is Modulated by α -Synuclein. *J. Biol. Chem.* **2002**, *277*, 23927–23933. [\[CrossRef\]](#) [\[PubMed\]](#)
17. Wilkaniec, A.; Lenkiewicz, A.M.; Czapski, G.A.; Jeśko, H.M.; Hilgier, W.; Brodzik, R.; Gąssowska-Dobrowolska, M.; Culmsee, C.; Adamczyk, A. Extracellular A-Synuclein Oligomers Induce Parkin S-Nitrosylation: Relevance to Sporadic Parkinson's Disease Etiopathology. *Mol. Neurobiol.* **2019**, *56*, 125–140. [\[CrossRef\]](#)
18. Wheeler, T.C.; Chin, L.-S.; Li, Y.; Roudabush, F.L.; Li, L. Regulation of Synaptophysin Degradation by Mammalian Homologues of Seven in Absentia. *J. Biol. Chem.* **2002**, *277*, 10273–10282. [\[CrossRef\]](#)
19. Liani, E.; Eyal, A.; Avraham, E.; Shemer, R.; Szargel, R.; Berg, D.; Bornemann, A.; Riess, O.; Ross, C.A.; Rott, R.; et al. Ubiquitylation of synphilin-1 and α -synuclein by SIAH and its presence in cellular inclusions and Lewy bodies imply a role in Parkinson's disease. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 5500–5505. [\[CrossRef\]](#)
20. Lee, J.T.; Wheeler, T.C.; Li, L.; Chin, L.S. Ubiquitination of α -synuclein by Siah-1 promotes α -synuclein aggregation and apoptotic cell death. *Hum. Mol. Genet.* **2007**, *17*, 906–917. [\[CrossRef\]](#)
21. Rott, R.; Szargel, R.; Haskin, J.; Shani, V.; Shainskaya, A.; Manov, I.; Liani, E.; Avraham, E.; Engelender, S. Monoubiquitylation of α -synuclein by seven in absentia homolog (SIAH) promotes its aggregation in dopaminergic cells. *J. Biol. Chem.* **2008**, *283*, 3316–3328. [\[CrossRef\]](#) [\[PubMed\]](#)
22. Shin, Y.; Klucken, J.; Patterson, C.; Hyman, B.T.; McLean, P.J. The co-chaperone carboxyl terminus of Hsp70-interacting protein (CHIP) mediates α -synuclein degradation decisions between proteasomal and lysosomal pathways. *J. Biol. Chem.* **2005**, *280*, 23727–23734. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Tetzlaff, J.E.; Putcha, P.; Outeiro, T.F.; Ivanov, A.; Berezovska, O.; Hyman, B.T.; McLean, P.J. CHIP targets toxic α -Synuclein oligomers for degradation. *J. Biol. Chem.* **2008**, *283*, 17962–17968. [\[CrossRef\]](#) [\[PubMed\]](#)
24. Kalia, L.V.; Kalia, S.K.; Chau, H.; Lozano, A.M.; Hyman, B.T.; McLean, P.J. Ubiquitylation of α -Synuclein by Carboxyl Terminus Hsp70-Interacting Protein (CHIP) Is Regulated by Bcl-2-Associated Athanogene 5 (BAG5). *PLoS ONE* **2011**, *6*, e14695. [\[CrossRef\]](#)
25. Gerez, J.A.; Prymaczok, N.C.; Rockenstein, E.; Herrmann, U.S.; Schwarz, P.; Adame, A.; Enchev, R.I.; Courtheoux, T.; Boersema, P.J.; Riek, R.; et al. A cullin-RING ubiquitin ligase targets exogenous α -synuclein and inhibits Lewy body-like pathology. *Sci. Transl. Med.* **2019**, *11*, 6722. [\[CrossRef\]](#)
26. Scheffner, M.; Kumar, S. Mammalian HECT ubiquitin-protein ligases: Biological and pathophysiological aspects. *Biochim. Biophys. Acta* **2014**, *1843*, 61–74. [\[CrossRef\]](#)
27. Mund, T.; Masuda-Suzukake, M.; Goedert, M.; Pelham, H.R. Ubiquitination of α -synuclein filaments by Nedd4 ligases. *PLoS ONE* **2018**, *13*, e0200763. [\[CrossRef\]](#)
28. Tofaris, G.K.K.; Hourez, R.; Jung, J.; Kim, K.P.; Goldberg, A.L. Ubiquitin ligase Nedd4 promotes α -synuclein degradation by the endosomal-lysosomal pathway. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 17004–17009. [\[CrossRef\]](#)

29. Sugeno, N.; Hasegawa, T.; Tanaka, N.; Fukuda, M.; Wakabayashi, K.; Oshima, R.; Konno, M.; Miura, E.; Kikuchi, A.; Baba, T.; et al. Lys-63-linked ubiquitination by E3 ubiquitin ligase Nedd4-1 facilitates endosomal sequestration of internalized α -synuclein. *J. Biol. Chem.* **2014**, *289*, 18137–18151. [[CrossRef](#)]
30. Davies, S.E.; Hallett, P.J.; Moens, T.; Smith, G.; Mangano, E.; Kim, H.T.; Goldberg, A.L.; Liu, J.-L.; Isacson, O.; Tofaris, G.K. Enhanced ubiquitin-dependent degradation by Nedd4 protects against α -synuclein accumulation and toxicity in animal models of Parkinson's disease. *Neurobiol. Dis.* **2014**, *64*, 79–87. [[CrossRef](#)]
31. Akkaya, B.G.; Zolnerchiks, J.K.; Ritchie, T.K.; Bauer, B.; Hartz, A.M.; Sullivan, J.A.; Linton, K.J. The multidrug resistance pump ABCB1 is a substrate for the ubiquitin ligase NEDD4-1. *Mol. Membr. Biol.* **2015**, *32*, 39–45. [[CrossRef](#)] [[PubMed](#)]
32. Platta, H.W.; Abrahamsen, H.; Thoresen, S.B.; Stenmark, H. Nedd4-dependent lysine-11-linked polyubiquitination of the tumour suppressor Beclin 1. *Biochem. J.* **2012**, *441*, 399–406. [[CrossRef](#)] [[PubMed](#)]
33. Shenoy, S.K.; Xiao, K.; Venkataramanan, V.; Snyder, P.M.; Freedman, N.J.; Weissman, A.M. Nedd4 Mediates Agonist-dependent Ubiquitination, Lysosomal Targeting, and Degradation of the β 2-Adrenergic Receptor. *J. Biol. Chem.* **2008**, *283*, 22166–22176. [[CrossRef](#)] [[PubMed](#)]
34. Simonin, A.; Fuster, D. Nedd4-1 and β -Arrestin-1 Are Key Regulators of Na⁺/H⁺ Exchanger 1 Ubiquitylation, Endocytosis, and Function. *J. Biol. Chem.* **2010**, *285*, 38293–38303. [[CrossRef](#)] [[PubMed](#)]
35. Harvey, K.F.; Kumar, S. Nedd4-like proteins: An emerging family of ubiquitin-protein ligases implicated in diverse cellular functions. *Trends Cell Biol.* **1999**, *9*, 166–169. [[CrossRef](#)]
36. Jiang, C.; Kawabe, H.; Rotin, D. The Ubiquitin Ligase Nedd4L Regulates the Na/K/2Cl Co-transporter NKCC1/SLC12A2 in the Colon. *J. Biol. Chem.* **2017**, *292*, 3137–3145. [[CrossRef](#)]
37. Liu, Q.; Zhang, S.; Sun, Z.; Guo, X.; Zhou, H. E3 ubiquitin ligase Nedd4 is a key negative regulator for non-canonical inflammasome activation. *Cell Death Differ.* **2019**, *26*, 2386–2399. [[CrossRef](#)]
38. Guo, J.; Wang, T.; Li, X.; Shallow, H.; Yang, T.; Li, W.; Xu, J.; Fridman, M.D.; Yang, X.; Zhang, S. Cell Surface Expression of Human Ether-a-go-go-related Gene (hERG) Channels Is Regulated by Caveolin-3 Protein via the Ubiquitin Ligase Nedd4-2*. *J. Biol. Chem.* **2012**, *287*, 33132–33141. [[CrossRef](#)]
39. Yang, B.; Gay, D.L.; Macleod, M.K.L.; Cao, X.; Hala, T.; Sweezer, E.M.; Kappler, J.; Marrack, P.; Oliver, P.M. Nedd4 augments the adaptive immune response by promoting ubiquitin-mediated degradation of Cbl-b in activated T cells. *Nat. Immunol.* **2008**, *9*, 1356–1363. [[CrossRef](#)]
40. Pham, N.; Rotin, D. Nedd4 regulates ubiquitination and stability of the guanine-nucleotide exchange factor CNrasGEF. *J. Biol. Chem.* **2001**, *276*, 46995–47003. [[CrossRef](#)]
41. Zhang, H.; Nie, W.; Zhang, X.; Zhang, G.; Li, Z.; Wu, H.; Shi, Q.; Chen, Y.; Ding, Z.; Zhou, X.; et al. NEDD4-1 regulates migration and invasion of glioma cells through CNrasGEF ubiquitination in vitro. *PLoS ONE* **2013**, *8*, e82789. [[CrossRef](#)] [[PubMed](#)]
42. Yang, B.; Kumar, S. Nedd4 and Nedd4-2: Closely related ubiquitin-protein ligases with distinct physiological functions. *Cell Death Differ.* **2010**, *17*, 68–77. [[CrossRef](#)] [[PubMed](#)]
43. Persaud, A.; Alberts, P.; Mari, S.; Tong, J.; Murchie, R.; Maspero, E.; Safi, F.; Moran, M.F.; Polo, S.; Rotin, D. Tyrosine phosphorylation of NEDD4 activates its ubiquitin ligase activity. *Sci. Signal.* **2014**, *7*, ra95. [[CrossRef](#)] [[PubMed](#)]
44. Bakkers, J.; Camacho-Carvajal, M.; Nowak, M.; Kramer, C.; Danger, B.; Hammerschmidt, M. Destabilization of $\Delta\alpha$ by Nedd4-Mediated Ubiquitination Ubc9-Mediated Sumoylation, and Its Implications on Dorsoventral Patterning of the Zebrafish Embryo. *Cell Cycle* **2005**, *4*, 790–800. [[CrossRef](#)]
45. Leonard, M.K.; Hill, N.T.; Grant, E.D.; Kadakia, M.P. Δ Np63 α represses nuclear translocation of PTEN by inhibition of NEDD4-1 in keratinocytes. *Arch. Derm. Res.* **2013**, *305*, 733–739. [[CrossRef](#)]
46. Woelk, T.; Oldrini, B.; Maspero, E.; Confalonieri, S.; Cavallaro, E.; Di Fiore, P.P.; Polo, S. Molecular mechanisms of coupled monoubiquitination. *Nat. Cell Biol.* **2006**, *8*, 1246–1254. [[CrossRef](#)]
47. Persaud, A.; Alberts, P.; Hayes, M.; Guettler, S.; Clarke, I.; Sicheri, F.; Dirks, P.; Ciruna, B.; Rotin, D. Nedd4-1 binds and ubiquitylates activated FGFR1 to control its endocytosis and function. *EMBO J.* **2011**, *30*, 3259–3273. [[CrossRef](#)]
48. Vittal, V.; Stewart, M.D.; Brzovic, P.S.; Klevit, R.E. Regulating the Regulators: Recent Revelations in the Control of E3 Ubiquitin Ligases. *J. Biol. Chem.* **2015**, *290*, 21244–21251. [[CrossRef](#)]
49. Blot, V.; Perugi, F.; Gay, B.; PréVost, M.-C.; Briant, L.; Tangy, F.D.R.; Abriel, H.; Staub, O.; DokhéLar, M.-C.; Pique, C. Nedd4.1-mediated ubiquitination and subsequent recruitment of Tsg101 ensure HTLV-1 Gag trafficking towards the multivesicular body pathway prior to virus budding. *J. Cell Sci.* **2004**, *117*, 2357–2367. [[CrossRef](#)]
50. Rost, M.; Döring, T.; Prange, R. γ 2-Adaptin, a ubiquitin-interacting adaptor, is a substrate to coupled ubiquitination by the ubiquitin ligase Nedd4 and functions in the endosomal pathway. *J. Biol. Chem.* **2008**, *283*, 32119–32130. [[CrossRef](#)]
51. Prange, R. Host factors involved in hepatitis B virus maturation, assembly, and egress. *Med. Microbiol. Immunol.* **2012**, *201*, 449–461. [[CrossRef](#)] [[PubMed](#)]
52. Wan, T.; Lei, Z.; Tu, B.; Wang, T.; Wang, J.; Huang, F. NEDD4 Induces K48-Linked Degradative Ubiquitination of Hepatitis B Virus X Protein and Inhibits HBV-Associated HCC Progression. *Front. Oncol.* **2021**, *11*, 625169. [[CrossRef](#)] [[PubMed](#)]
53. Huang, Z.; Choi, B.K.; Mujoo, K.; Fan, X.; Fa, M.; Mukherjee, S.; Owiti, N.; Zhang, N.; An, Z. The E3 ubiquitin ligase NEDD4 negatively regulates HER3/ErbB3 level and signaling. *Oncogene* **2015**, *34*, 1105–1115. [[CrossRef](#)] [[PubMed](#)]
54. Li, Y.; Zhang, L.; Zhou, J.; Luo, S.; Huang, R.; Zhao, C.; Diao, A. Nedd4 E3 ubiquitin ligase promotes cell proliferation and autophagy. *Cell Prolif.* **2015**, *48*, 338–347. [[CrossRef](#)]

55. Katz, M.; Shtiegman, K.; Tal-Or, P.; Yakir, L.; Mosesson, Y.; Harari, D.; Machluf, Y.; Asao, H.; Jovin, T.; Sugamura, K.; et al. Ligand-Independent Degradation of Epidermal Growth Factor Receptor Involves Receptor Ubiquitylation and Hgs, an Adaptor Whose Ubiquitin-Interacting Motif Targets Ubiquitylation by Nedd4. *Traffic* **2002**, *3*, 740–751. [[CrossRef](#)]
56. Chesarino, N.M.; McMichael, T.M.; Yount, J.S. E3 Ubiquitin Ligase NEDD4 Promotes Influenza Virus Infection by Decreasing Levels of the Antiviral Protein IFITM3. *PLoS Pathog.* **2015**, *11*, e1005095. [[CrossRef](#)]
57. Yan, C.; Zhao, M.; Li, S.; Liu, T.; Xu, C.; Liu, L.; Geng, T.; Gong, D. Increase of E3 ubiquitin ligase NEDD4 expression leads to degradation of its target proteins PTEN/IGF1R during the formation of goose fatty liver. *J. Anim. Sci.* **2020**, *98*, 270. [[CrossRef](#)]
58. Sun, L.; Amraei, R.; Rahimi, N. NEDD4 regulates ubiquitination and stability of the cell adhesion molecule IGPR-1 via lysosomal pathway. *J. Biomed. Sci.* **2021**, *28*, 731. [[CrossRef](#)]
59. Fukushima, T.; Yoshihara, H.; Furuta, H.; Kamei, H.; Hakuno, F.; Luan, J.; Duan, C.; Saeki, Y.; Tanaka, K.; Iemura, S.; et al. Nedd4-induced monoubiquitination of IRS-2 enhances IGF signalling and mitogenic activity. *Nat. Commun.* **2015**, *6*, 6780. [[CrossRef](#)]
60. Hu, Y.; Hong, X.Y.; Yang, X.F.; Ma, R.H.; Wang, X.; Zhang, J.F.; Feng, Q.; Li, X.G.; Sun, D.S.; Li, X.; et al. Inflammation-dependent ISG15 upregulation mediates MIA-induced dendrite damages and depression by disrupting NEDD4/Rap2A signaling. *Biochim. Biophys. Acta Mol. Basis Dis.* **2019**, *1865*, 1477–1489. [[CrossRef](#)]
61. Fan, J.-B.; Arimoto, K.-L.; Motamedchaboki, K.; Yan, M.; Wolf, D.A.; Zhang, D.-E. Identification and characterization of a novel ISG15-ubiquitin mixed chain and its role in regulating protein homeostasis. *Sci. Rep.* **2015**, *5*, 12704. [[CrossRef](#)] [[PubMed](#)]
62. Fan, J. Investigating the Crosstalk between Nedd4 Ubiquitin Ligases and PIAS3 SUMO Ligase. Ph.D. Thesis, University of London, London, UK, 2017.
63. Kumar, S.; Behera, A.; Saha, P.; Kumar Srivastava, A. The role of Krüppel-like factor 8 in cancer biology: Current research and its clinical relevance. *Biochem. Pharm.* **2021**, *183*, 114351. [[CrossRef](#)] [[PubMed](#)]
64. Sun, Y.; Cao, L.; Lin, J.T.; Yuan, Y.; Cao, Z.L.; Jia, J.D. Upregulated miRNA-1236-3p in osteosarcoma inhibits cell proliferation and induces apoptosis via targeting KLF8. *Eur. Rev. Med. Pharm. Sci.* **2019**, *23*, 6053–6061. [[CrossRef](#)]
65. Salah, Z.; Cohen, S.; Itzhaki, E.; Aqeilan, R.I. NEDD4 E3 ligase inhibits the activity of the Hippo pathway by targeting LATS1 for degradation. *Cell Cycle* **2013**, *12*, 3817–3823. [[CrossRef](#)]
66. Sun, A.; Wei, J.; Childress, C.; Shaw, J.H.t.; Peng, K.; Shao, G.; Yang, W.; Lin, Q. The E3 ubiquitin ligase NEDD4 is an LC3-interactive protein and regulates autophagy. *Autophagy* **2017**, *13*, 522–537. [[CrossRef](#)]
67. Liu, Z.; Huo, X.; Zhao, S.; Yang, J.; Shi, W.; Jing, L.; Li, W.; Li, Y.; Ma, L.; Gao, Y.; et al. Low density lipoprotein receptor class A domain containing 4 (LDLRAD4) promotes tumorigenesis of hepatic cancer cells. *Exp. Cell Res.* **2017**, *360*, 189–198. [[CrossRef](#)]
68. Ikeda, M.; Ikeda, A.; Longan, L.C.; Longnecker, R. The Epstein-Barr virus latent membrane protein 2A PY motif recruits WW domain-containing ubiquitin-protein ligases. *Virology* **2000**, *268*, 178–191. [[CrossRef](#)]
69. Xu, C.; Fan, C.D.; Wang, X. Regulation of Mdm2 protein stability and the p53 response by NEDD4-1 E3 ligase. *Oncogene* **2015**, *34*, 281–289. [[CrossRef](#)]
70. Sharma, P.; Murillas, R.; Zhang, H.; Kuehn, M.R. N4BP1 is a newly identified nucleolar protein that undergoes SUMO-regulated polyubiquitylation and proteasomal turnover at promyelocytic leukemia nuclear bodies. *J. Cell Sci.* **2010**, *123*, 1227–1234. [[CrossRef](#)]
71. Murillas, R.; Simms, K.S.; Hatakeyama, S.; Weissman, A.M.; Kuehn, M.R. Identification of developmentally expressed proteins that functionally interact with Nedd4 ubiquitin ligase. *J. Biol. Chem.* **2002**, *277*, 2897–2907. [[CrossRef](#)]
72. Hatstat, A.K.; Ahrendt, H.D.; Foster, M.W.; Mayne, L.; Moseley, M.A.; Englander, S.W.; McCafferty, D.G. Characterization of Small-Molecule-Induced Changes in Parkinson's-Related Trafficking via the Nedd4 Ubiquitin Signaling Cascade. *Cell Chem. Biol.* **2021**, *28*, 14–25.e19. [[CrossRef](#)] [[PubMed](#)]
73. Tardiff, D.F.; Jui, N.T.; Khurana, V.; Tambe, M.A.; Thompson, M.L.; Chung, C.Y.; Kamadurai, H.B.; Kim, H.T.; Lancaster, A.K.; Caldwell, K.A.; et al. Yeast Reveal a “Druggable” Rsp5/Nedd4 Network that Ameliorates -Synuclein Toxicity in Neurons. *Science* **2013**, *342*, 979–983. [[CrossRef](#)] [[PubMed](#)]
74. Xie, W.; Jin, S.; Wu, Y.; Xian, H.; Tian, S.; Liu, D.A.; Guo, Z.; Cui, J. Auto-ubiquitination of NEDD4-1 Recruits USP13 to Facilitate Autophagy through Deubiquitinating VPS34. *Cell Rep.* **2020**, *30*, 2807–2819.e2804. [[CrossRef](#)] [[PubMed](#)]
75. Quirit, J.G.; Lavrenov, S.N.; Poindexter, K.; Xu, J.; Kyauk, C.; Durkin, K.A.; Aronchik, I.; Tomasiak, T.; Solomatin, Y.A.; Preobrazhenskaya, M.N.; et al. Indole-3-carbinol (I3C) analogues are potent small molecule inhibitors of NEDD4-1 ubiquitin ligase activity that disrupt proliferation of human melanoma cells. *Biochem. Pharm.* **2017**, *127*, 13–27. [[CrossRef](#)] [[PubMed](#)]
76. Liu, P.Y.; Xu, N.; Malyukova, A.; Scarlett, C.J.; Sun, Y.T.; Zhang, X.D.; Ling, D.; Su, S.P.; Nelson, C.; Chang, D.K.; et al. The histone deacetylase SIRT2 stabilizes Myc oncoproteins. *Cell Death Differ.* **2013**, *20*, 503–514. [[CrossRef](#)]
77. Sakata, T.; Sakaguchi, H.; Tsuda, L.; Higashitani, A.; Aigaki, T.; Matsuno, K.; Hayashi, S. Drosophila Nedd4 regulates endocytosis of notch and suppresses its ligand-independent activation. *Curr. Biol.* **2004**, *14*, 2228–2236. [[CrossRef](#)]
78. Wilkin, M.B.; Carbery, A.M.; Fostier, M.; Aslam, H.; Mazaleyrat, S.L.; Higgs, J.; Myat, A.; Evans, D.A.; Cornell, M.; Baron, M. Regulation of notch endosomal sorting and signaling by Drosophila Nedd4 family proteins. *Curr. Biol.* **2004**, *14*, 2237–2244. [[CrossRef](#)]
79. Koncarevic, A.; Jackman, R.W.; Kandarian, S.C. The ubiquitin-protein ligase Nedd4 targets Notch1 in skeletal muscle and distinguishes the subset of atrophies caused by reduced muscle tension. *FASEB J.* **2007**, *21*, 427–437. [[CrossRef](#)]

80. Wang, L.; Zhu, B.; Wang, S.; Wu, Y.; Zhan, W.; Xie, S.; Shi, H.; Yu, R. Regulation of glioma migration and invasion via modification of Rap2a activity by the ubiquitin ligase Nedd4-1. *Oncol. Rep.* **2017**, *37*, 2565–2574. [\[CrossRef\]](#)
81. Zeng, T.; Wang, Q.; Fu, J.; Lin, Q.; Bi, J.; Ding, W.; Qiao, Y.; Zhang, S.; Zhao, W.; Lin, H.; et al. Impeded Nedd4-1-mediated Ras degradation underlies Ras-driven tumorigenesis. *Cell Rep.* **2014**, *7*, 871–882. [\[CrossRef\]](#)
82. Hyndman, B.D.; Crupi, M.J.F.; Peng, S.; Bone, L.N.; Rekab, A.N.; Lian, E.Y.; Wagner, S.M.; Antonescu, C.N.; Mulligan, L.M. Differential recruitment of E3 ubiquitin ligase complexes regulates RET isoform internalization. *J. Cell Sci.* **2017**, *130*, 3282–3296. [\[CrossRef\]](#) [\[PubMed\]](#)
83. Anindya, R.; Aygün, O.; Svejstrup, J.Q. Damage-induced ubiquitylation of human RNA polymerase II by the ubiquitin ligase Nedd4, but not Cockayne syndrome proteins or BRCA1. *Mol. Cell* **2007**, *28*, 386–397. [\[CrossRef\]](#) [\[PubMed\]](#)
84. Harreman, M.; Taschner, M.; Sigurdsson, S.; Anindya, R.; Reid, J.; Somesh, B.; Kong, S.E.; Banks, C.A.; Conaway, R.C.; Conaway, J.W.; et al. Distinct ubiquitin ligases act sequentially for RNA polymerase II polyubiquitylation. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 20705–20710. [\[CrossRef\]](#) [\[PubMed\]](#)
85. Canal, M.; Martín-Flores, N.; Pérez-Sisqués, L.; Romani-Aumedes, J.; Altas, B.; Man, H.-Y.; Kawabe, H.; Alberch, J.; Malagelada, C. Loss of NEDD4 contributes to RTP801 elevation and neuron toxicity: Implications for Parkinson's disease. *Oncotarget* **2016**, *7*, 58813–58831. [\[CrossRef\]](#)
86. Zhou, W.; Xu, J.; Zhao, Y.; Sun, Y. SAG/RBX2 is a novel substrate of NEDD4-1 E3 ubiquitin ligase and mediates NEDD4-1 induced chemosensitization. *Oncotarget* **2014**, *5*, 6746–6755. [\[CrossRef\]](#)
87. Aoh, Q.L.; Castle, A.M.; Hubbard, C.H.; Katsumata, O.; Castle, J.D. SCAMP3 Negatively Regulates Epidermal Growth Factor Receptor Degradation and Promotes Receptor Recycling. *Mol. Biol. Cell* **2009**, *20*, 1816–1832. [\[CrossRef\]](#)
88. Al Sorkhy, M.; Craig, R.; Market, B.; Ard, R.; Porter, L.A. The cyclin-dependent kinase activator, Spy1A, is targeted for degradation by the ubiquitin ligase NEDD4. *J. Biol. Chem.* **2009**, *284*, 2617–2627. [\[CrossRef\]](#)
89. Fifield, B.-A.; Qemo, I.; Kirou, E.; Cardiff, R.D.; Porter, L.A. The atypical cyclin-like protein Spy1 overrides p53-mediated tumour suppression and promotes susceptibility to breast tumourigenesis. *Breast Cancer Res.* **2019**, *21*, 1211. [\[CrossRef\]](#)
90. Lin, Q.; Dai, Q.; Meng, H.; Sun, A.; Wei, J.; Peng, K.; Childress, C.; Chen, M.; Shao, G.; Yang, W. The HECT E3 ubiquitin ligase NEDD4 interacts with and ubiquitylates SQSTM1 for inclusion body autophagy. *J. Cell Sci.* **2017**, *130*, 3839–3850. [\[CrossRef\]](#)
91. Murdaca, J.; Treins, C.; Monthouël-Kartmann, M.N.; Pontier-Bres, R.; Kumar, S.; Van Obberghen, E.; Giorgetti-Peraldi, S. Grb10 prevents Nedd4-mediated vascular endothelial growth factor receptor-2 degradation. *J. Biol. Chem.* **2004**, *279*, 26754–26761. [\[CrossRef\]](#)
92. Bhalla, V.; Daidié, D.e.; Li, H.; Pao, A.C.; LaGrange, L.P.; Wang, J.; Vandewalle, A.; Stockand, J.D.; Staub, O.; Pearce, D. Serum- and Glucocorticoid-Regulated Kinase 1 Regulates Ubiquitin Ligase Neural Precursor Cell-Expressed, Developmentally Down-Regulated Protein 4-2 by Inducing Interaction with 14-3-3. *Mol. Endocrinol.* **2005**, *19*, 3073–3084. [\[CrossRef\]](#) [\[PubMed\]](#)
93. Pohl, P.; Joshi, R.; Petrvalska, O.; Obsil, T.; Obsilova, V. 14-3-3-protein regulates Nedd4-2 by modulating interactions between HECT and WW domains. *Commun. Biol.* **2021**, *4*, 2419. [\[CrossRef\]](#) [\[PubMed\]](#)
94. Mackenzie, K.; Foot, N.J.; Anand, S.; Dalton, H.E.; Chaudhary, N.; Collins, B.M.; Mathivanan, S.; Kumar, S. Regulation of the divalent metal ion transporter via membrane budding. *Cell Discov.* **2016**, *2*, 16011. [\[CrossRef\]](#) [\[PubMed\]](#)
95. Shea, F.F.; Rowell, J.L.; Li, Y.; Chang, T.H.; Alvarez, C.E. Mammalian α arrestins link activated seven transmembrane receptors to Nedd4 family e3 ubiquitin ligases and interact with β arrestins. *PLoS ONE* **2012**, *7*, e50557. [\[CrossRef\]](#)
96. Bhalla, V.; Oyster, N.M.; Fitch, A.C.; Wijngaarden, M.A.; Neumann, D.; Schlattner, U.; Pearce, D.; Hallows, K.R. AMP-activated Kinase Inhibits the Epithelial Na⁺ Channel through Functional Regulation of the Ubiquitin Ligase Nedd4-2*. *J. Biol. Chem.* **2006**, *281*, 26159–26169. [\[CrossRef\]](#)
97. Hatanaka, T.; Hatanaka, Y.; Setou, M. Regulation of Amino Acid Transporter ATA2 by Ubiquitin Ligase Nedd4-2*. *J. Biol. Chem.* **2006**, *281*, 35922–35930. [\[CrossRef\]](#)
98. Fu, L.; Rab, A.; Tang, L.; Bebok, Z.; Rowe, S.M.; Bartoszewski, R.; Collawn, J.F. Δ F508 CFTR surface stability is regulated by DAB2 and CHIP-mediated ubiquitination in post-endocytic compartments. *PLoS ONE* **2015**, *10*, e0123131. [\[CrossRef\]](#)
99. Ahsan, M.K.; Figueroa-Hall, L.; Baratta, V.; Garcia-Milian, R.; Lam, T.T.; Hoque, K.; Salas, P.J.; Ameen, N.A. Glucocorticoids and serum- and glucocorticoid-inducible kinase 1 are potent regulators of CFTR in the native intestine: Implications for stress-induced diarrhea. *Am. J. Physiol. Gastrointest Liver Physiol.* **2020**, *319*, G121–G132. [\[CrossRef\]](#)
100. Caohuy, H.; Jozwik, C.; Pollard, H.B. Rescue of Δ F508-CFTR by the SGK1/Nedd4-2 Signaling Pathway*. *J. Biol. Chem.* **2009**, *284*, 25241–25253. [\[CrossRef\]](#)
101. Yamada, H.; Imajoh-Ohmi, S.; Haga, T. The high-affinity choline transporter CHT1 is regulated by the ubiquitin ligase Nedd4-2. *Biomed. Res.* **2012**, *33*, 1–8. [\[CrossRef\]](#)
102. Hryciw, D.H.; Ekberg, J.; Lee, A.; Lensink, I.L.; Kumar, S.; Guggino, W.B.; Cook, D.I.; Pollock, C.A.; Poronnik, P. Nedd4-2 Functionally Interacts with CIC-5: Involvement in Constitutive Albumin Endocytosis in Proximal Tubule Cells*. *J. Biol. Chem.* **2004**, *279*, 54996–55007. [\[CrossRef\]](#) [\[PubMed\]](#)
103. Nanami, M.; Pham, T.D.; Kim, Y.H.; Yang, B.; Sutliff, R.L.; Staub, O.; Klein, J.D.; Lopez-Cayuqueo, K.I.; Chambrey, R.; Park, A.Y.; et al. The Role of Intercalated Cell Nedd4-2 in BP Regulation, Ion Transport, and Transporter Expression. *J. Am. Soc. Nephrol.* **2018**, *29*, 1706–1719. [\[CrossRef\]](#) [\[PubMed\]](#)
104. Arévalo, J.C. Nedd4-2 regulation of voltage-gated ion channels: An update on structure–function relationships and the pathophysiological consequences of dysfunction. *J. Recept. Ligand Channel Res.* **2015**, *53*, 52534. [\[CrossRef\]](#)

105. Embark, H.M.; Böhmer, C.; Palmada, M.; Rajamanickam, J.; Wyatt, A.W.; Wallisch, S.; Capasso, G.; Waldegger, P.; Seyberth, H.W.; Waldegger, S.; et al. Regulation of CLC-Ka/barttin by the ubiquitin ligase Nedd4-2 and the serum- and glucocorticoid-dependent kinases. *Kidney Int.* **2004**, *66*, 1918–1925. [\[CrossRef\]](#)
106. Jentsch, T.J.; Pusch, M. CLC Chloride Channels and Transporters: Structure, Function, Physiology, and Disease. *Physiol. Rev.* **2018**, *98*, 1493–1590. [\[CrossRef\]](#)
107. Kim, Y.H.; Yoo, H.; Hong, A.R.; Kwon, M.; Kang, S.W.; Kim, K.; Song, Y. NEDD4L limits cAMP signaling through ubiquitination of CREB-regulated transcription coactivator 3. *FASEB J.* **2018**, *32*, 4053–4062. [\[CrossRef\]](#)
108. Vina-Vilaseca, A.; Sorkin, A. Lysine 63-linked Polyubiquitination of the Dopamine Transporter Requires WW3 and WW4 Domains of Nedd4-2 and UBE2D Ubiquitin-conjugating Enzymes*. *J. Biol. Chem.* **2010**, *285*, 7645–7656. [\[CrossRef\]](#)
109. Howitt, J.; Putz, U.; Lackovic, J.; Doan, A.; Dorstyn, L.; Cheng, H.; Yang, B.; Chan-Ling, T.; Silke, J.; Kumar, S.; et al. Divalent metal transporter 1 (DMT1) regulation by Ndfip1 prevents metal toxicity in human neurons. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 15489–15494. [\[CrossRef\]](#)
110. Ding, Y.; Zhang, Y.; Xu, C.; Tao, Q.-H.; Chen, Y.-G. HECT Domain-containing E3 Ubiquitin Ligase NEDD4L Negatively Regulates Wnt Signaling by Targeting Dishevelled for Proteasomal Degradation. *J. Biol. Chem.* **2013**, *288*, 8289–8298. [\[CrossRef\]](#)
111. Huang, X.; Chen, J.; Cao, W.; Yang, L.; Chen, Q.; He, J.; Yi, Q.; Huang, H.; Zhang, E.; Cai, Z. The many substrates and functions of NEDD4-1. *Cell Death Dis.* **2019**, *10*, 904. [\[CrossRef\]](#)
112. Boehmer, C.; Henke, G.; Schniepp, R.; Palmada, M.; Rothstein, J.D.; Bröer, S.; Lang, F. Regulation of the glutamate transporter EAAT1 by the ubiquitin ligase Nedd4-2 and the serum and glucocorticoid-inducible kinase isoforms SGK1/3 and protein kinase B. *J. Neurochem.* **2003**, *86*, 1181–1188. [\[CrossRef\]](#) [\[PubMed\]](#)
113. Boehmer, C.; Palmada, M.; Rajamanickam, J.; Schniepp, R.; Amara, S.; Lang, F. Post-translational regulation of EAAT2 function by co-expressed ubiquitin ligase Nedd4-2 is impacted by SGK kinases. *J. Neurochem.* **2006**, *97*, 911–921. [\[CrossRef\]](#) [\[PubMed\]](#)
114. Zhang, Y.; He, X.; Meng, X.; Wu, X.; Tong, H.; Zhang, X.; Qu, S. Regulation of glutamate transporter trafficking by Nedd4-2 in a Parkinson's disease model. *Cell Death Dis.* **2017**, *8*, e2574. [\[CrossRef\]](#) [\[PubMed\]](#)
115. Zhu, J.; Lee, K.Y.; Jewett, K.A.; Man, H.-Y.; Chung, H.J.; Tsai, N.-P. Epilepsy-associated gene Nedd4-2 mediates neuronal activity and seizure susceptibility through AMPA receptors. *PLoS Genet.* **2017**, *13*, e1006634. [\[CrossRef\]](#)
116. Albesa, M.; Grilo, L.S.; Gavillet, B.; Abriel, H. Nedd4-2-dependent ubiquitylation and regulation of the cardiac potassium channel hERG1. *J. Mol. Cell Cardiol.* **2011**, *51*, 90–98. [\[CrossRef\]](#) [\[PubMed\]](#)
117. Edinger, R.S.; Lebowitz, J.; Li, H.; Alzamora, R.; Wang, H.; Johnson, J.P.; Hallows, K.R. Functional Regulation of the Epithelial Na⁺ Channel by IκB Kinase-β Occurs via Phosphorylation of the Ubiquitin Ligase Nedd4-2*. *J. Biol. Chem.* **2009**, *284*, 150–157. [\[CrossRef\]](#) [\[PubMed\]](#)
118. Hallows, K.R.; Bhalla, V.; Oyster, N.M.; Wijngaarden, M.A.; Lee, J.K.; Li, H.; Chandran, S.; Xia, X.; Huang, Z.; Chalkley, R.J.; et al. Phosphopeptide Screen Uncovers Novel Phosphorylation Sites of Nedd4-2 That Potentiate Its Inhibition of the Epithelial Na⁺ Channel*. *J. Biol. Chem.* **2010**, *285*, 21671–21678. [\[CrossRef\]](#)
119. Ekberg, J.; Schuetz, F.; Boase, N.A.; Conroy, S.-J.; Manning, J.; Kumar, S.; Poronnik, P.; Adams, D.J. Regulation of the Voltage-gated K⁺ Channels KCNQ2/3 and KCNQ3/5 by Ubiquitination: Novel Role for Nedd4-2*. *J. Biol. Chem.* **2007**, *282*, 12135–12142. [\[CrossRef\]](#)
120. Jespersen, T.; Membrez, M.; Nicolas, C.S.; Pitard, B.; Staub, O.; Olesen, S.-P.; Baró, I.; Abriel, H. The KCNQ1 potassium channel is down-regulated by ubiquitylating enzymes of the Nedd4/Nedd4-like family. *Cardiovasc. Res.* **2007**, *74*, 64–74. [\[CrossRef\]](#)
121. Alzamora, R.; Gong, F.; Rondanino, C.; Lee, J.K.; Smolak, C.; Pastor-Soler, N.M.; Hallows, K.R. AMP-activated protein kinase inhibits KCNQ1 channels through regulation of the ubiquitin ligase Nedd4-2 in renal epithelial cells. *Am. J. Physiol. Ren. Physiol.* **2010**, *299*, F1308–F1319. [\[CrossRef\]](#)
122. Rougier, J.-S.; Bemmelen, M.X.v.; Bruce, M.C.; Jespersen, T.; Gavillet, B.; Apothéloz, F.; Cordonier, S.; Staub, O.; Rotin, D.; Abriel, H. Molecular determinants of voltage-gated sodium channel regulation by the Nedd4/Nedd4-like proteins. *Am. J. Physiol.-Cell Physiol.* **2005**, *288*, C692–C701. [\[CrossRef\]](#) [\[PubMed\]](#)
123. Laedermann, C.J.; Cachemaille, M.; Kirschmann, G.; Pertin, M.; Gosselin, R.D.; Chang, I.; Albesa, M.; Towne, C.; Schneider, B.L.; Kellenberger, S.; et al. Dysregulation of voltage-gated sodium channels by ubiquitin ligase NEDD4-2 in neuropathic pain. *J. Clin. Invest.* **2013**, *123*, 3002–3013. [\[CrossRef\]](#) [\[PubMed\]](#)
124. Gasser, A.; Cheng, X.; Gilmore, E.S.; Tyrrell, L.; Waxman, S.G.; Dib-Hajj, S.D. Two Nedd4-binding Motifs Underlie Modulation of Sodium Channel Nav1.6 by p38 MAPK*. *J. Biol. Chem.* **2010**, *285*, 26149–26161. [\[CrossRef\]](#) [\[PubMed\]](#)
125. Arroyo, J.P.; Lagnaz, D.; Ronzaud, C.; Vázquez, N.; Ko, B.S.; Moddes, L.; Ruffieux-Daidié, D.; Hausel, P.; Koesters, R.; Yang, B.; et al. Nedd4-2 modulates renal Na⁺-Cl⁻ cotransporter via the aldosterone-SGK1-Nedd4-2 pathway. *J. Am. Soc. Nephrol.* **2011**, *22*, 1707–1719. [\[CrossRef\]](#)
126. Rotin, D.; Staub, O. Nedd4-2 and the regulation of epithelial sodium transport. *Front. Physiol.* **2012**, *3*, 212. [\[CrossRef\]](#)
127. Bruce, M.C.; Kanelis, V.; Fouladkou, F.; Debonneville, A.; Staub, O.; Rotin, D. Regulation of Nedd4-2 self-ubiquitination and stability by a PY motif located within its HECT-domain. *Biochem. J.* **2008**, *415*, 155–163. [\[CrossRef\]](#)
128. Escobedo, A.; Gomes, T.; Aragón, E.; Martín-Malpartida, P.; Ruiz, L.; Macias, M.J. Structural basis of the activation and degradation mechanisms of the E3 ubiquitin ligase Nedd4L. *Structure* **2014**, *22*, 1446–1457. [\[CrossRef\]](#)
129. Zhao, C.; Jia, M.; Song, H.; Yu, Z.; Wang, W.; Li, Q.; Zhang, L.; Zhao, W.; Cao, X. The E3 Ubiquitin Ligase TRIM40 Attenuates Antiviral Immune Responses by Targeting MDA5 and RIG-I. *Cell Rep.* **2017**, *21*, 1613–1623. [\[CrossRef\]](#)

130. No, Y.R.; He, P.; Yoo, B.K.; Yun, C.C. Unique Regulation of Human Na⁺/H⁺ Exchanger 3 (NHE3) by Nedd4-2 Ligase That Differs from Non-primate NHE3s. *J. Biol. Chem.* **2014**, *289*, 18360–18372. [\[CrossRef\]](#)
131. Xu, D.; Wang, H.; You, G. An Essential Role of Nedd4-2 in the Ubiquitination, Expression, and Function of Organic Anion Transporter-3. *Mol. Pharm.* **2016**, *13*, 621–630. [\[CrossRef\]](#)
132. Xu, D.; Zhang, J.; Zhang, Q.; Fan, Y.; Liu, C.; You, G. PKC/Nedd4-2 Signaling Pathway Regulates the Cell Surface Expression of Drug Transporter hOAT1. *Drug Metab. Dispos.* **2017**, *45*, 887–895. [\[CrossRef\]](#) [\[PubMed\]](#)
133. Raikwar, N.S.; Vandewalle, A.; Thomas, C.P. Nedd4-2 interacts with occludin to inhibit tight junction formation and enhance paracellular conductance in collecting duct epithelia. *Am. J. Physiol.-Ren. Physiol.* **2010**, *299*, F436–F444. [\[CrossRef\]](#) [\[PubMed\]](#)
134. Snyder, P.M.; Olson, D.R.; Kabra, R.; Zhou, R.; Steines, J.C. cAMP and serum and glucocorticoid-inducible kinase (SGK) regulate the epithelial Na(+) channel through convergent phosphorylation of Nedd4-2. *J. Biol. Chem.* **2004**, *279*, 45753–45758. [\[CrossRef\]](#)
135. Wiemuth, D.; Lott, J.S.; Ly, K.; Ke, Y.; Teesdale-Spittle, P.; Snyder, P.M.; McDonald, F.J. Interaction of Serum- and Glucocorticoid Regulated Kinase 1 (SGK1) with the WW-Domains of Nedd4-2 Is Required for Epithelial Sodium Channel Regulation. *PLoS ONE* **2010**, *5*, e12163. [\[CrossRef\]](#) [\[PubMed\]](#)
136. Ingham, R.J.; Gish, G.; Pawson, T. The Nedd4 family of E3 ubiquitin ligases: Functional diversity within a common modular architecture. *Oncogene* **2004**, *23*, 1972–1984. [\[CrossRef\]](#)
137. Dieter, M.; Palmada, M.; Rajamanickam, J.; Aydin, A.; Busjahn, A.; Boehmer, C.; Luft, F.C.; Lang, F. Regulation of Glucose Transporter SGLT1 by Ubiquitin Ligase Nedd4-2 and Kinases SGK1, SGK3, and PKB. *Obes. Res.* **2004**, *12*, 862–870. [\[CrossRef\]](#)
138. Inoue, Y.; Imamura, T. Regulation of TGF- β family signaling by E3 ubiquitin ligases. *Cancer Sci.* **2008**, *99*, 2107–2112. [\[CrossRef\]](#)
139. Morén, A.; Imamura, T.; Miyazono, K.; Heldin, C.-H.; Moustakas, A. Degradation of the Tumor Suppressor Smad4 by WW and HECT Domain Ubiquitin Ligases. *J. Biol. Chem.* **2005**, *280*, 22115–22123. [\[CrossRef\]](#)
140. Kuratomi, G.; Komuro, A.; Goto, K.; Shinozaki, M.; Miyazawa, K.; Miyazono, K.; Imamura, T. NEDD4-2 (neural precursor cell expressed, developmentally down-regulated 4-2) negatively regulates TGF- β (transforming growth factor- β) signalling by inducing ubiquitin-mediated degradation of Smad2 and TGF- β type I receptor. *Biochem. J.* **2005**, *386*, 461–470. [\[CrossRef\]](#)
141. Gao, S.; Alarcón, C.; Sapkota, G.; Rahman, S.; Chen, P.-Y.; Goerner, N.; Macias, M.J.; Erdjument-Bromage, H.; Tempst, P.; Massagué, J. Ubiquitin Ligase Nedd4L Targets Activated Smad2/3 to Limit TGF- β Signaling. *Mol. Cell* **2009**, *36*, 457–468. [\[CrossRef\]](#)
142. Konkright, J.J.; Apsley, K.S.; Martin, E.P.; Ridsdale, R.; Rice, W.R.; Na, C.-L.; Yang, B.; Weaver, T.E. Nedd4-2-Mediated Ubiquitination Facilitates Processing of Surfactant Protein-C. *Am. J. Respir. Cell Mol. Biol.* **2010**, *42*, 181–189. [\[CrossRef\]](#) [\[PubMed\]](#)
143. Zeng, Y.; Feng, Z.; Liao, Y.; Yang, M.; Bai, Y.; He, Z. Diminution of microRNA-98 alleviates renal fibrosis in diabetic nephropathy by elevating Nedd4L and inactivating TGF- β /Smad2/3 pathway. *Cell Cycle* **2020**, *19*, 3406–3418. [\[CrossRef\]](#) [\[PubMed\]](#)
144. Sánchez-Sánchez, J.; Arévalo, J. A Review on Ubiquitination of Neurotrophin Receptors: Facts and Perspectives. *Int. J. Mol. Sci.* **2017**, *18*, 630. [\[CrossRef\]](#) [\[PubMed\]](#)
145. Arévalo, J.C.; Waite, J.; Rajagopal, R.; Beyna, M.; Chen, Z.-Y.; Lee, F.S.; Chao, M.V. Cell Survival through Trk Neurotrophin Receptors Is Differentially Regulated by Ubiquitination. *Neuron* **2006**, *50*, 549–559. [\[CrossRef\]](#)
146. Georgieva, M.V.; de Pablo, Y.; Sanchis, D.; Comella, J.X.; Llovera, M. Ubiquitination of TrkA by Nedd4-2 regulates receptor lysosomal targeting and mediates receptor signaling. *J. Neurochem.* **2011**, *117*, 479–493. [\[CrossRef\]](#)
147. He, Y.; Hryciw, D.H.; Carroll, M.L.; Myers, S.A.; Whitbread, A.K.; Kumar, S.; Poronnik, P.; Hooper, J.D. The Ubiquitin-Protein Ligase Nedd4-2 Differentially Interacts with and Regulates Members of the Tweety Family of Chloride Ion Channels*. *J. Biol. Chem.* **2008**, *283*, 24000–24010. [\[CrossRef\]](#)
148. Nazio, F.; Carinci, M.; Valacca, C.; Bielli, P.; Strappazzon, F.; Antonioli, M.; Ciccocanti, F.; Rodolfo, C.; Campello, S.; Fimia, G.M.; et al. Fine-tuning of ULK1 mRNA and protein levels is required for autophagy oscillation. *J. Cell Biol.* **2016**, *215*, 841–856. [\[CrossRef\]](#)
149. Oberfeld, B.; Ruffieux-Daidié, D.; Vitagliano, J.-J.; Pos, K.M.; Verrey, F.; Staub, O. Ubiquitin-specific protease 2-45 (Usp2-45) binds to epithelial Na⁺ channel (ENaC)-ubiquitylating enzyme Nedd4-2. *Am. J. Physiol.-Ren. Physiol.* **2011**, *301*, F189–F196. [\[CrossRef\]](#)
150. Roy, A.; Al-Qusairi, L.; Donnelly, B.F.; Ronzaud, C.; Marciszyn, A.L.; Gong, F.; Chang, Y.P.C.; Butterworth, M.B.; Pastor-Soler, N.M.; Hallows, K.R.; et al. Alternatively spliced proline-rich cassettes link WNK1 to aldosterone action. *J. Clin. Investig.* **2015**, *125*, 3433–3448. [\[CrossRef\]](#)
151. Lin, Q.; Wang, J.; Childress, C.; Sudol, M.; Carey, D.J.; Yang, W. HECT E3 ubiquitin ligase Nedd4-1 ubiquitinates ACK and regulates epidermal growth factor (EGF)-induced degradation of EGF receptor and ACK. *Mol. Cell Biol.* **2010**, *30*, 1541–1554. [\[CrossRef\]](#)
152. Chan, W.; Tian, R.; Lee, Y.-F.; Sit, S.T.; Lim, L.; Manser, E. Down-regulation of Active ACK1 Is Mediated by Association with the E3 Ubiquitin Ligase Nedd4-2*. *J. Biol. Chem.* **2009**, *284*, 8185–8194. [\[CrossRef\]](#) [\[PubMed\]](#)
153. Fan, C.D.; Lum, M.A.; Xu, C.; Black, J.D.; Wang, X. Ubiquitin-dependent regulation of phospho-AKT dynamics by the ubiquitin E3 ligase, NEDD4-1, in the insulin-like growth factor-1 response. *J. Biol. Chem.* **2013**, *288*, 1674–1684. [\[CrossRef\]](#) [\[PubMed\]](#)
154. Persaud, A.; Alberts, P.; Amsen, E.M.; Xiong, X.; Wasmuth, J.; Saadon, Z.; Fladd, C.; Parkinson, J.; Rotin, D. Comparison of substrate specificity of the ubiquitin ligases Nedd4 and Nedd4-2 using proteome arrays. *Mol. Syst. Biol.* **2009**, *5*, 333. [\[CrossRef\]](#) [\[PubMed\]](#)
155. Lee, I.-H.; Dinudom, A.; Sanchez-Perez, A.; Kumar, S.; Cook, D.I. Akt Mediates the Effect of Insulin on Epithelial Sodium Channels by Inhibiting Nedd4-2*. *J. Biol. Chem.* **2007**, *282*, 29866–29873. [\[CrossRef\]](#) [\[PubMed\]](#)

156. Scudder, S.L.; Goo, M.S.; Cartier, A.E.; Molteni, A.; Schwarz, L.A.; Wright, R.; Patrick, G.N. Synaptic strength is bidirectionally controlled by opposing activity-dependent regulation of Nedd4-1 and USP8. *J. Neurosci. Off. J. Soc. Neurosci.* **2014**, *34*, 16637–16649. [\[CrossRef\]](#)
157. Schwarz, L.A.; Hall, B.J.; Patrick, G.N. Activity-dependent ubiquitination of GluA1 mediates a distinct AMPA receptor endocytosis and sorting pathway. *J. Neurosci. Off. J. Soc. Neurosci.* **2010**, *30*, 16718–16729. [\[CrossRef\]](#)
158. Lin, A.; Hou, Q.; Jarzylo, L.; Amato, S.; Gilbert, J.; Shang, F.; Man, H.-Y. Nedd4-mediated AMPA receptor ubiquitination regulates receptor turnover and trafficking. *J. Neurochem.* **2011**, *119*, 27–39. [\[CrossRef\]](#)
159. Kim, J.E.; Lee, D.S.; Park, H.; Kim, T.H.; Kang, T.C. AMPA Receptor Antagonists Facilitate NEDD4-2-Mediated GRIA1 Ubiquitination by Regulating PP2B-ERK1/2-SGK1 Pathway in Chronic Epilepsy Rats. *Biomedicines* **2021**, *9*, 1069. [\[CrossRef\]](#)
160. Jewett, K.A.; Zhu, J.; Tsai, N.P. The tumor suppressor p53 guides GluA1 homeostasis through Nedd4-2 during chronic elevation of neuronal activity. *J. Neurochem.* **2015**, *135*, 226–233. [\[CrossRef\]](#)
161. Trimpert, C.; Wesche, D.; de Groot, T.; Pimentel Rodriguez, M.M.; Wong, V.; van den Berg, D.T.M.; Cheval, L.; Ariza, C.A.; Doucet, A.; Stagljar, I.; et al. NDFIP allows NEDD4/NEDD4L-induced AQP2 ubiquitination and degradation. *PLoS ONE* **2017**, *12*, e0183774. [\[CrossRef\]](#)
162. Claude, Eitelhuber, A.; Christian; Giallonardo, P.; Gegg, M.; Oller, H.; Seth; Krappmann, D.; Ueffing, M.; Lickert, H. Dlg3 Trafficking and Apical Tight Junction Formation Is Regulated by Nedd4 and Nedd4-2 E3 Ubiquitin Ligases. *Dev. Cell* **2011**, *21*, 479–491. [\[CrossRef\]](#)
163. Novellasademunt, L.; Kucharska, A.; Jamieson, C.; Prange-Barczynska, M.; Baulies, A.; Antas, P.; van der Vaart, J.; Gehart, H.; Maurice, M.M.; Li, V.S. NEDD4 and NEDD4L regulate Wnt signalling and intestinal stem cell priming by degrading LGR5 receptor. *EMBO J.* **2020**, *39*, e102771. [\[CrossRef\]](#) [\[PubMed\]](#)
164. Dinudom, A.; Harvey, K.F.; Komwatana, P.; Young, J.A.; Kumar, S.; Cook, D.I. Nedd4 mediates control of an epithelial Na⁺ channel in salivary duct cells by cytosolic Na⁺. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 7169–7173. [\[CrossRef\]](#) [\[PubMed\]](#)
165. Staub, O.; Dho, S.; Henry, P.; Correa, J.; Ishikawa, T.; McGlade, J.; Rotin, D. WW domains of Nedd4 bind to the proline-rich PY motifs in the epithelial Na⁺ channel deleted in Liddle's syndrome. *EMBO J.* **1996**, *15*, 2371–2380. [\[CrossRef\]](#) [\[PubMed\]](#)
166. Zhou, R.; Patel, S.V.; Snyder, P.M. Nedd4-2 catalyzes ubiquitination and degradation of cell surface ENaC. *J. Biol. Chem.* **2007**, *282*, 20207–20212. [\[CrossRef\]](#)
167. Kanelis, V.; Rotin, D.; Forman-Kay, J.D. Solution structure of a Nedd4 WW domain-ENaC peptide complex. *Nat. Struct. Biol.* **2001**, *8*, 407–412. [\[CrossRef\]](#)
168. Monami, G.; Emiliozzi, V.; Morrione, A. Grb10/Nedd4-mediated multiubiquitination of the insulin-like growth factor receptor regulates receptor internalization. *J. Cell Physiol.* **2008**, *216*, 426–437. [\[CrossRef\]](#)
169. Nakayama, M.; Berger, P. Coordination of VEGF receptor trafficking and signaling by coreceptors. *Exp. Cell Res.* **2013**, *319*, 1340–1347. [\[CrossRef\]](#)
170. Cheng, W.; Huang, P.C.; Chao, H.M.; Jeng, Y.M.; Hsu, H.C.; Pan, H.W.; Hwu, W.L.; Lee, Y.M. Glypican-3 induces oncogenicity by preventing IGF-1R degradation, a process that can be blocked by Grb10. *Oncotarget* **2017**, *8*, 80429–80442. [\[CrossRef\]](#)
171. Vélez, P.; Schwartz, A.B.; Iyer, S.R.; Warrington, A.; Fadool, D.A. Ubiquitin ligase Nedd4-2 modulates Kv1.3 current amplitude and ion channel protein targeting. *J. Neurophysiol.* **2016**, *116*, 671–685. [\[CrossRef\]](#)
172. Plant, P.J.; Correa, J.; Goldenberg, N.; Bain, J.; Batt, J. The inositol phosphatase MTMR4 is a novel target of the ubiquitin ligase Nedd4. *Biochem. J.* **2009**, *419*, 57–63. [\[CrossRef\]](#) [\[PubMed\]](#)
173. Nagpal, P.; Plant, P.J.; Correa, J.; Bain, A.; Takeda, M.; Kawabe, H.; Rotin, D.; Bain, J.R.; Batt, J.A.E. The ubiquitin ligase Nedd4-1 participates in denervation-induced skeletal muscle atrophy in mice. *PLoS ONE* **2012**, *7*, e46427. [\[CrossRef\]](#) [\[PubMed\]](#)
174. Lee, Y.K.; Sala, L.; Mura, M.; Rocchetti, M.; Pedrazzini, M.; Ran, X.; Mak, T.S.H.; Crotti, L.; Sham, P.C.; Torre, E.; et al. MTMR4 SNVs modulate ion channel degradation and clinical severity in congenital long QT syndrome: Insights in the mechanism of action of protective modifier genes. *Cardiovasc. Res.* **2021**, *117*, 767–779. [\[CrossRef\]](#) [\[PubMed\]](#)
175. Fotia, A.B.; Dinudom, A.; Shearwin, K.E.; Koch, J.-P.; Korbmacher, C.; Cook, D.I.; Kumar, S. The role of individual Nedd4-2 (KIAA0439) WW domains in binding and regulating epithelial sodium channels. *FASEB J.* **2003**, *17*, 70–72. [\[CrossRef\]](#) [\[PubMed\]](#)
176. Howitt, J.; Lackovic, J.; Low, L.H.; Naguib, A.; Macintyre, A.; Goh, C.P.; Callaway, J.K.; Hammond, V.; Thomas, T.; Dixon, M.; et al. Ndfip1 regulates nuclear Pten import in vivo to promote neuronal survival following cerebral ischemia. *J. Cell Biol.* **2012**, *196*, 29–36. [\[CrossRef\]](#)
177. Trotman, L.C.; Wang, X.; Alimonti, A.; Chen, Z.; Teruya-Feldstein, J.; Yang, H.; Pavletich, N.P.; Carver, B.S.; Cordon-Cardo, C.; Erdjument-Bromage, H.; et al. Ubiquitination regulates PTEN nuclear import and tumor suppression. *Cell* **2007**, *128*, 141–156. [\[CrossRef\]](#)
178. Wang, X.; Trotman, L.C.; Koppie, T.; Alimonti, A.; Chen, Z.; Gao, Z.; Wang, J.; Erdjument-Bromage, H.; Tempst, P.; Cordon-Cardo, C.; et al. NEDD4-1 is a proto-oncogenic ubiquitin ligase for PTEN. *Cell* **2007**, *128*, 129–139. [\[CrossRef\]](#)
179. Amodio, N.; Scrima, M.; Palaia, L.; Salman, A.N.; Quintiero, A.; Franco, R.; Botti, G.; Pirozzi, P.; Rocco, G.; De Rosa, N.; et al. Oncogenic role of the E3 ubiquitin ligase NEDD4-1, a PTEN negative regulator, in non-small-cell lung carcinomas. *Am. J. Pathol.* **2010**, *177*, 2622–2634. [\[CrossRef\]](#)
180. Drinjakovic, J.; Jung, H.; Campbell, D.S.; Strohlic, L.; Dwivedy, A.; Holt, C.E. E3 ligase Nedd4 promotes axon branching by downregulating PTEN. *Neuron* **2010**, *65*, 341–357. [\[CrossRef\]](#)

181. Fouladkou, F.; Landry, T.; Kawabe, H.; Neeb, A.; Lu, C.; Brose, N.; Stambolic, V.; Rotin, D. The ubiquitin ligase Nedd4-1 is dispensable for the regulation of PTEN stability and localization. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 8585–8590. [\[CrossRef\]](#)
182. Hsia, H.-E.; Kumar, R.; Luca, R.; Takeda, M.; Courchet, J.; Nakashima, J.; Wu, S.; Goebbels, S.; An, W.; Eickholt, B.J.; et al. Ubiquitin E3 ligase Nedd4-1 acts as a downstream target of PI3K/PTEN-mTORC1 signaling to promote neurite growth. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 13205. [\[CrossRef\]](#) [\[PubMed\]](#)
183. Giasson, B.I.; Murray, I.V.J.; Trojanowski, J.Q.; Lee, V.M.Y. A Hydrophobic Stretch of 12 Amino Acid Residues in the Middle of α -Synuclein Is Essential for Filament Assembly. *J. Biol. Chem.* **2001**, *276*, 2380–2386. [\[CrossRef\]](#) [\[PubMed\]](#)
184. Conway, K.A.; Harper, J.D.; Lansbury, P.T. Accelerated in vitro fibril formation by a mutant α -synuclein linked to early-onset Parkinson disease. *Nat. Med.* **1998**, *4*, 1318–1320. [\[CrossRef\]](#) [\[PubMed\]](#)
185. Chung, C.Y.; Khurana, V.; Auluck, P.K.; Tardiff, D.F.; Mazzulli, J.R.; Soldner, F.; Barua, V.; Lou, Y.; Frey, Y.; Cho, S.; et al. Identification and Rescue of α -Synuclein Toxicity in Parkinson Patient-Derived Neurons. *Science* **2013**, *342*, 983–987. [\[CrossRef\]](#) [\[PubMed\]](#)
186. Itier, J.M. Parkin gene inactivation alters behaviour and dopamine neurotransmission in the mouse. *Hum. Mol. Genet.* **2003**, *12*, 2277–2291. [\[CrossRef\]](#)
187. Dickens, R.A.; Frew, I.J.; House, C.M.; O'Bryan, M.K.; Holloway, A.J.; Haviv, I.; Traficante, N.; De Kretser, D.M.; Bowtell, D.D.L. The Ubiquitin Ligase Component Siah1a Is Required for Completion of Meiosis I in Male Mice. *Mol. Cell. Biol.* **2002**, *22*, 2294–2303. [\[CrossRef\]](#)
188. Frew, I.J.; Dickens, R.A.; Cuddihy, A.R.; Del Rosario, M.; Reinhard, C.; O'Connell, M.J.; Bowtell, D.D.L. Normal p53 Function in Primary Cells Deficient for Siah Genes. *Mol. Cell. Biol.* **2002**, *22*, 8155–8164. [\[CrossRef\]](#)
189. Frew, I.J.; Hammond, V.E.; Dickens, R.A.; Quinn, J.M.W.; Walkley, C.R.; Sims, N.A.; Schnall, R.; Della, N.G.; Holloway, A.J.; Digby, M.R.; et al. Generation and Analysis of Siah2 Mutant Mice. *Mol. Cell. Biol.* **2003**, *23*, 9150–9161. [\[CrossRef\]](#)
190. Schisler, J.C.; Rubel, C.E.; Zhang, C.; Lockyer, P.; Cyr, D.M.; Patterson, C. CHIP protects against cardiac pressure overload through regulation of AMPK. *J. Clin. Invest.* **2013**, *123*, 3588–3599. [\[CrossRef\]](#)
191. Moroishi, T.; Nishiyama, M.; Takeda, Y.; Iwai, K.; Keiichi. The FBXL5-IRP2 Axis Is Integral to Control of Iron Metabolism In Vivo. *Cell Metab.* **2011**, *14*, 339–351. [\[CrossRef\]](#)
192. Perry, W.L.; Hustad, C.M.; Swing, D.A.; O'Sullivan, T.N.; Jenkins, N.A.; Copeland, N.G. The itchy locus encodes a novel ubiquitin protein ligase that is disrupted in a18H mice. *Nat. Genet.* **1998**, *18*, 143–146. [\[CrossRef\]](#) [\[PubMed\]](#)
193. Yamashita, M.; Ying, S.-X.; Zhang, G.-M.; Li, C.; Cheng, S.Y.; Deng, C.-X.; Zhang, Y.E. Ubiquitin Ligase Smurf1 Controls Osteoblast Activity and Bone Homeostasis by Targeting MEKK2 for Degradation. *Cell* **2005**, *121*, 101–113. [\[CrossRef\]](#) [\[PubMed\]](#)
194. Narimatsu, M.; Bose, R.; Pye, M.; Zhang, L.; Miller, B.; Ching, P.; Sakuma, R.; Luga, V.; Roncari, L.; Attisano, L.; et al. Regulation of Planar Cell Polarity by Smurf Ubiquitin Ligases. *Cell* **2009**, *137*, 295–307. [\[CrossRef\]](#) [\[PubMed\]](#)
195. Zhao, D.; Zhi, X.; Zhou, Z.; Chen, C. TAZ antagonizes the WWP1-mediated KLF5 degradation and promotes breast cell proliferation and tumorigenesis. *Carcinogenesis* **2012**, *33*, 59–67. [\[CrossRef\]](#)
196. Zou, W.; Chen, X.; Shim, J.-H.; Huang, Z.; Brady, N.; Hu, D.; Drapp, R.; Sigrist, K.; Glimcher, L.H.; Jones, D. The E3 ubiquitin ligase Wwp2 regulates craniofacial development through mono-ubiquitylation of Goosecoid. *Nat. Cell Biol.* **2011**, *13*, 59–65. [\[CrossRef\]](#)
197. Kawabe, H.; Neeb, A.; Dimova, K.; Young, S.M., Jr.; Takeda, M.; Katsurabayashi, S.; Mitkovski, M.; Malakhova, O.A.; Zhang, D.E.; Umikawa, M.; et al. Regulation of Rap2A by the ubiquitin ligase Nedd4-1 controls neurite development. *Neuron* **2010**, *65*, 358–372. [\[CrossRef\]](#)
198. Boase, N.A.; Rychkov, G.Y.; Townley, S.L.; Dinudom, A.; Candi, E.; Voss, A.K.; Tsoutsman, T.; Semsarian, C.; Melino, G.; Koentgen, F.; et al. Respiratory distress and perinatal lethality in Nedd4-2-deficient mice. *Nat. Commun.* **2011**, *2*, 287. [\[CrossRef\]](#)
199. Leitz, D.; Duerr, J.; Dvornikov, D.; Szczygiel, M.; Konietzke, P.; Rotin, D.; Wielpütz, M.; Kawabe, H.; Beers, M.; Klingmüller, U.; et al. Development of pulmonary fibrosis in conditional Nedd4-2 deficient mice. *Eur. Respir. J.* **2018**, *52*, OA5364. [\[CrossRef\]](#)
200. Frank, C.; Pari, G.; Rossiter, J.P. Approach to diagnosis of Parkinson disease. *Can. Fam. Phys.* **2006**, *52*, 862–868.
201. Nalls, M.A.; Blauwendraat, C.; Vallerga, C.L.; Heilbron, K.; Bandres-Ciga, S.; Chang, D.; Tan, M.; Kia, D.A.; Noyce, A.J.; Xue, A.; et al. Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: A meta-analysis of genome-wide association studies. *Lancet Neurol.* **2019**, *18*, 1091–1102. [\[CrossRef\]](#)
202. Foo, J.N.; Chew, E.G.Y.; Chung, S.J.; Peng, R.; Blauwendraat, C.; Nalls, M.A.; Mok, K.Y.; Satake, W.; Toda, T.; Chao, Y.; et al. Identification of Risk Loci for Parkinson Disease in Asians and Comparison of Risk between Asians and Europeans. *JAMA Neurol.* **2020**, *77*, 746. [\[CrossRef\]](#) [\[PubMed\]](#)
203. Putz, U.; Howitt, J.; Lackovic, J.; Foot, N.; Kumar, S.; Silke, J.; Tan, S.S. Nedd4 family-interacting protein 1 (Ndfip1) is required for the exosomal secretion of Nedd4 family proteins. *J. Biol. Chem.* **2008**, *283*, 32621–32627. [\[CrossRef\]](#) [\[PubMed\]](#)
204. Kumar, S.; Harvey, K.F.; Kinoshita, M.; Copeland, N.G.; Noda, M.; Jenkins, N.A. cDNA cloning, expression analysis, and mapping of the mouse Nedd4 gene. *Genomics* **1997**, *40*, 435–443. [\[CrossRef\]](#) [\[PubMed\]](#)
205. Chen, H.; Ross, C.A.; Wang, N.; Huo, Y.; Mackinnon, D.F.; Potash, J.B.; Simpson, S.G.; McMahon, F.J.; Depaulo, J.J.R.; McInnis, M.G. NEDD4L on human chromosome 18q21 has multiple forms of transcripts and is a homologue of the mouse Nedd4-2 gene. *Eur. J. Hum. Genet.* **2001**, *9*, 922–930. [\[CrossRef\]](#)
206. Itani, O.A.; Campbell, J.R.; Herrero, J.; Snyder, P.M.; Thomas, C.P. Alternate promoters and variable splicing lead to hNedd4-2 isoforms with a C2 domain and varying number of WW domains. *Am. J. Physiol. Ren. Physiol.* **2003**, *285*, F916–F929. [\[CrossRef\]](#)

207. Itani, O.A.; Stokes, J.B.; Thomas, C.P. Nedd4-2 isoforms differentially associate with ENaC and regulate its activity. *Am. J. Physiol. Ren. Physiol.* **2005**, *289*, F334–F346. [\[CrossRef\]](#)
208. Ronzaud, C.; Loffing-Cueni, D.; Hausel, P.; Debonneville, A.; Malsure, S.R.; Fowler-Jaeger, N.; Boase, N.A.; Perrier, R.; Maillard, M.; Yang, B.; et al. Renal tubular NEDD4-2 deficiency causes NCC-mediated salt-dependent hypertension. *J. Clin. Investig.* **2013**, *2013*, 61110. [\[CrossRef\]](#)
209. Harvey, K.F.; Dinudom, A.; Cook, D.I.; Kumar, S. The Nedd4-like protein KIAA0439 is a potential regulator of the epithelial sodium channel. *J. Biol. Chem.* **2001**, *276*, 8597–8601. [\[CrossRef\]](#)
210. Kumar, S.; Tomooka, Y.; Noda, M. Identification of a set of genes with developmentally down-regulated expression in the mouse brain. *Biochem. Biophys. Res. Commun.* **1992**, *185*, 1155–1161. [\[CrossRef\]](#)
211. Fagerberg, L.; Hallström, B.M.; Oksvold, P.; Kampf, C.; Djureinovic, D.; Odeberg, J.; Habuka, M.; Tahmasebpour, S.; Danielsson, A.; Edlund, K.; et al. Analysis of the Human Tissue-specific Expression by Genome-wide Integration of Transcriptomics and Antibody-based Proteomics. *Mol. Cell. Proteom.* **2014**, *13*, 397–406. [\[CrossRef\]](#)
212. Dumitriu, A.; Latourelle, J.C.; Hadzi, T.C.; Pankratz, N.; Garza, D.; Miller, J.P.; Vance, J.M.; Foroud, T.; Beach, T.G.; Myers, R.H. Gene Expression Profiles in Parkinson Disease Prefrontal Cortex Implicate FOXO1 and Genes under Its Transcriptional Regulation. *PLoS Genet.* **2012**, *8*, e1002794. [\[CrossRef\]](#) [\[PubMed\]](#)
213. Plant, P.J.; Yeger, H.; Staub, O.; Howard, P.; Rotin, D. The C2 Domain of the Ubiquitin Protein Ligase Nedd4 Mediates Ca²⁺-dependent Plasma Membrane Localization. *J. Biol. Chem.* **1997**, *272*, 32329–32336. [\[CrossRef\]](#) [\[PubMed\]](#)
214. Goel, P.; Manning, J.A.; Kumar, S. NEDD4-2 (NEDD4L): The ubiquitin ligase for multiple membrane proteins. *Gene* **2015**, *557*, 1–10. [\[CrossRef\]](#) [\[PubMed\]](#)
215. Donovan, P.; Poronnik, P. Nedd4 and Nedd4-2: Ubiquitin ligases at work in the neuron. *Int. J. Biochem. Cell Biol.* **2013**, *45*, 706–710. [\[CrossRef\]](#) [\[PubMed\]](#)
216. Fotia, A.B.; Cook, D.I.; Kumar, S. The ubiquitin-protein ligases Nedd4 and Nedd4-2 show similar ubiquitin-conjugating enzyme specificities. *Int. J. Biochem. Cell Biol.* **2006**, *38*, 472–479. [\[CrossRef\]](#)
217. De Bie, P.; Ciechanover, A. Ubiquitination of E3 ligases: Self-regulation of the ubiquitin system via proteolytic and non-proteolytic mechanisms. *Cell Death Differ.* **2011**, *18*, 1393–1402. [\[CrossRef\]](#)
218. Mari, S.; Ruetalo, N.; Maspero, E.; Stoffregen, M.C.; Pasqualato, S.; Polo, S.; Wiesner, S. Structural and functional framework for the autoinhibition of Nedd4-family ubiquitin ligases. *Structure* **2014**, *22*, 1639–1649. [\[CrossRef\]](#)
219. Wang, J.; Peng, Q.; Lin, Q.; Childress, C.; Carey, D.; Yang, W. Calcium activates Nedd4 E3 ubiquitin ligases by releasing the C2 domain-mediated auto-inhibition. *J. Biol. Chem.* **2010**, *285*, 12279–12288. [\[CrossRef\]](#)
220. Mund, T.; Pelham, H.R.B. Control of the activity of WW-HECT domain E3 ubiquitin ligases by NDFIP proteins. *EMBO Rep.* **2009**, *10*, 501–507. [\[CrossRef\]](#)
221. Song, L.; Luo, Z.-Q. Post-translational regulation of ubiquitin signaling. *J. Cell Biol.* **2019**, *218*, 1776–1786. [\[CrossRef\]](#)
222. Noyes, N.C.; Hampton, B.; Migliorini, M.; Strickland, D.K. Regulation of Itch and Nedd4 E3 Ligase Activity and Degradation by LRAD3. *Biochemistry* **2016**, *55*, 1204–1213. [\[CrossRef\]](#) [\[PubMed\]](#)
223. Husnjak, K.; Dikic, I. EGFR trafficking: Parkin' in a jam. *Nat. Cell Biol.* **2006**, *8*, 787–788. [\[CrossRef\]](#) [\[PubMed\]](#)
224. Attali, I.; Tobelaim, W.S.; Persaud, A.; Motamedchaboki, K.; Simpson-Lavy, K.J.; Mashahreh, B.; Levin-Kravets, O.; Keren-Kaplan, T.; Pilzer, I.; Kupiec, M.; et al. Ubiquitylation-dependent oligomerization regulates activity of Nedd4 ligases. *EMBO J.* **2017**, *36*, 425–440. [\[CrossRef\]](#) [\[PubMed\]](#)
225. Shao, G.; Wang, R.; Sun, A.; Wei, J.; Peng, K.; Dai, Q.; Yang, W.; Lin, Q. The E3 ubiquitin ligase NEDD4 mediates cell migration signaling of EGFR in lung cancer cells. *Mol. Cancer* **2018**, *17*, 24. [\[CrossRef\]](#)
226. Tu, J.; Zhang, B.; Fang, G.; Chang, W.; Zhao, Y. Neddylation-mediated Nedd4-2 activation regulates ubiquitination modification of renal NBCe1. *Exp. Cell Res.* **2020**, *390*, 111958. [\[CrossRef\]](#)
227. Grimsey, N.J.; Narala, R.; Rada, C.C.; Mehta, S.; Stephens, B.S.; Kufareva, I.; Lapek, J.; Gonzalez, D.J.; Handel, T.M.; Zhang, J.; et al. A Tyrosine Switch on NEDD4-2 E3 Ligase Transmits GPCR Inflammatory Signaling. *Cell Rep.* **2018**, *24*, 3312–3323.e3315. [\[CrossRef\]](#)
228. Debonneville, C.; Flores, S.Y.; Kamynina, E.; Plant, P.J.; Tauxe, C.; Thomas, M.A.; Münster, C.; Chraïbi, A.; Pratt, J.H.; Horisberger, J.-D.; et al. Phosphorylation of Nedd4-2 by Sgk1 regulates epithelial Na⁺ channel cell surface expression. *EMBO J.* **2001**, *20*, 7052–7059. [\[CrossRef\]](#)
229. Snyder, P.M.; Olson, D.R.; Thomas, B.C. Serum and Glucocorticoid-regulated Kinase Modulates Nedd4-2-mediated Inhibition of the Epithelial Na⁺ Channel. *J. Biol. Chem.* **2002**, *277*, 5–8. [\[CrossRef\]](#)
230. Chandran, S.; Li, H.; Dong, W.; Krasinska, K.; Adams, C.; Alexandrova, L.; Chien, A.; Hallows, K.R.; Bhalla, V. Neural Precursor Cell-expressed Developmentally Down-regulated Protein 4-2 (Nedd4-2) Regulation by 14-3-3 Protein Binding at Canonical Serum and Glucocorticoid Kinase 1 (SGK1) Phosphorylation Sites. *J. Biol. Chem.* **2011**, *286*, 37830–37840. [\[CrossRef\]](#)
231. Nagaki, K.; Yamamura, H.; Shimada, S.; Saito, T.; Hisanaga, S.-I.; Taoka, M.; Isobe, T.; Ichimura, T. 14-3-3 Mediates Phosphorylation-Dependent Inhibition of the Interaction between the Ubiquitin E3 Ligase Nedd4-2 and Epithelial Na⁺ Channels. *Biochemistry* **2006**, *45*, 6733–6740. [\[CrossRef\]](#)
232. Zhou, R.; Snyder, P.M. Nedd4-2 phosphorylation induces serum and glucocorticoid-regulated kinase (SGK) ubiquitination and degradation. *J. Biol. Chem.* **2005**, *280*, 4518–4523. [\[CrossRef\]](#) [\[PubMed\]](#)

233. Sato, S.; Chiba, T.; Nishiyama, S.; Kakiuchi, T.; Tsukada, H.; Hatano, T.; Fukuda, T.; Yasoshima, Y.; Kai, N.; Kobayashi, K.; et al. Decline of striatal dopamine release in parkin-deficient mice shown by ex vivo autoradiography. *J. Neurosci. Res.* **2006**, *84*, 1350–1357. [[CrossRef](#)] [[PubMed](#)]
234. Kimura, T.; Kawabe, H.; Jiang, C.; Zhang, W.; Xiang, Y.-Y.; Lu, C.; Salter, M.W.; Brose, N.; Lu, W.-Y.; Rotin, D. Deletion of the ubiquitin ligase Nedd4L in lung epithelia causes cystic fibrosis-like disease. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 3216–3221. [[CrossRef](#)] [[PubMed](#)]
235. Lakso, M.; Pichel, J.G.; Gorman, J.R.; Sauer, B.; Okamoto, Y.; Lee, E.; Alt, F.W.; Westphal, H. Efficient in vivo manipulation of mouse genomic sequences at the zygote stage. *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 5860–5865. [[CrossRef](#)] [[PubMed](#)]
236. Shi, P.P.; Cao, X.R.; Sweezer, E.M.; Kinney, T.S.; Williams, N.R.; Husted, R.F.; Nair, R.; Weiss, R.M.; Williamson, R.A.; Sigmund, C.D.; et al. Salt-sensitive hypertension and cardiac hypertrophy in mice deficient in the ubiquitin ligase Nedd4-2. *Am. J. Physiol. Ren. Physiol.* **2008**, *295*, F462–F470. [[CrossRef](#)]
237. Dinudom, A.; Fotia, A.B.; Lefkowitz, R.J.; Young, J.A.; Kumar, S.; Cook, D.I. The kinase Grk2 regulates Nedd4/Nedd4-2-dependent control of epithelial Na⁺ channels. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 11886–11890. [[CrossRef](#)]
238. Pronin, A.N.; Morris, A.J.; Surguchov, A.; Benovic, J.L. Synucleins are a novel class of substrates for G protein-coupled receptor kinases. *J. Biol. Chem.* **2000**, *275*, 26515–26522. [[CrossRef](#)]
239. Howitt, J.; Gysbers, A.M.; Ayton, S.; Carew-Jones, F.; Putz, U.; Finkelstein, D.I.; Halliday, G.M.; Tan, S.-S. Increased Ndfip1 in the Substantia Nigra of Parkinsonian Brains Is Associated with Elevated Iron Levels. *PLoS ONE* **2014**, *9*, e87119. [[CrossRef](#)]
240. Guo, M.; Wang, J.; Zhao, Y.; Feng, Y.; Han, S.; Dong, Q.; Cui, M.; Tieu, K. Microglial exosomes facilitate α -synuclein transmission in Parkinson's disease. *Brain* **2020**, *143*, 1476–1497. [[CrossRef](#)]
241. Oliver, P.M.; Cao, X.; Worthen, G.S.; Shi, P.; Briones, N.; Macleod, M.; White, J.; Kirby, P.; Kappler, J.; Marrack, P.; et al. Ndfip1 Protein Promotes the Function of Itch Ubiquitin Ligase to Prevent T Cell Activation and T Helper 2 Cell-Mediated Inflammation. *Immunity* **2006**, *25*, 929–940. [[CrossRef](#)]
242. O'Leary, C.E.; Riling, C.R.; Spruce, L.A.; Ding, H.; Kumar, S.; Deng, G.; Liu, Y.; Seeholzer, S.H.; Oliver, P.M. Ndfip-mediated degradation of Jak1 tunes cytokine signalling to limit expansion of CD4⁺ effector T cells. *Nat. Commun.* **2016**, *7*, 11226. [[CrossRef](#)] [[PubMed](#)]
243. Behrends, C.; Sowa, M.E.; Gygi, S.P.; Harper, J.W. Network organization of the human autophagy system. *Nature* **2010**, *466*, 68–76. [[CrossRef](#)] [[PubMed](#)]
244. Wang, H.; Sun, R.Q.; Camera, D.; Zeng, X.Y.; Jo, E.; Chan, S.M.H.; Herbert, T.P.; Molero, J.C.; Ye, J.M. Endoplasmic reticulum stress up-regulates Nedd4-2 to induce autophagy. *FASEB J.* **2016**, *30*, 2549–2556. [[CrossRef](#)] [[PubMed](#)]
245. Conway, J.A.; Kramer, E.R. Is activation of GDNF/RET signaling the answer for successful treatment of Parkinson's disease? A discussion of data from the culture dish to the clinic. *Neural Regen. Res.* **2022**, *17*, 1462. [[CrossRef](#)]
246. Meng, B.; Lever, A.M.L. The Interplay between ESCRT and Viral Factors in the Enveloped Virus Life Cycle. *Viruses* **2021**, *13*, 324. [[CrossRef](#)]
247. Malakhova, O.A.; Zhang, D.-E. ISG15 Inhibits Nedd4 Ubiquitin E3 Activity and Enhances the Innate Antiviral Response*. *J. Biol. Chem.* **2008**, *283*, 8783–8787. [[CrossRef](#)]
248. Okumura, A.; Pitha, P.M.; Harty, R.N. ISG15 inhibits Ebola VP40 VLP budding in an L-domain-dependent manner by blocking Nedd4 ligase activity. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 3974–3979. [[CrossRef](#)]
249. Riederer, P.; Berg, D.; Casadei, N.; Cheng, F.; Classen, J.; Dresel, C.; Jost, W.; Krüger, R.; Müller, T.; Reichmann, H.; et al. α -Synuclein in Parkinson's disease: Causal or bystander? *J. Neural Transm.* **2019**, *126*, 815–840. [[CrossRef](#)]
250. Kramer, E.R.; Liss, B. GDNF-Ret signaling in midbrain dopaminergic neurons and its implication for Parkinson disease. *FEBS Lett.* **2015**, *589*, 3760–3772. [[CrossRef](#)]
251. Plaza-Menacho, I.; Barnouin, K.; Barry, R.; Borg, A.; Orme, M.; Chauhan, R.; Mouilleron, S.; Martínez-Torres, R.J.; Meier, P.; McDonald, N.Q. RET Functions as a Dual-Specificity Kinase that Requires Allosteric Inputs from Juxtamembrane Elements. *Cell Rep.* **2016**, *17*, 3319–3332. [[CrossRef](#)]
252. Conway, J.A.; Ince, S.; Black, S.; Kramer, E.R. GDNF/RET signaling in dopamine neurons in vivo. *Cell Tissue Res.* **2020**, *382*, 135–146. [[CrossRef](#)] [[PubMed](#)]
253. Kramer, E.R.; Aron, L.; Ramakers, G.M.J.; Seitz, S.; Zhuang, X.; Beyer, K.; Smidt, M.P.; Klein, R. Absence of Ret Signaling in Mice Causes Progressive and Late Degeneration of the Nigrostriatal System. *PLoS Biol.* **2007**, *5*, e39. [[CrossRef](#)] [[PubMed](#)]
254. Kowsky, S.; Pöppelmeier, C.; Kramer, E.R.; Falkenburger, B.H.; Kruse, A.; Klein, R.; Schulz, J.B. RET signaling does not modulate MPTP toxicity but is required for regeneration of dopaminergic axon terminals. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 20049–20054. [[CrossRef](#)] [[PubMed](#)]
255. Drinkut, A.; Tillack, K.; Meka, D.P.; Schulz, J.B.; Kügler, S.; Kramer, E.R. Ret is essential to mediate GDNF's neuroprotective and neuroregenerative effect in a Parkinson disease mouse model. *Cell Death Dis.* **2016**, *7*, e2359. [[CrossRef](#)] [[PubMed](#)]
256. Haglund, K.; Dikic, I. The role of ubiquitylation in receptor endocytosis and endosomal sorting. *J. Cell Sci.* **2012**, *125*, 265–275. [[CrossRef](#)] [[PubMed](#)]
257. Crupi, M.J.; Yoganathan, P.; Bone, L.N.; Lian, E.; Fetz, A.; Antonescu, C.N.; Mulligan, L.M. Distinct Temporal Regulation of RET Isoform Internalization: Roles of Clathrin and AP2. *Traffic* **2015**, *16*, 1155–1173. [[CrossRef](#)] [[PubMed](#)]
258. Richardson, D.S.; Rodrigues, D.M.; Hyndman, B.D.; Crupi, M.J.; Nicolescu, A.C.; Mulligan, L.M. Alternative splicing results in RET isoforms with distinct trafficking properties. *Mol. Biol. Cell* **2012**, *23*, 3838–3850. [[CrossRef](#)]

259. Grice, G.L.; Nathan, J.A. The recognition of ubiquitinated proteins by the proteasome. *Cell Mol. Life Sci.* **2016**, *73*, 3497–3506. [\[CrossRef\]](#)
260. Vecchione, A.; Marchese, A.; Henry, P.; Rotin, D.; Morrione, A. The Grb10/Nedd4 complex regulates ligand-induced ubiquitination and stability of the insulin-like growth factor I receptor. *Mol. Cell Biol.* **2003**, *23*, 3363–3372. [\[CrossRef\]](#)
261. Kwak, Y.D.; Wang, B.; Li, J.J.; Wang, R.; Deng, Q.; Diao, S.; Chen, Y.; Xu, R.; Masliah, E.; Xu, H.; et al. Upregulation of the E3 ligase NEDD4-1 by Oxidative Stress Degrades IGF-1 Receptor Protein in Neurodegeneration. *J. Neurosci.* **2012**, *32*, 10971–10981. [\[CrossRef\]](#)
262. Cao, X.R.; Lill, N.L.; Boase, N.; Shi, P.P.; Croucher, D.R.; Shan, H.; Qu, J.; Sweezer, E.M.; Place, T.; Kirby, P.A.; et al. Nedd4 controls animal growth by regulating IGF-1 signaling. *Sci. Signal.* **2008**, *1*, ra5. [\[CrossRef\]](#) [\[PubMed\]](#)
263. Giorgetti-Peraldi, S.; Murdaca, J.; Mas, J.C.; Van Obberghen, E. The adapter protein, Grb10, is a positive regulator of vascular endothelial growth factor signaling. *Oncogene* **2001**, *20*, 3959–3968. [\[CrossRef\]](#) [\[PubMed\]](#)
264. Hurley, J.H. Nipped in the bud: How the AMSH MIT domain helps deubiquitinate lysosome-bound cargo. *Structure* **2011**, *19*, 1033–1035. [\[CrossRef\]](#) [\[PubMed\]](#)
265. Shields, S.B.; Piper, R.C. How ubiquitin functions with ESCRTs. *Traffic* **2011**, *12*, 1306–1317. [\[CrossRef\]](#)
266. Roxrud, I.; Raiborg, C.; Pedersen, N.M.; Stang, E.; Stenmark, H. An endosomally localized isoform of Eps15 interacts with Hrs to mediate degradation of epidermal growth factor receptor. *J. Cell Biol.* **2008**, *180*, 1205–1218. [\[CrossRef\]](#)
267. Hasegawa, T.; Konno, M.; Baba, T.; Sugeno, N.; Kikuchi, A.; Kobayashi, M.; Miura, E.; Tanaka, N.; Tamai, K.; Furukawa, K.; et al. The AAA-ATPase VPS4 regulates extracellular secretion and lysosomal targeting of α -synuclein. *PLoS ONE* **2011**, *6*, e29460. [\[CrossRef\]](#)
268. Tanikawa, S.; Mori, F.; Tanji, K.; Kakita, A.; Takahashi, H.; Wakabayashi, K. Endosomal sorting related protein CHMP2B is localized in Lewy bodies and glial cytoplasmic inclusions in α -synucleinopathy. *Neurosci. Lett.* **2012**, *527*, 16–21. [\[CrossRef\]](#)
269. Kurashige, T.; Takahashi, T.; Yamazaki, Y.; Hiji, M.; Izumi, Y.; Yamawaki, T.; Matsumoto, M. Localization of CHMP2B-immunoreactivity in the brainstem of Lewy body disease. *Neuropathology* **2013**, *33*, 237–245. [\[CrossRef\]](#)
270. Jing, W.; Wang, G.; Cui, Z.; Xiong, G.; Jiang, X.; Li, Y.; Li, W.; Han, B.; Chen, S.; Shi, B. FGFR3 Destabilizes PD-L1 via NEDD4 to Control T-cell-Mediated Bladder Cancer Immune Surveillance. *Cancer Res.* **2022**, *82*, 114–129. [\[CrossRef\]](#)
271. Yu, T.; Calvo, L.; Anta, B.; López-Benito, S.; López-Bellido, R.; Vicente-García, C.; Tessarollo, L.; Rodríguez, R.E.; Arévalo, J.C. In vivo regulation of NGF-mediated functions by Nedd4-2 ubiquitination of TrkA. *J. Neurosci.* **2014**, *34*, 6098–6106. [\[CrossRef\]](#)
272. Yanpallear, S.; Wang, T.; Koh, D.C.; Quarta, E.; Fulgenzi, G.; Tessarollo, L. Nedd4-2 haploinsufficiency causes hyperactivity and increased sensitivity to inflammatory stimuli. *Sci. Rep.* **2016**, *6*, 32957. [\[CrossRef\]](#) [\[PubMed\]](#)
273. Melchior, B.; Nerrière-Daguin, V.; Laplaud, D.-A.; Remy, S.; Wiertlewski, S.; Neveu, I.; Naveilhan, P.; Meakin, S.; Brachet, P. Ectopic expression of the TrkA receptor in adult dopaminergic mesencephalic neurons promotes retrograde axonal NGF transport and NGF-dependent neuroprotection. *Exp. Neurol.* **2003**, *183*, 367–378. [\[CrossRef\]](#)
274. Ding, Y.-X.; Xia, Y.; Jiao, X.-Y.; Duan, L.; Yu, J.; Wang, X.; Chen, L.-W. The TrkB-Positive Dopaminergic Neurons are Less Sensitive to MPTP Insult in the Substantia Nigra of Adult C57/BL Mice. *Neurochem. Res.* **2011**, *36*, 1759–1766. [\[CrossRef\]](#) [\[PubMed\]](#)
275. Heldin, C.H.; Moustakas, A. Signaling Receptors for TGF- β Family Members. *Cold Spring Harb. Perspect. Biol.* **2016**, *8*, 22053. [\[CrossRef\]](#)
276. Vander Ark, A.; Cao, J.; Li, X. TGF- β receptors: In and beyond TGF- β signaling. *Cell. Signal.* **2018**, *52*, 112–120. [\[CrossRef\]](#)
277. Luo, S.X.; Huang, E.J. Dopaminergic Neurons and Brain Reward Pathways: From Neurogenesis to Circuit Assembly. *Am. J. Pathol.* **2016**, *186*, 478–488. [\[CrossRef\]](#)
278. Sorkina, T.; Miranda, M.; Dionne, K.R.; Hoover, B.R.; Zahniser, N.R.; Sorkin, A. RNA Interference Screen Reveals an Essential Role of Nedd4-2 in Dopamine Transporter Ubiquitination and Endocytosis. *J. Neurosci.* **2006**, *26*, 8195–8205. [\[CrossRef\]](#)
279. Efimova, E.V.; Gainetdinov, R.R.; Budygin, E.A.; Sotnikova, T.D. Dopamine transporter mutant animals: A translational perspective. *J. Neurogenet.* **2016**, *30*, 5–15. [\[CrossRef\]](#)
280. García-Tardón, N.; González-González, I.M.; Martínez-Villarreal, J.; Fernández-Sánchez, E.; Giménez, C.; Zafra, F. Protein Kinase C (PKC)-promoted Endocytosis of Glutamate Transporter GLT-1 Requires Ubiquitin Ligase Nedd4-2-dependent Ubiquitination but Not Phosphorylation. *J. Biol. Chem.* **2012**, *287*, 19177–19187. [\[CrossRef\]](#)
281. Yu, Z.; Liu, C.; Zhang, J.; Liang, Z.; You, G. Protein kinase C regulates organic anion transporter 1 through phosphorylating ubiquitin ligase Nedd4-2. *BMC Mol. Cell Biol.* **2021**, *22*, 53. [\[CrossRef\]](#)
282. Zhang, J.; Liu, C.; You, G. AG490, a JAK2-specific inhibitor, downregulates the expression and activity of organic anion transporter-3. *J. Pharm. Sci.* **2018**, *136*, 142–148. [\[CrossRef\]](#) [\[PubMed\]](#)
283. Malagelada, C.; Ryu, E.J.; Biswas, S.C.; Jackson-Lewis, V.; Greene, L.A. RTP801 Is Elevated in Parkinson Brain Substantia Nigral Neurons and Mediates Death in Cellular Models of Parkinson's Disease by a Mechanism Involving Mammalian Target of Rapamycin Inactivation. *J. Neurosci.* **2006**, *26*, 9996–10005. [\[CrossRef\]](#) [\[PubMed\]](#)
284. Canal, M.; Romani-Aumedes, J.; Martín-Flores, N.; Pérez-Fernández, V.; Malagelada, C. RTP801/REDD1: A stress coping regulator that turns into a troublemaker in neurodegenerative disorders. *Front. Cell Neurosci.* **2014**, *8*, 313. [\[CrossRef\]](#) [\[PubMed\]](#)
285. Grace, A.; Bunney, B. The control of firing pattern in nigral dopamine neurons: Single spike firing. *J. Neurosci.* **1984**, *4*, 2866–2876. [\[CrossRef\]](#) [\[PubMed\]](#)
286. Grace, A.; Bunney, B. The control of firing pattern in nigral dopamine neurons: Burst firing. *J. Neurosci.* **1984**, *4*, 2877–2890. [\[CrossRef\]](#)

-
287. Lammel, S.; Hetzel, A.; Häckel, O.; Jones, I.; Liss, B.; Roeper, J. Unique Properties of Mesoprefrontal Neurons within a Dual Mesocorticolimbic Dopamine System. *Neuron* **2008**, *57*, 760–773. [[CrossRef](#)]
288. Margolis, E.B.; Coker, A.R.; Driscoll, J.R.; Lemaître, A.-I.; Fields, H.L. Reliability in the Identification of Midbrain Dopamine Neurons. *PLoS ONE* **2010**, *5*, e15222. [[CrossRef](#)]
289. Drion, G.; Massotte, L.; Sepulchre, R.; Seutin, V. How Modeling Can Reconcile Apparently Discrepant Experimental Results: The Case of Pacemaking in Dopaminergic Neurons. *PLoS Comput. Biol.* **2011**, *7*, e1002050. [[CrossRef](#)]
290. Branch, S.Y.; Sharma, R.; Beckstead, M.J. Aging decreases L-type calcium channel currents and pacemaker firing fidelity in substantia nigra dopamine neurons. *J. Neurosci.* **2014**, *34*, 9310–9318. [[CrossRef](#)]
291. Rice, M.E.; Patel, J.C.; Cragg, S.J. Dopamine release in the basal ganglia. *Neuroscience* **2011**, *198*, 112–137. [[CrossRef](#)]
292. Van Wart, A.; Matthews, G. Impaired Firing and Cell-Specific Compensation in Neurons Lacking Nav1.6 Sodium Channels. *J. Neurosci.* **2006**, *26*, 7172–7180. [[CrossRef](#)] [[PubMed](#)]
293. Wang, Z.; Lin, Y.; Liu, W.; Kuang, P.; Lao, W.; Ji, Y.; Zhu, H. Voltage-Gated Sodium Channels Are Involved in Cognitive Impairments in Parkinson's Disease-like Rats. *Neuroscience* **2019**, *418*, 231–243. [[CrossRef](#)] [[PubMed](#)]
294. Hansen, H.H.; Ebbesen, C.; Mathiesen, C.; Weikop, P.; Rønn, L.C.; Waroux, O.; Scuvée-Moreau, J.; Seutin, V.; Mikkelsen, J.D. The KCNQ channel opener retigabine inhibits the activity of mesencephalic dopaminergic systems of the rat. *J. Pharm. Exp.* **2006**, *318*, 1006–1019. [[CrossRef](#)] [[PubMed](#)]
295. Cooper, E.C.; Harrington, E.; Jan, Y.N.; Jan, L.Y. M channel KCNQ2 subunits are localized to key sites for control of neuronal network oscillations and synchronization in mouse brain. *J. Neurosci.* **2001**, *21*, 9529–9540. [[CrossRef](#)] [[PubMed](#)]
296. Weber, Y.G.; Geiger, J.; Kämpchen, K.; Landwehrmeyer, B.; Sommer, C.; Lerche, H. Immunohistochemical analysis of KCNQ2 potassium channels in adult and developing mouse brain. *Brain Res.* **2006**, *1077*, 1–6. [[CrossRef](#)] [[PubMed](#)]
297. Martire, M.; D'Amico, M.; Panza, E.; Miceli, F.; Viggiano, D.; Lavergata, F.; Iannotti, F.A.; Barrese, V.; Preziosi, P.; Annunziato, L.; et al. Involvement of KCNQ2 subunits in [3H]dopamine release triggered by depolarization and pre-synaptic muscarinic receptor activation from rat striatal synaptosomes. *J. Neurochem.* **2007**, *102*, 179–193. [[CrossRef](#)]