

2022-07-08

Influenza vaccination reduced myocardial infarctions in UK older adults: a prior event rate ratio study

Streeter, Adam

<http://hdl.handle.net/10026.1/19409>

10.1016/j.jclinepi.2022.06.018

Journal of Clinical Epidemiology

Elsevier BV

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

IPTW diagnostic plots




Figure 1: Plots of weighted (by propensity scores in red) and unweighted standardised (blue) mean differences for balancing variables – 1997 - 1999 cohorts




Figure 2: Plots of weighted (by propensity scores in red) and unweighted standardised (blue) mean differences for balancing variables – 2000 - 2002 cohorts




Figure 3: Plots of weighted (by propensity scores in red) and unweighted standardised (blue) mean differences for balancing variables – 2003 - 2005 cohorts




Figure 4: Plots of weighted (by propensity scores in red) and unweighted standardised (blue) mean differences for balancing variables – 2006 - 2008 cohorts




Figure 5: Plots of weighted (by propensity scores in red) and unweighted standardised (blue) mean differences for balancing variables – 2009 - 2011 cohorts