

2022-09-01

Filling gaps via citizen science: Phyllorhiza punctata von Lendenfeld, 1884 (Cnidaria: Scyphozoa: Mastigiidae) in Cyprus (eastern Mediterranean Sea)

Kaminas, A

<http://hdl.handle.net/10026.1/19806>

10.3391/bir.2022.11.3.09

BiolInvasions Records

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Rapid Communication

Filling gaps via citizen science: *Phyllorhiza punctata* von Lendenfeld, 1884 (Cnidaria: Scyphozoa: Mastigiidae) in Cyprus (eastern Mediterranean Sea)

Alexandros Kaminas¹, Maria Shokouros-Oskarsson², Vasileios Minasidis^{1,3}, Joachim Langeneck⁴, Periklis Kleitou^{1,5}, Francesco Tiralongo^{6,7} and Fabio Crocetta^{8,*}

¹Sea, Environmental Organisation for the Preservation of the Aquatic Ecosystems, 54645 Thessaloniki, Greece

²Independent Researcher, Agiou Mama 91, 5295 Paralimni, Cyprus

³Department of Animal Production, Fisheries and Aquaculture, University of Patras, 30200 Mesolongi, Greece

⁴Dipartimento di Biologia, Università di Pisa, via Derna 1, I-56126 Pisa, Italy

⁵Marine & Environmental Research (MER) Lab., 202 Amathountos Avenue, Marina Gardens, Block B, Limassol 4533, Cyprus

⁶Department of Biological, Geological and Environmental Sciences, University of Catania, 95124 Catania, Italy

⁷Ente Fauna Marina Mediterranea, Scientific Organization for Research and Conservation of Marine Biodiversity, 96012 Avola, Italy

⁸Department of Integrative Marine Ecology, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Naples, Italy

*Corresponding author

E-mail: fabio.crocetta@szn.it

Citation: Kaminas A, Shokouros-Oskarsson M, Minasidis V, Langeneck J, Kleitou P, Tiralongo F, Crocetta F (2022) Filling gaps via citizen science: *Phyllorhiza punctata* von Lendenfeld, 1884 (Cnidaria: Scyphozoa: Mastigiidae) in Cyprus (eastern Mediterranean Sea). *BioInvasions Records* 11(3): 667–675, <https://doi.org/10.3391/bir.2022.11.3.09>

Received: 27 February 2022

Accepted: 7 April 2022

Published: 11 May 2022

Handling editor: Melih Ertan Çımar

Thematic editor: April Blakeslee

Copyright: © Kaminas et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International - CC BY 4.0).

OPEN ACCESS

Abstract

The Mediterranean Sea is one of the most heavily impacted areas by biological invasions worldwide, with over 650 non-indigenous species (NIS) reported as established in 2019. Information about the distribution of NIS in the basin is often fragmented or non-existent due to factors such as lack of finance, expertise, and appropriate surveillance systems. Citizen science might represent a reliable tool in monitoring the spread of NIS, being able to overcome the challenges of scientific monitoring. In the context of the citizen science project “Is it alien to you? Share it!!!”, we hereby first report the presence of the Australian spotted jellyfish *Phyllorhiza punctata* von Lendenfeld, 1884 in the marine waters of Cyprus and review its known distribution in the Mediterranean Sea. Although this taxon does not represent a direct threat to human health or tourism as it bears no painful sting, it already caused important ecological and economic damages in other regions of the world, and thus its further spread in Cypriot waters requires appropriate monitoring.

Key words: biological invasions, jellyfishes, Levant Sea, non-indigenous species (NIS), invasive species

Introduction

The Mediterranean Sea is one of the most heavily impacted areas by non-indigenous species (NIS) worldwide due to a combination of cumulative human stressors which include heavy maritime traffic and strong connectivity with other ecoregions through sea corridors (Coll et al. 2012; Katsanevakis et al. 2013). More than 650 NIS species were reported as established in the Mediterranean by 2019 (Zenetos and Galanidi 2020), including gelatinous planktonic species that often proliferate and create blooms that lead to substantial socioeconomic damage (Purcell et al. 2007;

Boero et al. 2009; Boero 2013). Among them, the cnidarian *Rhopilema nomadica* Galil, Spanier & Ferguson, 1990 and the ctenophore *Mnemiopsis leidyi* A. Agassiz, 1865 are causing the most harm in the Mediterranean Sea, with the former interacting with coastal trawling and purse-seine fishing in the southern and eastern coasts of the basin, and the latter threatening commercially important species such as the anchovy *Engraulis encrasicolus* (Linnaeus, 1758) in the Adriatic and the Black Sea (Knowler 2005; Purcell et al. 2007; Boero et al. 2009; Boero 2013; Malej et al. 2017). Notwithstanding that, little is known about other NIS jellyfishes, or they are still known based on scattered records despite possessing bright and distinctive colors, large sizes, or inhabiting the upper levels of the water column, or even a combination of all these features (e.g. Galil et al. 2010; Mamish et al. 2016; Langeneck et al. 2019; Douek et al. 2020; Mutlu et al. 2020; Badreddine and Bitar 2020).

The Australian spotted jellyfish *Phyllorhiza punctata* von Lendenfeld, 1884 is also encased in this statement. Considered as native from the western Pacific Ocean (Heeger et al. 1992; Graham et al. 2003), this species tolerates a broad range of salinity and temperature and can live in turbid waters around estuaries, lagoons, as well as in embayments and heavily populated areas (Rippingale and Kelly 1995). This taxon is characterized by a roughly semi-spherical umbrella, with an average diameter of 45–50 cm in adult individuals and eight broad and transparent branching mouth arms that culminate in large brown bundles of stinging cells, accentuated by white crystalline inclusions that appear as white spots (Hawaiian Biological Survey 2001). These conspicuous morphological characteristics make the species an apparent and easy target for visual detection, including by citizen scientists and non-specialists (Haddad and Nogueira Júnior 2006). However, while *P. punctata* is widely recorded as a NIS in the eastern Pacific and the western Atlantic, where its spread was presumably facilitated by ships through ballast water or fouling (review in Rosales-Catalán et al. 2021), it is still scarcely recorded in the Mediterranean Sea. In fact, although early records of *P. punctata* in the basin date back to 1965, when a single individual was found off Israel (Galil et al. 1990), and the species had already reached the western Mediterranean Sea (Boero et al. 2009; Gueroun et al. 2015), its presence in several Mediterranean countries (e.g. Lebanon, Libya, Syria, and Italy) is based so far on single or very few recent records, which are themselves often based on single specimens (Boero et al. 2009; Durgham 2011; Crocetta and Bariche in Dailianis et al. 2016; Deidun et al. 2017; Rizgalla and Crocetta 2020).

Within the framework of a citizen science project, we hereby first report the presence of *P. punctata* in Cyprus, thus filling a gap in the knowledge of its spread in the Mediterranean Sea, and review its known distribution in the basin.

Materials and methods

In 2016, the Greek environmental organization iSea launched an online project named “Is it Alien to you? Share it!!!” where citizen scientists could share sightings of marine organisms unknown to them. Citizens can submit their sightings with photographic material, videos, and other forms of observation, and can complement their records with auxiliary information, such as specimen size, number of specimens, depth of the sighting, location, date, as well as observation type (freediving, underwater photography, shore-based fishing, boat-based fishing, spearfishing, etc.). Following examination by experts, sightings are identified and assessed based on their validity and reliability. In the recent years, the project has further expanded to include a Google Form and a Facebook® group, with the aim of facilitating the data uploading process, and included several taxonomic experts on marine NIS that regularly provide their expertise in identifying the submitted observations (see Giovos et al. 2019 for further details).

When a sighting of potentially major scientific interest is submitted to the platform, a systematic literature review of records published in indexed and grey literature is also carried out through Google Scholar using appropriate keywords based on the species observed.

Results

Since its launch, the “Is it Alien to you? Share it!!!” project received 121 sightings related to native or NIS jellyfish. Specifically, scyphozoan taxa account for 4.23% of the total records, with numbers increasing during the last four years (less than 1% at the end of 2018: see Langeneck et al. 2019) and NIS that comprise ~ 18.2% of the sightings (Table 1). Among them, one pertained a specimen of a spotted jellyfish (Figure 1), found on the 6th of February 2022 by one of the authors (M.S.-O.) during a recreational dive activity. The animal was found drifting at a depth of around 6 m in the Bay of Da Costa (35.008288N; 34.061619E), Protaras, Cyprus. Ambient seawater temperature was 16 °C. No jellyfish aggregations or even single individuals were recorded in the immediate vicinity. The specimen was soon identified by the group experts as *P. punctata* based on its distinctive bell shape and coloration.

The bibliographic review conducted here yielded 20 records from the entire Mediterranean Sea. Among them, a record from Spain was originally reported in newspapers only (see Anonymous 2011), but then included in the scientific literature (Brotz and Pauly 2012; Galil 2012). As no confirmation was ever published by local researchers, and the newspapers never presented images of the alleged outbreak and of the species, we list it with caution as it may result to be incorrect. Results of the bibliographic review are summarized in Table 2 and depicted in a GIS-generated map in Figure 2.

Table 1. Records of native and non-indigenous Scyphozoan species collected in the present project, with: (i) species binomial name; (ii) N – cumulative number of records in the “Is it Alien to you? Share it!!!” project database; (iii) species status (N – native or NIS – non-indigenous species).

| Species | N | Status |
|---|----|--------|
| Family Cassiopeidae Tilesius, 1831 | | |
| <i>Cassiopea andromeda</i> (Forskål, 1775) | 6 | NIS |
| Family Cepheidae Agassiz, 1862 | | |
| <i>Cotylorhiza tuberculata</i> (Macri, 1778) | 24 | N |
| Family Rhizostomatidae Cuvier, 1800 | | |
| <i>Rhizostoma pulmo</i> (Macri, 1778) | 13 | N |
| <i>Rhopilema nomadica</i> Galil, Spanier & Ferguson, 1990 | 16 | NIS |
| Family Ulmaridae Haeckel, 1880 | | |
| <i>Aurelia</i> sp./spp. | 6 | N |
| Family Pelagiidae Gegenbaur, 1856 | | |
| <i>Chrysaora</i> cf. <i>achlyos</i> Martin et al. 1997 | 1 | NIS |
| <i>Chrysaora hysoscella</i> (Linnaeus, 1767) | 11 | N |
| <i>Pelagia noctiluca</i> (Forsskål, 1775) | 44 | N |



Figure 1. *Phyllorhiza punctata* from Da Costa Bay (Cyprus, eastern Mediterranean Sea). Photo Maria Shokouros-Oskarsson.

Discussion

Citizen science is expanding rapidly, as scientists, policymakers, and conservationists widely recognize citizens’ contributions in recording, analyzing, and responding species’ invasions worldwide (Thiel et al. 2014; Scyphers et al. 2015). In this view, the case of *P. punctata* in the Mediterranean Sea is also notable, as citizen science already proved to be crucial in tracking its spread in the Mediterranean Sea, and records after 2009 were mostly obtained through social media data mining, interviews with fishermen, and above all generalist citizen science projects or even specific campaigns focused on gelatinous species (Boero et al. 2009; Crocetta and Bariche in Dailianis et al. 2016; Deidun et al. 2017; Rabaoui and El Zrelli in Stamouli et al. 2017; Rizgalla and Crocetta 2020).

Table 2. Known records of *Phyllorhiza punctata* in the Mediterranean Sea arranged in temporal order, with: (i) country, locality, coordinates (latitude and longitude), and Marine Strategy Framework Directive (MSFD) subregion; (ii) sighting date (month and year, when known); (iii) number of individuals (N_IND); (iv) method of observation; (v) reference(s). Numbers (N) correspond to sampling locations shown in Figure 2. Abbreviations used: (i) MSFD: CM – central Mediterranean; EM – eastern Mediterranean; WM – western Mediterranean; (ii) Method: SI – scientific investigation; SS – scientific sampling; MC – museum collection; CS – citizen-science; SM – social media data mining; UNK – unknown. Note: * this outbreak is doubtful; moreover, it was incorrectly reported in the scientific literature as happened in 2010 instead of 2011 [Anonymous (2011); see also references in Brotz and Pauly (2012)].

| N | Country | Locality | Coordinates | MSFD | Date | N_IND | Method | Reference |
|----|---------|---|-----------------------------|------|------------------------------|----------|--------------------------|---|
| 1 | Israel | Beit Yanai | ~ 32.382379N, 34.860442E | EM | 1965 | 1 | SI (MC) | Galil et al. (1990) |
| 2 | Egypt | El-Arish | ~ 31.281151N, 33.865823E | EM | VII.1986 | 3 | stranded | Atta (1991) |
| 3 | Israel | Shikmona | 32.821626N, 34.950020E | EM | I.2005 | 1 | stranded (MC) | Galil et al. (2009) |
| 4 | Greece | Bay of Vlyho (Lefkada Island) | 38.684444N, 20.706389E | CM | IX.2005 and IX.2006 | 49 | SS | Abed-Navandi and Kikinger (2007) |
| 5 | Israel | Kiryat Yam (Haifa Bay) | 32.836109N, 34.627411E | EM | IX.2006 | 2 | UNK (MC) | Galil et al. (2009) |
| 6 | Greece | Igoumenitsa harbour | ~ 39.498769N, 20.261977E | CM | IX.2006 | > 50 | SS | Abed-Navandi and Kikinger (2007) |
| 7 | Israel | Ashdod | ~ 31.834615N, 34.541251E | EM | VI.2009 | 12 | UNK (bottom trawling) | Galil et al. (2009) |
| 8 | Italy | Cala Suaraccia (Sardinia Island) | 40.845110N, 9.707275E | WM | X.2009 | 1 | CS campaign | Boero et al. (2009) |
| 9 | Turkey | İskenderun Bay | 36.742500N, 36.178600E | EM | X.2010 | 1 | SS | Cevik et al. (2011) |
| 10 | Italy | Mar Piccolo (Gulf of Taranto) | 40.481395N, 17.235006E | CM | VII.2011 | 1 | UNK | Deidun et al. (2017) |
| 11 | *Spain | Orihuela (Alicante) | ~ 37.932455N, 0.709471W | WM | VII.2011 | outbreak | UNK | Anonymous (2011); Brotz and Pauly (2012); Galil (2012) |
| 12 | Syria | Lattakia | 35.524872N, 35.551716E | EM | VIII.2011 | 1 | SS | Durgham (2011) |
| 13 | Turkey | Sülüngür Lake | ~ 36.783950N, 28.652441E | EM | IX.2011 | 10 | SS | Gülşahin and Tarkan (2012) |
| 14 | Israel | from Caesarea to Nanariya | ~ 32.790511N, 34.866857E | EM | 2012–2017 | UNK | UNK | Mizrahi et al. (2021) |
| 15 | Tunisia | Bizerte Lagoon | ~ 37.194574N, 9.858965E | WM | VIII.2012 (to X.2013) | 1434 | SS | Gueroun et al. (2015) |
| 16 | Tunisia | between El Hicha and Zarrat (Gulf of Gabès) | ~ 34.149522N, 10.040956E | CM | since 2012 (to VIII.2017) | common | SS/interviews | Rabaoui and El Zrelli in Stamouli et al. (2017) |
| 17 | Libya | Al Dafiniyah | 32.443833N, 14.884028E | CM | I.2015 | > 1 | SM | Rizgalla and Crocetta (2020) |
| 18 | Lebanon | Sarafand | 33.439316N, 35.200751E | EM | X.2015 | 1 | SM | Crocetta and Bariche in Dailianis et al. (2016) |
| 19 | Malta | Salini salt pans (Malta Island) | 35.948328N, 14.425425E | CM | X–XI.2016 | > 50 | SS/CS project | Deidun et al. (2017) |
| | | St. Paul's Bay (Malta Island) | 35.949951N, 14.400052E | CM | XI.2016 | 1 | CS project | |
| | | Qawra Point (Malta Island) | 35.958673N, 14.426643E | CM | XI.2016 | 1 | CS project | |
| 20 | Egypt | Port Said harbour | 31.45N, 32.5E | EM | XI.2018 | 4 | stranded | Madkour et al. (2021) |
| 21 | Cyprus | Bay of DaCosta | 35.008288N, 34.061619E | EM | II.2022 | 1 | CS project | present paper |

The present record from the Bay of Da Costa constituted the first record of the species from Cyprus and thus fills a gap at a country level in the known invaded range of *P. punctata* in the Mediterranean. On one hand, it serves to reaffirm the importance of scientific data obtained through citizen science, but also raises questions regarding its possible pathway(s) of arrival. Galil et al. (2009) reported the existence of sexually mature specimens throughout the entire year, suggesting that established populations persist at least in certain regions of the Mediterranean Sea. This is in agreement with the high number of specimens reported in Greece and in




Figure 2. The known distribution of *Phyllorhiza punctata* in the Mediterranean Sea. Numbers within square brackets refer to references listed in Table 2. Red star refers to the present record. Note: the record [1] from Spain is doubtful.

Tunisia (Abed-Navandi and Kikinger 2007; Gueroun et al. 2015; Rabaoui and El Zrelli in Stamouli et al. 2017). Moreover, as the species is characterized by possessing an asexually reproducing benthic polyp and a sexually reproducing pelagic medusa, it may easily remain in the inconspicuous polyp phase for long periods before blooming when environmental conditions are favorable (Eldredge and Smith 2001). Therefore, as *P. punctata* has already invaded almost the entirety of the Mediterranean Sea in the last decades, it is possible that it was already present in Cyprus but went undetected due to absence of field studies. In agreement with that, a recent horizon scanning of invasive species in Cyprus scored as “very high” the likelihood of arrival of *P. punctata* (Peyton et al. 2019), whereas a recent study assessing the invasiveness of 45 jellyfish species in the Mediterranean ranked *P. punctata* even among the top three species (Killi et al. 2020). However, based on the available data, the possibility of a recent arrival in the area also cannot be excluded. This could have happened through adult passive drifting from nearby populations in the Levant Sea, but also with early life stages being transported in ballast waters or clinging to vessel hulls, as already speculated by Deidun et al. (2017) for records in the central Mediterranean Sea.

Apart from these uncertainties, the potential of establishing permanent large-scale populations of *P. punctata* in Cyprus, as well as the consequences this establishment may incur, should not be underestimated. In fact, although this taxon does not represent a direct threat to human health or tourism industry (Boero et al. 2009), it has already caused significant impacts to fisheries (i.e. clogging nets) and the ecosystem (i.e. large clearance rates of zooplankton) (Garcia 1990; Perry et al. 2000; Silveira and Cornelius 2000; Graham et al. 2003; Rosales-Catalán et al. 2021), and thus regular blooms of this species might constitute an additional stressor for local fisheries.

Acknowledgements

Two anonymous reviewers improved an earlier version of the manuscript.

Authors' contribution

Conceptualization, A.K. and F.C.; methodology, A.K. and F.C.; validation, J.L., F.T., and F.C.; investigation, A.K., M.S.-O., V.M., and P.K.; data curation, A.K., V.M., and F.C.; writing – original draft preparation, A.K. and F.C.; writing – review and editing, A.K., M.S.-O., V.M., J.L., P.K., F.T. and F.C.; supervision, F.C. All authors have read and agreed to the published version of the manuscript.

References

- Abed-Navandi D, Kikinger R (2007) First record of the tropical scyphomedusa *Phyllorhiza punctata* von Lendenfeld, 1884 (Cnidaria: Rhizostomeae) in the central Mediterranean Sea. *Aquatic Invasions* 2: 391–394, <https://doi.org/10.3391/ai.2007.2.4.7>
- Anonymous (2011) Australian spotted jellyfish, *Phyllorhiza punctata*, invade Spanish beaches. <https://www.news.com.au/travel/travel-updates/australian-spotted-jellyfish> (accessed 15 February 2022)
- Atta MM (1991) The Scyphomedusae of the Mediterranean coast of El Arish. *Journal of the Egyptian German Society of Zoology* 6(A): 251–256
- Badreddine A, Bitar G (2020) *Cotylorhiza erythraea* Stiasny, 1920 (Scyphozoa: Rhizostomeae: Cepheidae): a new lessepsian jellyfish in the Lebanese waters, the eastern Mediterranean Sea. *Journal of the Black Sea/Mediterranean Environment* 26(3): 321–328
- Boero F (2013) Review of jellyfish blooms in the Mediterranean and Black Sea. General Fisheries Commission for the Mediterranean Studies and Reviews 92, 53 pp
- Boero F, Putti M, Trainito E, Prontera E, Piraino S, Shiganova TA (2009) First records of *Mnemiopsis leidyi* (Ctenophora) from the Ligurian, Tyrrhenian and Ionian Seas (Western Mediterranean) and first record of *Phyllorhiza punctata* (Cnidaria) from the Western Mediterranean. *Aquatic Invasions* 4: 675–680, <https://doi.org/10.3391/ai.2009.4.4.13>
- Brotz L, Pauly D (2012) Jellyfish populations in the Mediterranean Sea. *Acta Adriatica* 53(2): 213–232
- Cevik C, Derici OB, Cevik F (2011) First record of *Phyllorhiza punctata* von Lendenfeld, 1884 (Scyphozoa: Rhizostomeae: Mastigiidae) from Turkey. *Aquatic Invasions* 6: S27–S28, <https://doi.org/10.3391/ai.2011.6.S1.006>
- Coll M, Piroddi C, Albouy C, Ben RaisLasram F, Cheung WW, Christensen V, Karpouzi S, Guilhaumon F, Mouillot D, Paleczny M, Palomares ML, Steenbeek J, Trujillo P, Watson R, Pauly D (2012) The Mediterranean Sea under siege: spatial overlap between marine biodiversity, cumulative threats and marine reserves. *Global Ecology and Biogeography* 21: 465–480, <https://doi.org/10.1111/j.1466-8238.2011.00697.x>
- Dailianis T, Akyol O, Babali N, Bariche M, Crocetta F, Gerovasileiou V, Chanem R, Gökoğlu M, Hasiotis T, Izquierdo-muñoz A, Julian D, Katsanevakis S, Lipej L, Mancini, E, Mytilineou CH, Ounifi Ben Amor K, Ozgül A, Ragkousis M, Rubio-Portillo E, Servello G, Sini M, Stamouli C, Steriotti A, Teker S, Tiralongo F, Trkov D (2016) New Mediterranean Biodiversity Records (July 2016). *Mediterranean Marine Science* 17: 608–626, <https://doi.org/10.12681/mms.1734>
- Deidun A, Sciberras J, Sciberras A, Gauci A, Balistreri P, Salvatore A, Piraino S (2017) The first record of the white-spotted Australian jellyfish *Phyllorhiza punctata* von Lendenfeld, 1884 from Maltese waters (western Mediterranean) and from the Ionian coast of Italy. *BioInvasions Records* 6: 119–124, <https://doi.org/10.3391/bir.2017.6.2.05>

- Douek J, Paz G, Rinkevich B, Gevili R, Galil BS (2020) First record of a non-native pelagiid jellyfish (Scyphozoa: Pelagiidae: *Chrysaora*) in the easternmost Mediterranean Sea. *BioInvasions Records* 9: 482–489, <https://doi.org/10.3391/bir.2020.9.3.04>
- Durgham H (2011) First records of *Phyllorhiza punctata* von Lendenfeld, 1884 (Cnidaria: Rhizostomeae) from the Mediterranean coast of Syria. *International Journal of Oceans and Oceanography* 5(2): 153–155
- Eldredge LG, Smith CM (2001) A Guidebook of Introduced Marine Species in Hawaii. Bishop Museum Technical Report 21, August, 2001, 21 pp
- Galil BS (2012) Truth and consequences: the bioinvasion of the Mediterranean Sea. *Integrative Zoology* 7: 299–311, <https://doi.org/10.1111/j.1749-4877.2012.00307.x>
- Galil BS, Spanier E, Ferguson WW (1990) The Scyphomedusae of the Mediterranean coast of Israel, including two Lessepsian migrants new to the Mediterranean. *Zoologische Mededelingen* 64(7): 95–105
- Galil BS, Shoal L, Goren M (2009) *Phyllorhiza punctata* von Lendenfeld, 1884 (Scyphozoa: Rhizostomeae: Mastigiidae) reappeared off the Mediterranean coast of Israel. *Aquatic Invasions* 4: 481–483, <https://doi.org/10.3391/ai.2009.4.3.6>
- Galil BS, Gershwin L-A, Douek J, Rinkevich B (2010) *Marivagia stellata* gen. et sp. nov. (Scyphozoa: Rhizostomeae: Cepheidae), another alien jellyfish from the Mediterranean coast of Israel. *Aquatic Invasions* 5: 331–340, <https://doi.org/10.3391/ai.2010.5.4.01>
- Garcia JR (1990) Population dynamics and production of *Phyllorhiza punctata* (Cnidaria: Scyphozoa) in Laguna Joyuda, Puerto Rico. *Marine Ecology Progress Series* 64: 243–251, <https://doi.org/10.3354/meps064243>
- Giovos I, Kleitou P, Poursanidis D, Batjakas I, Bernardi G, Crocetta F, Doumpas N, Kalogirou S, Kampouris TE, Keramidas I, Langeneck J, Maximidi M, Mitsou E, Stoilas VO, Tiralongo F, Romanidis-Kyriakidis G, Xentidis NJ, Zenetos A, Katsanevakis S (2019) Citizen-science for monitoring marine invasions and stimulating public engagement: a case project from the eastern Mediterranean. *Biological Invasions* 21: 3707–3721, <https://doi.org/10.1007/s10530-019-02083-w>
- Graham WM, Martin DL, Felder DL, Asper VL, Perry HM (2003) Ecological and economic implications of a tropical jellyfish invader in the Gulf of Mexico. *Biological Invasions* 5: 53–69, <https://doi.org/10.1023/A:1024046707234>
- Gueroun SKM, Kéfi-Daly Yahia O, Deidun A, Fuentes V, Piraino S, Daly Yahia MN (2015) First record and potential trophic impact of *Phyllorhiza punctata* (Cnidaria: Scyphozoa) along the north Tunisian coast (South Western Mediterranean Sea). *Italian Journal of Zoology* 82: 95–100, <https://doi.org/10.1080/11250003.2014.981306>
- Gülşahin N, Tarkan AN (2012) The first record of *Phyllorhiza punctata* von Lendenfeld, 1884 from the southern Aegean Coast of Turkey. *BioInvasions Records* 1: 41–44, <https://doi.org/10.3391/bir.2012.1.1.09>
- Haddad MA, Nogueira Júnior M (2006) Reappearance and seasonality of *Phyllorhiza punctata* von Lendenfeld (Cnidaria, Scyphozoa, Rhizostomeae) medusae in southern Brazil. *Revista Brasileira de Zoologia* 23: 824–831, <https://doi.org/10.1590/S0101-81752006000300030>
- Heeger T, Piatkowski U, Möller H (1992) Predation on jellyfish by the cephalopod *Argonauta argo*. *Marine Ecology Progress Series* 88: 293–296, <https://doi.org/10.3354/meps088293>
- Katsanevakis S, Zenetos A, Belchior C, Cardoso AC (2013) Invading European Seas: assessing pathways of introduction of marine aliens. *Ocean & Coastal Management* 76: 64–74, <https://doi.org/10.1016/j.ocecoaman.2013.02.024>
- Killi N, Tarkan AS, Kozic S, Copp GH, Davison PI, Vilizzi L (2020) Risk screening of the potential invasiveness of non-native jellyfishes in the Mediterranean Sea. *Marine Pollution Bulletin* 150: 110728, <https://doi.org/10.1016/j.marpolbul.2019.110728>
- Knowler D (2005) Reassessing the costs of biological invasion: *Mnemiopsis leidyi* in the Black Sea. *Ecological Economics* 52: 187–199, <https://doi.org/10.1016/j.ecolecon.2004.06.013>
- Langeneck J, Crocetta F, Doumpas N, Giovos I, Piraino S, Boero F (2019) First record of the non-native jellyfish *Chrysaora* cf. *achlyos* (Cnidaria: Pelagiidae) in the Mediterranean Sea. *BioInvasions Records* 8: 608–613, <https://doi.org/10.3391/bir.2019.8.3.17>
- Madkour F, Elnagar N, Killi N (2021) Re-occurrence of the invasive jellyfish *Phyllorhiza punctata* von Lendenfeld, 1884 in the coast of Egypt and spreading pattern in the Mediterranean. *Biharean Biologist* 15: 87–90
- Malej A, Tirelli V, Lučić D, Paliaga P, Vodopivec M, Goruppi A, Ancona S, Benzi M, Bettoso N, Camatti E, Ercolessi M, Ferrari CR, Shiganova T (2017) *Mnemiopsis leidyi* in the northern Adriatic: here to stay? *Journal of Sea Research* 124: 10–16, <https://doi.org/10.1016/j.seares.2017.04.010>
- Mamish S, Durgham H, Al-Masri MS (2016) First record of the new alien sea jelly species *Marivagia stellata* Galil and Gershwin, 2010 off the Syrian coast. *Marine Biodiversity Records* 9: 23, <https://doi.org/10.1186/s41200-016-0029-2>
- Mizrachi GA, Shemesh E, Mizrachi A, Tchernov D (2021) Comparative genetics of scyphozoan species reveals the geological history and contemporary processes of the Mediterranean Sea. *Ecology and Evolution* 11: 10303–10319, <https://doi.org/10.1002/ece3.7834>

- Mutlu E, Çağatay IT, Olguner MT, Emer Yilmaz H (2020) A new sea-nettle from the Eastern Mediterranean Sea: *Chrysaora pseudoocellata* sp. nov. (Scyphozoa: Pelagiidae). *Zootaxa* 4790: 229–244, <https://doi.org/10.11646/zootaxa.4790.2.2>
- Perry HM, Van Devender T, Graham D, Johnson K, Larsen K, Burke WD, Trigg C (2000) Diaphanous denizens from down under: first occurrence of *Phyllorhiza punctata* in Mississippi coastal waters. Gulf and Caribbean Fisheries Institute Annual Meeting, Biloxi Mississippi. November 6–10, 2000
- Peyton J, Martinou AF, Pescott OL, Demetriou M, Adriaens T, Arianoutsou M, Bazos I, Bean CW, Booy O, Botham M, Britton JR, Cervia JL, Charilaou P, Chartosia N, Dean HJ, Delipetrou P, Dimitriou AC, Dörflinger G, Fawcett J, Fyttis G, Galanidis A, Galil B, Hadjikyriakou T, Hadjistyli M, Ieronymidou C, Jimenez C, Karachle P, Kassinis N, Kerametsidis G, Kirschel ANG, Kleitou P, Kleitou D, Manolaki P, Michailidis N, Mountford JO, Nikolaou C, Papatheodoulou A, Payiatas G, Ribeiro F, Rorke SL, Samuel Y, Savvides P, Schafer SM, Tarkan AS, Silva-Rocha I, Top N, Tricarico E, Turvey K, Tziortzis I, Tzirkalli E, Verreycken H, Winfield IJ, Zenetos A, Roy HE (2019) Horizon scanning for invasive alien species with the potential to threaten biodiversity and human health on a Mediterranean island. *Biological Invasions* 21: 2107–2125, <https://doi.org/10.1007/s10530-019-01961-7>
- Purcell JE, Uye SI, Lo WT (2007) Anthropogenic causes of jellyfish blooms and their direct consequences for humans: a review. *Marine Ecology Progress Series* 350: 153–174, <https://doi.org/10.3354/meps07093>
- Silveira FL, Cornelius PFS (2000) New observations on medusae (Cnidaria, Scyphozoa, Rhizostomae) from the northeast and south Brazil. *Acta Biologica Leopoldensia* 22(1): 9–18
- Rippingale RJ, Kelly SJ (1995) Reproduction and survival of *Phyllorhiza punctata* (Cnidaria: Rhizostomae) in a seasonally fluctuating salinity regime in western Australia. *Marine Freshwater Research* 46: 1145–1151, <https://doi.org/10.1071/MF9951145>
- Rizgalla J, Crocetta F (2020) First record of *Phyllorhiza punctata* von Lendenfeld, 1884 (Cnidaria: Scyphozoa: Rhizostomeae) in Libya through social media data mining. *BioInvasions Records* 9: 490–495, <https://doi.org/10.3391/bir.2020.9.3.05>
- Rosales-Catalán L, Estrada-González MC, Rivera-Pérez C, Reza Sánchez MA, Gamero-Mora E, Morandini AC, Mendoza-Becerril MA (2021) Genetic and morphological evidence of the presence of *Phyllorhiza punctata* in the southwestern Gulf of California (NE Pacific Ocean). *Aquatic Invasions* 16: 637–652, <https://doi.org/10.3391/ai.2021.16.4.04>
- Scyphers SB, Powers SP, Akins JL, Drymon JM, Martin CW, Schobernd ZH, Schofield PJ, Shipp RL, Switzer TS (2015) The role of citizens in detecting and responding to a rapid marine invasion. *Conservation Letters* 8: 242–250, <https://doi.org/10.1111/conl.12127>
- Stamouli C, Akel EHK, Azzurro E, Bakiu R, Bas AA, Bitar G, Boyaci YÖ, Cakalli M, Corsini-Foka M, Crocetta F, Dragičević B, Dulčić J, Durucan F, Zrelli, RE, Erguden D, Filiz H, Giardina F, Giovos I, Gönülal O, Hemida F, Kassar A, Kondylatos G, Macali A, Mancini E, Ovalis P, Paladini De Mendoza F, Pavičić M, Rabaoui L, Rizkalla SI, Tiralongo F, Turan C, Vrdoljak D, Yapici S, Zenetos A (2017) New Mediterranean Biodiversity Records (December 2017). *Mediterranean Marine Science* 18: 534–556, <https://doi.org/10.12681/mms.15823>
- Thiel M, Penna-Díaz MA, Luna-Jorquera G, Salas S, Sellanes J, Stotz W (2014) Citizen scientists and marine research: volunteer participants, their contributions, and projection for the future. *Oceanography and Marine Biology: An Annual Review* 52: 257–314, <https://doi.org/10.1201/b17143-6>