Faculty of Science and Engineering

School of Geography, Earth and Environmental Sciences

2022-10-01

Post-rift geomorphological evolution of a passive continental margin (Paraiba region, northeastern Brazil): Insights from river profile and drainage divide analysis

Alves, FC

http://hdl.handle.net/10026.1/19955

10.1016/j.geomorph.2022.108384 Geomorphology Elsevier

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1	Title: Post-rift geomorphological evolution of a passive continental margin (Paraíba
2	region, northeastern Brazil): Insights from river profile and drainage divide analysis
3	Authors: Fabio Corrêa Alves ^{a,b*} , Martin Stokes ^b , Sarah J. Boulton ^b , Dilce de Fátima
4	Rossetti ^a , Márcio de Morisson Valeriano ^a
5	Authors address:
6	^a Earth Observation and Geoinformatics Division, National Institute for Space Research,
7	Av. dos Astronautas, 1758 - São José dos Campos -12245–970, São Paulo, Brazil
8	(alves.fabioc@gmail.com; dilce.rossetti@inpe.br; marcio.valeriano@inpe.br)
9	^b School of Geography, Earth and Environmental Sciences, Plymouth University, Devon
10	PL4 8AA, UK (M.Stokes@plymouth.ac.uk; sarah.boulton@plymouth.ac.uk)
11	(*) corresponding author

13 **1. Introduction**

14 A passive margin is a transition zone between the oceanic and the continental 15 lithosphere formed by rifting followed by seafloor spreading (Bradley, 2008). Low- and high-elevation passive margins have been commonly recognized (Gilchrist and 16 17 Summerfield, 1990; Japsen et al., 2012). Low-elevation passive margins comprise coastal plains that gradually rise to low inland surfaces. High-elevation passive margins, 18 also known as elevated passive continental margins (EPCMs), comprise high relief 19 20 escarpments traversed by incised rivers that separate coastal plains from high inland plateaus (Gilchrist and Summerfield, 1990; Japsen et al., 2012). EPCMs are common in 21 various bedrock lithologies, climatic conditions, and neotectonic settings, such as in 22 west Greenland, southwest Africa, southeast Australia, Norway and northeast Brazil 23 (Figure 1). 24

An assumption about EPCMs is that inland plateaus remain elevated after the 25 26 main rifting, with minimal post-rift deposition (Gilchrist and Summerfield, 1990; Ollier and Pain, 1997; Gallagher et al., 1998). Elevated plateau persistence along EPCMs has 27 been explained by lithospheric response to differential denudation between coastal and 28 inland flanks due to isostasy (e.g., Gilchrist and Summerfield, 1990). However, the 29 integration of offshore and onshore stratigraphic records with long-term denudation 30 insights from thermochronology (Cobbold et al., 2001; Japsen et al., 2012 and 31 references therein) suggests that EPCMs have multiple episodes of uplift, erosion and 32 subsidence long after rifting. Thus, they are not necessarily permanent topographic 33 34 highs. Conversely, short-term EPCM denudation histories using cosmogenic and/or geomorphological techniques indicate that erosion rates are low (i.e., a few tens of 35 meters per million years), suggesting stability during the Late Cenozoic (Bierman and 36 37 Caffee, 2001). In some margins, climate (Souza et al., 2019) or rock strength (Gallen, 2018) might be important controls of short-term EPCM erosional rates. However, the 38 39 late stage evolution of passive margins is generally not well understood, especially 40 along margins where cosmogenic data are limited or non-existent.

Potential new insights into passive margin development often include the 41 42 morphology of the margin escarpment and plateau, especially in geologically data-poor regions. These landscape elements respond to changes in climatic (weathering) and 43 base-level (surface / rock uplift, eustasy) boundary conditions by controlling the rates 44 45 and directions of river incision and drainage divide migration (Willett et al., 2014; Roy et al., 2015). Therefore, the investigation of rivers and their catchment drainage divides 46 has the potential to provide models of margin development and reveal the controls of 47 48 their morphological changes. Longitudinal profiles of bedrock rivers and their knickpoints provide information on landscape steady-state or transience (Kirby and 49

Whipple, 2012). Profile-derived metrics, such as the normalized steepness index $-k_{sn}$, 50 may reflect spatial or temporal variations in rock uplift rates, rock strength or climate 51 (e.g., orographic rainfall patterns; Wobus et al., 2006; Whipple et al., 2013). Recent 52 53 methods to access divide migration (Willett et al., 2014) can further be used to identify lithological, tectonic or climatic signals in fluvial settings (Goren et al., 2014; Bernard 54 55 et al., 2019; Zondervan et al., 2020). These geomorphological approaches, together with 56 geological information, have contributed with investigations aiming the development of fluvial landscapes, especially in tectonically active regions (Snyder et al., 2000; Wobus 57 et al., 2006; Kirby and Whipple, 2012; Willett et al., 2014). However, this approach is 58 59 still scarce in EPCMs (Gallen et al., 2013).

60 This study is focused on the South American EPCM. Our interest was to investigate the northeast sector of this margin, a Brazilian region of low-rate tectonics 61 (uplift rates of 0.1 mm/yr for the last ~122 ka: Pedoja et al., 2011), but with a long 62 record of moderate magnitude earthquakes (i.e., 5.2 mb body-wave magnitude), as 63 informed by instrumented records (Bezerra and Vita-Finzi, 2000, Bianchi et al., 2018; 64 Figure 1). Geological evidence suggests variations in the post-rift (i.e., Neogene and 65 Quaternary) tectonic styles and rates along this margin (Margues et al., 2014; Gandini et 66 al., 2014; Nogueira et al., 2015; Vasconcelos et al., 2019; Bezerra et al., 2020). This is 67 especially the case of the Paraíba region, the final bridge between the South American 68 and African plates that remained active up to the Early Cretaceous (Matos, 1992). 69 Notably, this region records an abundance of Quaternary deposits with seismically-70 triggered soft sediment deformation (Rossetti et al., 2011a, 2011b; Alves et al., 2019; 71 Andrades Filho et al., 2021) and some uplifted marginal marine deposits (Gandini et al., 72 73 2014). These deposits suggest that the relief in the Paraíba region may have developed under tectonic instability. As a result, a transient fluvial landscape with drainage 74

75 network reorganisation is expected, which might include incised bedrock channels, 76 knickpoints, unstable and mobile drainage divides. The investigation of this fluvial 77 landscape has the potential to inform on the geomorphological evolution of this and 78 similar passive continental margins elsewhere.

We explored whether variations in tectonic uplift during the Quaternary 79 promoted responses on transient fluvial landscape in the Paraíba region. This was 80 achieved by searching: 1) the long river profile geometries based on the types and 81 distribution of knickpoints, which allowed to explore spatial patterns and controls on 82 base-level lowering (i.e., rock strength, uplift, eustasy, capture); 2) the k_{sn} values and 83 their relationship with spatial variations in uplift rates and bedrock; 3) the fluvial 84 85 incision, a proxy for surface uplift (Kirby and Whipple, 2012), as the rivers of the 86 Paraíba region may have changed their incision dynamics due to shifts in the rates of tectonic uplift during the Quaternary; and 4) the mobility of the catchment drainage 87 divides and the dynamics of divide migration, which were achieved by analyzing their 88 89 potential relationship to variations in the tectonic uplift or spatial differences in bedrock erodibility. 90

91 **2.** Geological and geomorphological setting

92 **2.1.** The passive continental margin of northeastern Brazil

Northeastern Brazil shows two geographic trends of passive margin development formed during the Late Mesozoic breakup of Pangea: 1) an equatorial margin, with an E- to W-trending coast; and 2) an eastern margin with a ~ NNE- to SSW-trending coast (Figure 1). The eastern margin, focus of this study, is drained by NE-SW- or NW-SE-oriented rivers, such as the Paraíba River and lower course of the São Francisco River. Most rivers that drain this margin originate in highlands dominated by a complex of pre-rift Precambrian rocks (Almeida et al., 1981), bordered
to the north and east/northeast by Cretaceous and Cenozoic deposits of the coastal
lowlands (Figure 1). The relief changes gradually from the highlands to the coast and
lacks a well-defined escarpment.

103

Figure 1.

The highlands with pre-rift Precambrian rocks are from the São Francisco 104 105 Craton and Borborema Provinces (Almeida et al., 1981). These consist of Archean and Proterozoic rocks that separate the marginal sedimentary basins to the east from the 106 107 intracratonic Paleozoic Parnaíba Basin to the west (Figure 1). The Borborema Province was built by the collision between the São Luis/West Africa and São Francisco/Congo 108 cratons (West Gondwana amalgamation) during the Neoproterozoic Brasiliano/Pan-109 110 African orogeny (Almeida et al., 1981). These cratonic areas are formed by crystalline basement rocks of variable strengths (Almeida et al., 1981; Brito Neves et al., 2004). 111

112 The passive continental margin of northeastern Brazil formed during the Triassic-Jurassic to Early Cretaceous rifting between the South American and African 113 continents, which culminated with the opening of the South Atlantic Ocean (Matos, 114 115 1992). This event resulted in several sedimentary basins along the coast in both the marine and continental realms (Figure 1). The Paraíba Basin is one of these basins, 116 117 representing the last continental bridge between the South American and African plates (Matos, 1992). This basin was filled during the rifting and the post-rifting (Rossetti et 118 al., 2012). The rift deposition occurred in the Late Cretaceous and is recorded by 119 sandstones of the Beberibe Formation, limestones and calciferous sandstones of the 120 121 Itamaracá Formation, and limestones of the Gramame Formation (Barbosa et al., 2003). The post-rift stage is recorded by siliciclastic deposits of Miocene and Late Quaternary 122

ages, and corresponds to the Barreiras Formation and Post-Barreiras Sediments 123 (Barbosa et al., 2003; Rossetti et al., 2012). All these units are delimited by regional 124 unconformities. The unconformity between the Cretaceous and Neogene units are 125 particularly prominent, being marked by an erosional surface with a deep lateritic 126 paleosol horizon (Rossetti et al., 2012, 2013). This unconformity was formed during a 127 prolonged time of tectonic stability. Fault reactivation during the Neogene and 128 Quaternary (i.e., late post rift stage) created new space for sediment accommodation, 129 which was filled by the deposits of the Barreiras Formation and Post-Barreiras 130 Sediments (Rossetti et al., 2012). These deposits show evidence of syn-sedimentary 131 132 tectonic deformation, which indicates that the region remained tectonically unstable even after the main rifting (Rossetti et al., 2011b). 133

134 The Paraíba Basin is bounded by two major tectonic structures, the Pernambuco 135 Shear Zone to the south and the Mamanguape Fault, the latter a branch of the Patos Shear Zone to the north (Figure 1). These shear zones formed during the Neoproterozoic 136 137 Brasiliano/Pan-African orogens (Almeida et al., 1981; Bezerra et al., 2011) and 138 constitute prominent continental-scale Precambrian basement structures (Figure 1). They are E-W- and ENE-WSW-oriented shear zones, 1 to 5 km in width, and hundreds 139 of kilometers in length. The present-day instrumented earthquake records show a 140 141 considerable number of low to moderate earthquake magnitudes (up to 6.0 mR) near some shear zones, such as the Pernambuco Shear Zone, and in other areas of the 142 northeastern Brazilian margin (Figure 1; Bezerra and Vita-Finzi, 2000; Bianchi et al., 143 2018). 144

The pre-existing shear zones in this passive continental margin were reactivated during the rift stage and controlled the location, geometry and sedimentary fill of the marginal basins (e.g., Matos, 1992; Bezerra et al., 2011; 2014). Reactivations occurred

also in the post-rift phase (Neogene and Quaternary), when basin bounding faults were 148 displaced by strike-slip transpressive stress in concert with regional stress field changes, 149 which caused basin inversions (e.g., Araripe, Rio do Peixe and Potiguar basins; 150 Marques et al., 2014; Nogueira et al., 2015; Bezerra et al., 2020). However, the 151 northeastern Brazilian margin also records a set of ~ NE-SW-striking normal faults, 152 consistent with evolution models of transtensional strike-slip fault systems (e.g., 153 Bezerra et al., 2014, 2020). These normal faults serve as pathways for watersheds that 154 155 link the highlands to the Atlantic Ocean.

156 **2.2**

2.2. Geomorphology of the study area

Our study focuses a 165-km long and 132-km wide EPCM located in the Paraíba 157 region (Figures 1 and 2a). From north to south, ocean-facing catchments in this margin 158 159 are formed by large and generally SW- to NE-oriented trunk streams, such as the Curimataú, Camaratuba, Mamanguape, Miriri and low Paraíba rivers (Figure 2a). These 160 rivers form wide valleys, filled in the downstream sectors by late Holocene alluvial 161 162 sediments (Figure 2a). Trunk streams are fed by NNW-SSE or NE-SW oriented tributaries (Figure 3) that extend from the highlands to the coastal plains. These 163 tributaries are mainly bedrock rivers with thin alluvial deposits downstream. Many 164 tributaries of the Paraíba River are disrupted by normal faults of the Cariatá Graben to 165 the south (cf. Brito Neves et al., 2004; Bezerra et al., 2008). This is an asymmetrical 166 trough of ~ 40 km long, ~ 25 km wide, and 250–550 m deep (Bezerra et al., 2008). 167 Two main fault reactivation events of pre-Late Pleistocene and Late Pleistocene ages 168 controlled the development of this graben (Bezerra et al., 2008). To the north, tributaries 169 of the Mamanguape River are disrupted by the Patos Shear Zone, where the tectonic 170 171 effect is still poorly known (Figure 3).

Figure 2.

172

173

Figure 3.

The regional drainage divide bounds catchments to the west, some 140 km from 174 the coast (Figures 2a and 3) within the highland plateau. The plateau stands at altitudes 175 averaging 750 m a.s.l., and it is constituted by Precambrian crystalline basement rocks 176 (e.g., gneisses, migmatites, schists, and granites), and secondarily Paleogene sandstones 177 178 from the Serra do Martins Formation (Morais Neto et al., 2008; Figure 2a). A deeply dissected low elevation (< 200 m a.s.l.) surface, known as Sertaneja Depression (Costa 179 180 et al., 2020), occurs to the east of this plateau (Figure 2b). The low topography of this depression is locally interrupted by Cretaceous igneous rocks that stands at altitudes up 181 to 600 m (Figure 2a). The coastal lowlands are dominated by Miocene (Barreiras 182 183 Formation) and Late Quaternary (Post-Barreiras Formation) siliciclastic deposits (Rossetti et al., 2012), and secondarily Late Cretaceous sandstones and limestones 184 (Barbosa et al., 2003; Figure 2a). The Barreiras Formation, first attributed to essentially 185 continental environments, were reinterpreted as having deposits formed in various 186 coastal, mainly estuarine environments (see Rossetti et al. 2013 for a review). The Post-187 Barreiras Sediments (Rossetti et al., 2012) include two informal stratigraphic units dated 188 from the Late Pleistocene (Post-Barreiras 1 - PB1) and early/middle Holocene (Post-189 190 Barreiras 2 - PB2). PB1 consists of sandstones interbedded with mudstones and 191 conglomerates, formed by marine (Gandini et al., 2014) or alluvial (fluvial and gravitational) processes (Rossetti et al., 2012). PB2 is constituted by friable and 192 massive, mostly aeolian sands (Rossetti et al., 2012). Both the Miocene and Late 193 194 Quaternary deposits discordantly overly Precambrian units. The coastal lowlands consist of a table-shaped relief (Costa et al., 2020) and a domal relief (Alves et al., 195 2019; Figure 2a-b), which give way to an adjacent narrow coastal plain to the east. 196

The climate in the Paraíba region is currently humid tropical in the coast, with an 197 average annual precipitation of 1917 mm and an average temperature of 26.5 °C 198 (Carvalho et al., 2020). A semi-arid climate with prolonged droughts prevails inland. 199 200 The paleoclimate in this region remains elusive, but successive planation (erosional) surfaces have long been related to prevailing semi-arid conditions (King, 1967). 201 However, studies based on new methodologies, such as apatite fission track, 202 203 cosmogenic nuclides and remote sensing, disagree that climate has played a main role in 204 defining the landscape of this passive margin, and suggest instead a main neotectonic control (Bezerra et al., 2008; Japsen et al., 2012; Gandini et al., 2014; Alves et al., 2019 205 and references therein). 206

207 **3. Methods**

We used a digital elevation model (DEM) for the geomorphological analysis of tributary stream profiles and catchment drainage divides to investigate tectonic upliftinduced base-level changes and drainage network reorganization during the Quaternary of the Paraíba passive margin.

212 **3.1. DEM processing**

213 The study area was assessed using the 30-m spatial resolution Shuttle Radar 214 Topography Mission (SRTM) DEM derived from C-band interferometric synthetic aperture radar (InSAR) techniques (Farr et al., 2007). This product has greater ability to 215 216 reveal surface information than other DEMs (e.g., Boulton and Stokes, 2018), since the C-band (~ 5.6 cm) can penetrate through the dense clouds and the upper tree canopy, 217 218 typical in tropical regions (Henderson and Lewis, 1998) as in the case of the study area. 219 This DEM was filled using minimum topographic values from surrounding pixels to generate a continuous drainage network. The processed DEM was used as the basis to 220

extract long river profiles and topographic metrics of drainage divides using codes
available in ChiProfiler (Gallen and Wegmann, 2017), Topographic Analysis Kit (TAK,
Forte and Whipple, 2019) and DivideTools (Forte and Whipple, 2018). All these codes
are based on functions of TopoToolbox (Schwanghart and Scherler, 2014).

225 **3**

3.2. River profile analysis

A long profile is a plot of river channel elevation against distance (Hack, 1957). The geometry of long river profiles and the derivation of fluvial metrics have been regularly used in geomorphological research to assess rates of relative base-level fall due to variations in tectonics, river capture, sea-level, climate, or lithology (Wobus et al., 2006; Kirby and Whipple, 2012). This approach can benefit geomorphic investigations of bedrock rivers in landscapes where the extent and/or the scarcity of geological exposures make field-based investigations difficult, such as in the study area.

River profiles were extracted from the drainage network using routine DEM 233 234 techniques (e.g., flow direction and flow accumulation). A drainage area threshold (A_{cr}) 235 of 1 km² or higher was applied to the drainage network to eliminate upstream river segments related with debris-flow processes (Wobus et al., 2006; Kirby and Whipple, 236 237 2012). The river profile analysis was applied to 43 tributary streams (Figure 3) of third or higher orders (cf., Strahler, 1957). These orders were chosen because they are often 238 characterized by bedrock segments, as required for detachment-limited models of 239 240 erosion (Kirby and Whipple, 2012). The selected river tributaries flow across the elevated plateau and coastal lowlands, crossing a range of lithologies of varied ages and 241 242 pre-existing structures.

The passive margin river profiles were analyzed using the stream power incision
model (SPIM) (Kirby and Whipple, 2012; Lague, 2014). The generalized equation of

the SPIM can be described by an empirical power law relationship (Hack, 1957; Kirby
and Whipple, 2012) between channel slope (S) and upstream drainage area (A; a proxy
for discharge):

248

$$S = k_s A^{-\theta} \tag{1}$$

where, k_s is the channel steepness index (m^{2 θ}); and θ the channel concavity index 249 250 (dimensionless) (Snyder et al., 2000). In steady-state landscapes, where uplift equals 251 erosion, spatial variability in k_s values may reflect spatial or temporal changes in erosion and uplifting rates in reliefs with homogeneous lithological and climatic conditions 252 253 (Wobus et al., 2006; Kirby and Whipple, 2012). To compare k_s values from river profiles with different sizes, the normalized channel steepness index (k_{sn}) is obtained 254 based on a fixed reference concavity (θ_{ref}) (Wobus et al., 2006). Empirical studies have 255 shown that θ varies between 0.4–0.6, but a fixed reference concavity of 0.45 is often 256 used for characterizing steady-state channels in a variety of reliefs across the world 257 258 (e.g., Snyder et al., 2000; Wobus et al., 2006; Kirby and Whipple, 2012).

Parameters S and A can be extracted directly from DEMs, while k_{sn} and θ are 259 260 extracted from the linear regression between S and A on a log-log graph. However, the channel slope may enhance significant scatter with direct influence on the k_{sn} 261 calculation, since topographic data comprise noisy variations (microrelief, artifacts and 262 263 errors) that are propagated to the slope (first order derivative of elevation). To circumvent these issues and extract more reliable k_{sn} values, the integral method 264 (integration of drainage area along flow distance) has been proposed (Perron and 265 266 Royden, 2013). This approach rearranges equation 1 by replacing S for dz/dx, where dzis the elevation change and dx the distance along the channel. The integration is 267

268 performed in an upstream direction from a base-level (x_b) to a given point along the 269 channel (x) as follows:

270
$$z(x) = z(x_b) + \left(\frac{ks}{A_0^{\theta}}\right) \int_{x_b}^x \left(\frac{A_0}{A(x)}\right)^{\theta} dx$$
(2)

271 with the variable

272
$$\chi = \int_{x_b}^{x} \left(\frac{A_0}{A(x)}\right)^{\theta} dx$$
(3)

273 where, A_0 refers to a reference drainage area inserted to make the area term 274 dimensionless (c.f., Perron and Royden, 2013). Assuming steady-state landscapes, with 275 invariant bedrock erodibility, the longitudinal coordinate χ with dimensions of length 276 has a linear relationship with elevation z(x) (Perron and Royden, 2013; Willett et al., 277 2014; Whipple et al., 2017). In this situation, a given profile will appear as a straight 278 line in the χ -elevation space (also known as chi plot).

The k_{sn} calculation was performed using the integral method to identify regions 279 280 in the passive margin with spatial or temporal variability in rock uplift rates, climate or differences in bedrock erodibility. The integral method was calculated using an A_0 of 1 281 m^2 to ensure that the local slope of a transformed profile is the k_{sn} (Perron and Royden, 282 2013; Gallen and Wegmann, 2017). Exploratory analysis of log-log slope-area plots of 283 tributaries with graded stream profiles (concave shapes) in the study area revealed an 284 average concavity value of 0.41 (Supplementary Figure S1). This value, which is very 285 close to the standard concavity value of 0.45 (Wobus et al., 2006; Kirby and Whipple, 286 2012) was used in this study as basis to calculate the k_{sn} and allowed us to make 287 comparisons with previous works carried out in other similar EPCMs (Gallen, 2018; 288 Souza et al., 2019) and Brazilian intracontinental (Peifer et al., 2021) settings. Average 289

290 k_{sn} values were calculated for the tributaries by adjusting linear regression bounds in 291 profile segments above and below knickpoints; and for the entire drainage network 292 using stream segment lengths of 1 km, since k_{sn} values are strongly variable over short 293 distances.

The potential influence of rock strength on the spatial variability/magnitude of 294 k_{sn} values was investigated by averaging the k_{sn} of drainage network by rock types. In 295 the absence of other parameters (e.g., quantitative in situ rock mass strength, jointing, 296 and weathering data; Bursztyn et al., 2015), the average k_{sn} considers only the 297 lithological character of rocks (Bernard et al., 2019; Zondervan et al., 2020) and can be 298 299 used as a proxy for bedrock erodibility (Gallen, 2018). In this study, the various 300 geological units were simplified into a smaller number of lithological groups 301 (Supplementary Figure S2). According to literature-based descriptions of rock strength (e.g., Goudie, 2006), we defined three major rock groups in terms of relative importance 302 of resistance to fluvial erosion using the official geological map from the Brazilian 303 Geological Survey (CPRM) at 1:1000000 scale (http://geosgb.cprm.gov.br/geosgb): (i) 304 low resistant sedimentary rocks; (ii) moderately resistant metamorphosed rocks; and 305 (iii) highly resistant igneous and meta-igneous rock units (Table 1; Figure 6). The 306 307 average and standard deviation k_{sn} were extracted for each rock group based on a total of 10,000 random samples. These samples were distributed proportionally to the area of 308 each group based on their uneven distribution in the studied passive margin 309 (sedimentary rocks = 25.4 %; metamorphosed sediments = 5.3 %, and igneous and 310 meta-igneous rocks = 69.3 %). This analysis was not applied to 311 the alluvial/marine/aeolian deposits to avoid abrupt changes in k_{sn} related to depositional 312 channel segments. 313

314

Table 1.

In addition to the k_{sn} approach, rivers under non-steady-state conditions (uplift \neq 315 316 erosion) may produce unbalanced geomorphic features (transient features), such as convex-upward discontinuities (knickpoints; Kirby and Whipple, 2012). The study of 317 318 knickpoints along river profiles has been used as an indicator of rivers with a transient response due to rock strength controls or perturbations related to relative base-level 319 changes. These are caused by tectonics (e.g., increase in the fault slip rates), river 320 capture, sea-level fall and/or climatic variations. These factors can enhance or reduce 321 322 the efficiency of river incision (Kirby and Whipple, 2012; Whipple et al., 2013). We identified major knickpoints by visual interpretation of convex-upward deviations in 323 324 river profile shapes from the distance-elevation and γ -elevation plots. Knickpoints were classified into two end-member morphologies, vertical-step or slope-break (c.f., Haviv 325 et al., 2010 and Whipple et al., 2013), based on the recognition of spikes and breaks in 326 327 log-log slope-area graphs, respectively. Given the importance of knickpoints in 328 preserving information about the tectonic evolution of fluvial landscapes (Kirby and 329 Whipple, 2012), they were further characterized in terms of: spatial patterns between the 330 knickpoint locations and lithological contacts/pre-existing basement structures, distribution pattern, statistical relationships, formation and behavior. 331

332 The knickpoint analysis also included the application of two statistical hypothesis tests: the unpaired two-samples T-test (parametric test); and the Mann-333 Whitney U test (non-parametric test) (c.f., Noether, 1991; Venables and Smith, 2002). 334 335 These tests aimed to examine two clusters of knickpoints in the study area. To attend 336 the T-test assumptions, data normality analysis using the Shapiro-Wilk test and homogeneity of variances based on Levene's test were required (Venables and Smith, 337 2002). The parameters elevation, slope, relative relief and k_{sn} at a given knickpoint were 338 chosen because they may inform important aspects about the knickpoint origin and, 339

ultimately, the passive margin evolution. The relative relief was calculated varying the moving window radius (1, 3 and 5 km) to assess the windows sensitivity. Only the attributes that failed in the normality analysis (T-test) were submitted to the Mann-Whitney U test. The following null (h_0) and alternative (h_a) hypotheses were defined assuming a significance level (α) of 0.05: h_0 = the means/medians of the two groups of knickpoints are the same; h_a = the means/medians of the two groups of knickpoints are different.

The amount of fluvial incision was calculated for tributaries with slope-break 347 knickpoints. Fluvial incision was estimated from the reconstruction of channel profiles, 348 349 an approach typically applied to access the uplift history of a given area (Kirby and 350 Whipple, 2012). The calculations were made downstream of the slope-break 351 knickpoints in the lower reaches of the studied tributaries, that is, the region potentially adjusting to perturbations. Following the method implemented in the TAK (Forte and 352 Whipple, 2019), the river profiles downstream of the knickpoints were projected using 353 linear least squares fitting on the χ -elevation relationship, with a confidence interval of 354 95%. The estimated fluvial incision was then obtained by subtracting the elevations 355 from the projected (relict channel profile) and present-day profiles at the fault. The 356 357 position of the fault at the profile was defined based on the visual interpretation of river locations and major morphostructural lineaments/faults/shear zones in the study area. 358 The lineaments (204 in total with a main NE-SW trend) were mapped using SRTM-359 based shaded relief representations in different azimuth angles at a fixed 1:100.000 360 scale (Supplementary Figure S3). Known faults and shear zones were derived from 361 previous publications (Bezerra et al., 2008, 2014). 362

363

3.3

3.3. Assessment of divide mobility

Landscapes undergoing transient adjustments due to changes in boundary 364 conditions, such as those recording incised rivers, knickpoints and contrasting erosion 365 rates between adjacent drainage basins, may show drainage divide migration (Willett et 366 367 al., 2014). Active drainage divide migration has been recognized in a variety of fluvial landscapes as a result of many competing controls, such as lithology, where the 368 drainage divide moves towards highly resistant rocks (Bernard et al., 2019; Zondervan 369 et al., 2020), the spatial variability of precipitation (Goren et al., 2014), asymmetric 370 371 tectonic uplift (He et al., 2019), or large discrete river capture (Willett et al., 2014; Whipple et al., 2017). 372

The assessment of divide mobility has been performed by maps of the drainage 373 374 network coloured by χ (χ -maps). Significant variations in χ on opposite sides of a drainage divide (x-anomalies) may suggest divide migration (e.g., due to a river 375 capture), which is driven by differences in erosion rates of river channels (Willett et al., 376 2014). Alternatively, it has been suggested that divide instability analysis can be made 377 more reliable by interpreting cross-divide differences using topographic metrics, also 378 known as Gilbert metrics (i.e., elevation at the channel head, upstream relative relief 379 and upstream gradient), as these are topographic proxies for erosion rates (Whipple et 380 381 al., 2017; Forte and Whipple, 2018). In this study, the stability/mobility and direction of divide migration were investigated through the χ -map and Gilbert metrics extracted at 382 the channel heads. The χ -map was generated for complete drainage basins with outlets \geq 383 384 1 m a.s.l. based on the SRTM-DEM. For the sake of simplification, we analyzed only the drainage divides bounding major transverse fluvial systems. The drainage divide 385 was assumed to be stable if the mean on one side of the divide overlaps the uncertainty 386 387 of the mean on the opposite side; otherwise, the divide was assumed as mobile (c.f., Forte and Whipple, 2018). The direction of divide migration was defined following the 388

interpretations described in Willett et al. (2014), Whipple et al. (2017) and Forte andWhipple (2018).

- 391 4. River profiles and drainage divides
- 392 4.1. Knickpoints, *k*_{sn} and fluvial incision

The Paraíba region has tributaries up to ~117 km long and catchment areas up to 393 394 ~994 km² (Figure 3). Tributary streams crossing the Precambrian highlands are longer than those from the coastal lowlands (average lengths of \sim 34 km and 24 km, 395 396 respectively). These tributaries showed river profiles that can be qualitatively classified into two major groups according to their profile shapes shown in distance-elevation and 397 χ -elevation plots: (i) relatively graded profiles (concave-up profiles and linear χ -398 profiles); and (ii) ungraded profiles (convex-up river profiles and non-linear χ -profiles; 399 400 Figures 4a-f and 5a-f).

- 401 Figure 4.
- 402

Figure 5.

The first group of graded profiles comprises twenty-one tributaries, recorded mostly in the low-lying terrains of the passive margin, with profile elevations not exceeding 450 m a.s.l. (Figure 4a-f). These profiles have no knickpoints. Some small tributaries on flat terrains capped by Miocene deposits in the southeast sector of the study area show river profiles with convex segments, suggesting broader knickzones (rivers 26, 27, 28, and 29 in Figure 4f). However, these tributaries did not show clearly recognizable breaks or spikes in the log-log slope-area plots. The second group of ungraded profiles is represented by twenty-two tributaries (Figure 5a-f). These were recorded mainly in the Precambrian highlands of the passive continental margin to the west, and secondarily, in the coastal lowlands to the east. They showed: profile elevations up to 700 m a.s.l. (Table 2); and river profiles with widely scattered knickpoints (vertical-step and slope-break; Figure 5a-i).

416 We recorded a total of fourteen vertical-step knickpoints (green dots in Figure 5a-i). They occurred at or close to lithological contacts between Precambrian units (e.g., 417 knickpoints from rivers 1, 2, 5, 13 and 15 in Figure 6a and Supplementary Figure S4); 418 and in river segments downstream of slope-break knickpoints. These knickpoints 419 occurred at elevations between 73 and 507 m a.s.l., and revealed maximum k_{sn} values 420 421 almost twice as high as the slope-break knickpoints (Table 2). For instance, vertical-step knickpoints showed an average k_{sn} at knickpoint of 38.5 m^{0.9} and a maximum k_{sn} value 422 of 85.3 m^{0.9}, while slope-break knickpoints displayed an average k_{sn} at knickpoint of 423 27.4 m^{0.9} and a maximum k_{sn} value of 49.3 m^{0.9} (Table 2). 424

425

Figure 6.

426 The majority of the knickpoints were classified as slope-break type (purple dots 427 in Figure 5). They occurred across a range of elevations (65-676 m a.s.l.) and were 428 limited by downstream segments steeper than upstream (see k_{sn} ratio in Table 2). Most 429 rivers had one slope-break knickpoint each, but three rivers (rivers 5, 6, and 14) exhibited two slope-break knickpoints. Knickpoints from river 14 are likely to be part of 430 431 a single knickpoint, as they showed a similar morphology and are located close to each other across a distance of ~5 km. The slope-break knickpoints occurred generally 432 aligned across a \sim NNE-SSW-oriented belt over the highlands (Figure 6a), where they 433

lie upstream of the Patos Shear Zone to the north and upstream of the normal faults bounding the Cariatá Graben to the south (Figure 7). However, a small number of knickpoints were also observed closer to lithological contacts of the Precambrian units or between these and Miocene-Quaternary deposits (e.g., knickpoints from rivers 2 and 10 in Figure 6a and Supplementary Figure S4). Only the current position of the slopebreak knickpoints was recorded, but their behaviour is generally mobile over time (Crosby and Whipple, 2006; Berlin and Anderson, 2007; Boulton and Whittaker, 2009).

441

Figure 7.

The k_{sn} values ranged from 1 to 462 m^{0.9} with a right-skewed distribution and an 442 average k_{sn} of 15 m^{0.9} (Figure 6a). Higher k_{sn} values (> 20 m^{0.9}) were found in the 443 highlands (to the west) and dome-like reliefs (to the east), where the distribution of 444 relative relief values is high (Supplementary Figure S5). Lower k_{sn} values (< 20 m^{0.9}) 445 occurred mainly over the Precambrian low-lying and sedimentary table-like terrains. 446 The rock group average k_{sn} analysis showed a trend of an increased slope of channels 447 flowing over more resistant rock units (i.e., igneous and meta-igneous rocks of the 448 Precambrian basement), in contrast to low steepness channels that drain over less 449 resistant rock units (i.e., sedimentary rocks) (Figure 6b). Abrupt changes in k_{sn} values 450 owing to local differences in bedrock resistances were evident in the northwest and 451 northeast sectors of the study area, where pre-rift rock units of Precambrian age are in 452 453 contact with Paleogene and Miocene-Quaternary deposits, respectively (Figure 6a). However, local changes in k_{sn} were detected even in bedrock terrains of relatively 454 similar resistances, which cannot be explained only by lithological control. 455 456 Furthermore, this analysis showed a low variability of k_{sn} values among different rock groups (Figure 6b and Supplementary Figure S2), in general ranging from 10 to 25 m^{0.9}, 457 but with high uncertainties. 458

The distribution pattern of the knickpoints (slope-break and vertical-step) in the 459 460 study area is shown in Figure 8a. K_{sn} values downstream of knickpoints were higher than upstream for both slope-break and vertical-step morphologies, but with higher 461 462 uncertainties for the latter (Table 2). This situation was observed mainly for the tributary streams flowing over the Precambrian terrains of the study area, where 463 knickpoint elevations are high (Figure 8a). The statistical analysis between k_{sn} and 464 relative relief values at knickpoints showed moderate and weak positive linear 465 466 relationships for the slope-break ($r^2 = 0.61$) and vertical-step ($r^2 = 0.36$) morphologies, respectively (black/white dots in Figure 8b). The high coefficient of determination for 467 468 the slope-break knickpoints and their spatial correspondence with the structural fabric (i.e., upstream of known faults) of the study area suggest that they may have inherited 469 some tectonic influence. Therefore, subsequent analyses were carried out based only on 470 471 the slope-break knickpoints, given their potentially higher tectonic significance (Kirby 472 and Whipple, 2012; Whipple et al., 2013).

473

Figure 8.

We carried out some statistical analysis using information from the knickpoints 474 and their drainage basins to characterize the knickpoint formation (e.g., one or multiple 475 476 knickpoint clusters) and behavior (i.e., anchored in place or mobile). The regression analysis of the knickpoint elevation versus drainage area, and the knickpoint elevation 477 against the knickpoint distance from mouth showed weak relationships, with very poor 478 determination coefficients ($r^2 < 0.13$) when the slope-break knickpoints were analyzed 479 together (Figure 9a-b). In contrast, these relationships were strengthened when the 480 481 knickpoints from the north (Patos Shear Zone region) and south (Cariatá Graben region) of the passive continental margin were analyzed individually. The northern and southern 482 knickpoints showed positive ($r^2 = 0.43$ and 0.51) and negative ($r^2 = 0.42$ and 0.46) linear 483

relationships, respectively (Figure 9a-b). The statistical analysis between the average 484 basin gradient versus catchment area (not shown here) revealed a negative linear 485 relationship, with a moderate determination coefficient ($r^2 = 0.42$) only for the basins 486 located in the southern sector of the study area. These drainage basins have steeper 487 gradients than those from the northern sector, especially the small basins located closer 488 to the Cariatá Graben (Figure 7). The discrepancies with the knickpoint / drainage basin 489 relationships and the high range of slope-break knickpoint elevations led us to test the 490 491 hypothesis of two different clusters of knickpoints in the study area. The T-test showed that the means of the relative relief (radius of 3 and 5 km) of the southern knickpoints 492 493 are statistically different from the means of the northern ones (p-value < 0.05; Table 3), assuming an α of 0.05. The Mann Whitney U test did not show any statistical 494 differences in the slope of the knickpoints from both sectors (Table 3), although 495 496 limitations of dataset size could be an issue.

- 497 Figure 9.
- 498

Table 3.

The knickpoint retreat distance from the mouths and the downstream distances 499 500 from the drainage divides positively scaled with the total drainage area and drainage 501 area below and above the slope-break knickpoints, with strong r^2 coefficients (i.e., $r^2 > r^2$ 502 0.89; Figure 9c-e). Knickpoints from larger basins (e.g., from rivers 5 and 14, see Table 503 2) travelled further upstream than knickpoints from smaller basins (e.g., from rivers 3, 4 and 20). The drainage area upstream of knickpoints ranged from 4.2 to 238.3 km² and 504 from 2.4 to 125.3 km² for the northern and southern knickpoints, respectively. Hack's 505 506 exponents varied from 0.60-0.68 when all the knickpoints are analysed together; 0.62507 0.77 for the northern knickpoints; and 0.56-0.58 for the southern knickpoints (Figure
508 9c-e).

The amounts of fluvial incision derived from river profile segments downstream of slope-break knickpoints were 140 m on average (Table 2). Average fluvial incision was 156 m for tributaries over the Precambrian highlands and 43 m for tributaries crossing the coastal lowlands. This situation suggests spatially variable fluvial incision in the studied region, with higher incision amounts inland towards the elevated plateau formed by pre-rift Precambrian basement rocks.

515 **4.2. Drainage divide and divide mobility**

The main drainage divides of the study area extend from the elevated plateau to 516 the coastal lowlands, a total of 480 km in length (Figure 10a). In general, the drainage 517 divides have either NE-SW- or NW-SE-oriented segments and are at a maximum 125 518 km from the coastline. These were qualitatively divided into divide segments (D1-D8) 519 520 to assess local variations in divide migration (Figure 10a). From west to east, the 521 drainage divide segments traversed a range of lithologies (i.e., pre-rift basement rocks and post-rift depositional sequences) with contrasting erodibilities (Figures 2a and 10a). 522 523 The longest drainage divide segment (D5) is 153 km long and runs from 740 m a.s.l. across high resistant Precambrian units (igneous and meta-igneous rocks) and, 524 525 secondarily, low resistant Paleogene deposits to the west. Divide D5 decreased in elevation to ~ 350 m a.s.l., as it crossed moderate resistant metamorphic rocks to the 526 527 east. Divide D1 to the southwest crosses only high resistant Precambrian units, with 528 elevations ranging from \sim 500 to 140 m a.s.l. Divides D2-D3-D4 and D6-D7 to the southeast and northeast, respectively, occur between high to moderate resistant 529 Precambrian units and low resistant to erodible Miocene to Quaternary-deposits. D8 530

also crosses deposits formed since Miocene, but it occurs on domal reliefs withelevations up to 200 m a.s.l.

533

Figure 10.

Most divide segments (i.e., D1, D4, D5, D6 and D7) were stable, with no 534 significant cross-divide differences in the γ -map and Gilbert metrics (Figures 10a and 535 11a-d). For instance, D1, D4, D6, and D7 showed relatively equal average channel 536 537 elevation values of about 100 m a.s.l. (Figure 11a). They also displayed similar hillslope gradient values around 0.1 and relative relief values ranging from 40 to 50 m (Figure 538 539 11b-c). D5 showed the highest average elevation value up to 500 m a.s.l., the highest mean hillslope gradient, and relative relief of approximately 0.15 and 75 m, respectively 540 (Figure 11a-b-c). 541

542

Figure 11.

Among the drainage divide segments, only three (D2, D3, and D8; Figures 10a 543 and 11a-d) showed variations in χ and Gilbert metrics on both sides of the divide, 544 545 indicating divide migration. D8, located in a coastal lowland dome, had differences only 546 in the Gilbert metrics of hillslope gradient and relative relief. It also had the highest 547 average hillslope gradient and relative relief values (0.15 and 75 m, respectively), with 548 northward divide motion (Figure 11b-c-d). D2 and D3, located to the southeast, 549 revealed differences in both χ and Gilbert metrics, with relatively similar topographic 550 metrics (average channel elevation of ~ 100 m; hillslope gradient of ~ 0.05; relative relief of ~ 50 m; Figure 11a-c). γ and Gilbert metrics suggest divide movement to the 551 552 southeast for D2 and to the northeast for D3 (white arrows in Figure 10a and Figure 11). Divide movements follow the inland position of a high-elevation, chevron-shaped 553 plateau with elevations up to 170 m a.s.l. (Figures 2b and 10b). This plateau, not 554

included in previous geomorphological maps of northeastern Brazil (e.g., Costa et al., 2020), developed over low-resistant Miocene and Quaternary deposits that overly resistant Precambrian basement rocks. The plateau configuration is sustained by incised tributaries, some showing headwaters close to each other (Figure 10b). An ongoing river capture between a tributary of the Paraíba River left bank and the upper Miriri River was recorded in the lower part of this plateau (white dot in Figure 10b).

561 **5. Discussion**

The present-day rivers of the Paraíba region show a set of features compatible 562 with a transient landscape (Kirby and Whipple, 2012; Willett et al., 2014). Evidence 563 includes non-linear χ -profiles bounded by knickpoints, bedrock river channels with 564 spatially variable k_{sn} , and mobile drainage divide segments. These features were often 565 566 recognized in erosional reliefs of tectonically active collisional mountain landscapes (Snyder et al., 2000; Wobus et al., 2006; Kirby and Whipple, 2012), rather than in the 567 generally tectonically stable EPCMs (e.g., Gallen et al., 2013). The transient features 568 recorded in the Precambrian highlands and coastal lowlands indicate that the Paraíba 569 region had a topographic rejuvenation in the recent geological time due to fluvial 570 disturbances. The significance of such transient features and the potential controlling 571 factors that triggered the transient response in the late development of this passive 572 573 continental margin are discussed below.

574 5.1. Dynamics and control on divide migration

575 The resulting χ -map and Gilbert metrics did not show many topographic or 576 geometric changes in the fluvial channels and drainage divides of the studied seaward-577 dipping catchments. However, the migration of divides D2 and D3 (Figures 10 and 11) 578 in the southeastern sector is due a lithological strength contrast. This is proposed

because these divides occur in a plateau sustained by less resistant Miocene-Quaternary 579 580 sedimentary units in contact with Precambrian basement rocks. However, the topographic expression of this plateau may not have formed primarily due to a 581 582 lithological control. This plateau is topographically similar to the Shillong Plateau of northeast India (Strong et al., 2019). The Shillong Plateau resulted from the erosion of 583 horizontally layered lithologies with contrasting resistances. This condition re-exposed 584 a basement paleosurface, as proposed by numerical modelling of erosion (soft rocks 585 586 over hard rocks; Forte et al., 2016). However, we interpret that the studied plateau was formed by a young tectonic uplift, i.e., probably due to the Late Pleistocene reactivation 587 588 of the normal fault bounding the left margin of the Cariatá Graben (Bezerra et al., 2008), rather than the exhumation of bedrocks with contrasting erodibilities. This is 589 evidenced by: i) the plateau covered by younger deposits (i.e., Late Quaternary Post-590 591 Barreiras Sediments) at higher elevations (i.e., up to 200 m a.s.l.) than adjacent 592 Precambrian rocks (Figures 2b and 7); and ii) incised tributaries to the east of the 593 plateau draining across a high-elevation Quaternary dome (Alves et al., 2019). Thus, the 594 most likely is that tectonic uplift would have elevated basement rocks and overlying marine deposits, followed by fluvial incision and partial dissection of both the plateau 595 and the dome. 596

597 Divide mobility to the southeast of the studied region is also supported by the 598 current river capture between the left-bank tributary of the Paraíba River and the upper 599 Miriri River (Figure 10). The headwater of the Paraíba River tributary at a lower base-600 level records accelerated headward erosion. This is evidenced by channel deepening and 601 valley widening towards an elevated dome, where the Miriri River headwater is located. 602 Once this fluvial connection has been completed, the less active Miriri River tributary 603 will entrench south through the Paraíba Basin and abandon some of its uppermost 604 stretches. Similar river capture has been recorded in other EPCMs, such as in the 605 Ethiopian rift margin of northeast Africa (Giachetta and Willett, 2018). In the study 606 area, the river capture will be of low magnitude and will result in the local re-607 organization of the drainage network.

608

5.2. Characterization of slope-break knickpoints

Slope-break knickpoints are presumed to propagate upstream at constant vertical 609 rates, so they should occur at similar elevations (Neimann et al., 2001). This situation 610 611 would be expected, for instance, in landscapes with invariant climatic, rock uplift and bedrock erodibility conditions, and where the drainage basins have similar geometric 612 613 characteristics and incision dynamics. However, this expected transient response is generally rare in nature due to the variability of these factors and absence of an earlier 614 615 steady-state that might disperse elevations during the knickpoint retreat (Boulton and Whittaker, 2009; Kirby and Whipple, 2012; Whipple et al., 2013). The range of 616 knickpoint elevations (i.e., from 65 to 676 m a.s.l.) in the study area, especially in the 617 618 Precambrian highlands (Figures 5, 6a and 7), and the weak statistical relationships including drainage area versus knickpoint elevation ($r^2 = 0.06$), and knickpoint elevation 619 versus knickpoint distance from the mouth ($r^2 = 0.13$; Figure 9a-b), suggest either 620 knickpoints with distinct formation or natural variability as a result of a single 621 formation. Such conditions were documented in several other landscapes (e.g., Crosby 622 and Whipple, 2006; Berlin and Anderson, 2007; Boulton and Whittaker, 2009; Boulton 623 624 et al., 2014).

Basin gradient seems to explain, at least in part, the differences in statistical relationships (positive and negative) between the two knickpoint groups (southern and northern sectors) in the study area, which was higher for the southern basins. The 628 differences of means between the two knickpoint groups only for the relative relief parameter (Table 3) indicated by the unpaired two-samples T-test, were expected given 629 the high topographic heights of the graben from the southern sector. Alternatively, such 630 631 differences in the relative relief are perhaps due to technical reasons, such as the configuration of the search window (e.g., Dibiase et al., 2010). The catchment 632 dimensions might have been exceeded when larger radius thresholds were used (i.e., 3 633 634 and 5 km). The statistical similarities among most topographic parameters are perhaps 635 indicative that the lowest knickpoints in each river are from a unique base-level fall. This event followed the Miocene, because some slope-break knickpoints were found on 636 deposits of this age (Figures 2 and 6a). A previous simulations stated that fluvial 637 disturbances are perceived generally in the order of 1–5 Myr time interval (Whittaker 638 639 and Boulton, 2012). Thus, the fluvial disturbances in the study area occurred most likely 640 in the Late Quaternary. This is also supported by tributaries incising into dome-like reliefs during this time. However, the migration of some knickpoints from most 641 642 upstream catchment locations (e.g., rivers 5 and 6) is related to an earlier transient stage 643 of the rivers.

The regression analysis pointing to the strong scalings ($r^2 > 0.89$) between the 644 645 knickpoint horizontal distances and the drainage areas (Figure 9) suggests knickpoints not anchored in place. This observation is in line with records of horizontal knickpoint 646 retreat as primarily a function of the drainage area (e.g., Crosby and Whipple, 2006; 647 648 Berlin and Anderson, 2007; Boulton et al., 2014). The high variability of Hack's exponents by comparing the Cariatá Graben region (south) and the Patos Shear Zone 649 region (north) is more likely due to the sinuosity of some large tributaries. Hence, the 650 651 high sinuosity of the northern tributaries (e.g., rivers 5 and 6 in Figure 3) downstream of the knickpoints may have inflated part of the deviation of Hack's exponent ($L = 1.4A^{0.6}$; 652

Hack, 1957), as verified in other studies (Smart and Surkan, 1967; Willemin, 2000). However, the wide range of drainage areas upstream of knickpoints (i.e., from 2.4 to 238 km²) confirms that they are not pinned to a specific drainage area threshold, but are actively migrating upstream (Figure 9). The high scaling between the k_{sn} and the relative relief values (Figure 8b) further evidences that the tributaries are responding to a transient fluvial incision, with fluvial channels still adjusting to the new boundary conditions.

660

5.3. Control on knickpoints, k_{sn} variations, and fluvial incision

661 Transient features in fluvial landscapes, such as knickpoints and spatial variability in k_{sn} values, are indicative of base-level changes in response to boundary 662 conditions. The knickpoint formation might be due to the exposure of rocks with 663 664 differential erodibility, large river captures, and either spatial or temporal changes in past climate, sea-level and uplift rates (Wobus et al., 2006; Kirby and Whipple, 2012, 665 Gallen, 2018). Our topographic analysis showed that some vertical-step knickpoints 666 were lithologically controlled, as revealed by their: high scattering in the regression 667 analysis (Figure 8b); absence of a clear spatial correspondence between some 668 669 knickpoints and the structural fabric of the passive continental margin; and location close to lithological contacts of variable rock resistances (e.g., knickpoints from rivers 1 670 and 2 in Figure 6a and Supplementary Figure S4). Vertical-step knickpoints may reflect 671 local contrasting lithologies, as shown in previous numerical modeling (e.g., Forte et al., 672 673 2016).

674 However, the high number of vertical-step knickpoints with significant increase 675 in k_{sn} values downstream (Figure 8a) is a typical characteristic of slope-break 676 knickpoints (Whipple et al., 2013). These knickpoints indicate the influence of a

regional-scale factor. The uneven distribution of k_{sn} values (i.e., ranging from 8 to 462 677 $m^{0.9}$) in the highlands with Precambrian rocks to the west, where slope-break 678 knickpoints follow a broad ~NNE-SSW- trend (see dashed blue lines in Figure 6a), is 679 680 intriguing. Spatial variability of k_{sn} values in some EPCMs, such as in SE Brazil and eastern North America, was related to rock strength (Gallen, 2018; Peifer et al., 2021) or 681 climatic control (Souza et al., 2019). However, in the highlands of the study area, the 682 683 variability of k_{sn} values occurs across lithologies of relatively similar resistance (Figure 6a). In addition, climatic conditions, such as precipitation rates, do not vary 684 significantly from the coast to the passive margin high relief zone (Carvalho et al., 685 686 2020). Thus, these factors could not have altered the channel dynamics, neither they could explain the regional k_{sn} variations in the study area. 687

Large river captures may alter the fluvial channel dynamics and produce 688 knickpoints that propagate upstream (Whipple et al., 2017). However, the only ongoing 689 river capture identified in the study area (Figure 10) is of a low magnitude, and it could 690 691 not explain all knickpoints formed across the study area. The influence of eustatic sealevel lowering on knickpoints and k_{sn} variations is possible, particularly during the last 692 glaciation maximum (ca. 23 and 18 ka), when the global sea-level reached 80-120 m 693 below the modern sea level (e.g., Cutler et al., 2003; Clark et al., 2009). The extent to 694 which the sea level controls inland base-level and fluvial incision is still uncertain, but 695 100-150 km upstream from the coast has generally been proposed (Blum and Törnqvist, 696 2000). The uppermost tributaries in the study area are located up to 125 km from the 697 698 coast (Figure 8a), i.e., within the range of sea level influence. However, sea-level fall 699 alone could not explain the increased fluvial incision (i.e., 156 m on average) in bedrock rivers on the Precambrian highlands (Table 2). In addition, a sustained base-level 700 lowering longer than 100 ka is needed for the incision of tributary streams and 701

migration of knickpoints (e.g., Whittaker and Boulton, 2012). Furthermore, the ~120 m
global sea-level drop advocated for the Last Glaciation Maximum was fast, i.e., in the
scale of orbital cycles of up to 100 ka (Cutler et al., 2003; Clark et al., 2009). Thus, its
influence in the base-level changes and knickpoint propagation in this passive
continental margin is unlikely.

We propose that the Late Quaternary tectonics controlled the formation of many 707 708 knickpoints in the study area. Despite some differences in the proposed rates, several passive margins worldwide record increased rock uplift rates long after the rifting phase 709 710 (Japsen et al., 2012; Pedoja et al., 2011; Gandini et al., 2014; Tribaldos et al., 2017), 711 with multiple and regionalized uplift events during the Late Quaternary. For instance, a compilation of palaeoshoreline data (e.g., marine terraces) on a global scale with 712 713 emphasis on the last interglacial stage (130-115 ka) showed that most coastal segments of passive margins were raised relative to sea-level due to an increase in the mean 714 compression of the lithosphere (Pedoja et al., 2011). For the coastal segments to the 715 716 north and south of the study area, they found an average uplift rate of only 0.1 mm/yr. However, a study integrating ichnological and sedimentary facies proposed a higher 717 uplift rate of 0.63 mm/yr for the Paraíba region in the last 60 ka (Gandini et al., 2014). 718 The rise of Late Pleistocene shallow marine deposits at ~38 m a.s.l reported by these 719 authors is compatible with the fluvial incision averaging 43 m (Table 2) of some 720 tributaries from the coastal lowlands of the study area. However, the low incision (140 721 722 m on average) of the rivers in this instance suggests a low-magnitude uplifting (Table 723 2), which resulted in limited topographic changes compared to those from tectonically active settings (Snyder et al., 2000; Wobus et al., 2006; Kirby and Whipple, 2012). 724

A trend of increased uplift rates in the last 30 Myr, with an increased peak since
Middle Miocene (Figure 12), was recently documented for the Precambrian highlands

(Borborema Province) west of the study area, based on inverse modelling of river 727 728 profiles (Tribaldos et al., 2017). In addition, the time scale for fluvial disturbances is generally lower than 5 Myr (Whittaker and Boulton, 2012). These data, together with 729 730 the uplifting of Late Quaternary marine deposits in the Paraíba region (Gandini et al., 2014), led to relate the knickpoint migration and the knickpoints at higher elevations in 731 the Precambrian highlands (Table 2 and Figures 8a) to variations in Late Quaternary 732 uplift rates. The displacement of recent alluvial deposits with offsets up to 100 m 733 734 following the main drainage courses in the Cariatá Graben region (Fonsêca et al., 2020) is consistent with uplifting timing. 735

736

Figure 12.

The differential erosion of the studied relief is probably due to large rivers that 737 738 drained the Precambrian terrains, as these rivers have higher potential of erosion than the smaller and less erosive rivers from the coastal lowlands to the east. However, 739 740 extensive fracturing during and after the pre-rift stage could have enhanced the 741 likelihood of erosion in the low-lying Precambrian rocks to the west. A numerical 742 modelling of reliefs traversed by shear zones compared to the studied Precambrian terrains has indicated that fluvial incision rates close to areas with joints, fractures and 743 744 faults are orders of magnitude faster than in adjacent intact bedrocks (Roy et al., 2015). In the instance of the study area, many tributary streams are (Figure 3) aligned 745 746 according to regional E-W, NE-SW and NW-SE structural trends (Bezerra et al., 2011). For instance, the Cariatá Graben subsided due to the reactivation of normal faults during 747 the Late Quaternary (Bezerra et al., 2008). In addition, the reactivation of normal faults 748 749 and major dextral strike-slip mylonitic belts are recorded along the Patos and Pernambuco shear zones due to transpressional/transtensional deformation (Bezerra et 750 al., 2014, 2020; Nogueira et al., 2015). These pre-existing structures were reactivated 751

several times during the post-rift development of the northeastern Brazilian passive 752 753 continental margin, with the youngest reactivation occurring in the Quaternary (e.g., Bezerra et al., 2008; Marques et al., 2014; Nogueira et al., 2015). Tectonic reactivation 754 755 along the Patos Shear Zone raised the Araripe Plateau (Marques et al., 2014) and Rio do Peixe Basin (Nogueira et al., 2015), located to the west of the study area, in tens of 756 757 meters. In addition, the Pernambuco Shear Zone is a present-day well-known active seismogenic zone that concentrate a high number of low to moderate magnitude 758 759 earthquakes (e.g., Lima Neto et al., 2013; Figure 1). Thus, we propose that the study area was uplifted due to the reactivation of pre-existing structural trends. The tectonic 760 761 reactivation is likely to have triggered the incision wave that migrated upstream into the drainage systems to produce the mobile knickpoints and incised river channels in the 762 763 highlands. This interpretation agrees with the fact that the knickpoints lie upstream of 764 the Patos Shear Zone to the north and upstream of normal faults of the Cariatá Graben to the south (Figure 7). 765

766

5.4. Mechanisms for the passive margin uplift in the Paraíba region

767 There is no general agreement on the mechanisms that cause regional uplifting of passive continental margins. Main hypotheses are mantle forcing (Cox, 1989) and 768 769 increased lithosphere compression due to plate interactions (Leroy et al., 2004). The latter produces stresses with far-field effects even thousands of kilometres far from the 770 stress source (e.g., at plate boundaries). Stresses produced at the plate boundaries are 771 transmitted across the lithosphere and are accommodated by pre-existing weakness 772 773 zones of the crust, such as shear zones. These zones are eventually reactivated to 774 produce differential uplifts (Holdsworth et al., 1997). Several works have demonstrated that far-field stresses can be responsible for intraplate faulting and seismicity in passive 775 margins and intracontinental settings (Zoback, 1992; Assumpção and Araújo, 1993; Hill 776

et al., 1995; Pedoja et al., 2011; Japsen et al., 2012). For instance, the rise of the Andean
mountain range and the continuous lateral growth of the Andean Plateau caused a local
compression that increased the regional compression of the entire South American
platform (Assumpção and Araújo, 1993). Similarly, the long distance (i.e., 3500 km)
transmission of compressional stresses through the lithosphere caused basin inversions
in southeast Australia (Hill et al., 1995).

783 Intraplate compression due to motions of the Mid-Atlantic Ridge (MAR) pushing towards the west, and of the Andean Chain pushing towards the east, was 784 785 interpreted as the main mechanism for post-rift tectonic reactivation and inversion of 786 many sedimentary basins in the northeastern Brazilian passive continental margin (Marques et al., 2014; Nogueira et al., 2015; Vasconcelos et al., 2019; Bezerra et al., 787 2020). These studies proposed that tectonic inversion of the sedimentary basins in this 788 passive margin would have occurred during three main phases of Andean tectonism: 789 Peruvian (80-65 Myr), Incaic (45-28 Myr), and Quechuan (22-0 Myr; Coutand et al., 790 791 2001; Garzione et al., 2008). Whilst the oldest compressional event is difficult to describe due to the long-term erosion (Marques et al., 2014), the record of 792 normal/reverse faults and folds in Miocene and Late Quaternary deposits in this margin 793 794 (Rossetti et al., 2011a, 2011b; Bezerra et al., 2014; Alves et al., 2019; Andrades Filho et al., 2021) agrees with the influence of the Quechuan compressional event. Post-rift 795 (Miocene and Quaternary) tectonic deformation, including both extension and 796 compression, is compatible with the present-day stress-field documented in northeastern 797 Brazil, that is, E-W compression and N-S oriented extension within a strike-slip 798 799 deformational regime (Bezerra et al., 2014).

800 The timing of knickpoint formation in the Paraíba region, presumably of Late 801 Quaternary age, matches with the Quechuan tectonic phase, which occurred when the Andean Plateau was exceptionally high (Garzione et al., 2008; see paleoelevations in Figure 12). The synchronicity between this compressional event and the fluvial disturbances observed in the study area suggests a causative link between far-field stress-induced tectonic uplifting and base-level lowering.

806 6. Conclusion

DEM-based river profile and divide migration analyses helped to decode the 807 post-rift geomorphic history of the South American EPCM in the Paraíba region. These 808 809 methods have the potential to inform short-term events in the evolution of EPCMs, 810 particularly in regions with limited or lack of field-based geological data. The results of the topographic analysis showed that the overall drainage routing pattern of the Paraíba 811 812 region is sustained by the broad configuration of the passive continental margin, with a 813 highland plateau composed of pre-rift bedrocks of varied strengths and tectonic fabrics. In contrast to proposals of stable passive margins over the last million years, this study 814 shows that the late development of the studied EPCM was marked by increased local 815 816 tectonics during the Late Quaternary. This event changed the topographic gradient and likely accelerated the erosion, as informed by a number of transient fluvial features, 817 818 such as: mobile knickpoints; incised fluvial channels with increased k_{sn} ; and divide migration, many of which recorded over lithologies with similar resistances at or close 819 to the uplifted zones. Spatial variations in the uplift rates, potentially driven by far-field 820 compressive stresses concentrated in the shear zones, resulted in base-level lowering, 821 822 headward erosion, and drainage divide upstream migration, with drainage reorganizations taking place along the drainage divides. These results support previous 823 824 studies that the topographic development and final stages of evolution of the eastern South America passive continental margin had a tectonic contribution, at least in its 825 northeastern sector. 826

827 Acknowledgements

This study was financed by the Coordination for the Improvement of Higher 828 Education Personnel - Institutional Program for Internationalization (CAPES-PRINT) 829 830 within the Satellite Applications for Sustainable Development project from the National 831 Institute for Space Research - INPE in Brazil. The School of Geography, Earth and Environmental Sciences of the University of Plymouth is acknowledged for providing 832 833 the workstation for data processing. We would like to thank Jesse R. Zondervan for helping the rock strength and divide migration analyses and for the scientific 834 discussions, which helped to improve an earlier version of this work. Authors are 835 836 scholarship holders of the Council for Scientific and Technological Development 307943/2019-4 (M.M.V.), 163159/2020-3 processes 837 (CNPq), (F.C.A.) and 301343/2018-7 (D.F.R.). Finally, we thank the two anonymous reviewers and the editor, 838 Achim Beylich, for comments that improved this article. 839

840 **References**

- Almeida, F.F.M., Hasui, Y., Brito Neves, B.B., Fuck, R.A., 1981. Brazilian structural
- 842 provinces: An introduction. Earth-Science Reviews. 17, 1–29.
- 843 https://doi.org/10.1016/0012-8252(81)90003-9
- 844
- Alves, F.C., Rossetti, D.F., Valeriano, M.M., Andrades Filho, C.O., 2019. Neotectonics
- 846 in the South American passive margin: Evidence of Late Quaternary uplifting in the
- northern Paraiba Basin (NE Brazil). Geomorphology. 325, 1-16.
- 848 https://doi.org/10.1016/j.geomorph.2018.09.028
- 849
- Andrades Filho, C.O., Rossetti, D.F., Bezerra, F.H.R., 2021. The unsteady post-rift
 stage of the South American passive margin based on the tectono-sedimentary evolution
- stage of the South American passive margin based on the tectono-sedimentof the onshore Paraíba Basin, NE Brazil. Quaternary International.
- 853 https://doi.org/10.1016/j.quaint.2020.10.051
- 854
- Assumpção, M., Araújo, M., 1993. Effect of the Altiplano-Puna plateau, South
- America, on the regional intraplate stresses. Tectonophysics. 221, 475-496.
- 857 https://doi.org/10.1016/0040-1951(93)90174-I
- 858
- Barbosa, J.A., Souza, E., Lima Filho, M., Neumann, V.A., 2003. Estratigrafia da Bacia
 Paraíba: uma reconsideração. Estud. Geol. 13, 89–108.
- 861

Berlin, M.M., Anderson, R.S., 2007. Modeling of knickpoint retreat on the Roan 862 Plateau, western Colorado. Journal of Geophysical Research: Earth Surface. 112, 1–16. 863 864 https://doi.org/10.1029/2006JF000553 865 Bernard, T., Sinclair, H.D., Gailleton, B., Mudd, S.M., Ford, M., 2019. Lithological 866 867 control on the post-orogenic topography and erosion history of the Pyrenees. Earth and 868 Planetary Science Letters. 518, 53-66. https://doi.org/10.1016/j.epsl.2019.04.034 869 Bezerra, F.H., Vita-Finzi, C., 2000. How active is a passive margin? Palaeoseismicity in 870 Northeastern Brazil. Geology. 28, 591-594. https://doi.org/10.1130/0091-871 7613(2000)28%3C591:HAIAPM%3E2.0.CO;2 872 873 874 Bezerra, F.H., Brito Neves, B.B., Corrêa, A.C., Barreto, A.M., Suguio, K., 2008. Late 875 Pleistocene tectonic-geomorphological development within a passive margin - The 876 Cariatá trough, northeastern Brazil. Geomorphology. 97, 555-582. https://doi.org/10.1016/j.geomorph.2007.09.008 877 878 879 Bezerra, F.H., Nascimento, A.F., Ferreira, J.M., Nogueira, F.C., Fuck, R.A., Neves, 880 B.B., Sousa, M.O., 2011. Review of active faults in the Borborema Province, Intraplate South America - Integration of seismological and paleoseismological data. 881 Tectonophysics. 510, 269–290. https://doi.org/10.1016/j.tecto.2011.08.005 882 883 Bezerra, F.H., Rossetti, D.F., Oliveira, R.G., Medeiros, W.E., Neves, B.B., Balsamo, F., 884 Nogueira, F.C., Dantas, E.L., Andrades Filho, C.O., Góes, A.M., 2014. Neotectonic 885 886 reactivation of shear zones and implications for faulting style and geometry in the continental margin of NE Brazil. Tectonophysics. 614, 78-90. 887 888 https://doi.org/10.1016/j.tecto.2013.12.021 889 890 Bezerra, F.H., Castro, D.L., Maia, R.P., Sousa, M.O., Moura-Lima, E.N., Rossetti, D.F., Bertotti, G., Souza, Z.S., Nogueira, F.C., 2020. Postrift stress field inversion in the 891 892 Potiguar Basin, Brazil - Implications for petroleum systems and evolution of the equatorial margin of South America. Marine and Petroleum Geology.111, 88-104. 893 https://doi.org/10.1016/j.marpetgeo.2019.08.001 894 895 896 Bianchi, M.B., M. Assumpção, M.P., Rocha, J.M., Carvalho, P.A., Azevedo, S.L., Fontes, F.L., Dias, J.M., Ferreira, A.F., Nascimento, M.V., Ferreira, Costa, I.S.L., 2018. 897 898 The Brazilian Seismographic Network (RSBR): Improving Seismic Monitoring in 899 Brazil. Seism. Res. Lett. 89, 452-457. 900 doi: 10.1785/0220170227 901 Bierman, P.R., Caffee, M., 2001. Slow rates of rock surface erosion and sediment 902 903 production across the Namib desert and escarpment, Southern Africa. American Journal 904 of Science. 301, 326-358. https://doi.org/10.2475/ajs.301.4-5.326 905 906 Blum, M.D., Törnqvist, T.E., 2000. Fluvial responses to climate and sea-level change: a 907 review and look forward. Sedimentology. 47, 2-48. https://doi.org/10.1046/j.1365-908 3091.2000.00008.x 909 910 Boulton, S.J., Whittaker, A.C., 2009. Quantifying the slip rates, spatial distribution and 911 evolution of active normal faults from geomorphic analysis: field examples from an

- oblique-extensional graben, southern Turkey. Geomorphology. 104, 299–316.
- 913 https://doi.org/10.1016/j.geomorph.2008.09.007
- 914
- Boulton, S.J., Stokes, M., Mather, A.E., 2014. Transient fluvial incision as an indicator
- of active faulting and Plio-Quaternary uplift of the Moroccan high Atlas.
- 917 Tectonophysics. 633, 16–33. https://doi.org/10.1016/j.tecto.2014.06.032
- 918
- Boulton, S.J., Stokes, M., 2018. Which DEM is best for analyzing fluvial landscape
- development in mountainous terrains?. Geomorphology. 310, 168-187.
 https://doi.org/10.1016/j.geomorph.2018.03.002
- 921 922
- Bradley, D.C., 2008. Passive margins through earth history. Earth-Science Reviews.
 91,1-26. https://doi.org/10.1016/j.earscirev.2008.08.001
- 925
 926 Brito Neves, B.B., Riccomini, C., Fernandes, T.M.G., Santanna, L.G., 2004. O sistema tafrogênico terciário do saliente oriental nordestino na Paraíba: um legado Proterozóico.
 928 Revista Brasileira de Geociências. 1, 127–134.
- 929
- Bursztyn, N., Pederson, J.L., Tressler, C., Mackley, R.D., Mitchell, K.J., 2015. Rock
 strength along a fluvial transect of the Colorado Plateau quantifying a fundamental
 control on geomorphology. Earth and Planetary Science Letters. 429, 90–100.
 https://doi.org/10.1016/j.epsl.2015.07.042
- 934
- Carvalho, A.A., Montenegro, A.A.A., Silva, H.P., Lopes, I., Morais, J.E.F., Silva,
 T.G.F., 2020. Trends of rainfall and temperature in Northeast Brazil. Revista Brasileira
 de Engenharia Agrícola e Ambiental. 24, 15–23. https://doi.org/10.1590/18071929/agriambi.v24n1p15-23
- 939
- Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B., Mitrovica,
 J.X., Hostetler, S.W., McCABE, M., 2009. The last glacial maximum. Science. 325,
 710–714. https://doi.org/10.1126/science.1172873
- 943
- Cobbold, P.R., Meisling, K.E., Mount, V.S., 2001. Reactivation of an obliquely rifted
 margin, Campos and Santos basins, southeastern Brazil. AAPG Bulletin. 85, 19251944. https://doi.org/10.1306/8626D0B3-173B-11D7-8645000102C1865D
- 947
 948 Costa, L.R.F., Maia, R.P., Barreto, L.L., Sales, V.C.C., 2020. Geomorfologia do
 949 nordeste setentrional brasileiro: uma proposta de classificação. Revista Brasileira de
 950 Geomorfologia. 21, 185-208. http://dx.doi.org/10.20502/rbg.v21i1.1447
- 951
- Coutand, I., Cobbold, P.R., Urreiztieta, M., Gautier, P., Chauvin, A., Gapais, D.,
 Rossello, E.A., López-Gamundí, O., 2001. Style and history of Andean deformation,
- 954 Puna plateau, northwestern Argentina. Tectonics. 20, 210–234.
- 955 https://doi.org/10.1029/2000TC900031
- 956
- Cox, K.G., 1989. The role of mantle plumes in the development of continental drainage
 patterns. Nature. 342, 873–877.
- 959

- 960 Crosby, B.T., Whipple, K.X., 2006. Knickpoint initiation and distribution within fluvial networks: 236 waterfalls in the Waipaoa River, North Island, New Zealand. 961 962 Geomorphology. 82, 16–38. https://doi.org/10.1016/j.geomorph.2005.08.023 963 Cutler, K.B., Edwards, R.L., Taylor, F.W., Cheng, H., Adkins, J., Gallup, C.D., Cutler, 964 965 P.M., Burr, G.S., Bloom, A.L., 2003. Rapid sea-level fall and deep-ocean temperature 966 change since the last interglacial period. Earth and Planetary Science Letters. 206, 253-271. https://doi.org/10.1016/S0012-821X(02)01107-X 967 968 969 DiBiase, R.A., Whipple, K.X., Heimsath, A.M., Ouimet, W.B., 2010. Landscape form and millennial erosion rates in the San Gabriel Mountains, CA. Earth and 970 971 Planetary Science Letters. 289, 134–144. https://doi.org/10.1016/j.epsl.2009.10.036 972 973 Farr, T.G., Rosen, P.A., Caro, E., Crippen, R., Duren, R., Hensley, S., Kobrick, M., 974 Paller, M., Rodriguez, E., Roth, L., Seal, D., Shaffer, S., Shimada, J., Umland, J., Werner, M., Oskin, M., Burbank, D., Alsdorf, D., 2007. The Shuttle radar topography 975 mission. Rev. Geophys. 45, 1-33. https://doi.org/10.1029/2005RG000183 976 977 978 Fonsêca, D.N., Corrêa, A.C.B., Tavares, B.A.C., Lira, D.R., Barros, A.C.M., Mutzenberg, D.S., 2020. Coupling of tectonic factors and precipitation 979 980 variability as a driver of Late Quaternary aggradation in Northeast Brazil. Earth Surface 981 Processes and Landforms. 45, 3525-3539. 982 Forte, A.M., Yanites, B.J., Whipple, K.X., 2016. Complexities of landscape evolution 983 during incision through layered stratigraphy with contrasts in rock strength. Earth 984 Surface Processes and Landforms. 41, 1736–1757. https://doi.org/10.1002/esp.3947 985 986 987 Forte, A.M., Whipple, K.X., 2018. Criteria and tools for determining drainage divide 988 stability. Earth and Planetary Science Letters. 493, 102–117. https://doi.org/10.1016/j.epsl.2018.04.026 989 990 Forte, A.M., Whipple, K.X., 2019. Short communication: The Topographic Analysis 991 992 Kit (TAK) for TopoToolbox. Earth Surface Dynamics. 7, 87–95. 993 https://doi.org/10.5194/esurf-7-87-2019 994 995 Gallagher, K., Brown, R., Johnson, C., 1998. Fission track analysis and its applications 996 to geological problems. Annual Review of Earth and Planetary Sciences. 26, 519–572. 997 https://doi.org/10.1146/annurev.earth.26.1.519 998 999 Gallen, S.F., Wegmann, K.W., Bohnenstiehl, D.W.R., 2013. Miocene rejuvenation of topographic relief in the southern Appalachians. GSA Today. 23, 4-10. 1000 10.1130/GSATG163A.1 1001 Gallen, S.F., Wegmann, K.W., 2017. River profile response to normal fault growth and 1002 1003 linkage: An example from the Hellenic forearc of south-central Crete, Greece. Earth Surface Dynamics. 5, 161–186. https://doi.org/10.5194/esurf-5-161-2017 1004 1005 1006 Gallen, S.F., 2018. Lithologic controls on landscape dynamics and aquatic species 1007 evolution in post-orogenic mountains. Earth and Planetary Science Letters. 493, 150-160. https://doi.org/10.1016/j.epsl.2018.04.029 1008
- 1009

1010 1011	Gandini, R., Rossetti, D.F., Netto, R.G., Bezerra, F.H.R., Góes, A.M., 2014. Neotectonic evolution of the Brazilian northeastern continental margin based on
1012	sedimentary facies and ichnology. Ouaternary Research (United States). 82, 462–472.
1013	https://doi.org/10.1016/j.yares.2014.07.003
1014	
1015	Garzione, C.N., Hoke, G.D., Libarkin, J.C., Withers, S., Macfadden, B., Eiler, J.,
1016	Ghosh P. Mulch A 2008 Rise of the Andes Science 320 1304-1307
1017	https://doi.org/10.1126/science.1148615
1012	
1010	Giachetta E Willett S.D. 2018 Effects of River Capture and Sediment Flux on the
1019	Evolution of Plateaus: Insights From Numerical Modeling and River Profile Analysis in
1020	the Upper Plue Nile Catebrant Journal of Geophysical Pessarah: Earth Surface, 123
1021	1187 1217 https://doi.org/10.1020/2017/E004252
1022	1187-1217. https://doi.org/10.1029/2017JF004252
1023	Cilchnist A.B. Symmetricial M.A. 1000 Differential demydation and flammal instances
1024	Glichrist, A.R., Summerfield, M.A., 1990. Differential denudation and flexural isostasy
1025	in formation of rifted-margin upwarps. Nature. 346, 739-742.
1026	
1027	Gómez, J., Schobbenhaus, C. & Montes, N.E., compilers. 2019. Geological Map of
1028	South America 2019. Scale 1:5 000 000. Commission for the Geological Map of the
1029	World (CGMW), Colombian Geological Survey and Geological Survey of Brazil. Paris.
1030	
1031	Goren, L., Willett, S.D., Herman, F., Braun, J., 2014. Coupled numerical-analytical
1032	approach to landscape evolution modeling. Earth Surface Processes and Landforms. 39,
1033	522-545. https://doi.org/10.1002/esp.3514
1034	
1035	Goudie, A.S., 2006. The Schmidt Hammer in geomorphological research. Progress in
1036	Physical Geography. 30, 703-718. https://doi.org/10.1177%2F0309133306071954
1037	
1038	Hack, J.T., 1957. Studies of Longitudinal Stream Profiles in Virginia and Maryland.
1039	U.S.G.S. Professional Paper 294-B, pp. 1–91.
1040	
1041	Haviv, I., Enzel, Y., Whipple, K. X., Zilberman, E., Matmon, A., Stone, J., and Fifield,
1042	K. L. 2010. Evolution of vertical knickpoints (waterfalls) with resistant caprock:
1043	Insights from numerical modeling, J. Geophys. Res., 115, F03028.
1044	https://doi.org/10.1029/2008JF001187
1045	
1046	He, C., Rao, G., Yang, R., Hu, L., Yao, O., Yang, C.J., 2019, Divide migration in
1047	response to asymmetric unlift. Insights from the Wula Shan horst North China
1048	Geomorphology 339 44-57 https://doi.org/10.1016/j.geomorph.2019.04.024
1040	Geomorphology. 359, 44 97. https://doi.org/10.1010/j.geomorph.2019.04.024
1045	Henderson F.M. Lewis A. I. 1998 Principles and Applications of Imaging Radar
1050	Manual of remote sensing New York: John Wiley and Sons
1051	Hill K C Hill K A Cooper C T O'Sulliven A L O'sulliven D P Dichardson
1052	M L 1005 Inversion around the Deeg Deein SE Australia In L C Duchanon & D C
1053	M.J., 1995. Inversion around the bass basin, SE Australia. In J. G. Buchanali, & F. G.
1054	buchanan (Eus.), Dasin inversion. Geological Society, London, Special Publications 88,
1055	323-347.
1056	
1057	Holdsworth, K.E., Butler, C.A., Koberts, A.M., 1997. The recognition of reactivation
1058	auring continental deformation. Journal of the Geological Society, London. 154, 73–78.
1059	

- 1060 Japsen, P., Chalmers, J.A., Green, P.F., Bonow, J.M., 2012. Elevated, passive
- 1061 continental margins: Not rift shoulders, but expressions of episodic, post-rift burial and1062 exhumation. Global and Planetary Change. 90-91, 73–86.
- 1063 https://doi.org/10.1016/j.gloplacha.2011.05.004
- 1064
- 1065 King, L.C., 1967. The Morphology of the Earth. Oliver and Boyd, Edinburgh.
- 1066
- Kirby, E., Whipple, K.X., 2012. Expression of active tectonics in erosional landscapes.
 Journal of Structural Geology. 44, 54–75. https://doi.org/10.1016/j.jsg.2012.07.009
- 1069
 1070 Lague, D., 2014. The stream power river incision model: evidence, theory and beyond.
 1071 Earth Surface Processes and Landforms. 39, 38–61. https://doi.org/10.1002/esp.3462
- 1072
 1073 Leroy, M., Dauteuil, O., Cobbold, P.R., 2004. Incipient shortening of a passive margin:
 1074 The mechanical roles of continental and oceanic lithospheres. Geophysical Journal
 1075 International. 159, 400–411. <u>https://doi.org/10.1111/j.1365-246X.2004.02400.x</u>
- 1076
- Lima Neto, H.C., Ferreira, J.M., Bezerra, F.H.R., Assumpção, M.S., Do Nascimento,
 A.F., Sousa, M.O.L., Menezes, E.A.S., 2013. Upper crustal earthquake swarms in São
 Caetano: reactivation of the Pernambuco shear zone and trending branches in intraplate
 Brazil. Tectonophysics 608, 804–811. https://doi.org/10.1016/j.tecto.2013.08.001
- 1081
 1082 Marques, F.O., Nogueira, F.C., Bezerra, F.H., Castro, D.L., 2014. The Araripe Basin in
 1083 NE Brazil: An intracontinental graben inverted to a high-standing horst.
- 1084 Tectonophysics. 630, 251–264. https://doi.org/10.1016/j.tecto.2014.05.029 1085
- 1086 Matos, R.M.D., 1992. The Northeast Brazilian Rift System. Tectonics. 11, 766–791.
- 1087
 1088 Morais Neto, J.M., Green, P.F., Karner, G.D., Alkmim, F.F., 2008. Age of the Serra do
 1089 Martins Formation, Borborema Plateau, northeastern Brazil: constraints from
 1090 apatite and zircon fission track analysis. Bol. Geoc. Petrobras 16, 23–52.
- Neimann, J.D., Gasparini, N.M., Tucker, G.E., Bras, R.L., 2001. A quantitative
 evaluation of Playfair's law and its use in testing long-term stream erosion models.
 Earth Surface Processes and Landforms. 26, 1317–1332.
- 1095

- 1096 Noether, G.E., 1991. Introduction to Statistics: The Nonparametric Way. Stanford, CA:1097 Springer Science Business Media, LLC.
- 1098
- 1099 Nogueira, F.C., Marques, F.O., Bezerra, F.H., Castro, D.L., Fuck, R.A., 2015.
- 1100 Cretaceous intracontinental rifting and post-rift inversion in NE Brazil: Insights from
- the Rio do Peixe Basin. Tectonophysics. 644, 92–107.
- 1102 https://doi.org/10.1016/j.tecto.2014.12.016
- 1103
- Ollier, C.D., Pain, C.F., 1997. Equating the basal unconformity with the palaeoplain: amodel for passive margins. Geomorphology. 19, 1-15.
- 1106
- 1107 Pedoja, K., Husson, L., Regard, V., Cobbold, P.R., Ostanciaux, E., Johnson, M.E.,
- 1108 Kershaw, S., Saillard, M., Martinod, J., Furgerot, L., Weill, P., Delcaillau, B., 2011.

Relative sea-level fall since the last interglacial stage: Are coasts uplifting worldwide? 1109 Earth-Science Reviews. 108, 1–15. https://doi.org/10.1016/j.earscirev.2011.05.002 1110 1111 Peifer, D., Persano, C., Hurst, M.D., Bishop, P., Fabel, D., 2021. Growing topography 1112 due to contrasting rock types in a tectonically dead landscape. Earth Surf. Dyn. 9, 167-1113 1114 181. https://doi.org/10.5194/esurf-9-167-2021 1115 1116 Perron, J.T., Royden, L., 2013. An integral approach to bedrock river profile analysis. Earth Surface Processes and Landforms. 38, 570–576. https://doi.org/10.1002/esp.3302 1117 1118 Roy, S.G., Koons, P.O., Upton, P., Tucker, G.E., 2015. The influence of crustal strength 1119 fields on the patterns and rates of fluvial incision. J. Geophys. Res. Earth Surf., 120: 1120 1121 275-299. https://doi.org/10.1002/2014JF003281 1122 Rossetti, D.F., Bezerra, F.H.R., Góes, A.M., Valeriano, M.M., Andrades Filho, C.O., 1123 Mittani, J.C.R., Tatumi, S.H., Brito Neves, B.B., 2011a. Late Quaternary sedimentation 1124 in the Paraíba Basin, Northeastern Brazil: Landform, sea level and tectonics in Eastern 1125 South America passive margin. Palaeogeography, Palaeoclimatology, Palaeoecology. 1126 1127 300, 191–204. https://doi.org/10.1016/j.palaeo.2010.12.026 1128 Rossetti, D.F., Bezerra, F.H.R., Góes, A.M., Neves, B.B.B., 2011b. Sediment 1129 1130 deformation in Miocene and post-Miocene strata, Northeastern Brazil: Evidence for paleoseismicity in a passive margin. Sedimentary Geology. 235, 172–187. 1131 https://doi.org/10.1016/j.sedgeo.2010.02.005 1132 1133 1134 Rossetti, D.F., Góes, A.M., Bezerra, F.H., Valeriano, M.M., Brito Neves, B.B., Ochoa, F.L., 2012. Contribution to the stratigraphy of the onshore Paraíba basin, Brazil. Anais 1135 da Academia Brasileira de Ciencias. 84, 313-333. https://doi.org/10.1590/S0001-1136 1137 37652012005000026 1138 1139 Rossetti, D.F., Bezerra, F.H., Dominguez, J.M.L., 2013. Late Oligocene-Miocene transgressions along the equatorial and eastern margins of Brazil. Earth-Sci. Rev. 123, 1140 87-112. https://doi.org/10.1016/j.earscirev.2013.04.005. 1141 1142 1143 Schwanghart, W., Scherler, D., 2014. Short Communication: TopoToolbox 2 -MATLAB-based software for topographic analysis and modeling in Earth surface 1144 1145 sciences. Earth Surface Dynamics. 2, 1-7. https://doi.org/10.5194/esurf-2-1-2014. 1146 Smart, J.S., Surkan, A.J., 1967. The relation between mainstream length and area in 1147 drainage basins. Water Resources Research. 3, 963–974. 1148 1149 Snyder, N.P., Whipple, K.X., Tucker, G.E., Merritts, D.J., 2000. Stream profiles in the 1150 Mendocino triple junction region, northern California. GSA Bulletin. 112, 1250-1263. 1151 1152 Strahler, A.N., 1957. Quantitative analysis of watershed geomorphology. Trans. Am. 1153 Geophys. Union 38, 913–920 1154 1155 1156 Strong, C. M., Attal, M., Mudd, S.M., Sinclair, H.D., 2019. Lithological control on the geomorphic evolution of the Shillong Plateau in Northeast India. Geomorphology. 330, 1157 133-150. https://doi.org/10.1016/j.geomorph.2019.01.016 1158

1159	
1160	Souza, D. H., Stuart, F.M., Rodés, A., Pupim, F.N., Hackspacher, P.C., 2019. Controls
1161	on the erosion of the continental margin of southeast Brazil fromcosmogenic10Be in
1162	river sediments. Geomorphology. 330: 163-176.
1163	https://doi.org/10.1016/j.geomorph.2019.01.020
1164	
1165	Tribaldos, R.V., White, N.J., Roberts, G.G., Hoggard, M.J., 2017. Spatial and temporal
1166	uplift history of South America from calibrated drainage analysis. Geochemistry,
1167	Geophysics, Geosystems. 18, 2321–2353. https://doi.org/10.1002/2017GC006909
1168	
1169	Vasconcelos, D.L., Bezerra, F.H., Medeiros, W.E., Castro, D.L., Clausen, O.R., Vital,
1170	H., Oliveira, R.G., 2019. Basement fabric controls rift nucleation and postrift basin
1171	inversion in the continental margin of NE Brazil. Tectonophysics. 751, 23-40.
1172	https://doi.org/10.1016/j.tecto.2018.12.019
1173	
1174	Venables WN, Smith DM. 2002. An Introduction to R. Network Theory, Bristol, UK
1175	
1176	Whipple, K.X., DiBiase, R., Crosby, B., 2013. Bedrock rivers. In: Shroder, J. (Ed.),
1177	Treatise on Geomorphology, vol. 9. Academic Press, San Diego, CA, pp. 550–573.
1178	
1179	Whipple, K.X., Forte, A.M., Dibiase, R.A., Gasparini, N.M., Ouimet, W.B., 2017.
1180	Timescales of landscape response to divide migration and drainage capture:
1181	Implications for the role of divide mobility in landscape evolution. Journal of
1182	Geophysical Research: Earth Surface. 122, 248–273.
1183	https://doi.org/10.1002/2016JF003973
1184	
1185	Whittaker, A.C., Boulton, S.J., 2012. Tectonic and climatic controls on
1186	knickpoint retreat rates and landscape response times. Journal of Geophysical
1187	Research: Earth Surface. 117, 1–19. https://doi.org/10.1029/2011JF002157
1188	
1189	Willemin, J.H., 2000. Hack's law: Sinuosity, convexity, elongation. Water Resources
1190	Research. 36, 3365–3374. https://doi.org/10.1029/2000WR900229
1191	
1192	Willett, S.D., Mccoy, S.W., Taylor Perron, J., Goren, L., Chen, C.Y., 2014. Dynamic
1193	reorganization of River Basins. Science. 343, 1–19.
1194	https://doi.org/10.1126/science.1248/65
1195	
1196	Wobus, C., Whipple, K.X., Kirby, E., Snyder, N., Johnson, J., Spyropolou, K., Crosby,
1197	B., Sheehan, D., 2006. Tectonics from topography: Procedures, promise, and pitfalls.
1198	Special Paper of the Geological Society of America. 398, 55–74.
1199	$\frac{1}{10000000000000000000000000000000000$
1200	Zondervan, J.K., Stokes, M., Boulton, S.J., Telfer, M.W., Matner, A.E., 2020. Rock
1201	strength and structural controls on fluvial erodibility: implications for drainage divide
1202	mobility in a collisional mountain belt. Earth and Planetary Science Letters. 538,
1203	110221. https://doi.org/10.1010/j.epsi.2020.116221
1204	Zahoole M. L. 1002 First and second order rotteres of stress in the lither three The
1205	ZOUACK, WI. L., 1992. FIRST- and second-order patterns of stress in the hunosphere: The World Stress Man Project, I. Goophys. Box. 07, 11702
1205	wond Suess Map Project. J. Geophys. Res. $9/$, 11/03.
1207	nups.//doi.org/10.1029/92JD00152

1210 Figure Captions

1211 1212

Figure 1. Continental scale geological framework and epicentral distribution in the 1213 1214 northeastern Brazilian passive continental margin. Shear zones: Pa = Patos; Pe = Pernambuco. Sedimentary basins: Ar = Araripe; Ig = Iguatú; Ja = Jatobá; Po = Potiguar; 1215 Pn = Parnaíba; RP = Rio do Peixe; SA = Sergipe-Alagoas; Tu = Tucano; Pe = 1216 1217 Pernambuco; and Pb = Paraíba. Geological compartments and structures modified from the Brazilian Geological Survey (CPRM) database (http://geosgb.cprm.gov.br/geosgb) 1218 and the geological map of South America at 1:5.000.000 scale (Gómez et al., 2019). 1219 1220 Instrumental earthquake data (white circles) available for the period of 1955 to 2020 was simplified from the Brazilian Seismic Catalog (Bianchi et al., 2018; 1221 http://moho.iag.usp.br/eq/bulletin/). 1222

Figure 2. Geological context and regional topography of the study area. (a) Geology of the passive continental margin and location of main trunk streams. Trunk rivers: Cu = Curimataú; Ca = Camaratuba; Ma = Mamanguape; Mi = Miriri; Pb = Paraíba. Shaded relief derived from the SRTM-DEM. Geological units simplified from the CPRM geological map at 1:1000000 scale (<u>http://geosgb.cprm.gov.br/geosgb</u>). (b) Swath topographic profile derived from the SRTM-DEM, with the regional morphology of the passive continental margin.

1230 Figure 3. Location of the tributary streams investigated in the study area and major tectonic structures. Pa-sz = Patos shear zone. Trunk streams: Cu = Curimataú; Ca = 1231 Camaratuba; Ma = Mamanguape; Mi = Miriri; and Pb = Paraíba River. Dashed black 1232 1233 line = inferred fault in the basin. RDD = Regional drainage divide. Geological classes geological 1:1000000 1234 simplified from the map at scale from CPRM 1235 (http://geosgb.cprm.gov.br/geosgb). The location of the shear zone and graben-related
1236 normal faults based on Bezerra et al. (2008, 2014).

Figure 4. River profiles of tributary streams from the study area without apparent knickpoints (see location of numbered tributaries in Figure 3). (a-c) River profiles in distance-elevation plots. (d-f) River profiles in χ -elevation plots. Trunk streams with knickpoints are also indicated for comparison. Trunk streams: Cu = Curimataú; Ma = Mamanguape; Pb = Paraiba. River profiles were coloured by their corresponding geology. River profile segments related to artifacts (lakes/reservoirs) are shown only in the distance-elevation graphs for the sake of simplification.

Figure 5. River profiles of tributary streams from the study area showing knickpoints 1244 (slope-break and vertical-step). (a-c) River profiles in distance-elevation plots. (d-f) 1245 1246 River profiles in χ -elevation plots (see the location of numbered tributaries in Figure 3). Trunk streams: Cu = Curimataú; Ma = Mamanguape; Pb = Paraíba. River profile 1247 segments related to artifacts (lakes/reservoirs) shown only in distance-elevation plots 1248 for simplification. River profiles coloured by their corresponding geology. (g-h-i) Log-1249 log slope-area (black dots) and χ -elevation plots with examples of profiles bounded by 1250 vertical-step (g), slope-break (h), and both (i). A_{cr} = critical drainage area. 1251

Figure 6. K_{sn} spatial distribution and rock strength analysis. (a) Drainage network coloured by k_{sn} values, with the location of knickpoints, drainage basins, and spatial contacts of rock groups of similar relative resistance to erosion (see Table 1 for definition of rock groups). K_{sn} values were classified by quantiles due to their skewed distribution. Major trunk streams: Cu = Curimataú; Ca = Camaratuba; Ma = Mamanguape; Mi = Miriri; Pb = Paraíba. RDD = Regional drainage divide. Dashed blue line = inferred area of high relief. (b) Average k_{sn} and standard deviation values for 1259 each rock group (Sr = sedimentary rocks; Mr = metamorphic rocks; and Imr = igneous
1260 and meta-igneous rocks). The number of samples is displayed besides each error bar.

Figure 7. 3D-view of the EPCM based on the SRTM-DEM (vertical exaggeration of 1261 30×), with the location of tributaries, slope-break knickpoints and main tectonic 1262 1263 structures. Purple dot = slope-break knickpoint (numbered tributaries); blue line = tributary streams. HEP = High-elevation plateau; LRS = Low-relief surface; HECP = 1264 high-elevation, chevron-shaped plateau; DR = dome-like relief; CP = coastal plain. 1265 Solid black line = normal fault; dashed black line = dextral strike-slip fault. Dashed 1266 white line = general limits between basement rocks to the west and sedimentary covers 1267 to the east. The location of the Patos shear zone and faults bounding the Cariatá Graben 1268 1269 was based on Bezerra et al. (2008, 2014).

Figure 8. Statistic analysis of knickpoints from the study area. (a) Distribution of average k_{sn} values upstream and downstream of knickpoints and knickpoint elevations from the coastline (E-W). (b) Linear regression between k_{sn} at knickpoint (slope-break and vertical-step) and relative relief.

Figure 9. Statistic analysis of the slope-break knickpoints. Linear regressions: (a) drainage area versus knickpoint elevation; (b) knickpoint elevation versus knickpoint distance from the mouth; (c) total drainage area versus knickpoint distance from the mouth; (d) drainage area above knickpoint versus knickpoint distance from the divide; and (e) drainage area below knickpoint versus knickpoint distance from the mouth. The distributions in c-d-e were fitted by a power-law function in the general form of Hack's law (L = kA^h; Hack, 1957).

1281 Figure 10. χ-map for the complete drainage basins of the passive continental margin 1282 (outlets \geq 1m asl) and the 3D-view of an ongoing river capture. Trunk streams in a: Cu 1283 = Curimataú; Ca = Camaratuba; Ma = Mamanguape; Mi = Miriri; Pb = Paraíba. White 1284 arrow = χ -anomaly. Dashed red line = lithological contacts between basement rocks 1285 (west) and sedimentary deposits (east). RDD = Regional drainage divide. Shaded relief 1286 based on SRTM-DEM. (b) SRTM-DEM 3D-view of an ongoing river capture between 1287 the upper Miriri River and a tributary of the Paraíba River left margin (vertical 1288 exaggeration = 30×). White dot in b = possible capture point.

1289 Figure 11. Cross-divide statistics (average and standard deviation) and delta values based on Gilbert metrics and γ , including the estimated direction of the divide migration 1290 1291 (see the location of divide segments in Figure 10a). (a) Channel elevation; (b) channel 1292 gradient; (c) relative relief at channel head. Red and black lines represent the sides of 1293 the drainage divides. The main geological units of each divide segment are presented on the top of the graph in a. Pb = Precambrian basement rocks; Pg = Paleogene unit; M =1294 Miocene unit; Qa = Quaternary unit. (d) Normalized cross-divide delta values (c.f., 1295 Forte and Whipple, 2018) and uncertainties (standard deviation) for each of the four 1296 1297 metrics derived from headwater channels, illustrating the stable and mobile divide segments in the passive continental margin. 1298

1299 Figure 12. Predicted average uplift rates for the Borborema Province in northeastern Brazil and palaeoelevation estimates for the central Andean Plateau. Black line and grey 1300 1301 area with ± 1 error band = average predicted uplift rates over the last 70 Myr based on inverse river profile modelling. Purple bar = palaeoelevation estimates (vertical axis to 1302 1303 the left) over the past 30 Myr based on carbonate oxygen isotopes. The main phases of Andean orogeny are also indicated above the diagram (i.e., Peruvian: 80-65 Myr, Incaic: 1304 1305 45-28 Myr and Quechuan: 22-0 Myr). Uplift rate and palaeoelevation data simplified from Tribaldos et al. (2017) and Garzione et al. (2008), respectively. Phases of Andean 1306 Orogeny based on Coutand et al. (2001). 1307



Fig.1.







Fig. 3.







Elevation (m)



10 km





Fig.9.



Fig.10.



Fig.11.



Table 1. Summary of the main lithologies exposed in the study area, including theirages and groups according to their relative resistance to erosion.

Age	Main	Geological unit name	Relative resistance to
	lithologies		erosion
*Quaternary	Sandstones	Post-Barreiras	
Miocene	Sandstones and mudstones	Barreiras	Low resistant
Paleogene	Sandstones	Serra do Martins	sedimentary
Late Cretaceous	Sandstones and limestones	Beberibe and Gramame	rocks
Early Cretaceous	Phonolith and rhyolite	Itapororoca	Igneous, more or less massive, highly resistant
Neoproterozoic	Schists	Seridó	Metamorphic rocks, moderately resistant
Neoproterozoic, Mesoproterozoic and Paleoproterozoic	Gneisses, Migmatites and Granites	Suíte Várzea Alegre, Serrinha - Pedro Velho - units 1-2-3-4, Cabaceiras, Salgadinho, Serra de Jabitacá, Sertânea, Recanto - Riacho do Forno, São Caetano and Campina Grande, Esperança, Monte de Gameleiras, Serra Redonda, São Lourenço, Sumé, and Dona Inês and Caxexa plutons	Igneous and meta- igneous, more or less massive, highly resistant

1338 * after Rossetti et al. (2012).

1344	Table 2. Summary of the main characteristics of tributary streams bounded by
1345	knickpoints (slope-break and vertical-step), including fluvial metrics and estimates of
1346	fluvial incision.

River number	River length (km)	Drainage area (km ²)	Elev (m)	Dfm (km)	Dfd (km)	Area up (km ²)	Area down (km ²)	$\begin{array}{c} K_{sn} \operatorname{up} \\ (\mathrm{m}^{0.9}) \end{array}$	±	K_{sn} down (m ^{0.9})	±	K_{sn} at kp (m ^{0.9})	K _{sn} ratio	Fluvial incision (m)
Slope-bre	ak knickj	point				(kiii)	(kiii)			(111)		(111)		(111)
2	16.0	70.6	490	14.2	3.4	4.2	66.4	13.7	0.5	92.3	2.7	12.3	6.7	135
3	9.9	30.7	183	5.0	5.7	7.3	23.4	13.2	0.8	80.2	4.1	13.8	6.1	45
4	13.7	46.5	65	10.0	5.4	9.4	37.1	4.8	1.7	24.6	1.2	12.6	5.1	20
5	117.4	950.6	676	112.0	7.5	9.5	941.1	6.7	0.1	18.2	0.1	24.7	2.7	100
5	117.4	950.6	423	68.6	51.0	238.3	712.3	18.2	0.1	163.6	16.3	28.3	9.0	60
6	48.8	319.4	483	45.7	5.7	7.4	312.0	7.8	0.3	17.0	0.9	19.2	2.2	370
6	48.8	319.4	419	40.8	10.5	17.3	302.1	17.0	0.9	36.5	1.7	41.2	2.1	270
7	33.9	198.9	484	20.6	16.5	51.6	147.3	12.2	0.1	61.9	1.5	31.4	5.1	180
8	43.2	200.1	291	32.2	13.4	34.8	165.3	17.6	0.3	58.6	0.7	49.2	3.3	110
10	16.0	53.3	91	8.2	4.7	8.8	44.5	7.1	0.4	58.9	4.1	9.9	8.3	50
11	28.8	92.0	125	25.6	5.2	5.2	86.8	12.4	0.7	27.7	2.3	10.4	2.2	60
13	77.9	422.3	250	42.6	37.4	125.3	297.0	24.6	0.2	174.3	7.5	35.3	7.1	75
14	94.0	993.8	281	86.8	9.7	14.7	979.1	7.6	0.1	45.4	1.4	28.7	5.9	112
14	94.0	993.8	239	84.8	11.7	18.7	975.1	7.6	0.1	45.4	1.4	22.6	5.9	90
15	64.9	349.3	265	42.7	25.4	120.8	228.5	17.1	0.2	67.2	2.9	19.2	3.9	80
16	38.0	126.4	289	35.6	4.2	3.6	122.8	35.4	0.8	106.7	10.3	23.6	3.0	150
18	29.3	162.9	300	25.3	6.2	7.7	155.2	15.1	0.7	55.9	3.2	38.1	3.7	115
19	33.7	123.4	401	29.9	6.1	8.6	114.9	6.3	0.3	81.5	4.0	20.8	13.0	240
20	13.4	46.5	404	11.2	4.0	4.8	41.7	59.0	4.7	88.1	8.8	46.5	1.5	280
21	22.0	51.2	499	17.9	6.4	6.9	44.3	16.1	0.6	220.8	10.9	49.3	13.7	290
22	20.4	89.3	384	19.1	3.2	2.4	86.9	63.3	2.0	94.1	3.1	39.4	1.5	120
Vertical-s	step knick	point												
1	64.7	666.4	442	56.7	9.7	32.6	633.8	17.3	0.2	70.8	3.6	38.3	4.1	
1	64.7	666.4	371	54.0	12.4	39.5	626.9	70.8	3.6	137.8	6.0	42	1.9	
1	64.7	666.4	231	40.7	25.7	121.7	544.7	20.7	0.2	81.5	7.7	37.9	3.9	
2	16.0	70.6	324	4.5	13.2	63.8	6.7	10.5	0.3	40.4	2.4	34.5	3.8	
5	117.4	950.6	209	44.5	75.0	408.9	541.7	48.5	1.3	498.2	15.8	85.3	10.3	
6	48.8	319.4	317	35.5	15.8	38.8	280.6	36.5	1.7	494.5	16.6	46.9	13.5	
7	33.9	198.9	304	7.6	29.6	173.4	25.5	61.9	1.5	155.2	6.8	60.4	2.5	
9	10.2	46.8	342	5.9	7.2	16.2	30.6	14.3	1.0	263.6	7.1	54.3	18.4	
12	16.7	51.6	75	14.3	4.5	3.1	48.5	6.9	0.9	19.2	0.7	7.2	2.8	
13	77.9	422.3	507	70.8	9.2	15.3	407.0	19.0	0.2	78.2	2.5	20.4	4.1	
15	64.9	512.1	174	25.6	42.5	261.5	250.6	21.0	0.3	101.0	5.2	27.1	4.8	
17	31.7	190.4	73	28.6	4.7	5.8	184.6	11.4	0.5	19.3	0.4	13.2	1.7	
19	33.7	123.4	311	24.3	11.7	36.9	86.6	18.0	0.5	93.1	9.6	24.3	5.2	
20	13.4	46.5	246	6.7	8.6	25.6	20.9	41.3	3.2	171.9	6.4	48.5	4.2	

1347	Elev = Knickpoint elevation; Dfm = knickpoint distance from mouth; Dfd = Knickpoint
1348	distance from divide; Area up = drainage area upstream of knickpoint; Area down =
1349	drainage area downstream of knickpoint; k_{sn} up = k_{sn} upstream of knickpoint; k_{sn} down =
1350	k_{sn} downstream of knickpoint; k_{sn} at kp = k_{sn} at knickpoint.
1351	
1352	
1353	
1354	
1355	
1356	
1357	
1358	
1359	
1360	
1361	
1362	
1363	
1364	

- Table 3. Summary of the unpaired two-samples T and Mann-Whitney U statistical tests
- comparing topographic parameters of slope-break knickpoints recorded in the northern

1367	(N) and souther	n (S) sectors	of the studied	passive	continental	margin.
1307	(1) und bouiner	$\Pi(S)$ sectors	of the studied	pubbive	continental	margin

Unpaired two-samples T-test									
				Shapiro-	Levene	Two	Two		
Topographic	Sector	Average	Standard	Wilk	test:	sample t-	sample		
parameter	Sector	Tiverage	deviation	test: p-	p-	test:	t-test:		
				value	value	t-statistic	p-value		
Flavation	Ν	339.1	200.0	0.337	0.004	0 1 1 0	0.007		
Elevation	S	331.2	85.5	0.168	0.004	0.119	0.907		
* Relative	N	139.1	41.9	0.471	0.003	1.640	0.124		
relief	S	186.7	82.5	0.098	0.005	-1.040	0.124		
** Relative	Ν	208.5	66.8	0.575	0.013	2 610	0.021		
relief	S	326.3	128.0	0.223	0.015	-2.010			
*** Relative	Ν	277.6	106.0	0.122	0.501	-3.060	0.006		
relief	S	421.7	109.0	0.074	0.391				
K	N	28.7	18.1	0.099	0.036	1.620	0.124		
Λ_{Sn}	S	46.1	29.1	0.362	0.030	-1.030			
Mann-Whitne	ey U test								
						Mann	Mann		
Topographic	Sector	Median	Interquartile			Whitney	Whitney		
parameter	Sector	wieulan	range			test: W-	test: p-		
						statistic	value		
Slope	N	2.71	1.9	-	-	17	0 5065		
Stope	S	2.74	0.9	-	-	+/	0.3903		

Relative relief with window radius of 1 (*), 3 (**) and 5 km (***).