Faculty of Science and Engineering

School of Biological and Marine Sciences

2022-12-08

Ecological niche modeling of the Macedonian mouse, Mus macedonicus (Mammalia, Rodentia), under climate change conditions

Dermitas, S

http://hdl.handle.net/10026.1/20044

10.1163/22244662-bja10044 Israel Journal of Ecology and Evolution Brill Academic Publishers

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1	Ecological niche modeling	of the Macedonian mouse.	Mus macedonicus (Mammalia.	Rodentia)	
_	needed in the mouth of the second					

- 2 under climate change conditions
- 3
- 4 Sadik Demirtaş^{1*}, Nevran E. Gündüz¹, David T. Bilton^{2,3}
- 5
- 6 ¹Department of Biology, Faculty of Arts and Sciences, Ondokuz Mayis University, Samsun, Turkey. E-
- 7 mail: sadikd@omu.edu.tr ORCID NO: 0000-0003-0859-7887
- 8 ¹Department of Biology, Faculty of Arts and Sciences, Ondokuz Mayis University, Samsun, Turkey. E-
- 9 mail: eakman@omu.edu.tr ORCID NO: 0000-0001-5777-470X
- 10 ²Marine Biology and Ecology Research Centre, School of Biological and Marine Sciences, University of
- 11 Plymouth, Plymouth PL4 8AA, Devon, UK. E-mail: D.Bilton@plymouth.ac.uk ORCID NO: 0000-0003-
- 12 1136-0848
- ³Department of Zoology, University of Johannesburg, PO Box 524, Auckland Park, Johannesburg 2006,
- 14 Republic of South Africa
- 15

18 Abstract

19 Mus macedonicus Petrov & Ružic, 1983, also known as the Macedonian mouse or Balkan short-tailed 20 mouse, lives in the southern Balkans and the Middle East. While this species is common in Mediterranean 21 ecosystems and is listed as "least concern" by the IUCN, little is known about how its distribution may 22 shift with climate change. This study explores the 'species' potential distribution in three different periods: 23 during the Last Glacial Maximum (LGM) and under current and future climate scenarios, using 24 Maximum Entropy modelling. Modelling was based on 137 georeferenced occurrence records from 25 Macedonia, Bulgaria, Greece, Turkey, Syria, Lebanon, Israel, Iran, Georgia, and Russia and ten 26 bioclimatic variables from the WorldClim database. We show that a combination of precipitation and 27 temperature variables appear to shape the geographical range of the Macedonian mouse and that its 28 predicted distribution during the LGM is consistent with its survival in multiple refugia, as suggested by 29 previous genetic studies. Modelled future distributions are subtly but significantly different from the

^{16 *}Corresponding author. E-mail: sadikd@omu.edu.tr

- current, with population losses and gains in different regions. Our results provide a sound framework for
 future studies on this model species' range dynamics, suggesting that the overall geographical range of *M*.
 macedonicus is relatively stable in the long term.
- 33

Keywords: Bioclimatic variables, Ecological niche modeling, Jackknife test, MaxEnt, Mediterranean
 climate zone

36

37 Introduction

38 Mus macedonicus Petrov & Ružic, 1983 (the Macedonian mouse or Balkan short-tailed mouse) is found 39 in the southern Balkans, Anatolia, Transcaucasia, northwest Iran, and the Levant (Macholán et al. 2007; 40 Macholán 1999). This species is common in Mediterranean climate zones and is rated as "least concern" 41 in the International Union for Conservation of Nature (IUCN) Red List of Threatened Species (Kryštufek 42 & Vohralík 2016). The Macedonian mouse is a mesic species living in habitats containing areas with 43 medium and relatively constant humidity (Haim et al. 1999). As a common species, the Macedonian 44 mouse is also relatively eurytopic, inhabiting various habitats, including long grass, bushes, cultivated 45 lands, woodland borders, and stream banks (Auffray et al. 1990). 46 Molecular studies (Orth et al. 2002; Kryštufek & Vohralík 2016) have revealed that M. macedonicus 47 appears to have persisted in several separate refugia (western Georgia or eastern Azerbaijan, southern 48 Turkey, or around the Euphrates and Tigris and the Levant) during the Last Glacial Maximum (LGM) and 49 the species would appear to be an excellent model organism for studies of refugia and colonization 50 patterns after the LGM in the region. However, to date, no attempt has been made to model the species' 51 potential distribution based on climatic data in the LGM, an approach that could be most illuminating. 52 Indeed, studies on bioclimatically suitable areas and the ecological requirements of *M. macedonicus* are 53 extremely limited, and no attempt has been made to assess how this model species may respond to 54 ongoing anthropogenic climate change. 55 The Köppen–Geiger climate classification system is a widely used ecological niche modeling (ENM) 56 study. This system is divided into five major groups: Tropical (A), dry (B), temperate (C), continental

- 57 (D), and polar (E) (Köppen 1884) in which second and third letters further sub-divide the climate group
- 58 based on seasonal precipitation type and the level of temperature patterns, respectively. Mediterranean

59 climate zone is warm temperate with moderate and relatively constant levels of humidity, and warm

summer; represented as C and D in the Köppen-Geiger classification (Peel et al. 2007; Beck et al. 2018),

61 making it suitable for Macedonian mouse. According to the IUCN distribution map, the known

62 distribution range of *Mus macedonicus* have a total of six sub-climate zones in which Csa (hot-summer

63 Mediterranean), Csb (warm-summer Mediterranean), Cfb (warm-summer humid Mediterranean), Dsa

64 (Mediterranean-influenced hot-summer continental), Dsb (Mediterranean-influenced warm-summer

65 continental), Dfb (Mediterranean-influenced warm-summer humid continental).

66 ENM has been widely used to predict the distribution of taxa across given geographical regions, using 67 climatic variables (Peterson et al. 1999; Soberón & Peterson 2005; Soberón 2010; Peterson et al. 2011). 68 This method is based on the fact that if the species or population's geographic location is known, the 69 potential distribution area at unsampled locations in the past, current, or future can be estimated using 70 environmental variables. Maximum Entropy modeling (MaxEnt) (Phillips et al. 2006; 2017) is one of the 71 most widely used approaches in ENM, producing a spatial habitat suitability map and evaluating the 72 significance of individual bioclimatic variables in shaping a 'species' range. Although several studies have 73 examined morphological, karyological, and genetic variation in *M. macedonicus*, as well as its 74 phylogenetic position within the genus (Ivantcheva & Cassaing 1999; Gündüz et al. 2000; Orth et al. 75 2002; Colak et al. 2006; Macholán et al. 2007; 2008; Rajabi & Azizi 2013; Abi-Said & Karam 2017), 76 very little information, is available about the factors shaping the 'species' range (Auffray et al. 1990; Abi-77 Said & Karam 2017). This study uses a large, mainly genetically-based occurrence dataset for the 78 Macedonian mouse to explore the importance of bioclimatic variables in determining the 'species' range. 79 We explore the past, current, and future potential geographical distribution of the species, addressing the 80 following questions: 1. What are the key bioclimatic variables affecting the distribution of species today? 81 2. What was the potential distribution of *M. macedonicus* in Europe and Asia during the Last Glacial 82 Maximum? 3. Is the modelled potential distribution during the LGM compatible with multiple refugia, 83 suggested by molecular studies? 4. How will M. macedonicus respond to future changes under different 84 climate scenarios?

85 Materials and Methods

86 Species Data Source

87 Occurrence records of *M. macedonicus* were obtained from literature sources (Gündüz et al. 2000; Orth et

- 88 al. 2002; van der Wal et al. 2003; Macholán et al. 2007; 2008; Gündürü 2008; Rajabi & Azizi 2013; Abi-
- 89 Said & Karam 2017; unpublished data from Demirtaş et al. 2022) as well as TRAMEM
- 90 (http://www.tramem.org), and our field observations which together added a significant number of new
- 91 Turkish records for the species (supplementary material, Table SI). All literature records used in this
- 92 study were based on genetic identifications and cranial measurements since *M. macedonicus* can be
- 93 confused with *Mus musculus domesticus* Rutty, 1772 on external morphological characters alone.
- 94 TRAMEM records with habitat information and field observations were based on external morphology
- 95 and habitat type but are considered reliable because we avoided using occurrence records in the human
- 96 settlements and cultivated areas where they coexist with other wild mammals such as Microtus sp.,
- 97 Apodemus sp., and Crocidura sp. Macholán (1999) revealed the Macedonian mouse prefers mainly open
- 98 places with tall and dense vegetation such as grasses, reeds, and bushes, whereas it does not prefer forests
- and human settlements. To correct for sampling bias and ensure high geographical heterogeneity (Pearson
- 100 et al. 2006; Fourcade et al. 2014), multiple presence records within 20 km of each other were spatially
- 101 filtered and removed. Our complete filtered dataset contained 137 georeferenced occurrences (Figure 1).
- 102 Geographic Distance Matrix Generator version 1.2.3 (http://biodiversityinformatics.amnh.org) and
- 103 Google Maps (http://maps.google.com) were used to calculate and verify the occurrence records and the
- 104 spatial geographic distances between localities.



Figure 1. Known distribution of *Mus macedonicus*. Black dots represent occurrence records from our
dataset, the yellow dotted area indicating the 'species' distribution according to IUCN. Note that some

- 108 records fall outside this distribution, particularly in Iran.
- 109 Environmental Variables
- 110 We used bioclimatic variables from WorldClim archive version 1.4 (www.worldclim.org), including a set
- 111 of free climate data with a high resolution (~1 km²), to reconstruct past (LGM, approximately 22 kya),
- 112 current (1960-1990) and future (average 2070) distribution scenarios for *M. macedonicus*. The
- 113 Community Climate System Model (CCSM version 4) was used to predict potential distributions for past
- and future periods. Future climate scenarios used for ENM were representative concentration pathways
- 115 RCP 4.5 (medium greenhouse gas emission scenario) and RCP 8.5 (maximum greenhouse gas emission
- scenario), as presented in the Fifth Assessment Report (IPCC5) of the Intergovernmental Panel on
- 117 Climate Change (IPCC, 2014).
- 118 GeoTIFF raster files were downloaded and converted to ".ascii" format using SDM toolbox version 2.4
- 119 (http://www.sdmtoolbox.org, Brown 2014; Brown et al. 2017) in ArcGIS version 10.2
- 120 (https://www.esri.com). Data processing, mapping, and other file conversions were conducted using
- 121 ArcGIS and DIVA-GIS version 7.5 software (https://www.diva-gis.org). The ".ascii" files were clipped to
- 122 include only 19° to 52° east and 30° to 47° north. Highly correlated independent variables were removed
- from these files to eliminate multicollinearity (which can lead to overfitting problems) (Elith et al. 2011;

124 Dormann et al. 2013; Tanner et al. 2017; Feng et al. 2019) using Pearson correlation coefficients (r >

125 0.90) in the SDM toolbox. A resulting dataset of ten bioclimatic variables was employed for niche

126 modelling (see Table I).

127 Modelling Methods and Statistical Analyses

128 MaxEnt version 3.4.1 (Phillips et al. 2004; 2017) was used to calculate the percentage contribution and

129 permuted importance of the ten selected bioclimatic variables. MaxEnt analyses were conducted with

130 cloglog output format, ten random replicate analyses with the cross-validation method, maximum training

131 presence threshold, random seed, and fade by clamping subsets; all other options were set as defaults. The

132 performance of ENM was tested by receiver operating characteristic analyses (ROC) (Swets 1979) using

- 133 AUC (area under the curve) for ten replicate runs to determine the discriminating capacity of the model.
- 134 The closer the AUC test value is to 1, the better the distinction i.e., the model is sensitive and

descriptive; models with AUC test values between 0.75-1.00 are also considered applicable as they have

high predictive power (Swets 1979; 1988). If the AUC value is less than 0.5, the model is not sensitive

137 enough and is considered unsuitable (Phillips & Dudík 2008). Moreover, we calculated the percentage of

suitable and unsuitable habitats based on the maximum test sensitivity plus specificity threshold (MSS).

139 In addition, the Jack-knife test (Shcheglovitova & Anderson 2013) option in MaxEnt was used to evaluate

- 140 the relative importance of each environmental variable in the model, again with ten replicate runs.
- 141 Results

142 Model Perfomance and Key Bioclimatic Variables

AUC scores for all models were above 0.85 (see Table I), indicating good accuracy. We found that under
the current climate scenario, 70.3% of the habitat was unsuitable and 29.7% suitable. In contrast, all the
remaining scenarios had less suitable habitats than the current one, ranging from 22.8% to 28.0% (Table
I).

147

Table I. AUC values with standard deviation, MSS values, and the percentage of suitable and unsuitablehabitats for LGM, present, and two future climate scenarios.

Distribution	AUC±Sd	MSS	Suitable Habitat	Unsuitable
scenario			(%)	Habitat (%)
LGM	0.865±0.036	0.4277	22.9	77.1

Current	0.859±0.018	0.2908	29.7	70.3
RCP 4.5	0.874 ± 0.042	0.3007	28.0	72.0
RCP 8.5	0.873±0.023	0.3886	22.8	77.2

151 Table II shows the ten bioclimatic variables predicting the current potential distribution of *M*.

152 *macedonicus* across Europe and Asia. Of the bioclimatic variables with a high percentage contribution,

153 BIO12 annual precipitation (41.1%), BIO11 mean temperature of coldest quarter (18.4%), and BIO9

154 mean temperature of the driest quarter (18.3%) appeared to play significant roles in determining the

distribution of *M. macedonicus*. Moreover, bioclimatic variables with the highest permutation importance

- 156 were BIO11 (35.3%), BIO12 (13.7%), and BIO9 (11.9%).
- 157
- 158 Table II

159 Percentage contribution (PC) and permutation importance (PI) values for selected bioclimatic variables.

Abbreviation and variables	РС	PI
BIO1 = Annual mean temperature (°C)	1.8	3.4
BIO2 = Mean diurnal range (mean of monthly (max temp - min temp)) (°C)	0.9	1.5
BIO8 = Mean temperature of the wettest quarter ($^{\circ}$ C)	8.6	7.5
BIO9 = Mean temperature of the driest quarter (°C)	18.3	11.9
BIO11 = Mean temperature of the coldest quarter (°C)	18.4	35.3
BIO12 = Annual precipitation (mm)	41.1	13.7
BIO13 = Precipitation of the wettest month (mm)	0.8	10.4
BIO14 = Precipitation of the driest month (mm)	4.7	7
BIO15 = Precipitation seasonality (coefficient of variation) (mm)	1.2	3.8
BIO19 = Precipitation of the coldest quarter (mm)	4.3	5.6

160

161 According to the Jackknife test (Figure 2), the bioclimatic variable with the highest gain when isolated

162 was BIO19 precipitation of the coldest quarter, and this variable alone provided the most useful

163 information. Moreover, BIO11 mean temperature of the coldest quarter, was the bioclimatic variable

164 whose exclusion produced the largest decrease in gain; this variable contained the most information not

165 incorporated by other variables. These variables suggest that *M. macedonicus* is highly affected by

166 rainfall and temperature during the year's coldest quarter.



Figure 2. Jackknife test for variable importance in the Macedonian mouse. Blue bars = each predictor
variable alone, teal blue bars: how much the total gain is diminished without the given predictor variable,

and red bar: all bioclimatic variables.

172 Current Potential Geographical Distribution of M. macedonicus

173 The results of ENM revealed that *M. macedonicus* prefers to be present in areas where the annual mean

temperature (BIO1) is between 4.63-22.09°C (mean 13.57°C), and annual precipitation (BIO12) between

175 281-1197 mm (mean 593 mm). Additionally, the Macedonian mouse is found in geographical regions

176 where BIO9 is up to 29.16 °C, BIO11 is up to 14.18 °C, and BIO19 is up to 679 mm °C. According to

177 occurrence records, *M. macedonicus* was recorded at altitudes ranging from -195 m to 1965 m (mean 567

- 178 m) above sea level. ENM analyses also revealed that *M. macedonicus* has a potential distributional area
- 179 between latitudes 30.5° to 44.1°N and longitudes 19.3° to 51.4°E under current climatic conditions
- 180 (Figure 3). The distribution map provided by the IUCN has *M. macedonicus* distributed between latitudes
- 181 31.3° to 43.0° N and longitudes 19.3° to 49.3° E (Figure 1).
- 182 The potential distribution preference of species was inferred by comparing potential ecological niches to
- 183 the Köppen-Geiger climate classification. Therefore, the geographical range of species can be generalised
- as mainly Csa (hot-summer), Csb (warm-summer), and partially Dsb (warm-summer humid continental)
- under the Köppen-Geiger classification, temperate in the coast and inner regions and continental on high
- 186 plateaus.



187

Figure 3. Ecological niche model showing the potential geographical distribution of the Macedonian
mouse under current bioclimatic conditions. Color gradient indicates the logistic probability of species
occurrence, increasing from blue (low probability) to red (high probability). Open circles indicate
occurrence records.

193 Past Potential Geographical Distribution of M. macedonicus

194 The predicted potential geographical distribution during the LGM was subtly but significantly different 195 from the current distribution (Figure 4). In the LGM model, 77.1% of the habitat was unsuitable, and 196 22.9% was suitable, suggesting a smaller area of suitable habitat for the species than today. The predicted 197 distribution in the LGM of the species can be generalised as mainly Csa and Csb climate types. Unlike 198 today, ENM inferred that the Macedonian mouse did not prefer the Dsb climate type. Namely, it preferred 199 lower altitude and temperate areas instead of high plateau and cool areas throughout the LGM. Moreover, 200 Csa and Csb sub-climate zones contained more suitable habitats than today. ENM pointed out two evident 201 corridors that allowed the spread of the species. These were a northern corridor reaching the Caucasus 202 along the Black Sea coast in the north and a southern corridor running through northern Iraq to 203 northwestern Iran. While the south corridor is vague, the north is indistinct in the current period. ENM 204 also showed the niche suitability is more favorable, and the predicted distribution is wider than it is in the



- 207 Figure 4. Ecological niche model showing the potential geographical distribution of the Macedonian
- 208 mouse under LGM bioclimatic conditions. Color gradient indicates the logistic probability of species
- 209 occurrence, increasing from blue (low probability) to red (high probability).

210 Future Potential Geographical Distribution of M. macedonicus

- 211 The predicted potential geographical distributions under RCP 4.5 and RCP 8.5 climate change scenarios
- for 2070 are shown in Figures 5 and 6. Both scenarios closely resemble each other. However, in the future
- model with RCP 4.5, 72.0% of the habitat was unsuitable and 28.0% suitable, broadly similar to today. In
- the future model with RCP 8.5, 77.2% of the habitat was unsuitable and 22.8% suitable, significantly
- 215 lower than today. The results of both models showed that the predicted distributions partially overlapped
- 216 with the current model compared to LGM model. Nevertheless, the niche suitability was different inner
- 217 Anatolian Peninsula and along the Aegean Sea and Black Sea coasts. Moreover, the niche suitability had
- 218 increased in northwest Iran and shifted to more eastern areas including in eastern Georgia and central
- 219 Azerbaijan. The two corridors detected in the past model have entirely disappeared.



220 🖌

Figure 5: Ecological niche model showing the potential geographical distribution of the Macedonian

222 mouse under future bioclimatic conditions (RCP 4.5). Color gradient indicates the logistic probability of

- species occurrence, increasing from blue (low probability) to red (high probability).
- 224



- 225
- Figure 6: Ecological niche model showing the potential geographical distribution of the Macedonian
- 227 mouse under future bioclimatic conditions (RCP 8.5). Color gradient indicates the logistic probability of
- 228 species occurrence, increasing from blue (low probability) to red (high probability).
- 229 Discussion

230 ENM provides information about the potential geographical distribution of species using several

bioclimatic variables, representing a 'species' ecological niche within a set of environmental dimensions

232 (Rotenberry et al. 2006). A species realised niche, or geographical range is often smaller than its potential

233 ones due to historical factors, competition, predation, and the realized environment (i.e., current

conditions) across localities (Pulliam 2000; Anderson & Martínez-Meyer 2004). Better knowledge of the

ature of organismal niches is critical for both present and future management and conservation, but such

studies are lacking for most taxa. This is the first study to employ ENM to explore the requirements and

237 distribution of the Macedonian mouse, making several important contributions to our understanding of

this species's ecology and population history.

239 Our MaxEnt results strongly suggest that precipitation is an important variable shaping the range of this 240 species and that the Macedonian mouse is adapted to biomes with a particular annual precipitation regime 241 characterised by warm, hot, dry summers and cool, wet winters. The mean temperature of the driest and 242 coldest quarters of the year also appeared crucial and are likely related to the physiological requirements 243 of *M. macedonicus*, including its capacity for natural hypothermia (Schubert et al. 2010). Studies 244 conducted on the house mouse (M. musculus domesticus), a close relative of the Macedonian mouse, 245 indicate that M. m. domesticus exhibits hypothermic responses to overnight fasting (Hudson & Scott 246 1979) and low ambient temperatures (Tomlinson et al. 2007). Moreover, physiological studies on the 247 house mouse have shown that daily torpor is not restricted to starvation conditions and depends on the 248 environment (Schubert et al. 2010). Our Jackknife test results support the ENM analyses, suggesting that 249 physiological activity may shape the distribution of the species.

The results of ENM based on current bioclimatic variables demonstrate that the Macedonian mouse is aspecies adapted to relatively mesic habitats in the Mediterranean climate zone, with moderate and fairly

constant humidity and warm-hot temperature levels. Studies of nonshivering thermogenesis capacity in

the Macedonian house also conclude that it is a mesic species (Haim et al. 1999; Shabtay et al. 2000). M.

- 254 macedonicus often occupy the same habitat as other mesic species, such as Apodemus flavicollis
- 255 (Melchior, 1834), Apodemus uralensis (Pallas, 1811), and Apodemus mystacinus (Danford & Alston,
- 256 1877). Moreover, the Macedonian mouse is also a relatively eurytopic species, coexisting with similar
- 257 taxa such as Apodemus sylvaticus (Linnaeus, 1758) and Apodemus witherbyi (Thomas, 1902) (Kryštufek

258 & Vohralik 2009).

259 Some of our confirmed occurrence data are outside the range boundaries suggested by the IUCN, 260 particularly in northwest Iran, northeast Bulgaria, and the Dagestan region of Russia. On the other hand, 261 there are no occurrence records of *M. macedonicus* in many apparently climatic suitable regions (see 262 Figure 3), including in the west (Albania), north (Crimean peninsula and Russia's Black Sea coast), and 263 south (Cyrenaica region of northern Libya, Peloponnesos peninsula, Crete and Cyprus) of the 'species' 264 distributional area. A combination of demographic history, ecological and dispersal barriers, and 265 competition may explain the absence of the species in these areas. In much of the Balkans and Crimea, 266 Mus spicilegus Petényi, 1882 may outcompete M. macedonicus, the two species rarely being sympatric in 267 contact zones (Orth et al. 2002). The absence of *M. macedonicus* from Cyrenaica may be due to a failure 268 of the species to disperse to this relatively isolated pocket of suitable habitat. Still, it may also be down to 269 the area occupied by Mus spretus Lataste, 1883, another potential competitor. On Cyprus and Crete Mus 270 cypriacus (Cucchi et al. 2006) and Acomys minous (Bate, 1906) are found, respectively. Again the 271 absence of *M. macedonicus* may result from the 'species' failure to reach these islands and/or the presence 272 of ecologically similar species. It is worth noting that the Macedonian mouse is known from some islands 273 in the Aegean, showing that it is able to cross some marine barriers and establish populations in the 274 absence of close competitors. The remaining two regions (Crimea and Russian Black Sea Coast and 275 Peloponnesos) may have been affected by topographic barriers such as high mountains and water masses 276 or historical factors such as floods, forest fires, and glaciation, all of which would have prevented the 277 spread of *M. macedonicus*. 278 Our LGM range prediction is compatible with previous studies on molecular and morphological variation 279 in M. macedonicus (Hewitt 1996; 1999; Orth et al. 2002; Çolak et al. 2006; Macholán et al. 2007),

280 which suggest that separate glacial refugia were located in the south of the Caucasus and the Black Sea,

southern Turkey and the Levant. Populations from the Levant have been considered to belong to a

separate subspecies (*Mus macedonicus spretoides*) (Orth et al. 2002), which diverged from *M*.

283 macedonicus macedonicus ca. 300,000 BP (Macholán et al. 2007). Time to a most recent common

ancestor for *M. macedonicus macedonicus* populations in Europe, and Asia has been suggested to be in

- the order of 160,000 years, meaning that population subdivision over multiple glacial cycles has been
- essential in generating the genetic diversity seen in this mouse today (Macholán et al. 2007). Our
- 287 bioclimatic model for the LGM provides an important tool for further exploration of the genetic structure

288 of Macedonian mouse populations, suggesting, as it does, other areas where the species is likely to have 289 persisted during glacial episodes. Our ENM results showed that the LGM model significantly differed 290 from the present and future models. Because the sub-climate type in LGM was different from today. 291 Although we showed that the Levant population had a broader distribution in the LGM period, it did not 292 provide enough knowledge to indicate a separate subspecies. However, this foresight may become 293 prominent with more regional analyzes. The two corridors indicated by the LGM map show that the 294 Caucasian and Iranian populations separated from the main population after LGM. Moreover, this 295 separation will be evident in the near future. However, other climate methods and molecular data need to 296 support this prediction. Future climate change scenarios showed significant geographical range 297 differences between the current and future periods, especially in the central Anatolian Peninsula, the 298 Aegean Sea and Black Sea coasts, and northwest Iran. These regions are expected to be drier and 299 experience less precipitation under continued global climate change. However, these analyses also 300 identified several potentially suitable future areas including in eastern Georgia and central Azerbaijan. 301 Such regions were located at lower altitudes than those in western Georgia and northern Azerbaijan, 302 which become unsuitable for *M. macedonicus* under the future IPCC scenarios. Under both scenarios, the 303 range of the Macedonian mouse is likely to retreat in more arid regions such as central Anatolia, the 304 Turkey-Syria border region, and northwest Iran, but it seems likely to expand its distribution range into 305 central Iran and Azerbaijan. Such an observation for this mesic species is consistent with the suggestion 306 that coniferous forests are likely to undergo a contraction in the eastern Mediterranean over the same 307 period (Zeydanlı et al. 2011), and predicted reductions in the potential distributional areas of a variety of 308 taxa in the region in the future (Perktaş et al. 2015; Gür et al. 2017; Malekian & Sadeghi 2019; Örücü 309 2019; Walas et al. 2019). 310 In summary, our study has evaluated habitat suitability and the potential geographical distribution of M.

311 *macedonicus* throughout its range in the Balkans, Anatolia, the Levant, Russia, Transcaucasia, and Iran

312 using Maximum Entropy Modelling based on bioclimatic variables and known occurrence records.

313 Models for past (LGM), current, and future climatic scenarios were robust and pointed to the importance

of a suite of precipitation and temperature variables in determining the 'species' range. Modelled

315 distribution during the LGM is consistent with the view that the species survived cold periods in multiple

316 glacial refugia and suggests the existence of additional refugial areas to those proposed to date based on

317	genetic studies.	Shifts in the	'species'	range are	predicted	under both	modelled f	uture climate	change
	generie staares.		opeeres.		p10010000				

318 scenarios, with population losses and gains suggested across its distribution. Our findings provide a sound

framework to explore Macedonian mouse range dynamics in more detail and highlight the relatively

- 320 limited shifts in the distribution of this species during changing climates something which is in sharp
- 321 contrast to the situation with similar taxa further north in Eurasia.
- 322 Acknowledgments
- 323 We would like to thank Dr. I. Gündüz (Ondokuz Mayis University, Samsun, Turkey) for information
- about the species occurrence in Turkey.
- 325 Declarations
- 326 Conflict of interest
- 327 The authors declare that they have no conflict of interest
- 328 Ethical approval
- 329 This article does not contain any studies with human and animal subjects performed by any of the authors
- 330 References
- 331 Abi-Said MR, Karam SS (2017) Morphological Cranial Study and Habitat Preference of Mus
- 332 macedonicus (Petrov & Ruzic, 1983) (Mammalia: Rodentia) in Lebanon. Jordan J. Biol. Sci 10:235–237.
- Anderson RP, Martínez-Meyer E (2004) Modeling 'species' geographic distributions for preliminary
- 334 conservation assessments: an implementation with the spiny pocket mice (Heteromys) of Ecuador. Biol.
- 335 Conserv 116:167–179. https://doi.org/10.1016/S0006-3207(03)00187-3
- 336 Auffray JC, Tchernov E, Bonhomme F, Heth G, Simson S, Nevo E (1990) Presence and ecological
- 337 distribution of *Mus* "spretoides" Mus musculus domesticus in Israel: Circum Mediterranean vicariance in
- the genus Mus. Z. Saugetierkd 55: 1–10.
- 339 Beck HE, Zimmermann NE, McVicar TR, Vergopolan N, Berg A, Wood EF (2018) Present and future
- 340 Köppen-Geiger climate classification maps at 1-km resolution. Sci. Data 5:180214.
- 341 https://doi.org/10.1038/sdata.2018.214
- 342 Bilton DT, Mirol PM, Mascheretti S, Fredga K, Zima J, Searle JB (1998) Mediterranean Europe as an
- area of endemism for small mammals rather than a source for northwards postglacial colonisation. Proc R
- 344 Soc Lond B Biol Sci 265:1219–1226. https://doi.org/10.1098/rspb.1998.0423

- Brown JL (2014) SDMtoolbox: a python-based GIS toolkit for landscape genetic, biogeographic, and
- 346 species distribution model analyses. Methods Ecol. Evol 5(7):694–700. https://doi.org/10.1111/2041-

347 210X.12200

- 348 Brown JL, Bennett JR, French CM (2017) SDMtoolbox 2.0: the next generation Python-based GIS toolkit
- for landscape genetic, biogeographic, and species distribution model analyses. PeerJ. 5:e4095.
- 350 https://doi.org/10.7717/peerj.4095
- 351 Çolak E, Yiğit N, Sözen M, Çolak R, Özkurt Ş, Kankılıç T, Kankılıç T (2006) The morphological
- analysis of *Mus domesticus* and *Mus macedonicus* (Mammalia: Rodentia) in Turkey. Turk J Zool 30:309-
- **353** 317.
- 354 Dormann CF, Elith J, Bacher S, Carré GCG, García-Márquez JR, Gruber B, Lafourcade B, Leitao PJ,
- 355 Münkemüller T, McClean CJ, Osborne PE, Reneking B, Schröder B, Skidmore AK, Zurell, D,
- 356 Lautenbach S (2013) Collinearity: A review of methods to deal with it and a simulation study evaluating
- 357 their performance. Ecography 36(1):27–46. https://doi.org/10.1111/j.1600-0587.2012.07348.x
- 358 Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE, Yates CJ (2011) A statistical explanation of MaxEnt for
- 359 ecologists. Divers Distrib 17(4):3–57. https://doi.org/10.1111/j.1472-4642.2010.00725.x
- 360 Feng X, Park DS., Liang Y, Pandey R, Papeş M (2019) Collinearity in ecological niche modeling:
- 361 Confusions and challenges. Ecol Evol 9(18):10365–10376. https://doi.org/10.1002/ece3.5555
- 362 Fourcade Y, Engler JO, Rödder D, Secondi J (2014) Mapping species distributions with MAXENT using
- 363 a geographically biased sample of presence data: A performance assessment of methods for correcting
- 364 sampling bias. PLoS ONE 9(5): e97122. https://doi.org/10.1371/journal.pone.0097122
- 365 Gündürü U (2008) Allozymic variations of the genus Mus L., 1758 (Mammalia: Rodentia) in Western
- 366 Black Sea and Thrace. Ankara University, Graduate School of Natural and Applied Sciences, Ankara,
- 367 Turkey.
- 368 Gündüz İ, Tez C, Malikov V, Vaziri A, Polyakov AV, Searle JB (2000) Mitochondrial DNA and
- 369 chromosomal studies of wild mice (*Mus*) from Turkey and Iran. Heredity 84:458–467.
- 370 Gür H, Perktaş U, Gür MK (2017) Do climate-driven altitudinal range shifts explain the intraspecific
- diversification of a narrow ranging montane mammal, Taurus ground squirrels? Mammal Res 63(2):1–15.
- 372 https://doi.org/10.1007/s13364-017-0347-8

- 373 Haim A, Shabtay A, Arad Z (1999) Thermoregulatory and metabolic responses to photoperiod
- 374 manipulations of the Macedonian mouse *Mus macedonicus*, a post-fire invader. J. Therm. Biol 24:279–

375 286.

- Hewitt GM (1996) Some genetic consequences of the ice ages, and their role in divergence and
- 377 speciation. Biol J Linn Soc Lond 58(3):247–276. https://doi.org/10.1006/bijl.1996.0035
- Hewitt GM (1999) Post-glacial re-colonization of European biota. Biol J Linn Soc Lond 68(1-2):87–112.
- 379 https://doi.org/10.1006/bijl.1999.0332
- 380 Hudson JW, Scott IM (1979) Daily torpor in the laboratory mouse, *Mus musculus* var. albino. Physiol
- 381 Zool 52:205–218. https://doi.org/10.1086/physzool.52.2.30152564
- 382 IPCC (2014) Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the
- 383 Fifth Assessment Report of the Intergovernmental Panel on Climate Change. In Pachauri RK, Meyer LA
- (eds) IPCC, Geneva, Switzerland, pp 151.
- 385 Ivantcheva A, Cassaing J (1999) Male-female interactions and socio-sexual isolation of *Mus spicilegus*
- toward two other species of east-European mice. M. macedonicus and M. m. musculus. Comptes Rendus
- 387 Académie Des Sciences, Paris, Sciences Vie, 322:597–605.
- 388 Köppen W (1884) Die Wärmezonen der Erde, nach der Dauer der heissen, gemässigten und kalten Zeit
- 389 und nach der Wirkung der Wärme auf die organische Welt betrachtet (The thermal zones of the earth
- according to the duration of hot, moderate and cold periods and to the impact of heat on the organic
- 391 world). Meteorol. Z. 20(3): 351–360. https://doi.org/10.1127/0941-2948/2011/105
- 392 Kryštufek B, Vohralik V. (2009) Mammals of Turkey and Cyprus. Rodentia II. Cricetinae, Muridae,
- 393 Spalacidae, Calomyscidae, Capromyidae, Hystricidae, Castoridae. Annales Majora, Koper.
- 394 Kryštufek B, Vohralík V (2016) Mus macedonicus (errata version published in 2017). The IUCN Red List
- 395 of Threatened Species 2016: e.T13966A115117069. https://dx.doi.org/10.2305/IUCN.UK.2016-
- **396** 3.RLTS.T13966A22404035.en
- 397 Macholán M (1999) Mus macedonicus Petrov & Ružic, 1983. In: Mitchell-Jones AJ, Amori J,
- Bogdanowicz W et al. (eds) The atlas of European mammals. London: Academic Press, pp 284–285.
- 399 Macholán M, Mikula O, Vohralík V (2008) Geographic phenetic variation of two eastern-Mediterranean
- 400 non-commensal mouse species, *Mus macedonicus* and *M. cypriacus* (Rodentia: Muridae) based on

- 401 traditional and geometric approaches to morphometrics. Zool Anz 247(1):67–80.
- 402 https://doi.org/10.1016/j.jcz.2007.07.003
- 403 Macholán M, Vyskočilová M, Bonhomme F, Kryštufek B, Orth A, Vohralík, V (2007) Genetic variation
- 404 and phylogeography of free-living mouse species (genus *Mus*) in the Balkans and the Middle East. Mol
- 405 Ecol 16(22): 4774–88. https://doi.org/10.1111/j.1365-294X.2007.03526.x
- 406 Malekian M, Sadeghi M (2019) Predicting impacts of climate change on the potential distribution of two
- 407 interacting species in the forests of western Iran. Meteorol Appl 27(1):e1800.
- 408 https://doi.org/10.1002/met.1800
- 409 Orth A, Auffray JC, Bonhomme F (2002) Two deeply divergent mitochondrial clades in the wild mouse
- 410 *Mus macedonicus* reveal multiple glacial refuges south of Caucasus. Heredity 89:353–357.
- 411 Örücü ÖK (2019) Prediction of future and current distribution of *Phoenix theophrasti* Gr. with using
- 412 MaxEnt model and its utilization for planting design. Turk J For 20(3):274–283.
- 413 https://doi.org/10.18182/tjf.613205
- 414 Pearson RG, Thuiller W, Araújo MB, Martinez-Meyer E, Brotons L, Mcclean C, Miles L, Segurado P,
- 415 Dawson TP, Lees DC (2006) Model based uncertainty in species range prediction. J. Biogeogr 33:1704–
- 416 1711. https://doi.org/10.1111/j.1365-2699.2006.01460.x
- 417 Peel MC, Finlayson BL, McMahon TA (2007) Updated world map of the Köppen–Geiger climate
- 418 classification. Hydrol. Earth Syst. Sci. 11(5):1633–1644. https://doi.org/10.5194/hess-11-1633-2007
- 419 Perktaş U, Gür H, Sağlam İK, Quintero E (2015) Climate-driven range shifts and demographic events
- 420 over the history of Kruper's Nuthatch Sitta krueperi. Bird Study 62(1):14–28.
- 421 https://doi.org/10.1080/00063657.2014.977220
- 422 Peterson AT, Soberón J & Sánchez-Cordero V (1999). Conservatism of ecological niches in evolutionary
- 423 time. Science, 285:1265–1267. http://dx.doi.org/10.1126/ science.285.5431.1265
- 424 Phillips SJ, Anderson RP, Dudík M, Schapire RE, Blair ME (2017) Opening the black box: an open-
- 425 source release of Maxent. Ecography 40(7):887–893. https://doi.org/10.1111/ecog.03049
- 426 Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species geographic
- 427 distributions. Ecol Modell 190: 231–259. https://doi.org/10.1016/j.ecolmodel.2005.03.026
- 428 Phillips SJ, Dudík M (2008) Modeling of species distributions with Maxent: New extensions and a
- 429 comprehensive evaluation. Ecography 31:161–175. https://doi.org/10.1111/j.0906-7590.2008.5203.x

- 430 Phillips SJ, Dudík M, Schapire RE (2004) A maximum entropy approach to species distribution
- 431 modeling. In Proceedings of the Twenty-First International Conference on Machine Learning. Banff, AB:
- 432 ACM Press, pp. 655–662.
- 433 Phillips SJ, Dudík M, Schapire RE (2017) MaxEnt software for modeling species niches and
- 434 distributions. Version 3.4.1. Available from:
- 435 https://biodiversityinformatics.amnh.org/open_source/maxent
- 436 Pulliam HR (2000) On the relationship between niche and distribution. Ecol Lett 3:349–361.
- 437 https://doi.org/10.1046/j.1461-0248.2000.00143.x
- 438 Rajabi MH and Azizi V (2013) Phylogeographical study of short-tailed mouse species (Mus macedonicus
- 439 Petrov & Ruzic, 1983) in North West of Iran. J Anim Res. 26(3):289–297.
- 440 Rotenberry JT, Preston KL, Knick ST (2006) GIS-based niche modeling for mapping species habitat.
- 441 Ecology 87:1458–1464. http://dx.doi.org/10.1890/0012-9658(2006)87[1458:gnmfms]2.0.co;2
- 442 Schubert KA, Boerema AS, Vaanholt LM, de Boer SF, Strijkstra AM, Daan S (2010) Daily torpor in
- 443 mice: high foraging costs trigger energy-saving hypothermia. Biol Lett 6(1):132–135.
- 444 https://doi.org/10.1098/rsbl.2009.0569
- 445 Shabtay A, Haim A, Arad Z (2000) Metabolic rate and thermoregulation in the Macedonian mouse Mus
- 446 *macedonicus*. Isr. J. Zool 46:305–312.
- 447 Shcheglovitova M, Anderson RP (2013) Estimating optimal complexity for ecological niche models: a
- 448 jackknife approach for species with small sample sizes. Ecol Model 269:9–17.
- 449 https://doi.org/10.1016/j.ecolmodel.2013.08.011
- 450 Swets JA (1979) ROC analysis applied to the evaluation of medicalimaging techniques. Invest Radiol
- 451 14:109–121. https://doi.org/10.1097/00004424-197903000-00002
- 452 Swets JA (1988) Measuring the accuracy of diagnostic systems. Science 240(4857):1285–1293.
- 453 https://doi.org/10.1126/science.3287615
- 454 Tanner EP, Papeş M, Elmore RD, Fuhlendorf SD, Davis CA (2017) Incorporating abundance information
- 455 and guiding variable selection for climate-based ensemble forecasting of species' distributional shifts.
- 456 PLoS ONE:12, e0184316. https://doi.org/10.1371/journal.pone.0184316
- 457 van der Wal A, Willemsen J, de Wit J (2003) Mammal Survey Eastern Rhodopes Bulgaria. Uitgave van
- 458 de Vereniging voor Zoogdierkunde en Zoogdierbescherming (VZZ).

- 459 Walas Ł, Sobierajska K, Ok T, Dönmez AA, Kanoğlu SS, Dagher-Kharrat MB, Douaihy B, Romo A,
- 460 Stephan J, Jasińska AK, Boratyński A (2019) Past, present, and future geographic range of an oro-
- 461 Mediterranean Tertiary relict: The juniperus drupacea case study. Reg Environ Change 19:1507–1520.
- 462 https://doi.org/10.1007/s10113-019-01489-5
- 463 Zeydanlı U, Turak A, Bilgin C, Kınıkoğlu Y, Yalçın S, Doğan H (2011) İklim Değişikliği ve Ormancılık:
- 464 Modellerden Uygulamaya. Ankara: Doğa Koruma Merkezi
- 465

Figure Legends

467

468 Figure 1. Known distribution of *Mus macedonicus*. Black dots represent occurrence records from our

- dataset, the yellow dotted area indicating the 'species' distribution according to IUCN. Note that some
- 470 records fall outside this distribution, particularly in Iran.
- 471
- 472 Figure 2. Jackknife test for variable importance in the Macedonian mouse. Blue bars each predictor
 473 variable alone, teal blue bars how much the total gain is diminished without the given predictor variable,
 474 and red bar all bioclimatic variables.
- 475

476 Figure 3. Ecological niche model showing the potential geographical distribution of the Macedonian

477 mouse under current bioclimatic conditions. Color gradient indicates the logistic probability of species

478 occurrence, increasing from blue (low probability) to red (high probability). Open circles indicate

479 occurrence records.

480

481 Figure 4. Ecological niche model showing the potential geographical distribution of the Macedonian

482 mouse under LGM bioclimatic conditions. Color gradient indicates the logistic probability of species

483 occurrence, increasing from blue (low probability) to red (high probability).

484

- 485 Figure 5: Ecological niche model showing the potential geographical distribution of the Macedonian
- 486 mouse under future bioclimatic conditions (RCP 4.5). Color gradient indicates the logistic probability of

487 species occurrence, increasing from blue (low probability) to red (high probability).

488

489 Figure 6: Ecological niche model showing the potential geographical distribution of the Macedonian

- 490 mouse under future bioclimatic conditions (RCP 8.5). Color gradient indicates the logistic probability of
- 491 species occurrence, increasing from blue (low probability) to red (high probability).
- 492

493 Supplementary Material

494 Table S1. Occurrence records of Macedonian mouse