Faculty of Health: Medicine, Dentistry and Human Sciences

School of Health Professions

2021-10

# Benign paroxysmal positional vertigo in acute traumatic brain injury patients data from a multi-centre prospective randomised feasibility study

#### Smith, R

http://hdl.handle.net/10026.1/20068

10.1016/j.jns.2021.117727 Journal of the Neurological Sciences Elsevier BV

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

### **Background and aims**

Benign paroxysmal positional vertigo (BPPV) is the most common peripheral vestibular diagnosis in acute TBI. However, there is a paucity of acute prospective data regarding the most effective treatment, optimal time to treat and recurrence. Given such uncertainties we aimed to investigate the feasibility of assessing and treating BPPV in acute TBI.

# Methods

Participants were recruited from two major trauma centres in London, UK. Inclusion criteria included those over the age of 18, with a closed head injury (as noted by CT scan) and were an inpatient on a major trauma or outlying ward. Importantly, subjective dizziness was not a prerequisite for inclusion. Following consent, patients were assessed for BPPV by trained ward therapists. Those with BPPV were randomly allocated to one of three interventions (repositioning manoeuvres, Brandt Daroff exercises or advice) and were followed up at two time intervals.

# Results

95 patients have consented to take part in the study. Of those, BPPV was diagnosed in twenty eight (mean age 50 years; 20 males, 8 females). 62% of patients had a moderate-severe head injury. 14 (50%) had unilateral BPPV, of which 71% had right sided posterior canal BPPV. 12 patients (43%) had bilateral posterior canal BPPV.

### Conclusions

This is the first data from a prospective study investigating BPPV in acute TBI. Initial data suggests just under a third of patients with acute TBI have BPPV. Interestingly, a large number of patients had bilateral BPPV. Data from this study will inform a more definitive randomised controlled trial.