

2022-11

Artificial reefs in the Arabian Gulf: Benefits, challenges and recommendations for policy-makers

Bartholomew, A

<http://hdl.handle.net/10026.1/20402>

10.1016/j.rsma.2022.102723

Regional Studies in Marine Science

Elsevier

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Artificial Reefs in the Arabian Gulf: Benefits, Challenges and Recommendations for Policy-Makers

Aaron Bartholomew^{1*}, John A. Burt², Louise B. Firth³

¹ Department of Biology, Chemistry and Environmental Science
American University of Sharjah
P.O. Box 26666
Sharjah, United Arab Emirates

² Center for Genomics and Systems Biology
New York University – Abu Dhabi
PO Box 129188
Abu Dhabi, United Arab Emirates

³ School of Biological and Marine Sciences
University of Plymouth
Drake Circus, Plymouth
PL48AA, United Kingdom

*Corresponding author: abartholomew@aus.edu

Abstract

Artificial reef (AR) deployment has increased dramatically in the Arabian Gulf in recent years, and will likely continue as Gulf nations continue to develop their coastlines and expand fisheries. Unfortunately, there is little publicly-available information about AR programs in the Gulf, including information about management goals and program success. ARs can provide economic, social and ecological benefits, but they also have underappreciated risks associated with them. Benefits include increasing short-term catch rates for fisheries, increasing tourism, enhancing and protecting biodiversity and providing ecological services. Risks include exacerbating regional overfishing in the long-term, facilitating the spread of invasive species, altering benthic habitat around the AR, contributing to marine pollution and creating habitats that are “sinks” for larval fish. This paper provides recommendations for managers that are considering whether to initiate AR programs in the Arabian Gulf. Deployment of ARs should not be used as an excuse to allow the destruction or degradation of natural habitat, since ARs are not surrogates of natural habitat. Managers should define clear, explicit management goals in the planning stages of the reef project, and then design and deploy the reef to meet those particular goals. Managers should also set quantifiable objectives for each goal, and implement long-term monitoring programs to determine whether the reef is successful in meeting its goals. Finally, managers should disseminate the results of the monitoring program and share “lessons learned”. Implementation of these recommendations will help to guide future sustainable AR programs in the Arabian Gulf and elsewhere.

Keywords Artificial reef, Arabian Gulf, Management, Goals, Benefits, Recommendations

1 Introduction

1.1 Pressures on Arabian Gulf Coastal Ecosystems

A mosaic of highly productive and diverse marine ecosystems border the Arabian Gulf, providing critical goods and services to coastal populations (Vaughan et al. 2019). These ecosystems encompass highly diverse habitats in a region that is typically recognized for its comparatively low-diversity terrestrial deserts (Sheppard et al. 1992, Burt 2014). Mangrove forests, mudflats and sabkha (salt flat) ecosystems support extensive coastal and subtidal food webs (Sheppard et al. 1992, Hegazy 1998, Barth and Böer 2002). Important nearshore subtidal ecosystems include coral reefs, seagrass meadows, algal beds and oyster reefs that cover extensive areas of seafloor (Vaughan et al. 2019). Economically, fisheries are the second-most important economic sector in the Gulf after oil and gas, and these marine habitats are critical for numerous commercially-important species (Siddeek et al. 1999, Van Lavieren et al. 2011, Vaughan et al. 2019). These ecosystems are also important for tourism (Nouri et al. 2008, Ryan et al. 2012, Blignaut et al. 2016), they protect against coastal erosion (Erftemeijer et al. 2020) and they effectively sequester carbon (Campbell et al. 2015, Schile et al. 2017, Rabaoui et al. 2019).

Nations bordering the Arabian Gulf have population growth rates that are almost twice the global average, and most of this growth has occurred in cities bordering coasts (Van Lavieren et al. 2011). This rapid population growth, along with global climate change, has put immense pressure on all coastal ecosystems of the Gulf, and has resulted in widespread habitat degradation and loss (Sale et al. 2011). Threats to Gulf coastal ecosystems include dredging, land reclamation and related development

(Sheppard et al. 2010; Sale et al. 2011; Burt 2014), water pollution and desalination-brine discharge (Freije 2015; Alshahri 2017; Le Quesne et al. 2021) and eutrophication leading to hypoxia (De Verneil et al. 2021; Lachkar et al. 2022), harmful algal blooms and mass die-offs of fishes and other marine organisms (Bauman et al. 2010; Chatziefthimiou et al. 2021). Furthermore, the Gulf fishing industry has expanded rapidly in recent decades to support the growing human populations. The combination of increased fishing pressure and the degradation or loss of critical ecosystems that support fisheries has led to dramatic declines in regional fish stocks (Al-Abdulrazzak 2015; Buchanan et al. 2019; EAD 2021). Unfortunately, fisheries management practices and law enforcement are not adequate to reverse these trends in many parts of the Gulf (Siddeek et al. 1999; Al-Abdulrazzak 2015; Aljasmí 2017).

Global climate change is exacerbating the effects of ecosystem degradation and overfishing in the Gulf, and will continue to have widespread negative impacts in the future (Burt et al. 2021; Lincoln et al. 2021). Some coastal ecosystems in the Gulf are threatened by “coastal squeeze” due to rising sea levels (Melville-Rea et al. 2021). Coral bleaching events associated with elevated sea surface temperatures are increasingly frequent, resulting in reduced coral cover, enhanced risk of regional extinctions and reduced reef complexity (Riegl and Purkis 2015; Riegl et al. 2018; Burt et al. 2019). Reef-associated fish are impacted by this habitat loss (Buchanan et al. 2016, 2019), and global warming may also affect fish size, growth, fecundity and susceptibility to disease (Donelson et al. 2010; Brandl 2020; D’Agostino et al. 2021). Overall, the Gulf represents a highly vulnerable marine environment that is challenged by a wide array of local and global factors that threaten to further degrade the integrity of its coastal ecosystems.

1.2 Management Responses to Environmental Change in the Gulf

Gulf leaders have become increasingly aware of both the importance of coastal ecosystems and the threats to them, and various marine conservation efforts have been instituted in recent years (Khan 2007; Hamza and Munawar 2009; Lamine et al. 2020). Marine environmental management actions have included establishing marine protected areas (Al-Cibahy et al. 2012; Van Lavieren and Klaus 2013), adopting ecosystem-based management approaches (Burt et al. 2017; Mateos-Molina et al. 2021), developing more sustainable fishing practices (Grandcourt 2012), enhancing environmental regulations and strengthening environmental impact assessment processes (Naser 2015). Gulf nations have also adopted various regional and international frameworks to improve sustainable development practices, address climate change, prevent pollution and protect biodiversity, including the Regional Organization for Protection of Marine Environment (ROPME) (Nadim et al. 2008; Lincoln et al. 2021), the United Nations 2030 Sustainable Development Agenda and many others (Al Saidi 2021).

1.3 Artificial Reefs as a Tool for Marine Management in the Gulf

Better conservation management, stronger regulations and stricter enforcement will help slow the degradation of marine ecosystems, but there is also a need for habitat restoration and enhancement efforts to compensate for habitat that has been degraded or lost (Burt and Bartholomew 2019; Alghunaim et al. 2020; Erftemeijer et al. 2020). Artificial reefs (ARs) are a management tool that can be deployed for various ecological and economic benefits in the Gulf. To assess the current status of artificial reef use and performance, we performed a systematic literature review to explore patterns and trends in AR-related research in the Arabian Gulf region. We searched Google Scholar for “artificial reef” combined with “Arabian Gulf”, “Persian Gulf” and the names of each country of the Gulf, and we included “gray literature” sources. Some of the Gulf AR programs are described in detail, and

information about these reefs is included in Table 1. In other cases, ARs are only mentioned very briefly. These are listed below by country, but are not included in Table 1. (Kuwait: Maghsoudlou et al. 2008), (Bahrain: Shams 2002), (Qatar: Abdel-Moati et al. 2006; Maghsoudlou et al. 2008; Richer 2008; Rosciszewski-Dodgson and Cirella 2022), (Oman: Alhabsi 2013), (Iran: Ajdari and Ajdari 2016) (United Arab Emirates: Salahuddin 2006; EAD 2016; Gibling 2013; Mateos-Molina et al. 2020). In some cases, one or more management goals were stated for the AR program, and this information was incorporated into Figure 1. In several cases no management goals were explicitly stated. Figure 1 shows that artificial reefs have been deployed in the Gulf for a variety of purposes, including: promoting fisheries (Downing et al. 1985; Harris and Schroder 2001; Ajdari et al. 2011; Azhdari et al. 2012; Ajdari and Ajdari 2016; Loughland and Bass 2016), enhancing biodiversity (Salahuddin 2006; Hopkins 2007; Al-Cibahy et al. 2009; Loughland and Bass 2016), promoting dive tourism (Harris and Schroder 2001; Gibling 2013), providing additional reef habitat to compensate for lost natural hard bottom habitat (EAD 2016; Mohamed 2021), preventing trawling (Shams 2002) and creating hard bottom habitat to transplant corals that would otherwise be destroyed by development (Deb et al. 2014). Most of these are common goals for artificial reef projects worldwide, but the use of artificial reefs to support large-scale coral transplantation efforts in Qatar (Deb et al. 2014; Rosciszewski-Dodgson and Cirella 2022) is unusual.

For many Gulf AR programs there is little publicly-available information available, and no stated management goals. When stated, management goals are often imprecise, and in most cases there is insufficient monitoring after deployment to determine if any stated goals were met. Marine managers also usually do not account for possible negative consequences of the AR program, so no efforts are made to prevent or mitigate these problems (Feary et al. 2011). Without explicitly-stated goals and adequate monitoring, managers may claim their reef projects are successful, when in fact they are not meeting management objectives (Baine 2001). AR programs may even exacerbate some of the existing problems managers are trying to fix, or create new unanticipated problems (Grossman et al. 1997; Bulleri and Airoidi 2005; Morley et al. 2008).

The purposes of this paper are threefold. Firstly, we summarize the various ways in which ARs have been used as tools for marine management globally and in the Gulf, and we discuss possible benefits associated with their use. Secondly, we highlight various risks associated with ARs if they are not properly designed, deployed and managed. Thirdly, we use this information to propose six policy recommendations for managers considering AR programs, in order to minimize risks and maximize benefits of future AR programs in the Gulf.

2 Uses and Benefits Associated with ARs

2.1 Economic and Cultural Uses and Benefits

This section of the paper is summarized in Fig. 2.

ARs have been deployed successfully to promote various economic interests. ARs have been deployed in the Gulf (Table 1; Ajdari and Ajdari 2016) and elsewhere (Delmendo 1990; Zhang et al. 2005) to attract commercially-important species of fish, or to provide them with habitat and refuge spaces (Charbonnel et al. 2002; Leitao et al. 2008a). This can benefit local fisheries in the short term, but there is a long-standing debate about whether ARs “produce” new fish biomass for local fisheries or simply attract existing fish to them (*sensu* “attraction versus production” Bohnsack and Sutherland 1985; Grossman et al. 1997; Roa-Ureta et al. 2019). Efforts to design and create reefs that produce fish have

1
2
3
4 been less successful than reefs that simply attract fish, although there are instances where ARs seem to
5 contribute to increased net production of fish, even with the increased fishing pressure associated with
6 the ARs (Roa-Ureta et al. 2019). ARs can increase fish production by providing refuges from predation, if
7 refuge habitat is limiting (Simon et al. 2011; Roa-Ureta et al. 2019; Folpp et al. 2020), or by providing
8 habitat for planktivorous fish that would not live in an area without the reef (Rilov and Benayahu 2002;
9 Champion et al. 2015; Smith et al. 2016). ARs can also increase production by providing suitable
10 breeding habitat (Pondella et al. 2002; Kim et al. 2011). ARs can increase feeding opportunities, leading
11 to increased growth rates and more fish biomass production (Fabi et al. 2006; Love et al. 2007; Cresson
12 et al. 2014). ARs can also alter water circulation patterns to improve larval recruitment to an area (Cenci
13 et al. 2011) or provide suitable recruitment habitat for planktonic fish larvae that would otherwise die
14 because they would be transported to regions lacking hard-bottom habitat (Emery et al. 2006).
15
16
17
18

19 Improving the design of ARs may increase the likelihood that they produce fish, rather than just
20 attract fish. Recommendations include creating vertically-oriented reefs (Rilov and Benayahu 2002), and
21 creating reefs with void spaces that are the appropriate size to provide refuge spaces for fish, especially
22 juvenile fish that normally have high mortality rates (Charbonnel et al. 2002; Patranella et al. 2017; Burt
23 and Bartholomew 2019). Placing reefs “downstream” of natural reefs in areas without natural hard-
24 bottom habitat may provide settlement sites for planktonic fish larvae that would otherwise not survive
25 (Carr and Hixon 1997).
26
27
28

29 ARs can also support bivalve fisheries by providing attachment sites and increasing production
30 (Steimle et al. 2002; Macreadie et al. 2011; Lipcius and Burke 2018), and ARs can be designed to support
31 aquaculture for benthic invertebrates (Fabi et al. 1989; Xu et al. 2017). ARs have also been used
32 successfully to promote marine biology education (Seidelin et al. 2018), to create surfing locations (Black
33 and Mead 2001) and to promote dive tourism (Table 1; Salahuddin 2006; Gibling 2013; Belhassen et al.
34 2017). Dive tourism associated with some ARs, like large shipwrecks, can bring substantial economic
35 benefits to local economies (Broughton 2012).
36
37
38

39 *2.2 Ecological Uses and Benefits*

40 This section of the paper is summarized in Fig. 3.

41
42 ARs have also been used to promote various ecological goals. There are examples of large-scale
43 artificial marine structures that have had significant positive ecological effects, including producing fish
44 larvae (Stephens and Pondella 2002), increasing biodiversity (Loughland and Bass 2016), protecting
45 valuable natural habitat from trawling (Charbonnel and Bachet 2010) creating hard-bottom settlement
46 sites for ecological-valuable species like corals and kelp (Burt et al. 2011; Schroeter et al. 2015) and
47 creating transplantation sites for coral that would otherwise be destroyed by development projects
48 (Table 1; Rosciszewski-Dodgson and Cirella 2022). There are also many successful examples of “greening
49 of gray infrastructure”, which refers to ecoengineering coastal infrastructure to enhance its ecological
50 benefits (Firth et al. 2014; Airoidi et al. 2021; O’Shaughnessy et al. 2020). ARs may be used to increase
51 biodiversity as they support various reef-associated fish and invertebrates (Maghsoudlou et al. 2008;
52 Mousavi et al. 2015; EAD 2016). ARs provide attachment sites for habitat-forming species of macroalgae,
53 bivalves, corals, etc. which contribute to increased biodiversity as well (Farinas-Franco and Roberts
54 2014; Alghunaim et al. 2020; Rosciszewski-Dodgson and Cirella 2022). The organisms associated with
55 ARs can also provide ecological services such as water filtration (Chojnacki and Ceronik 1996; Israel et al.
56 2017) and carbon sequestration (Hall et al. 2011; Pi-Hai et al. 2014). ARs have also been successfully
57
58
59
60
61
62
63
64
65

used to prevent bottom trawling, which protects critical benthic habitat from destruction (Shams 2002; Relini et al. 2007; Ramm et al. 2021), and can help prevent overfishing (Polovina 1990).

ARs are often deployed to provide ecological benefits as a part of mitigation plans for coastal development projects or accidental destruction of natural habitats (Richer 2008; Levrel et al. 2012; Al-Horani and Khalaf 2013). The “mitigation hierarchy” is used by managers to reduce negative impacts of development projects. The first step in the hierarchy is to avoid negative impacts, followed by reducing negative impacts, followed by providing offsets for any remaining unavoidable impacts (Cabrera and Castro 2018). ARs have been repeatedly shown to support diverse and abundant communities of fish and invertebrates. In some cases, diversity and abundances can be the same or higher on ARs compared with natural hard bottom habitat in the Gulf (Al Cibahy et al. 2009; Azhdari et al. 2012, Burt et al. 2013; Ajdari and Ajdari 2016) and elsewhere (Perkol-Finkel and Benayahu 2004; Fowler and Booth 2012; Folpp et al. 2013). With appropriate design, implementation and management, ARs could be deployed as part of the wider actions of the mitigation hierarchy to achieve a “net gain” in biodiversity. Gulf countries do not have adequate management policies for biodiversity offsetting, unfortunately (GIBOP 2019).

It is important to note, however, that ARs typically support very different communities of benthic invertebrates (Burt et al. 2009; Sedano et al. 2019; Hill et al. 2021) and fish (Komyakova et al. 2019; Jones et al. 2020; Mohamed 2021) than surrounding soft-bottom habitats and natural reefs. Paxton et al. (2020a), in contrast, found that artificial reefs generally support similar fish densities and diversities compared with natural reefs. Since ARs often support different communities than natural habitats, using ARs to mitigate destruction of natural habitat can be problematic, because they may be a poor substitutes for the lost biodiversity and ecosystem functions associated with natural habitats (Carvalho et al. 2013; Sanabria-Fernandez et al. 2018; Hill et al. 2021), particularly when the reefs are newly deployed and are in the early stages of colonization and succession (Walker and Schlacher 2014). The creation of ARs should not be used as an excuse to allow the destruction of existing natural habitat through so-called “green-washing” (Firth et al. 2020). Unfortunately, governmental policies may incentivize creation of new habitat, like ARs, for mitigation purposes over preservation of natural habitats (Levrel et al. 2012). At best, small-scale ARs may have limited ecological benefits with significant costs to deploy them. At worst, ARs can have a variety of unintended negative consequences, and have been used as an excuse to allow the destruction of economically valuable and ecologically important natural habitats.

3 Unintended Negative Consequences of ARs

This section of the paper is summarized in Fig. 4.

3.1 Opportunity Costs – ARs May Not be the Best Use of Limited Conservation Resources

Although there are certainly examples of large-scale ARs that have significant positive ecological effects, most AR deployments are small-scale, and may be too small relative to natural-occurring hard bottom habitat to create significant ecological benefits (DeMartini et al. 1989; Pitcher et al. 2002), as the effects of an individual reef may only extend a short distance from the edge of the reef itself (Scott et al. 2015; Smith et al. 2017; Reeds et al. 2018). These AR projects can be costly and time-consuming to create and deploy (Pioch and Doumenge 2010; Levrel et al. 2012; Bayraktarov et al. 2019), and in many cases the time, effort and money spent on ARs would be better spent on alternative conservation projects, such as protecting, restoring or enhancing natural habitat (McCreless and Beck 2016; Morris et

al. 2018; Airoidi et al 2021). In the mitigation hierarchy, deploying ARs can be used as an offset with positive impacts. Using offsets is a “last resort”, however, and it is preferable to avoid and minimize negative environmental impacts or restore natural habitat instead (Cabrera and Castro 2018).

3.2 Attraction of Fish that Exacerbates Overfishing

The most widely-cited and important negative economic consequence of deploying ARs is that they can contribute to unsustainably high fisheries catch rates, leading to a decline in regional fish stocks and overfishing over the medium- to long-term (Bohnsack 1989; Chou 1997; Grossman et al. 1997). This occurs because ARs often attract more fish from surrounding habitats than they produce themselves, which can contribute to decreased fish abundance in nearby natural habitats (Matthews 1985; Simon et al. 2011). ARs are effective at attracting and concentrating commercially-important species, including highly migratory species (Pears and Williams 2005; Shroepfer and Szedlmayer 2006), and numerous studies around the world have recorded increased short term catch rates and economic gains associated with ARs for both commercial and recreational fishers (Korea: Kim et al. 1994; Taiwan: Lin and Su 1994; Philippines: Chou 1997; Portugal: Santos and Monteiro 1998; Iran: Azhdari et al. 2012; India: Kasim et al. 2013; Australia: Keller et al. 2017). In the long term, these increased catch rates can contribute to regional overfishing, however (see paragraph below). ARs may improve catch rates of certain species but not others, and ARs may not affect overall fisheries catches in a region (Polovina 1989; Polovina and Sakai 1989; Sun et al. 2017). In some cases, ARs may even negatively affect regional fishing income (Islam et al. 2014).

ARs are often deployed to be easily accessible, and the combination of short travel times to known, fixed locations with high densities of commercially-important species means that ARs usually concentrate fishers as well. (McGlennon and Branden 1994; Watanuki and Gonzales 2006; Keller et al. 2016). High concentrations of fish and fishers around an AR can contribute to regional overfishing in the long term (Polovina 1990; Whitmarsh et al. 2008; Simon et al. 2011), particularly in countries with poor fisheries management (Watanuki and Gonzales 2006), unless regulations restricting fishing at ARs are implemented and enforced (Pears and Williams 2005). Concerns about ARs contributing to the decline of vulnerable commercial species has caused the government of South Australia, for example, to discourage any further deployment of ARs (Pears and Williams 2005).

3.3 ARs as Sink Habitat for Larval Fish

ARs may intercept fish larvae that would have otherwise settled in superior natural nursery habitats, and thus contribute to lower regional recruitment. ARs, particularly those with high vertical relief, tend to have high densities of large, predatory fish species (Paxton et al. 2020b), which can lead to low juvenile and prey fish survivorship on these reefs (Herrera et al. 2002; Leitao et al. 2008b; Folpp et al. 2011) compared with natural habitats. If this occurs, ARs would be a “sink” habitat that contribute to a decline in the regional abundance of prey species (Carr and Hixon 1997; Levrel et al. 2012).

3.4 Halos of Altered Benthic Habitat Around ARs

ARs usually support different faunal communities than the surrounding natural habitats, particularly if they are deployed in soft sediments (Bulleri 2005). ARs may also alter currents and sediment movement. Both the altered currents and the reef-associated fauna contribute to a “halo” of altered benthic habitat surrounding ARs (Herrera et al. 2002). Several studies have noted that the high

1
2
3
4 densities of predatory fish associated with ARs alter the benthic prey community around the reef
5 (Bortone et al. 1998; Herrera et al. 2002; Reeds et al. 2018) and high densities of herbivorous fish may
6 deplete vegetation around ARs (Edelist and Spanier 2009). Alternatively, high densities of fauna on ARs
7 may lead to increased nutrient deposition that spurs the growth of vegetation and alters the vegetation
8 community around the AR (Alevizon 2002; Dewsbury and Fourqurean 2010). ARs can also alter the local
9 hydrology and can cause increased sediment erosion around the reef in high-energy environments
10 (Herrera et al. 2002; Raineault et al. 2013) or increased sediment deposition in other cases (Martin et al.
11 2005; Cenci et al. 2011). These changes in sedimentation can, in turn, affect the benthic community
12 around the reef (Danovaro et al. 2002; Herrera et al. 2002; Raineault et al. 2013).

13 14 15 16 *3.5 Facilitating the Spread of Invasive Species*

17
18 ARs may act as stepping stones for the spread of hard-bottom invasive species (Bulleri and
19 Airoidi 2005; Adams et al. 2014; Firth et al. 2016), particularly if the ARs are deployed in soft sediment
20 habitats, such that these invasive species would not become established without the ARs (Sheehy and
21 Vik 2010; Airoidi et al. 2015). ARs can also facilitate the spread of invasives when deployed in hard-
22 bottom habitats, however, particularly if the invasive species are more successful on the artificial
23 structures compared with natural hard-bottom habitats (Dafforn et al. 2012). After invasive species
24 become established on artificial structures, they can then successfully invade nearby natural hard-
25 bottom habitats (Glasby et al. 2007; Broughton 2012; Simkanin et al. 2012). There are several examples
26 where ARs have clearly contributed to the spread of invasive species, including a green algae, ascidians,
27 a coral and a caprellid amphipod in the Mediterranean (Bulleri and Airoidi 2005; Ros et al. 2013;
28 Salomidi et al. 2013; Airoidi et al. 2015), a green algae in Chile (Neill et al. 2006) and a blenny, a coral,
29 mussels and an ascidian in the Gulf of Mexico (Sheehy and Vik 2010). The supplemental material of
30 Clarke et al. (2020) lists more than 30 benthic or demersal invasive species that have been found in the
31 Gulf that could potentially become associated with ARs. These include species of macroalgae, tunicates,
32 barnacles, nudibranchs, bryozoans and demersal fish. There are several design features that can reduce
33 the prevalence of invasive species on ARs, including: using natural rock and shell as substrates while
34 avoiding wood, concrete, rope and other artificial materials, avoiding steep vertical relief, reducing
35 shading on the reef and “seeding” the reef with native fouling species soon after the reef is deployed
36 (Firth et al. 2016; Dafforn 2017).

37 38 39 40 41 42 43 *3.6 Waste Dumping and Destruction of Natural Habitat*

44
45 Some “AR projects” may be little more than glorified trash dumping (with lower disposal costs)
46 in which cars, boats, airplanes, tires, construction waste, decommissioned oil rigs, etc. are sunk to the
47 bottom as “habitat” for marine life (Chou 1997; Firth et al. 2020). Decommissioning oil rigs may cost
48 several million dollars, for example, and oil companies can reduce their disposal costs with “rigs-to-
49 reefs” programs (Broughton 2012). ARs made from haphazardly deployed “scrap” material support
50 lower fish abundance and diversity compared with purpose-built ARs that are carefully deployed (Brock
51 and Norris 1989). These scrap materials are generally not built to withstand strong waves and currents,
52 and they may break apart and be transported surprisingly long distances by storms. Hurricanes have
53 transported relatively low-density AR materials, like concrete pipes, steel cubes, automobiles, tires and
54 recycled plastic 1 – 2 km, or further, away from their original location (Bell and Hall 1994; Turpin and
55 Bortone 2002; Morley et al. 2008). Even materials like sections of steel radio towers and vessels can be
56 moved several hundred meters in a hurricane (Bell and Hall 1994; Blair et al. 1994; Turpin and Bortone
57
58
59
60
61
62
63
64
65

2002). In some cases, scrap transported from ARs may end up damaging nearby natural habitat. An AR in Florida was created using millions of tires that moved several kilometers in some cases, and damaged nearby natural coral habitat. This necessitated costly cleanup efforts to remove some of the tires (Morley et al. 2008). Low-density materials, like automobiles, tires and shipping containers, should not be used as ARs where there is a possibility of strong storms, and the state of Florida, for example, has outlawed the use of these materials (Broughton 2012). Even relatively stable scrap artificial structures, like vessels, should be placed at least several hundred meters away from natural hard bottom habitats in regions susceptible to strong storms.

3.7 Pollution Released from ARs

ARs associated with waste materials may also release various pollutants associated with fuels, lubricants, batteries, metals, asbestos and paints, including toxic antifouling paints, unless they are thoroughly decontaminated before deployment. Examples of pollutants associated with ARs include polychlorinated biphenyls (PCBs) associated with vessels (Dodrill et al. 2011), heavy metals associated with paint (Broughton 2012; Gaylarde et al. 2021), heavy metals from tires (Collins et al. 2002) and toxic antifouling paint associated with vessels. This antifouling paint can reduce the densities of sessile invertebrates as well as the fish that would normally feed on them (Wendt et al. 1989; Walker et al. 2007; Szedlmayer and Miller 2018). Concrete associated with construction waste has a high pH (~13), and may be a poor substrate for recruitment of sessile organisms (Sella and Perkol-Finkel 2015).

4 Recommendations for Improved AR Policy in the Arabian Gulf

This section is summarized in Fig. 5

Gulf countries have developed ultra-modern, cosmopolitan cities, and Gulf leaders have clear visions for the future of their nations. Gulf countries are increasingly focusing on sustainability and effective environmental management as a part of their visions (Burt and Bartholomew 2019). Below we provide general recommendations that decision makers and managers should adopt in order to enhance the effectiveness of future AR programs in the Gulf.

4.1 Avoid 'greenwashing' with ARs

ARs provide additional hard-bottom habitat that can support diverse communities of fish and invertebrates. The communities that develop on artificial reefs are usually distinct from natural communities, however, so ARs should not be considered replacement surrogates for natural habitats that are degraded or destroyed by coastal development projects (Firth et al. 2020; Watchorn et al. 2022). Deployment of artificial reefs should not be used as an excuse to allow the destruction of natural habitat. Coastal management policies should promote the preservation, restoration and enhancement of natural ecosystems whenever possible. These policies will often be less costly than ARs and more successful in supporting sustainability-related management objectives.

4.2 List explicit goals for the AR program

The management goals for any AR project should be clearly defined at the earliest planning stage of the project, as these goals underpin all subsequent work on the AR (Spieler et al. 2001; Seaman 2007). ARs can be used to fulfill a variety management goals, and different goals would be supported by

different AR materials, designs, deployment patterns and site locations. Reef programs with multiple goals should list and assess each goal independently, and some goals may be incompatible to an extent (fishing and dive tourism, for example).

4.3 Define quantifiable criteria for assessing the AR's performance in meeting each goal

AR projects should have specific, quantifiable benchmarks associated with each goal, so that managers can judge AR performance and determine whether the AR project can be considered successful or not (Spieler et al. 2001; Becker et al. 2018). It is insufficient to say that the reef should simply increase fish density or coral percent coverage, for example. Multiple variables can be measured to determine if some goals are being met. For example, if a goal is enhancing dive tourism, managers could measure AR fish abundances, number of visitors to the reef, increased dive shop revenues, etc. These measurable objectives should be devised early on in any AR program, and should be re-evaluated after engaging relevant stakeholders, before moving on to later phases of the project (Pears and Williams 2005).

4.4 Design and deploy the ARs to meet the intended goals

ARs have been constructed using many different materials, they come in a wide variety of shapes and sizes, they may have different surface features and attachments and they have been deployed in a variety of configurations in many different locations and habitats around the world (Tickell et al. 2019). Only a subset of these will support the intended goals of the project, however, and managers should read published accounts of regional AR programs to determine what design features and deployment techniques have worked in the past to achieve similar goals. There are government-issued guidelines available for artificial reef planning, materials, construction and deployment that are available (Lukens and Selberg. 2004; Lindberg and Seaman 2011; Fabi et al. 2015). Decision makers should also consult with regional stakeholders during the planning stages and afterwards, so that these stakeholders can input their knowledge and they feel included, both of which would contribute to the success of the AR program (Clarke et al. 2002; Sayer and Wilding 2002).

4.5 Develop a robust monitoring program to assess criteria in meeting goals

It is critical to implement a scientific monitoring program after deployment, to determine the impacts of the reef and whether the goals of the AR program are met (Ramm et al. 2021). The monitoring program should ideally be implemented by professional marine ecologists associated with independent, non-commercial organizations. Factors such as the sampling design (e.g. Before-After, Control-Impact), amount of replication, the extent of sampling in space and time, the appropriate methods for sampling different species and data analyses need to be considered in-depth. Monitoring should be long-term and systematic, because the communities associated with ARs will change rapidly soon after deployment, but can continue to change for decades (or longer) (Perkol-Finkel and Benayahu 2005; Nicoletti et al. 2007). ARs are often expensive and time-consuming to deploy, and monitoring requires relatively little cost and effort by comparison, but unfortunately long-term monitoring is rare (Ramm et al. 2021). Monitoring data will be a "good return on investment" because it can help guide decision makers about whether and how to deploy more ARs in the future.

4.6 Publicize the results of the AR program

The final stage in any AR program should be to evaluate its effectiveness, and share the results and 'lessons learned' publicly. Managers should report information about the reef itself (Ramm et al. 2021), management goals and the data from the long-term monitoring program. Ideally, information about the costs and work hours associated with the AR program should be presented, so managers can assess whether the outcomes created by the AR are worth the money and effort. All this information will help decision makers to determine whether to deploy additional ARs, and how to improve them if more are deployed. By sharing this information publicly, marine managers can demonstrate their commitment to promoting environmental sustainability and help other managers develop successful AR programs in the future (Ramm et al. 2021).

5 Acknowledgements

This manuscript originated from a project supported by the Sheikh Saud bin Saqr Al Qasimi Foundation for Policy Research.

JAB was supported by a grant from Tamkeen to New York University Abu Dhabi through the Water Research Center (WRC) (grant number CG007), by a grant from Arabian Center for Climate and Environmental Sciences (ACCESS) (grant number CG009) and by grant G1902 from SwissRe to the NYUAD CITIES research center.

6 References

- Abdel-Moati, M., 2006. Coral Reef Conservation in Qatar, presented at Marine Conservation Forum. EWS-WWF, Abu Dhabi, UAE.
- Adams, T., Miller, R., Aleynik, D., Burrows, M., 2014. Offshore marine renewable energy devices as stepping stones across biogeographical boundaries. *J. Appl. Ecol.* 51, 330-338. <https://doi.org/10.1111/1365-2664.12207>.
- Airoidi, L., Beck, M., Firth, L., Bugnot, A., Steinberg, P., Dafforn, K., 2021. Emerging solutions to return nature to the urban ocean. *Ann. Rev. Mar. Sci.*, 13, 445-477. <https://doi.org/10.1146/annurev-marine-032020-020015>.
- Airoidi, L., Turon, X., Perkol-Finkel, S., Rius, M., 2015. Corridors for aliens but not for natives: effects of marine urban sprawl at a regional scale. *Divers. Distrib.* 21, 755-768. <https://doi.org/10.1111/ddi.12301>.
- Ajdari, D., Ajdari, Z., 2016. Effect of marine artificial reefs on improvement and development of sea environment in Kish Island, presented at International Conference on Coasts, Ports and Marine Structures (ICOPMAS), Tehran, IR, 16, 449-450.
- Ajdari D, Hajirezaee S, Ajdari Z., 2011. Fish colonization of artificial reefs in the Persian Gulf (Bandar Lengeh-Iran): Reef designs and seasonal changes of fish abundance. *Int. J. Fish. Aquac.* 3, 204-212.

- Al-Abdulrazzak, D., Zeller, D., Belhabib, D., Tesfamichael, D., Pauly, D., 2015. Total marine fisheries catches in the Persian/Arabian Gulf from 1950 to 2010. *Reg. Stud. Mar. Sci.* 2, 28-34. <https://doi.org/10.1016/j.rsma.2015.08.003>.
- Al-Cibahy, A., Al-Khalifa, K., Böer, B. and Samimi-Namin, K., 2012. Conservation of marine ecosystems with a special view to coral reefs in the Gulf, in: Riegl, B., Purkis, S. (Eds.), *Coral reefs of the Gulf*. Springer, Dordrecht, NL, pp. 337-348.
- Al-Cibahy, A., Grandcourt, E. and Bugla, I., 2009. Report on evaluation of eco-reefs installed in Marawah Marine Biosphere Reserve. Environment Agency-Abu Dhabi, Abu Dhabi, UAE, 17.
- Al-Horani, F., Khalaf, M., 2013. Developing artificial reefs for the mitigation of man-made coral reef damages in the Gulf of Aqaba, Red Sea: coral recruitment after 3.5 years of deployment. *Mar. Biol. Res.* 9, 749-757. <https://doi.org/10.1080/17451000.2013.765582>.
- Al-Saidi, M., 2021. Regional Environmental Cooperation: The (Lost) Potential for a Sustainable Future in the Arabian/Persian Gulf, in: Standish K., Devere H., Suazo A., Rafferty R. (Eds.), *The Palgrave Handbook of Positive Peace*. Palgrave Macmillan, Singapore, pp. 814-833. https://doi.org/10.1007/978-981-15-3877-3_44-1.
- Alevizon, W., 2002. Enhanced seagrass growth and fish aggregations around Bahamian patch reefs: The case for a functional connection. *Bull. Mar. Sci.* 70, 957-966.
- Alghunaim, A., Yamamoto, T., Madhusoodhanan, R., Polikarpov, I., Chen, W., Al-Said, T., Al-Zekri, W., Alenezi, Y., Al-Yamani, F., 2020. Evaluating the potential area and environmental key factors for Sargassum bed restoration in highly turbid waters of the northwestern Arabian Gulf. *Mar. Pollut. Bull.*, 153, p.110949. <https://doi.org/10.1016/j.marpolbul.2020.110949>.
- Alhabsi, M., 2013. Involvement of fishermen in fisheries co-management in Oman. *Greener J. Agric. Sci.* 3, 341-354.
- Aljasmi, H., 2017. Assessment of the environmental management practices to combat overfishing in the UAE, in: Anwar S., El Sergany M., Ankit A. (Eds.), *Innovation Arabia 10: Health and Environment Conference Proceedings*, Hamdan Bin Mohammed Smart University, Dubai, UAE, pp. 298-305.
- Alshahri, F., 2017. Heavy metal contamination in sand and sediments near to disposal site of reject brine from desalination plant, Arabian Gulf: Assessment of environmental pollution. *Environ. Sci. Pollut. Res.* 24,1821-1831. <https://doi.org/10.1007/s11356-016-7961-x>.
- Azhdari, H., Ibrahim, M., Arshad, A., Shohaimi, S., Ajdari, Z., Ibrahim, Z., 2012. The effect of artificial reefs on fish assemblage versus natural sites in the Bandar Lengeh-Iran. *Iran. J. Fish. Sci.* 11, 1-12.
- Baine, M., 2001. Artificial reefs: a review of their design, application, management and performance. *Ocean Coast. Manag.* 44, 241-259. [https://doi.org/10.1016/S0964-5691\(01\)00048-5](https://doi.org/10.1016/S0964-5691(01)00048-5).
- Barth, H., Böer, B. (Eds.), 2002. *Sabkha Ecosystems: Volume I: The Arabian Peninsula and Adjacent Countries*. Springer Science and Business Media, Dordrecht, NL.

1
2
3
4 Bauman, A., Burt, J., Feary, D., Marquis, E., Usseglio, P., 2010. Tropical harmful algal blooms: An
5 emerging threat to coral reef communities?. *Mar. Poll. Bull.* 60, 2117-2122.
6 <https://doi.org/10.1016/j.marpolbul.2010.08.015>.
7
8 Bayraktarov, E., Stewart-Sinclair, P., Brisbane, S., Boström-Einarsson, L., Saunders, M., Lovelock, C.,
9 Possingham, H., Mumby, P., Wilson, K., 2019. Motivations, success, and cost of coral reef
10 restoration. *Restor. Ecol.* 27, 981-991. <https://doi.org/10.1111/rec.12977>.
11
12
13 Becker, A., Taylor, M., Folpp, H., Lowry, M., 2018. Managing the development of artificial reef systems:
14 The need for quantitative goals. *Fish Fish.*, 19, 740-752. <https://doi.org/10.1111/faf.12288>.
15
16 Belhassen, Y., Rousseau, M., Tynyakov, J., Shashar, N., 2017. Evaluating the attractiveness and
17 effectiveness of artificial coral reefs as a recreational ecosystem service. *J. Environ. Manage.* 203, 448-
18 456. <https://doi.org/10.1016/j.jenvman.2017.08.020>.
19
20
21 Bell, M., Hall, W., 1994. Effects of Hurricane Hugo on South Carolina's marine artificial reefs. *Bull. Mar.*
22 *Sci.* 55, 836-847.
23
24 Black, K., Mead, S., 2001. Design of the Gold Coast reef for surfing, public amenity and coastal
25 protection: surfing aspects. *J. Coast. Res. special issue* 28, 115-130.
26
27 Blair, S., McIntosh, T., Mostkoff, B., 1994. Impacts of Hurricane Andrew on the offshore reef systems of
28 central and northern Dade County, Florida. *Bull. Mar. Sci.* 54, 961-973.
29
30
31 Blignaut, J., Mander, M., Inglesi-Lotz, R., Glavan, J., Parr, S., 2016. The amenity value of Abu Dhabi's
32 coastal and marine resources to its beach visitors. *Ecosyst. Serv.*, 19, 32-41.
33 <https://doi.org/10.1016/j.ecoser.2016.04.005>.
34
35 Bohnsack, J.A., 1989. Are high densities of fishes at artificial reefs the result of habitat limitation or
36 behavioral preference?. *Bull. Mar. Sci.* 44, 631-645.
37
38 Bohnsack, J., Sutherland, D., 1985. Artificial reef research: a review with recommendations for future
39 priorities. *Bull. Mar. Sci.* 37, 11-39.
40
41
42 Bortone, S., Cody, R., Turpin, R., Bundrick, C., 1998. The impact of artificial-reef fish assemblages on their
43 potential forage area. *Ital. J. Zool.* 65, 265-267. <https://doi.org/10.1080/11250009809386830>.
44
45 Brandl, S., Johansen, J., Casey, J., Tornabene, L., Morais, R., Burt, J., 2020. Extreme environmental
46 conditions reduce coral reef fish biodiversity and productivity. *Nat. Commun.* 11, 3832.
47 <https://doi.org/10.1038/s41467-020-17731-2>
48
49
50 Brock, R., Norris, J., 1989. An analysis of the efficacy of four artificial reef designs in tropical waters. *Bull.*
51 *Mar. Sci.* 44, 934-941.
52
53 Broughton, K. 2012. Office of National Marine Sanctuaries Science Review of Artificial Reefs. Marine
54 Sanctuaries Conservation Series ONMS-12-05, U.S. Department of Commerce, National Oceanic and
55 Atmospheric Administration, Office of National Marine Sanctuaries, Silver Spring, MD.
56
57
58 Buchanan, J., Krupp, F., Burt, J., Feary, D., Ralph, G., Carpenter, K., 2016. Living on the edge: Vulnerability
59 of coral-dependent fishes in the Gulf. *Mar. Poll. Bull.* 105, 480-488.
60 <https://doi.org/10.1016/j.marpolbul.2015.11.033>.
61
62
63
64
65

Buchanan, J., Ralph, G., Krupp, F., Harwell, H., Abdallah, M., Abdulqader, E., Al-Husaini, M., Bishop, J., Burt, J., Choat, J., Collette, B., Feary, D., Hartmann, S., Iwatsuki, Y., Kaymaram, F., Larson, H., Matsuura, K., Motomura, H., Munroe, T., Russell, B., Smith-Vaniz, W. Williams, J., Carpenter, K., 2019. Regional extinction risks for marine bony fishes occurring in the Persian/Arabian Gulf. *Biol. Conserv.* 230, 10-19. <https://doi.org/10.1016/j.biocon.2018.11.027>.

Bulleri, F., 2005. The introduction of artificial structures on marine soft-and hard-bottoms: ecological implications of epibiota. *Environ. Conserv.* 32, 101-102. <https://doi.org/10.1017/S0376892905002183>.

Bulleri, F., Airoidi, L., 2005. Artificial marine structures facilitate the spread of a non-indigenous green alga, *Codium fragile* ssp. *tomentosoides*, in the north Adriatic Sea. *J. Appl Ecol.* 42, 1063-1072. <https://doi.org/10.1111/j.1365-2664.2005.01096.x>.

Burt, J., 2014. The environmental costs of coastal urbanization in the Arabian Gulf. *City.* 18, 760-770. <https://doi.org/10.1080/13604813.2014.962889>

Burt, J., Bartholomew, A., 2019. Towards more sustainable coastal development in the Arabian Gulf: Opportunities for ecological engineering in an urbanized seascape. *Mar. Pollut. Bull.* 142, 93-102. <https://doi.org/10.1016/j.marpolbul.2019.03.024>.

Burt, J., Bartholomew, A. Sale, P., 2011. Benthic development on large-scale engineered reefs: a comparison of communities among breakwaters of different age and natural reefs. *Ecol. Eng.* 37, 191-198. <https://doi.org/10.1016/j.ecoleng.2010.09.004>.

Burt, J., Bartholomew, A., Usseglio, P., Bauman, A. and Sale, P., 2009. Are artificial reefs surrogates of natural habitats for corals and fish in Dubai, United Arab Emirates?. *Coral Reefs* 28, 663-675. <https://doi.org/10.1007/s00338-009-0500-1>.

Burt, J., Ben-Hamadou, R., Abdel-Moati, M., Fanning, L., Kaitibie, S., Al-Jamali, F., Range, P., Saeed, S., Warren, C., 2017. Improving management of future coastal development in Qatar through ecosystem-based management approaches. *Ocean Coast. Manag.* 148, 171-181. <https://doi.org/10.1016/j.ocecoaman.2017.08.006>.

Burt, J., Feary, D., Cavalcante, G., Bauman, A., Usseglio, P., 2013. Urban breakwaters as reef fish habitat in the Persian Gulf. *Mar. Poll. Bull.* 72. 342-350. <https://doi.org/10.1016/j.marpolbul.2012.10.019>.

Burt, J., Paparella, F., Al-Mansoori, N., Al-Mansoori, A., Al-Jailani, H., 2019. Causes and consequences of the 2017 coral bleaching event in the southern Persian/Arabian Gulf. *Coral Reefs* 38, 567-589. <https://doi.org/10.1007/s00338-019-01767-y>.

Burt, J., Range, P., Claereboudt, M., Al-Mealla, R., Salimi, P., Salimi, M., Ben-Hamadou, R., Bolouki, M., Bouwmeester, J., Taylor, O., Wilson S., 2021. Status and trends of coral reefs of the ROPME Sea Area, in: Souter, D., Planes, S., Wicquart, J., Logan, M., Obura, D., Staub, F. (Eds.), *Status of Coral Reefs of the World: 2020, Global Coral Reef Monitoring Network and United Nations Environment Programme* Lausanne, CH, pp. 74-81

Cabrera, E., Castro, J., 2018 *Guidance on Using the Mitigation Hierarchy to Avoid Impacts of Projects and Activities*. Commonwealth of the Northern Mariana Islands, Bureau of Environmental and Coastal Quality.

- Campbell, J., Lacey, E., Decker, R., Crooks, S., Fourqurean, J., 2015. Carbon storage in seagrass beds of Abu Dhabi, United Arab Emirates. *Estuar. Coast.* 38, 242-251. <https://doi.org/10.1007/s12237-014-9802-9>.
- Carr, M., Hixon, M., 1997. Artificial reefs: the importance of comparisons with natural reefs. *Fisheries* 22, 28-33. [https://doi.org/10.1577/1548-8446\(1997\)022%3C0028:ARTIOC%3E2.0.CO;2](https://doi.org/10.1577/1548-8446(1997)022%3C0028:ARTIOC%3E2.0.CO;2).
- Carvalho, S., Moura, A., Cúrdia, J., da Fonseca, L., Santos, M., 2013. How complementary are epibenthic assemblages in artificial and nearby natural rocky reefs?. *Mar. Environ. Res.* 92, pp.170-177. <https://doi.org/10.1016/j.marenvres.2013.09.013>.
- Cenci, E., Pizzolon, M., Chimento, N., Mazzoldi, C., 2011. The influence of a new artificial structure on fish assemblages of adjacent hard substrata. *Estuar., Coast. Shelf Sci.* 91,133-149. <https://doi.org/10.1016/j.ecss.2010.10.009>.
- Champion, C., Suthers, I., Smith, J., 2015. Zooplanktivory is a key process for fish production on a coastal artificial reef. *Mar. Ecol. Prog. Ser.* 541, 1-14. DOI: <https://doi.org/10.3354/meps11529>.
- Charbonnel, E., Bachet, F., 2010. Artificial reefs in the Cote Bleue Marine Park: Assessment after 25 years of experiments and scientific monitoring, in: Ceccaldi, H., Dekeyser I., Girault M., Stora G. (Eds.), *Global Change: Mankind-Marine Environment Interactions*, Springer, Dordrecht, NL, pp. 73-79.
- Charbonnel, E., Serre, C., Ruitton, S., Harmelin, J., Jensen, A., 2002. Effects of increased habitat complexity on fish assemblages associated with large artificial reef units (French Mediterranean coast). *ICES J. Mar. Sci.* 59(suppl), S208-S213. <https://doi.org/10.1006/jmsc.2002.1263>.
- Chatziefthimiou, A., Banack, S., Metcalf, J., 2021. Harmful Algal and Cyanobacterial Harmful Algal Blooms in the Arabian Seas: Current Status, Implications, and Future Directions, in: Jawad L. (Ed.), *The Arabian Seas: Biodiversity, Environmental Challenges and Conservation Measures*. Springer, Cham., pp.
- Chojnacki, J., Ceronik, E., 1996. Artificial reefs in the Pomeranian Bay (Southern Baltic) as biofiltration sites, in: Andrushaitis, A. (Ed.) *Proceedings of the 13th Symposium of the Baltic Marine Biologists*, Riga University, Riga, LV, pp. 169-172.
- Chou, L., 1997. Artificial reefs of Southeast Asia-Do they enhance or degrade the marine environment?. *Environ. Monit. Assess.* 44, 45-52. <https://doi.org/10.1023/A:1005759818050>.
- Clarke, S., Leung Wai-Yin, A., Mak, Y., Kennish, R., Haggan, N., 2002. Consultation with local fishers on the Hong Kong artificial reefs initiative. *ICES J. Mar. Sci.*, 59(suppl), S171-S177. <https://doi.org/10.1006/jmsc.2002.1274>.
- Clarke, S., Vilizzi, L., Lee, L., Wood, L., Cowie, W., Burt, J., Mamiit, R., Ali, H., Davison, P., Fenwick, G., Harmer, R., 2020. Identifying potentially invasive non-native marine and brackish water species for the Arabian Gulf and Sea of Oman. *Glob. Change Biol.* 26, 2081-2092. <https://doi.org/10.1111/gcb.14964>.
- Collins, K., Jensen, A., Mallinson, J., Roenelle, V., Smith, I., 2002. Environmental impact assessment of a scrap tyre artificial reef. *ICES J. Mar. Sci.* 59(suppl), S243-S249. <https://doi.org/10.1006/jmsc.2002.1297>.

- Cresson, P., Ruitton, S., Harmelin-Vivien, M., 2014. Artificial reefs do increase secondary biomass production: mechanisms evidenced by stable isotopes. *Mar. Ecol. Prog. Ser.* 509, 15-26.
<https://doi.org/10.3354/meps10866>
- D'Agostino, D., Burt, J., Santinelli, V., Vaughan, G., Fowler, A., Reader, T., Taylor, B., Hoey, A., Cavalcante, G., Bauman, A., Feary, D., 2021. Growth impacts in a changing ocean: insights from two coral reef fishes in an extreme environment. *Coral Reefs* 40, 433-446. <https://doi.org/10.1007/s00338-021-02061-6>.
- Dafforn, K., 2017. Eco-engineering and management strategies for marine infrastructure to reduce establishment and dispersal of non-indigenous species. *Manag. Biol. Invasion.* 8, 153-161.
<https://doi.org/10.3391/mbi.2017.8.2.03>.
- Dafforn, K., Glasby, T., Johnston, E., 2012. Comparing the invasibility of experimental "reefs" with field observations of natural reefs and artificial structures. *PLoS One*, 7, e38124.
<https://doi.org/10.1371/journal.pone.0038124>.
- Danovaro, R., Gambi, C., Mazzola, A., Mirto, S., 2002. Influence of artificial reefs on the surrounding infauna: analysis of meiofauna. *ICES J. Mar. Sci.* 59(suppl), S356-S362.
<https://doi.org/10.1006/jmsc.2002.1223>.
- De Verneil, A., Burt, J., Mitchell, M., Paparella, F., 2021. Summer Oxygen Dynamics on a Southern Arabian Gulf Coral Reef. *Front. Mar. Sci.* p.1676. <https://doi.org/10.3389/fmars.2021.781428>
- Deb, K., McCarthy, A., Harkanson, B., 2014. Coral relocation as habitat mitigation for impacts from the Barzan Gas Project pipeline construction, offshore Qatar. Presented at SPE International Conference on Health, Safety, and Environment, Long Beach, CA. <https://doi.org/10.2118/168369-MS>.
- Delmendo, M., 1990. A review of artificial reefs development and use of fish aggregating devices (FADs) in the ASEAN region. In: Pietersz, V. (Ed.), *Symposium on Artificial Reefs and Fish Aggregating Devices as Tools for the Management and Enhancement of Marine Fishery Resources*. Regional Office for Asia and the Pacific, United Nations Food and Agriculture Organisation, Bangkok, TH, pp. 116-141
- DeMartini, E., Roberts, D., Anderson, T., 1989. Contrasting patterns of fish density and abundance at an artificial rock reef and a cobble-bottom kelp forest. *Bull. Mar. Sci.* 44, 881-892.
- Dewsbury, B., Fourqurean, J., 2010. Artificial reefs concentrate nutrients and alter benthic community structure in an oligotrophic, subtropical estuary. *Bull. Mar. Sci.* 86, 813-829.
<https://doi.org/10.5343/bms.2009.1066>.
- Dodrill, J., K. Mille, B. Horn, R. Turpin. 2011. Progress report summarizing the reef fish sampling, PCB analysis results and visual monitoring associated with the Oriskany Reef, a decommissioned former Navy aircraft carrier sunk in 2006 as an artificial reef in the Northeastern Gulf of Mexico off Pensacola, Florida. Florida Fish and Wildlife Conservation Commission.
- Donelson, J., Munday, P., McCormick, M., Pankhurst, N., Pankhurst, P., 2010. Effects of elevated water temperature and food availability on the reproductive performance of a coral reef fish. *Mar. Ecol. Prog. Ser.* 401, 233-243. <https://doi.org/10.3354/meps08366>.
- Downing, N., Tubb, R., El-Zahr, C., McClure, R., 1985. Artificial reefs in Kuwait, northern Arabian Gulf. *Bull. Mar. Sci.* 37, 157-178.

1
2
3
4 EAD, 2016. Marine and Coastal Environment of Abu Dhabi Emirate, United Arab Emirates, Abu Dhabi
5 Global Environmental Data Initiative, Abu Dhabi, UAE.
6
7 EAD, 2021. Sustainable fisheries policy (2021-2025), Environment Agency Abu Dhabi, Abu Dhabi, UAE.
8
9 Edelist, D., Spanier, E., 2009. Influence of Levantine artificial reefs on the fish assemblage of the
10 surrounding seabed. *Mediterr. Mar. Sci.* 10, 35-54. <https://doi.org/10.12681/mms.121>.
11
12 Emery, B.M., L. Washburn, M.S. Love, M.M. Mishimoto, J.C. Ohlmann. 2006. Do oil and gas platforms off
13 California reduce recruitment of bocaccio (*Sebastes paucispinis*) to natural habitat? An analysis based on
14 trajectories derived from high-frequency radar. *Fish. Bull.* 104, 391-400.
15
16 Erftemeijer, P., Agastian, T., Yamamoto, H., Cambridge, M., Hoekstra, R., Toms, G., Ito, S., 2020.
17 Mangrove planting on dredged material: three decades of nature-based coastal defence along a
18 causeway in the Arabian Gulf. *Mar. Freshw. Res.* 71, 1062-1072. <https://doi.org/10.1071/MF19289>.
19
20 Fabi, G., Fiorentini, L., Giannini, S., 1989. Experimental shellfish culture on an artificial reef in the Adriatic
21 Sea. *Bull. Mar. Sci.* 44, 923-933.
22
23 Fabi, G., Manoukian, S., Spagnolo, A., 2006. Feeding behavior of three common fishes at an artificial reef
24 in the northern Adriatic Sea. *Bull. Mar. Sci.* 78, 39-56.
25
26 Fabi, G., Scarcella, G., Spagnolo, A., Bortone, S., Charbonnel, E., Goutayer, J., Haddad, N., Lök, A.,
27 Trommelen, M., 2015. Practical guidelines for the use of artificial reefs in the Mediterranean and the
28 Black Sea. General Fisheries Commission for the Mediterranean. Studies and Reviews, 96.
29
30 Fariñas-Franco, J., Roberts, D., 2014. Early faunal successional patterns in artificial reefs used for
31 restoration of impacted biogenic habitats. *Hydrobiologia*, 727, 75-94.
32
33 Feary, D., Burt, J., Bartholomew, A., 2011. Artificial marine habitats in the Arabian Gulf: review of
34 current use, benefits and management implications. *Ocean Coast. Manag.* 54, 742-749.
35 <https://doi.org/10.1016/j.ocecoaman.2011.07.008>.
36
37 Firth, L., Airoidi, L., Bulleri, F., Challinor, S., Chee, S., Evans, A., Hanley, M., Knights, A., O'Shaughnessy, K.,
38 Thompson, R., Hawkins, S., 2020. Greening of grey infrastructure should not be used as a Trojan horse to
39 facilitate coastal development. *J. Appl. Ecol.* 57, 1762-1768. <https://doi.org/10.1111/1365-2664.13683>.
40
41 Firth, L., Knights, A., Bridger, D., Evans, A., Mieszkowska, N., Moore, P., O'Connor, N., Sheehan, E.,
42 Thompson, R., Hawkins, S., 2016. Ocean sprawl: challenges and opportunities for biodiversity
43 management in a changing world. *Oceanogr. Mar. Biol.* 54, pp.189-262.
44
45 Firth, L., Thompson, R., Bohn, K., Abbiati, M., Airoidi, L., Bouma, T., Bozzeda, F., Ceccherelli, V.,
46 Colangelo, M., Evans, A., Ferrario, F., 2014. Between a rock and a hard place: environmental and
47 engineering considerations when designing coastal defence structures. *Coast. Eng.* 87, pp.122-135.
48 <https://doi.org/10.1016/j.coastaleng.2013.10.015>.
49
50 Folpp, H., Lowry, M., Gregson, M., Suthers, I., 2011. Colonization and community development of fish
51 assemblages associated with estuarine artificial reefs. *Braz. J. Oceanogr.* 59, 55-67.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Folpp, H., Lowry, M., Gregson, M., Suthers, I., 2013. Fish assemblages on estuarine artificial reefs:
5 natural rocky-reef mimics or discrete assemblages?. PLoS One 8, p.e63505.
6 <https://doi.org/10.1371/journal.pone.0063505>.
7
8
9 Folpp, H., Schilling, H., Clark, G., Lowry, M., Maslen, B., Gregson, M., Suthers, I., 2020. Artificial reefs
10 increase fish abundance in habitat-limited estuaries. J. Appl. Ecol. 57, 1752-1761.
11 <https://doi.org/10.1111/1365-2664.13666>.
12
13 Fowler, A., Booth, D., 2012. How well do sunken vessels approximate fish assemblages on coral reefs?
14 Conservation implications of vessel-reef deployments. Mar. Biol. 159, 2787-2796.
15 <https://doi.org/10.1007/s00227-012-2039-x>.
16
17
18 Freije, A., 2015. Heavy metal, trace element and petroleum hydrocarbon pollution in the Arabian Gulf. J.
19 Assoc. Arab Univ. Basic Appl. Sci. 17, 90-100. <https://doi.org/10.1016/j.jaubas.2014.02.001>.
20
21 Gaylarde, C., Neto, J., da Fonseca, E., 2021. Paint fragments as polluting microplastics: A brief
22 review. Mar. Poll. Bull., 162, p.111847. <https://doi.org/10.1016/j.marpolbul.2020.111847>.
23
24 Gibling, C., 2013. Construction process and post-construction impacts of the Palm Jumeirah in Dubai,
25 United Arab Emirates. Prototype, 1.
26
27 GIBOP 2019. Global inventory of biodiversity offset policies (GIBOP). International Union for Conservation
28 of Nature, The Biodiversity Consultancy, Durrell Institute of Conservation and Ecology.
29 <https://portals.iucn.org/offsetpolicy/> (accessed 18 Dec 2021).
30
31
32 Glasby, T., Connell, S., Holloway, M., Hewitt, C., 2007. Nonindigenous biota on artificial structures: could
33 habitat creation facilitate biological invasions?. Mar. Biol. 151, 887-895.
34 <https://doi.org/10.1007/s00227-006-0552-5>.
35
36
37 Grandcourt, E., 2012. Reef fish and fisheries in the Gulf, in: Riegl B., Purkis, S. (Eds.), Coral reefs of the
38 Gulf. Springer, Dordrecht, NL, 127-161.
39
40 Grossman, G., Jones, G., Seaman Jr., W., 1997. Do artificial reefs increase regional fish production? A
41 review of existing data. Fisheries 22, 17-23.
42 [https://doi.org/10.1577/1548-8446\(1997\)022%3C0017:DARIRF%3E2.0.CO;2](https://doi.org/10.1577/1548-8446(1997)022%3C0017:DARIRF%3E2.0.CO;2)
43
44
45 Hall, S., Risinger, J., Farlow, J., 2011. Ecological engineering of artificial oyster reefs to enhance carbon
46 sequestration via the algae-oyster complex. Presented at American Society of Agricultural and Biological
47 Engineers, Louisville, KY.
48
49
50 Hamza, W., Munawar, M., 2009. Protecting and managing the Arabian Gulf: Past, present and
51 future. Aquat. Ecosyst. Health Manag., 12, 429-439. <https://doi.org/10.1080/14634980903361580>.
52
53 Harris C., Schroder, T., 2001. UAE Underwater Explorer. Explorer Publishing, Dubai, UAE.
54
55 Hegazy, A., 1998. Perspectives on survival, phenology, litter fall and decomposition, and caloric content
56 of *Avicennia marina* in the Arabian Gulf region. J. Arid Environ. 40, 417-429.
57 <https://doi.org/10.1006/jare.1998.0457>
58
59
60
61
62
63
64
65

- Herrera, R., Espino, F., Garrido, M., Haroun, R., 2002. Observations on fish colonization and predation on two artificial reefs in the Canary Islands. *ICES J. Mar. Sci.* 59(suppl), S69-S73.
<https://doi.org/10.1006/jmsc.2002.1291>.
- Hill, C., Lymperaki, M., Hoeksema, B., 2021. A centuries-old manmade reef in the Caribbean does not substitute natural reefs in terms of species assemblages and interspecific competition. *Mar. Pollut. Bull.* 169, p.112576. <https://doi.org/10.1016/j.marpolbul.2021.112576>.
- Hopkins, C., 2007. Dubai pilot project: test report for 'Reef Systems' at the Palm Jumeirah. AquaMarine Advisors, Astorp, Sweden.
- Islam, G., Noh, K., Sidique, S., Noh, A., 2014. Economic impact of artificial reefs: A case study of small scale fishers in Terengganu, Peninsular Malaysia. *Fish. Res.*, 151, 122-129.
<https://doi.org/10.1016/j.fishres.2013.10.018>.
- Israel, D., Gallo, C., Angel, D., 2017. Benthic artificial reefs as a means to reduce the environmental effects of cod mariculture in Skutulsfjörður, Iceland. *Mar. Biodivers.* 47, 405-411.
<https://doi.org/10.1007/s12526-016-0490-5>.
- Jones, S., Asher, J., Boland, R., Kanenaka, B., Weng, K., 2020. Fish biodiversity patterns of a mesophotic-to-subphotic artificial reef complex and comparisons with natural substrates. *Plos one* 15, e0231668.
<https://doi.org/10.1371/journal.pone.0231668>.
- Kasim, H., Rao, G., Rajagopalan, M., Vivekanandan, E., Mohanraj, G., Kandasamy, D., Muthiah, P., Jagadis, I., Gopakumar, G., Mohan, S., 2013. Economic performance of artificial reefs deployed along Tamil Nadu coast, South India. *Indian J. Fish.* 60, 1-8. <http://eprints.cmfri.org.in/id/eprint/9370>.
- Keller, K., Steffe, A., Lowry, M., Murphy, J., Smith, J., Suthers, I., 2017. Estimating the recreational harvest of fish from a nearshore designed artificial reef using a pragmatic approach. *Fish. Res.* 187, 158-167. <https://doi.org/10.1016/j.fishres.2016.11.022>.
- Keller, K., Steffe, A., Lowry, M., Murphy, J., Suthers, I., 2016. Monitoring boat-based recreational fishing effort at a nearshore AR with a shore-based camera. *Fish. Res.* 181, 84-92.
<https://doi.org/10.1016/j.fishres.2016.11.022>.
- Khan, N., 2007. Multiple stressors and ecosystem-based management in the Gulf. *Aquat. Ecosyst. Health Manag.* 10, 259-267. <https://doi.org/10.1080/14634980701551168>.
- Kim, C., Lee, J., Park, J., 1994. Artificial reef designs for Korean coastal waters. *Bull. Mar. Sci.* 55, 858-866.
- Kim, C., Lee, S., Cha, H., Yang, J., Son, Y., 2011. A case study of artificial reefs in fisheries management enhancement of sandfish, *Arctoscopus japonicus*, by artificial reefs in the eastern waters of Korea, in: Bortone, S., Brandini, F., Fabi, G., Otake, S. (Eds.), *Artificial Reefs in Fisheries Management*. CRC Press. Boca Raton, FL, USA, pp.111-124.
- Komyakova, V., Chamberlain, D., Jones, G., Swearer, S., 2019. Assessing the performance of ARs as substitute habitat for temperate reef fishes: Implications for reef design and placement. *Sci. Total Environ.* 668, 139-152. <https://doi.org/10.1016/j.scitotenv.2019.02.357>

Lachkar, Z., Mehari, M., Lévy, M., Paparella, F., Burt, J., 2022. Recent expansion and intensification of hypoxia in the Arabian Gulf and its drivers. *Front. Mar. Sci.*, p. 1616.
<https://doi.org/10.3389/fmars.2022.891378>

Lamine, E., Mateos-Molina, D., Antonopoulou, M., Burt, J., Das, H., Javed, S., Muzaffar, S., Giakoumi, S., 2020. Identifying coastal and marine priority areas for conservation in the United Arab Emirates. *Biodivers. Conserv.* 29, 2967-2983. <https://doi.org/10.1007/s10531-020-02007-4>.

Le Quesne, W., Fernand, L., Ali, T., Andres, O., Antonopoulou, M., Burt, J., Dougherty, W., Edson, P., El Kharraz, J., Glavan, J., Mamiit, R., 2021. Is the development of desalination compatible with sustainable development of the Arabian Gulf?. *Mar. Pollut. Bull.* 173, 112940.
<https://doi.org/10.1016/j.marpolbul.2021.112940>.

Leitao, F., Santos, M., Erzini, K., Monteiro, C., 2008a. Fish assemblages and rapid colonization after enlargement of an artificial reef off the Algarve coast (Southern Portugal). *Mar. Ecol.* 29, 435-448.
<https://doi.org/10.1111/j.1439-0485.2008.00253.x>.

Leitao, F., Santos, M.N., Erzini, K. and Monteiro, C.C., 2008b. The effect of predation on artificial reef juvenile demersal fish species. *Mar. Biol.* 153, 1233-1244. <https://doi.org/10.1007/s00227-007-0898-3>.

Levrel, H., Pioch, S., Spieler, R., 2012. Compensatory mitigation in marine ecosystems: which indicators for assessing the “no net loss” goal of ecosystem services and ecological functions?. *Mar. Policy* 36, 1202-1210. <https://doi.org/10.1016/j.marpol.2012.03.004>.

Lin, J., Su, W., 1994. Early phase of fish habitation around a new artificial reef off southwestern Taiwan. *Bull. Mar. Sci.* 55, 1112-1121.

Lincoln, S., Buckley, P., Howes, E., Maltby, K., Pinnegar, J., Ali, T., Alosairi, Y., Al-Ragum, A., Baglee, A., Balmes, C., Hamadou, R., Burt, J., Claereboudt, M., Glavan, J., Mamiit, R., Naser, H., Sedighi, O., Shokri, M., Shuhaibar, B., Wabnitz, C., Le Quesne, W., 2021. A regional review of marine and coastal impacts of climate change on the ROPME sea area. *Sustain.* 13, 13810. <https://doi.org/10.3390/su132413810>.

Lindberg, W., Seaman W., (Eds). 2011. *Guidelines and Management Practices for Artificial Reef Siting, Use, Construction, and Anchoring in Southeast Florida*. Florida Department of Environmental Protection. Miami, FL.

Lipcius, R., Burke, R., 2018. Successful recruitment, survival and long-term persistence of eastern oyster and hooked mussel on a subtidal, artificial restoration reef system in Chesapeake Bay. *PloS one* 13, e0204329. <https://doi.org/10.1371/journal.pone.0204329>.

Loughland, R., Bass, W., 2016. *Artificial reefs enhancing marine biodiversity in the Arabian Gulf*, Dimensions International, 20-25.

Love, M.S., E. Brothers, D.M. Schroeder, W.H. Lenarz. 2007. Ecological performance of young-of-the-year blue rockfish (*Sebastes mystinus*) associated with oil platforms and natural reefs in California as measured by daily growth rates. *Bull. Mar. Sci.* 80, 147-157.

Lukens, R., Selberg, C., 2004. *Guidelines for marine artificial reef materials*. Atlantic and Gulf States Marine Fisheries Commissions.

- Macreadie, P., Fowler, A., Booth, D., 2011. Rigs-to-reefs: will the deep sea benefit from artificial habitat?. *Front. Ecol. Environ.* 9, 455-461. <https://doi.org/10.1890/100112>.
- Maghsoudlou, A., Araghi, P., Wilson, S., Taylor, O., Medio, D., 2008. Status of coral reefs in the ROPME sea area (The Persian Gulf, Gulf of Oman, and Arabian Sea), in: Wilkinson, C., (Ed.), *Status of Coral Reefs of the World: 2008*, Australian Institute of Marine Science, Townsville, AU pp.79-90.
- Martin, D., Bertasi, F., Colangelo, M., de Vries, M., Frost, M., Hawkins, S., Macpherson, E., Moschella, P., Satta, M., Thompson, R., Ceccherelli, V., 2005. Ecological impact of coastal defence structures on sediment and mobile fauna: evaluating and forecasting consequences of unavoidable modifications of native habitats. *Coast. Eng.* 52, 1027-1051. <https://doi.org/10.1016/j.coastaleng.2005.09.006>.
- Mateos-Molina, D., Antonopoulou, M., Baldwin, R., Bejarano, I., Burt, J., García-Charton, J., Al-Ghais, S., Walgamage, J., Taylor, O., 2020. Applying an integrated approach to coastal marine habitat mapping in the north-western United Arab Emirates. *Mar. Environ. Res.* 161, p.105095. <https://doi.org/10.1016/j.marenvres.2020.105095>.
- Mateos-Molina, D., Lamine, E., Antonopoulou, M., Burt, J., Das, H., Javed, S., Judas, J., Khan, S., Muzaffar, S., Pilcher, N., Rodriguez-Zarate, C., 2021. Synthesis and evaluation of coastal and marine biodiversity spatial information in the United Arab Emirates for ecosystem-based management. *Mar. Pollut. Bull.* 167, 112319. <https://doi.org/10.1016/j.marpolbul.2021.112319>
- Matthews, K., 1985. Species similarity and movement of fishes on natural and artificial reefs in Monterey Bay, California. *Bull. Mar. Sci.* 37, 252-270.
- McCreless, E., Beck, M., 2016. Rethinking our global coastal investment portfolio. *J. Ocean Coast. Econ.* 3, 6. <https://doi.org/10.15351/2373-8456.1069>.
- McGlennon, D., Branden, K., 1994. Comparison of catch and recreational anglers fishing on artificial reefs and natural seabed in Gulf St. Vincent, South Australia. *Bull. Mar. Sci.* 55, 510-523.
- Melville-Rea, H., Eayrs, C., Anwahi, N., Burt, J., Holland, D., Samara, F., Paparella, F., Al Murshidi, A., Al-Shehhi, M., Holland, D., 2021. A Roadmap for Policy-Relevant Sea-Level Rise Research in the United Arab Emirates. *Front. Mar. Sci.* 907. <https://doi.org/10.3389/fmars.2021.670089>.
- Mohamed, M., 2021. Assessment of temporal variation of fish assemblages between natural and artificial reefs MS thesis, Qatar University, Doha, QA.
- Morley, D., Sherman, R., Jordan, L., Banks, K., Quinn, T., Spieler, R., 2008. Environmental enhancement gone awry: characterization of an artificial reef constructed from waste vehicle tires, in: Brebbia, C., (Ed.) *Environmental Problems in Coastal Regions VII*, WIT press, Wessex, UK, 73-87.
- Morris, R., Konlechner, T., Ghisalberti, M., Swearer, S., 2018. From grey to green: Efficacy of eco-engineering solutions for nature-based coastal defence. *Glob. Change Biol.* 24, 1827-1842. <https://doi.org/10.1111/gcb.14063>.
- Mousavi, S., Danehkar, A., Shokri, M., Poorbagher, H., Azhdari, D., 2015. Site selection for artificial reefs using a new combine Multi-Criteria Decision-Making (MCDM) tools for coral reefs in the Kish Island–Persian Gulf. *Ocean Coast. Manag.* 111, pp.92-102. <https://doi.org/10.1016/j.ocecoaman.2015.03.004>.

- Nadim, F., Bagtzoglou, A., Iranmahboob, J., 2008. Coastal management in the Persian Gulf region within the framework of the ROPME programme of action. *Ocean Coast. Manag.* 51, 556-565. <https://doi.org/10.1016/j.ocecoaman.2008.04.007>.
- Naser, H., 2015. The role of environmental impact assessment in protecting coastal and marine environments in rapidly developing islands: The case of Bahrain, Arabian Gulf. *Ocean Coast. Manag.* 104, 159-169. <https://doi.org/10.1016/j.ocecoaman.2014.12.009>.
- Neill, P., Alcalde, O., Faugeton, S., Navarrete, S., Correa, J., 2006. Invasion of *Codium fragile* ssp. *tomentosoides* in northern Chile: a new threat for *Gracilaria* farming. *Aquaculture* 259, 202-210. <https://doi.org/10.1016/j.aquaculture.2006.05.009>.
- Nicoletti, L., Marzioletti, S., Paganelli, D., Ardizzone, G., 2007. Long-term changes in a benthic assemblage associated with artificial reefs. *Hydrobiologia*, 580, 233-240. <https://doi.org/10.1007/s10750-006-0450-3>.
- Nouri, J., Danehkar, A., Sharifipour, R., 2008. Evaluation of ecotourism potential in the northern coastline of the Persian Gulf. *Environ. Geo.* 55, 681-686. <https://doi.org/10.1007/s00254-007-1018-x>.
- O'Shaughnessy, K., Hawkins, S., Evans, A., Hanley, M., Lunt, P., Thompson, R., Francis, R., Hoggart, S., Moore, P., Iglesias, G., Simmonds, D., 2020. Design catalogue for eco-engineering of coastal artificial structures: a multifunctional approach for stakeholders and end-users. *Urban Ecosyst.* 23, 431-443. <https://doi.org/10.1007/s11252-019-00924-z>.
- Patranella, A., Kilfoyle, K., Pioch, S., Spieler, R., 2017. Artificial reefs as juvenile fish habitat in a marina. *J. Coast. Res.* 33, 1341-1351. <https://doi.org/10.2112/JCOASTRES-D-16-00145.1>.
- Paxton, A., Newton, E., Adler, A., Van Hoeck, R., Iversen Jr., E., Taylor, J., Peterson, C., Silliman, B., 2020b. Artificial habitats host elevated densities of large reef-associated predators. *PloS one* 15, e0237374. <https://doi.org/10.1371/journal.pone.0237374>.
- Paxton, A., Shertzer, K., Bacheler, N., Kellison, G., Riley, K., Taylor, J., 2020a. Meta-analysis reveals artificial reefs can be effective tools for fish community enhancement but are not one-size-fits-all. *Front. Mar. Sci.* 7, p.282. <https://doi.org/10.3389/fmars.2020.00282>.
- Pears, R., Williams, D., 2005. Potential effects of artificial reefs on the Great Barrier Reef: background paper. CRC Reef Research Centre, Townsville, AU.
- Perkol-Finkel, S., Benayahu, Y., 2004. Community structure of stony and soft corals on vertical unplanned artificial reefs in Eilat (Red Sea): comparison to natural reefs. *Coral Reefs* 23, 195-205. <https://doi.org/10.1007/s00338-004-0384-z>.
- Perkol-Finkel, S., Benayahu, Y., 2005. Recruitment of benthic organisms onto a planned artificial reef: shifts in community structure one decade post-deployment. *Mar. Environ. Res.*, 59, 79-99. <https://doi.org/10.1016/j.marenvres.2004.03.122>.
- Pi-hai, G., Jiao, L., Chang-tao, G., Meng-Jie, L., Chao, L., 2014. Estimation and experiment of carbon sequestration by oysters attached to the enhancement artificial reefs in Laizhou Bay, Shandong, China. *Yingyong Shengtai Xuebao* 25(10).

- Pioch, S., Doumenge, J., 2010. Summary of French artificial reefs immersions since 1968, sites, volumes, types and costs. In: Ceccaldi, H., Dekeyser, I., Girault, M., Stora G., (Eds.), *Global Change: Mankind-Marine Environment Interactions* Springer, Dordrecht, NL, pp. 125-127. https://doi.org/10.1007/978-90-481-8630-3_22.
- Pitcher, T., Buchary, E., Hutton, T., 2002. Forecasting the benefits of no-take human-made reefs using spatial ecosystem simulation. *ICES J. Mar. Sci.* 59(suppl), S17-S26. <https://doi.org/10.1006/jmsc.2002.1185>.
- Polovina, J., 1989. Artificial reefs: nothing more than benthic fish aggregators. *Reports of California Cooperative Oceanic Fisheries Investigations*, 30, pp.37-39.
- Polovina, J., 1990. Assessment of biological impacts of artificial reefs and FADs. In: Pietersz, V. (Ed.), *Symposium on Artificial Reefs and Fish Aggregating Devices as Tools for the Management and Enhancement of Marine Fishery Resources*. Regional Office for Asia and the Pacific, United Nations Food and Agriculture Organisation, Bangkok, TH, pp.258-263.
- Polovina, J., Sakai, I., 1989. Impacts of artificial reefs on fishery production in Shimamaki, Japan. *Bull. Mar. Sci.* 44, 997-1003.
- Pondella, D., Stephens Jr., J., Craig, M., 2002. Fish production of a temperate artificial reef based on the density of embiotocids (Teleostei: Perciformes). *ICES J. Mar. Sci.* 59(suppl), S88-S93. <https://doi.org/10.1006/jmsc.2002.1219>.
- Rabaoui, L., Cusack, M., Saderne, V., Krishnakumar, P., Lin, Y., Shemsi, A., El Zrelli, R., Arias-Ortiz, A., Masqué, P., Duarte, C., Qurban, M., 2020. Anthropogenic-induced acceleration of elemental burial rates in blue carbon repositories of the Arabian Gulf. *Sci. Total Environ.* 719, p.135177. <https://doi.org/10.1016/j.scitotenv.2019.135177>.
- Raineault, N., Trembanis, A., Miller, D., Capone, V., 2013. Interannual changes in seafloor surficial geology at an artificial reef site on the inner continental shelf. *Cont. Shelf Res.* 58, 67-78. <https://doi.org/10.1016/j.csr.2013.03.008>.
- Ramm, L., Florisson, J., Watts, S., Becker, A., Tweedley, J., 2021. Artificial reefs in the Anthropocene: a review of geographical and historical trends in their design, purpose, and monitoring. *Bull. Mar. Sci.* 97, 699-728. <https://doi.org/10.5343/bms.2020.0046>.
- Reeds, K., Smith, J., Suthers, I., Johnston, E., 2018. An ecological halo surrounding a large offshore artificial reef: sediments, infauna, and fish foraging. *Mar. Environ. Res.* 141, 30-38. <https://doi.org/10.1016/j.marenvres.2018.07.011>.
- Relini, G., Relini, M., Palandri, G., Merello, S., Beccornia, E. 2007. History, ecology and trends for artificial reefs of the Ligurian sea, Italy, in: Relini, G., Ryland, J. (Eds.) *Biodiversity in Enclosed Seas and Artificial Marine Habitats*. *Developments in Hydrobiology*, vol 193. Springer, Dordrecht, NL, pp. 193-217. https://doi.org/10.1007/978-1-4020-6156-1_17
- Richer, R., 2009. Conservation in Qatar: Impacts of Increasing Industrialization. Occasional Paper. Centre for International and Regional Studies, Georgetown University of Foreign Service in Qatar, p. 23.

- Riegl, B., Johnston, M., Purkis, S., Howells, E., Burt, J., Steiner, S., Sheppard, C., Bauman, A., 2018. Population collapse dynamics in *Acropora downingi*, an Arabian/Persian Gulf ecosystem-engineering coral, linked to rising temperature. *Glob. Chang. Biol.* 24, 2447-2462. <https://doi.org/10.1111/gcb.14114>.
- Riegl, B., Purkis, S., 2015. Coral population dynamics across consecutive mass mortality events. *Glob. Chang. Biol.* 21, 3995-4005. <https://doi.org/10.1111/gcb.13014>.
- Rilov, G., Benayahu, Y., 2002. Rehabilitation of coral reef-fish communities: the importance of artificial-reef relief to recruitment rates. *Bull. Mar. Sci.* 70, 185-197.
- Roa-Ureta, R., Santos, M., Leitão, F., 2019. Modelling long-term fisheries data to resolve the attraction versus production dilemma of artificial reefs. *Ecol. Modell.* 407, 108727. <https://doi.org/10.1016/j.ecolmodel.2019.108727>.
- Ros, M., Vázquez-Luis, M., Guerra-García, J., 2013. The role of marinas and recreational boating in the occurrence and distribution of exotic caprellids (Crustacea: Amphipoda) in the Western Mediterranean: Mallorca Island as a case study. *J. Sea Res.* 83, 94-103. <https://doi.org/10.1016/j.seares.2013.04.004>.
- Rosciszewski-Dodgson, M., Cirella, G., 2022. The Pearl's Social and Environmental Failures: Development Challenges, in: Cirella G. (Ed.), *Human Settlements*. Springer, Singapore, pp. 215-234. https://doi.org/10.1007/978-981-16-4031-5_12.
- Ryan, C., Ninov, I., Aziz, H., 2012. Ras Al Khor—Eco-tourism in constructed wetlands: Post modernity in the modernity of the Dubai landscape. *Tour. Manag. Perspect.* 4, 185-197. <https://doi.org/10.1016/j.tmp.2012.08.006>.
- Salahuddin B., 2006. The marine environmental impacts of artificial island construction: Dubai, UAE. MS thesis, Duke University, Durham, NC.
- Sale, P., Feary, D., Burt, J., Bauman, A., Cavalcante, G., Drouillard, K., Kjerfve, B., Marquis, E., Trick, C., Usseglio, P., Van Lavieren, H., 2011. The growing need for sustainable ecological management of marine communities of the Persian Gulf. *Ambio*, 40, 4-17. <https://doi.org/10.1007/s13280-010-0092-6>.
- Salomidi, M., Katsanevakis, S., Issaris, Y., Tsiamis, K., Katsiaras, N., 2013. Anthropogenic disturbance of coastal habitats promotes the spread of the introduced scleractinian coral *Oculina patagonica* in the Mediterranean Sea. *Biol. Invasions* 15, 1961-1971. <https://doi.org/10.1007/s10530-013-0424-0>.
- Sanabria-Fernandez, J., Lazzari, N., Riera, R., Becerro, M.A., 2018. Building up marine biodiversity loss: artificial substrates hold lower number and abundance of low occupancy benthic and sessile species. *Mar. Environ. Res.* 140, pp.190-199. <https://doi.org/10.1016/j.marenvres.2018.06.010>.
- Santos, M., Monteiro, C., 1998. Comparison of the catch and fishing yield from an artificial reef system and neighbouring areas off Faro (Algarve, south Portugal). *Fish. Res.* 39, 55-65. [https://doi.org/10.1016/S0165-7836\(98\)00169-6](https://doi.org/10.1016/S0165-7836(98)00169-6).
- Sayer, M., Wilding, T., 2002. Planning, licensing, and stakeholder consultation in an artificial reef development: the Loch Linnhe reef, a case study. *ICES J. Mar. Sci.*, 59(suppl), S178-S185. <https://doi.org/10.1006/jmsc.2002.1270>.

- Schile, L., Kauffman, J., Crooks, S., Fourqurean, J., Glavan, J., Megonigal, J., 2017. Limits on carbon sequestration in arid blue carbon ecosystems. *Ecol. Appl.* 27, 859-874.
<https://doi.org/10.1002/eap.1489>.
- Schroepfer, R., Szedlmayer, S., 2006. Estimates of residence and site fidelity for red snapper *Lutjanus campechanus* on artificial reefs in the northeastern Gulf of Mexico. *Bull. Mar. Sci.* 78, 93-101.
- Schroeter, S., Reed, D., Raimondi, P., 2015. Effects of reef physical structure on development of benthic reef community: a large-scale artificial reef experiment. *Mar. Ecol. Prog. Ser.* 540, pp.43-55.
<https://doi.org/10.3354/meps11483>.
- Scott, M., Smith, J., Lowry, M., Taylor, M., Suthers, I., 2015. The influence of an offshore artificial reef on the abundance of fish in the surrounding pelagic environment. *Mar. Freshw. Res.* 66, 429-437.
<https://doi.org/10.1071/MF14064>.
- Seaman, W., 2007. Artificial habitats and the restoration of degraded marine ecosystems and fisheries. *Hydrobiologia* 580, 143-155. DOI 10.1007/s10750-006-0457-9.
- Sedano, F., Florido, M., Rallis, I., Espinosa, F., Gerovasileiou, V., 2019. Comparing sessile benthos on shallow artificial versus natural hard substrates in the Eastern Mediterranean Sea. *Mediterr. Mar. Sci.* 20 (Special Issue 4), 688-702. <https://doi.org/10.12681/mms.17897>.
- Seidelin, L., Wahlberg, M., Holmer, M., 2018. Using live-stream video from an artificial reef to increase interest in marine biology. *J. Mar. Sci. Eng.* 6, 47. <https://doi.org/10.3390/jmse6020047>.
- Sella, I., Perkol-Finkel S., 2015. Blue is the new green—ecological enhancement of concrete based coastal and marine infrastructure. *Ecol. Eng.* 84, 260-272. <https://doi.org/10.1016/j.ecoleng.2015.09.016>.
- Shams, A., 2002. Some environmental aspects of aquaculture and fisheries activities in the Kingdom of Bahrain, presented at the Workshop on Environmentally Friendly Aquaculture and Fisheries Practices, Hurghada, EG.
- Sheehy, D., Vik, S., 2010. The role of constructed reefs in non-indigenous species introductions and range expansions. *Ecol. Eng.* 36, 1-11. <https://doi.org/10.1016/j.ecoleng.2009.09.012>.
- Sheppard, C., Al-Husiani, M., Al-Jamali, F., Al-Yamani, F., Baldwin, R., Bishop, J., Benzoni, F., Dutrieux, E., Dulvy, N., Durvasula, S., Jones, D., 2010. The Gulf: a young sea in decline. *Mar. Pollut. Bull.* 60, 3-38.
<https://doi.org/10.1016/j.marpolbul.2009.10.017>.
- Sheppard, C., Price, A., Roberts, C., 1992. *Marine Ecology of the Arabian region: Patterns and Processes in Extreme Tropical Environments*. Academic Press, Toronto, CA.
- Siddeek, M., Fouda, M., Hermosa Jr., G., 1999. Demersal fisheries of the Arabian Sea, the Gulf of Oman and the Arabian Gulf. *Estuar. Coast. Shelf Sci.* 49, 87-97. [https://doi.org/10.1016/S0272-7714\(99\)80013-4](https://doi.org/10.1016/S0272-7714(99)80013-4).
- Simkanin, C., Davidson, I., Dower, J., Jamieson, G., Therriault, T., 2012. Anthropogenic structures and the infiltration of natural benthos by invasive ascidians. *Mar. Ecol.* 33, 499-511.
<https://doi.org/10.1111/j.1439-0485.2012.00516.x>.

- Simon, T., Pinheiro, H., Joyeux, J., 2011. Target fishes on artificial reefs: evidences of impacts over nearby natural environments. *Sci. Total Environ.* 409, 4579-4584.
<https://doi.org/10.1016/j.scitotenv.2011.07.057>.
- Smith, J., Cornwell, W., Lowry, M., Suthers, I., 2017. Modelling the distribution of fish around an artificial reef. *Mar. Freshw. Res.* 68, 1955-1964. <https://doi.org/10.1071/MF16019>.
- Smith, J., Lowry, M., Champion, C., Suthers, I., 2016. A designed artificial reef is among the most productive marine fish habitats: new metrics to address 'production versus attraction'. *Mar. Biol.* 163, 1-8. <https://doi.org/10.1007/s00227-016-2967-y>.
- Spieler, R., Gilliam, D., Sherman, R., 2001. Artificial substrate and coral reef restoration: what do we need to know to know what we need. *Bull. Mar. Sci.*, 69, 1013-1030.
- Steimle, F., Foster, K., Kropp, R. and Conlin, B., 2002. Benthic macrofauna productivity enhancement by an artificial reef in Delaware Bay, USA. *ICES J. Mar. Sci.* 59(suppl), S100-S105.
<https://doi.org/10.1006/jmsc.2002.1268>.
- Stephens Jr., J., Pondella, D., 2002. Larval productivity of a mature artificial reef: the ichthyoplankton of King Harbor, California, 1974–1997. *ICES J. Mar. Sci.* 59(suppl), S51-S58.
<https://doi.org/10.1006/jmsc.2002.1189>.
- Sun, P., Liu, X., Tang, Y., Cheng, W., Sun, R., Wang, X., Wan, R., Heino, M., 2017. The bio-economic effects of artificial reefs: mixed evidence from Shandong, China. *ICES J. Mar. Sci.* 74, 2239-2248.
<https://doi.org/10.1093/icesjms/fsx058>.
- Szedlmayer, S., Miller, D., 2018. A comparison of fish and epibenthic assemblages on artificial reefs with and without copper-based, anti-fouling paint. *American Fisheries Society Symposium* 86, 47-67.
- Tickell, S., Sáenz-Arroyo, A., Milner-Gulland, E., 2019. Sunken worlds: The past and future of human-made reefs in marine conservation. *BioScience*, 69, 725-735. <https://doi.org/10.1093/biosci/biz079>.
- Turpin, R., Bortone, S., 2002. Pre-and post-hurricane assessment of artificial reefs: evidence for potential use as refugia in a fishery management strategy. *ICES J. Mar. Sci.* 59(suppl), S74-S82.
<https://doi.org/10.1006/jmsc.2002.1191>.
- Van Lavieren, H., Burt, J., Feary, D., Cavalcante, G., Marquis, E., Benedetti, L., Trick, C., Kjerfve, B., and Sale, P., 2011. Managing the growing impacts of development on fragile coastal and marine ecosystems: Lessons from the Gulf. UNU-INWEH, Hamilton, CA.
- Van Lavieren, H., Klaus, R., 2013. An effective regional Marine Protected Area network for the ROPME Sea Area: Unrealistic vision or realistic possibility?. *Mar. Pollut. Bull.* 72, 389-405.
<https://doi.org/10.1016/j.marpolbul.2012.09.004>.
- Vaughan, G., Al-Mansoori, N., Burt, J., 2019. The Arabian Gulf, in: Sheppard, C. (Ed.), *World Seas: An Environmental Evaluation*. Elsevier Science, Amsterdam, NL. pp. 1-23.
- Walker, S., Schlacher, T., 2014. Limited habitat and conservation value of a young artificial reef. *Biodivers. Conserv.* 23, 433-447. <https://doi.org/10.1007/s10531-013-0611-4>.

- Walker, S., Schlacher, T., Schlacher-Hoenlinger, M., 2007. Spatial heterogeneity of epibenthos on artificial reefs: fouling communities in the early stages of colonization on an East Australian shipwreck. *Mar. Ecol.* 28, 435-445. <https://doi.org/10.1111/j.1439-0485.2007.00193.x>.
- Watanuki, N., Gonzales, B., 2006. The potential of artificial reefs as fisheries management tools in developing countries. *Bull. Mar. Sci.* 78, 9-19.
- Watchorn, D., Cowan, M., Driscoll, D., Nimmo, D., Ashman, K., Garkaklis, M., Wilson, B., Doherty, T., 2022. Artificial habitat structures for animal conservation: design and implementation, risks and opportunities. *Front. Ecol. Environ.*, 20, 301-309. <https://doi.org/10.1002/fee.2470>.
- Wendt, P., Knott, D., Van Dolah, R., 1989. Community structure of the sessile biota on five artificial reefs of different ages. *Bull. Mar. Sci.* 44, 1106-1122.
- Whitmarsh, D., Santos, M., Ramos, J., Monteiro, C., 2008. Marine habitat modification through artificial reefs off the Algarve (southern Portugal): An economic analysis of the fisheries and the prospects for management. *Ocean Coast. Manag.* 51, 463-468. <https://doi.org/10.1016/j.ocecoaman.2008.04.004>.
- Xu, Q., Zhang, L., Zhang, T., Zhang, X., Yang, H., 2017. Functional groupings and food web of an artificial reef used for sea cucumber aquaculture in northern China. *J. Sea Res.* 119, 1-7. <https://doi.org/10.1016/j.seares.2016.10.005>.
- Zhang, H., Zhu, K., Tang, J., 2005. The effect of fishery resource enhancement in the artificial reef area in the Haizhou Gulf. *Marine Fisheries*, 27, 38-43.

Table

Table 1. A summary of AR programs in the Arabian Gulf that are adequately described in the publicly-available literature, including information about the reef, the stated management goals and the reported results.

Country / reference(s)	Reef Information	Management Goals	Results of the AR Program
Kuwait / Downing et al. 1985	- 3 car tire reefs - 25 m ² x 0.8 m or 2.4 m high	Test if reefs attract fish	- ARs quickly attracted commercially important species. - 2.4 m reef attracted more fish biomass
Saudi Arabia / Loughland and Bass 2016	- Multiple 1 m ³ cubical concrete modules at 24 sites. - 1 "mega site" with central and peripheral modules connected by limestone basket spokes	Create habitat to enhance fisheries and biodiversity	Reefs became "thriving new ecosystems" with a variety of fish and other reef organisms.
Qatar / Deb et al. 2014	Almost 2800 coral colonies were transplanted to 550 limestone boulders and 213 low-relief concrete modules	Transplant corals away from pipeline sites	Initial transplant monitoring indicated: - 5% coral mortality, and similar health to reference corals - Corals support reef fauna - Coral recruitment
Qatar / Mohamed 2021	Concrete orb-shaped reef modules deployed in clusters	Compensate for lost natural reefs	ARs had lower fish species richness but higher fish densities than a natural reef.
Iran / Ajdari et al. 2011 Azhdari et al. 2012	- 1.5 x 1.2 domed concrete modules, 1.5 x 1.4 pyramid concrete modules and concrete construction waste deployed in different combinations - Each reef type combination replicated 3 times - Each replicate had 16 pieces of artificial reef material	Test if reefs increase fish catch	Reefs with all 3 materials performed best: - They had higher catches than 2 other reef types in Ajdari et al. 2011. - They had higher catches than other reef types and natural habitat in Azhdari et al. 2012. - Other reef types did not have higher catches than natural habitat
United Arab Emirates / Hopkins 2007	- 3 reefs deployed, each with: - A central concrete cylinder 1.4 m diameter x 2.5 m tall - 14 rows of plastic pipes with 11 pipes in each row radiating from the cylinder	Test if reefs enhance biodiversity	After two years, the reefs were covered by fouling organisms, including corals, and attracted a variety of fish.

United Arab Emirates / Al-Cibahy et al. 2009	144 ceramic branching modules deployed in 4 reefs	Test if reefs enhance biodiversity	In less than 1 year, artificial reefs had: - Higher fish biomass and slightly higher fish species richness - Higher coral % coverage compared with natural reference sites
United Arab Emirates / Harris and Schroder 2001	Various vessels sunk by fishers, private companies or the government to create ARs	Attract fish	- These wrecks attract a variety of fish, including commercially-important species. - They are used by recreational divers
United Arab Emirates / Harris and Schroder 2001	3000 1m ³ concrete construction blocks deployed in a 30 x 100 m area.	Create an AR for dive tourism	- Supports a variety of sessile invertebrates, including corals, and abundant fish.

Management goals of Gulf artificial reef projects





Fig. 1 Management goals of 13 Arabian Gulf AR projects with publicly-available stated goals. One project had more than one goal, and both are included in this chart. *Note to editors: this figure should be in color for online readers, but grayscale for print publication.*

Economic benefits of artificial reefs

Fisheries

Increase availability and catch
of commercial species



Produce more fish biomass by:

Providing predation refuges for juvenile fish

Creating recruitment habitat, if there is none "downstream"

Altering circulation to encourage settlement of recruits

Creating suitable breeding habitats

Increasing fish growth rates (more feeding opportunities)

Provide attachment sites for commercial bivalves

Tourism

Create dive sites



Create surfing locations

Marine Biology Education



Fig. 2 Infographic depicting possible economic benefits of artificial reefs.

Ecological benefits of artificial reefs

Biodiversity

Create distinct hard-bottom communities

Create attachment sites for habitat-forming corals, algae, bivalves, etc.



Coral transplantation sites



Ecological services

Water filtration by reef organisms

Carbon sequestration by reef organisms



Prevent trawling to protect habitat




Fig. 3 Infographic depicting possible ecological benefits of artificial reefs.

Possible negative consequences of artificial reefs

Opportunity costs

Time and money spent on artificial reefs could be better spent on different conservation / management projects instead.



Artificial reefs may be “sink” habitat

Juvenile fish that settle on artificial reefs may suffer high predation. They may survive better in natural habitat.



Overfishing

Artificial reefs attract fish and fishers. Increased catch rates on artificial reefs can contribute to overfishing.



“Halos” around artificial reefs

The benthic habitat and community around the reef will change due to altered currents and the organisms on the reef.




Facilitating the spread of invasives

Artificial reefs may allow invasive hard-bottom species to spread quicker and colonize new areas.



Contamination

Some waste materials used to create artificial reefs may be contaminated with heavy metals, PCB's and anti-fouling paint.



Waste disposal and habitat destruction

Trash dumping may masquerade as “reef creation”. Waste material can be transported to natural habitats by storms.



Fig. 4 Infographic depicting possible negative consequences of artificial reefs.




Fig. 5 Flowchart illustrating the policy recommendations in this paper for Gulf AR projects.

Declaration of competing interest

None of the authors has any competing interests to declare.

Note to the editors: I did click on the tool to complete a competing interests statement, provided on the website: <https://www.elsevier.com/journals/regional-studies-in-marine-science/2352-4855/guide-for-authors>

This took me to this website: <https://declarations.elsevier.com/home>

This website was a “dead-end” for me, however, as it did not link to any shared statement that I could access. Sorry, I tried.