

2017-04

Cascading of high salinity bottom waters from the Arabian/Persian Gulf to the northern Arabian Sea

Shapiro, Georgy

<http://hdl.handle.net/10026.1/20418>

EGU General Assembly Conference abstracts 19, 7366

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.



Cascading of high salinity bottom waters from the Arabian/Persian Gulf to the northern Arabian Sea

Georgy Shapiro (1), Fred Wobus (1), Vladimir Solovyev (2), Xavier Francis (1), Patrick Hyder (3), Feng Chen (1), and Muhammad Asif (1)

(1) University of Plymouth, School of Marine Science and Engineering, Plymouth, Devon, United Kingdom (gshapiro@plymouth.ac.uk), (2) Aqualog LTD, Moscow, Russia, (3) Met Office, Fitzroy Road, Exeter, Devon, United Kingdom

Cascading (aka shelf convection) is a specific type of buoyancy driven current in which dense water is formed over the continental shelf and then descends down the slope to a greater depth. The cascades of dense water down continental slopes provide a mechanism for shelf–ocean exchange in many parts of the world’s oceans (Shapiro et al, 2003). Dense water is formed on the shelf by a number of processes, with high evaporation, limited river discharge and low precipitation being the major processes in warm climates (Ivanov et al, 2004). The formation and outflow of high salinity waters in the near-bottom layer of the Arabian/Persian Gulf is an example of dense water cascading (Bower et al 2000). Despite of its importance for the self-cleaning and the state of the marine ecosystem in the Arabian/Persian Gulf, the properties of the outflow have so far mainly been analysed using climatologically averaged data or observations of a limited set of parameters (mainly temperature), see (Bower et al 2000).

In this paper we study the dynamics of the flow using a comprehensive set of observational data (temperature, salinity velocity and turbidity profiles) obtained during the GRASP (Gulf Reconnaissance And Selective Profiling) observational campaign in the Gulf of Oman, which are complemented by the results of numerical modelling of the area using a number of 3D ocean models, and some ARGO T/S profiles. The GRASP measurements were carried out using an Aqualog climbing moored profiler, which was equipped with a Seabird CTD sensor, a Nortek Aquadopp current meter and a Seapoint turbidity meter. The Ocean circulation models used in the study include PGM4 and IND12 (UK Met Office); and AS20 and AG60 (University of Plymouth). All models are based on NEMO (Nucleus for European Modelling of the Ocean) codebase with a resolution from 9 km down to 1.8 km. The models were calibrated and validated against ARGO float profiles in the area.

The study revealed the mesoscale and sub-mesoscale circulation patterns of the outflow, their spatial and temporal variability over time scales from a few days to seasonal.

References

- Shapiro, G.I.; Huthnance, J.M.; Ivanov, V.V.. 2003 Dense water cascading off the continental shelf. *Journal of Geophysical Research*, 108 (C12). 3390.10.1029/2002JC001610
- Ivanov, V.V.; Shapiro, G.I.; Huthnance, J.M.; Aleynik, D.L.; Golovin, P.N.. 2004 Cascades of dense water around the world ocean. *Progress in Oceanography*, 60 (1). 47-98.10.1016/j.pocean.2003.12.002
- Bower, A. S., H. D. Hunt and J. Price, 2000. Character and Dynamics of the Red Sea and Persian Gulf Outflows. *Journal of Geophysical Research - Oceans*, Vol. 105, No. C3, pp. 6387-6414.