

2022-08

Matrix compression along isogenic blocks

Belton, Alexander

<http://hdl.handle.net/10026.1/20465>

10.1007/s44146-022-00023-0

Acta Scientiarum Mathematicarum

Springer Science and Business Media LLC

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.



Matrix compression along isogenic blocks

ALEXANDER BELTON, DOMINIQUE GUILLOT, APOORVA KHARE,
and MIHAI PUTINAR

Abstract. A matrix-compression algorithm is derived from a novel isogenic block decomposition for square matrices. The resulting compression and inflation operations possess strong functorial and spectral-permanence properties. The basic observation that Hadamard entrywise functional calculus preserves isogenic blocks has already proved to be of paramount importance for thresholding large correlation matrices. The proposed isogenic stratification of the set of complex matrices bears similarities to the Schubert cell stratification of a homogeneous algebraic manifold. An array of potential applications to current investigations in computational matrix analysis is briefly mentioned, touching concepts such as symmetric statistical models, hierarchical matrices and coherent matrix organization induced by partition trees.

1. Introduction

1.1. Prelude

In a previous paper [4] we were concerned with the study of entrywise positivity preservers acting on the cone of $N \times N$ positive semidefinite matrices. By a classical observation of Loewner, as developed by Horn in [16, Theorem 1.2], if the real-valued function f defined on the positive half line preserves positive semidefiniteness for all $N \times N$ matrices via entrywise Hadamard calculus (for brevity, we say that f *preserves positivity*), then f is necessarily of class $C^{(N-3)}$ and has non-negative derivatives up to this order. Moreover, the derivatives of order up to $N - 1$ are non-negative if they exist. In the case where f is known to be analytic, preserving positivity only for rank-one $N \times N$ matrices already implies that the first N non-zero Taylor coefficients of f are strictly positive. In particular, this holds for f a polynomial. Henceforth A^{ok} denotes the entrywise (fractional) power of a matrix A .

Article history: received 30.12.2021, revised 17.3.2022, accepted 18.3.2022.

AMS Subject Classification: 65F55; 65F45, 15A86, 47A60, 14M15.

Key words and phrases: structured matrix, Hadamard calculus, matrix compression, conditional expectation.

The central result in our previous work [4] (subsequently refined by one of us with Tao [19]) is a closed form for the greatest lower bound for the only possible negative coefficient c_M in the linear pencil

$$p(A) = c_0 A^{o n_0} + c_1 A^{o n_1} + \dots + c_{N-1} A^{o n_{N-1}} + c_M A^{o M}, \tag{1.1}$$

with real powers $n_0 < \dots < n_{N-1} < M$ lying in $\{0, 1, 2, \dots\} \cup [N - 2, \infty)$, such that $p(A)$ is positive semidefinite for any positive semidefinite $N \times N$ matrix A with entries in a specific domain in \mathbb{C} . The threshold value for the coefficient c_M was obtained via a combinatorial formula involving Schur polynomials.

A more natural matrix-theoretic approach to obtain the greatest lower bound for c_M in (1.1) is to study the spectrum of the Rayleigh quotient R of two quadratic forms,

$$R = R(A, \mathbf{u}) := \frac{\mathbf{u}^* A^{o M} \mathbf{u}}{\mathbf{u}^* (c_0 A^{o n_0} + c_1 A^{o n_1} + \dots + c_{N-1} A^{o n_{N-1}}) \mathbf{u}}. \tag{1.2}$$

Finding the maximum of R with respect to both \mathbf{u} and A indeed yields the critical threshold for preserving positivity as mentioned above; we refer the reader to [4, Section 4] and [19, Proposition 11.2]. One first obtains a closed-form expression for $R_A := \sup_{\mathbf{u} \in \mathbb{R}^N} R(A, \mathbf{u})$ and then maximizes with respect to A . However, a significant difficulty arises when using this approach:

the function $A \mapsto R_A$ is not continuous on the set of $N \times N$ positive semidefinite matrices.

As shown in [4], discontinuities occur precisely where constant-block structures emerge in A . The discovery of this phenomenon led the authors to investigate, in [4, Section 5], the simultaneous kernel $\bigcap_{k \geq 0} A^{o k}$ of the Hadamard powers of an arbitrary positive semidefinite matrix A , and a related Schubert cell-type stratification of the set of all positive semidefinite matrices of a given size.

A series of remarkable properties of this stratification is unveiled in the present work. In order to facilitate access to the concepts and notation in subsequent sections, we start by explaining informally the main ideas with the help of a simple example. Consider the 6×6 real symmetric matrix

$$A := \begin{pmatrix} 1 & 1 & 2 & -3 & 2 & 1 \\ 1 & 1 & 2 & -3 & 2 & 1 \\ 2 & 2 & 6 & -5 & 6 & 2 \\ -3 & -3 & -5 & 10 & -5 & -3 \\ 2 & 2 & 6 & -5 & 6 & 2 \\ 1 & 1 & 2 & -3 & 2 & 1 \end{pmatrix}.$$

Each row and column of A is constant on the sets in the partition

$$\pi = \{\{1, 2, 6\}, \{3, 5\}, \{4\}\}$$

of the index set $\{1, 2, 3, 4, 5, 6\}$. This observation gives at once some null vectors for the matrix A : those annihilated by the pattern of the partition, $(1, 1, 0, 0, 0, 1)$, $(0, 0, 1, 0, 1, 0)$ and $(0, 0, 0, 1, 0, 0)$. In fact, the kernel of every positive Hadamard power of A , including A itself, is precisely this set of vectors.

Given any matrix $A' \in \mathbb{C}^{N \times N}$, there is a unique coarsest partition π' of $\{1, \dots, N\}$ such that A' is constant on the blocks determined by π' . We denote by $\mathcal{S}_{\pi'}$ the set of all matrices having this constant-block structure. This naturally yields a stratification of the space of complex matrices,

$$\mathbb{C}^{N \times N} = \bigsqcup_{\pi'} \mathcal{S}_{\pi'}, \tag{1.3}$$

where the disjoint union is taken over all partitions of $\{1, \dots, N\}$. We refer to each set $\mathcal{S}_{\pi'}$ as a *stratum*, and call the above decomposition the *isogenic block stratification* of $\mathbb{C}^{N \times N}$.

We explore several applications of the stratification below. A natural operation is the compression $\Sigma_{\pi}^{\downarrow}$ of each constant block to a single entry: with A and π as above,

$$\Sigma_{\pi}^{\downarrow}(A) = \begin{pmatrix} 1 & 2 & -3 \\ 2 & 6 & -5 \\ -3 & -5 & 10 \end{pmatrix}.$$

This straightforward transformation has some remarkable and very useful properties. For example, the spectrum of $D_{\pi}^{1/2} \Sigma_{\pi}^{\downarrow}(A) D_{\pi}^{1/2}$ has the same non-zero eigenvalues, counting multiplicities, as the original matrix A , where the matrix $D_{\pi} = \text{diag}(3, 2, 1)$ simply reflects the size of the blocks of π . The spectrum of A has 3 additional zero eigenvalues, corresponding to its kernel.

An even more striking use of compression is for the computation of the Moore–Penrose generalized inverse: with A as above, this is

$$A^{\dagger} = \frac{1}{36} \begin{pmatrix} 140 & 140 & -30 & 96 & -30 & 140 \\ 140 & 140 & -30 & 96 & -30 & 140 \\ -30 & -30 & 9 & -18 & 9 & -30 \\ 96 & 96 & -18 & 72 & -18 & 96 \\ -30 & -30 & 9 & -18 & 9 & -30 \\ 140 & 140 & -30 & 96 & -30 & 140 \end{pmatrix}.$$

In general, there is a fast method to compute the pseudo-inverse of a block matrix:

$$A^{\dagger} = \Sigma_{\pi}^{\uparrow} (D_{\pi}^{-1} \Sigma_{\pi}^{\downarrow}(A)^{-1} D_{\pi}^{-1}), \tag{1.4}$$

where the inflation map Σ_{π}^{\uparrow} expands a matrix into one with blocks which are constant for the index-set partition π .

While the isogenic block structure considered in the present article arose from the entrywise matrix operations which preserve positivity, and is highly relevant to them, its simplicity and versatility invites the exploration of its potential utility beyond pure mathematics, to other areas of much current interest. Compression, fast computation and analysis of structured matrices are of great importance at present. For example, large positive semidefinite matrices make an important appearance in the analysis of big data, where they occur as covariance or correlation matrices of random vectors. In such settings, the rows and columns of the matrix are ordered according to the ordering of the underlying variables. Entrywise operations such as thresholding are natural in that setting (see [23], for example), as are grouping and averaging variables, and the related block operations on the corresponding matrices. We conclude the paper by indicating several such applications.

On the other hand, two notable connections with established areas of mathematics should be mentioned. The partition (1.3) of $\mathbb{C}^{N \times N}$ into subsets \mathcal{S}_π – and more generally, as in Proposition 2.6 – is akin to the stratification of the flag variety in *Schubert calculus*. It too emerges from a group action – of $G^{N \times N}$ acting entrywise on $\mathbb{C}^{N \times N}$.

Second, the matrix compression and inflation operations we focus on are a very specific instance of a *conditional expectation* map. Originating in probability theory, conditional expectations were generalized to the non-commutative setting of operator algebras with hard to underestimate benefits. From the ample bibliography on the subject we only mention an inspired, very recent survey [6] which presents Blecher and Read's extension of positivity beyond the well-known C^* -algebraic setting.

In the present work, aimed at a large audience of practitioners of matrix analysis, we do not pursue in detail these two paths.

1.2. Summary of contents

In Section 2, we study the stratification of complex matrices with blocks that lie in single orbits under the action of a multiplicative subgroup G of the group of units \mathbb{C}^\times , which generalises the case $G = \{1\}$ discussed above. In Section 3, we show that the same stratification naturally emerges from studying simultaneous entrywise powers or functions of matrices that are positive semidefinite; in fact, we do not even require the positivity of determinants of size 4 or more.

In Section 4, we investigate the compression $\Sigma_\pi^\downarrow(A)$, which collapses every block of the matrix A to a single entry, equal to the average over that block. Compression and its right inverse, inflation, provide an effective tool for operating inside each stratum or its closure. Section 5 contains results that describe how compression and inflation relate to spectra and the functional calculus. These exploit the observation that a weighted version of the compression map

provides a $*$ -isomorphism of unital C^* algebras between the stratum given by a partition with m elements and $\mathbb{C}^{m \times m}$.

The final section, Section 6, is devoted to providing examples and links to other areas of matrix analysis. Although our study was prompted by the structure of matrices arising in the statistics of big data, the isogenic stratification we introduce here is relevant to sparse-matrix compression procedures and related fast computational tools. Indeed, the averaging on isogenic blocks we propose is a simple and efficient method of eliminating the redundancy of operations involving this class of structured matrices. The rapidly developing theory of *hierarchical matrices* [7, 12, 14] is a natural framework where our study finds deep resonances. A second setting where isogenic stratifications are very natural is that of *coherent matrix organization* [9–11, 20]. An immediate consequence of our technique is the quick computation of spectra and singular values for structured matrices; other possible applications are the efficient solution of large linear systems and certifying the stability of evolution semigroups.

1.3. Acknowledgements

The authors extend their thanks to the International Centre for Mathematical Sciences, Edinburgh, where part of this work was carried out, and to an anonymous referee, for their helpful comments. D.G. was partially supported by a University of Delaware Research Foundation grant, by a Simons Foundation collaboration grant for mathematicians, and by a University of Delaware Research Foundation Strategic Initiative grant. A.K. was partially supported by Ramanujan Fellowship SB/S2/RJN-121/2017, MATRICS grant MTR/2017/000295, and SwarnaJayanti Fellowship grants SB/SJF/2019-20/14 and DST/SJF/MS/2019/3 from SERB and DST (Govt. of India), by grant F.510/25/CAS-II/2018(SAP-I) from UGC (Govt. of India), by a Tata Trusts gravel grant, and by a Young Investigator Award from the Infosys Foundation. M.P. was partially supported by a Simons Foundation collaboration grant.

1.4. List of symbols

We collect below some notation introduced and used throughout the text.

- $\overline{D}(0, \rho)$ is the closed disc in \mathbb{C} with radius ρ centered at the origin, S^1 is the unit circle in \mathbb{C} and \mathbb{C}^\times is the set of non-zero complex numbers; more generally, the group of units for a unital commutative ring R is denoted by R^\times .
- $\mathbf{1}_{M \times N}$ is the $M \times N$ matrix with each entry equal to 1, whereas Id_N is the $N \times N$ identity matrix.
- $f[A]$ is the matrix obtained by applying the function f to each entry of the matrix A .

- A^{α} is the matrix obtained from A by taking the α th power of each entry, whenever this is well defined. We set $0^0 := 1$.
- A^\dagger is the Moore–Penrose pseudo-inverse of A .
- (Π_N, \prec) is the poset of partitions of $\{1, \dots, N\}$, where $\pi' \prec \pi$ if π is a refinement of π' .
- D_π is the $m \times m$ diagonal matrix $\text{diag}(|I_1|, \dots, |I_m|)$, for any partition $\pi = \{I_1, \dots, I_m\} \in \Pi_N$.
- $\Sigma_\pi^\downarrow(A)$ is the $m \times m$ matrix obtained by averaging the $N \times N$ matrix A over each block determined by the m -element partition π .
- $\Theta_\pi^\downarrow(A)$ is the weighted $m \times m$ compression $D_\pi^{1/2} \Sigma_\pi^\downarrow(A) D_\pi^{1/2}$ of the $N \times N$ matrix A .
- $\Sigma_\pi^\uparrow(B)$ is the $N \times N$ matrix given by inflating each entry of the $m \times m$ matrix B to a constant block subordinate to the m -element partition π .
- $\Theta_\pi^\uparrow(B)$ is the weighted $N \times N$ inflation $\Sigma_\pi^\uparrow(D_\pi^{-1/2} B D_\pi^{-1/2})$ of the $m \times m$ matrix B .
- $\mathcal{P}_N(\Omega)$ is the set of $N \times N$ positive semidefinite matrices with entries in the set Ω .

2. Isogenic stratification of complex matrices

In order to define the isogenic stratification, we begin with a summary of results appearing in our earlier work.

Theorem 2.1. ([4, Theorems 5.1 and 5.8]) *Fix a multiplicative subgroup $G \subset \mathbb{C}^\times$, an integer $N \geq 1$, and a non-zero matrix $A \in \mathcal{P}_N(\mathbb{C})$.*

(1) *Suppose $\{I_1, \dots, I_m\}$ is a partition of $\{1, \dots, N\}$ satisfying the following two conditions.*

(a) *Each diagonal block A_{I_j} of A is a submatrix having rank at most one, and $A_{I_j} = \mathbf{u}_j \mathbf{u}_j^*$ for a unique $\mathbf{u}_j \in \mathbb{C}^{|I_j|}$ with first entry $\mathbf{u}_{j,1} \in [0, \infty)$.*

(b) *The entries of each diagonal block A_{I_j} lie in a single G -orbit.*

Then there exists a unique matrix $C = (c_{ij})_{i,j=1}^m$ such that $c_{ij} = 0$ unless $\mathbf{u}_i \neq 0$ and $\mathbf{u}_j \neq 0$, and A is a block matrix with

$$A_{I_i \times I_j} = c_{ij} \mathbf{u}_i \mathbf{u}_j^* \quad (1 \leq i, j \leq m).$$

Moreover, the entries of each off-diagonal block of A also lie in a single G -orbit. Furthermore, the matrix $C \in \mathcal{P}_m(\overline{D}(0, 1))$, and the matrices A and C have equal rank.

(2) *Consider condition (c).*

(c) *The diagonal blocks of A have maximal size, that is, each diagonal block is not contained in a larger diagonal block that has rank one.*

There exists a partition $\{I_1, \dots, I_m\}$ such that (a), (b) and (c) hold, and such a partition is unique up to relabelling of the indices.

- (3) Suppose (a)–(c) hold and $G = \mathbb{C}^\times$. Then the off-diagonal entries of C lie in the open disc $D(0, 1)$.
- (4) If $G \subset S^1$, then blocks in a single G -orbit automatically have rank at most one.

Theorem 2.1 naturally leads to Schubert-cell type stratifications of the cone $\mathcal{P}_N(\mathbb{C})$, and some properties of these stratifications were studied in [4]. Here, we explore a coarser form of partitioning on the whole set of $N \times N$ complex matrices, based solely on G -equivariance.

Definition 2.2. We denote by (Π_N, \prec) the poset of all *partitions* of the set $\{1, \dots, N\}$, ordered so that $\pi' \prec \pi$ if and only if π is a refinement of π' .

The poset Π_N is a lattice. Given $\pi, \pi' \in \Pi_N$, let $\pi \wedge \pi'$ and $\pi \vee \pi'$ denote the meet and join of π and π' , respectively: see [27, Example 3.10.4], but note that our ordering is opposite to the one employed there. For example, the meet and join of the partitions

$$\pi = \{\{1, 2\}, \{3, 4\}, \{5\}\} \quad \text{and} \quad \pi' = \{\{1, 3, 4\}, \{2\}, \{5\}\}$$

are

$$\pi \wedge \pi' = \{\{1, 2, 3, 4\}, \{5\}\} \quad \text{and} \quad \pi \vee \pi' = \{\{1\}, \{2\}, \{3, 4\}, \{5\}\}.$$

The lattice Π_N has maximum element $\pi_\vee := \{\{1\}, \dots, \{N\}\}$ and minimum element $\pi_\wedge := \{\{1, \dots, N\}\}$.

Theorem 2.3. Fix integers $M, N \geq 1$, a unital commutative ring R , and a multiplicative subgroup G of units in R^\times . Given any matrix $A \in R^{M \times N}$, there exist unique minimal (that is, coarsest) partitions $\pi_{\min} = \{I_1, \dots, I_m\} \in \Pi_M$ and $\varpi_{\min} = \{J_1, \dots, J_n\} \in \Pi_N$ such that the entries of the block submatrix $A_{I_i \times J_j}$ lie in a single G -orbit, for all $i \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$.

Now suppose $M = N$. If either A is symmetric, or $R = \mathbb{C}$, $G \subset S^1$, and A is Hermitian, then $\pi_{\min} = \varpi_{\min}$.

Proof. The non-trivial part is to establish uniqueness. To do so, we claim that if (π_1, ϖ_1) and $(\pi_2, \varpi_2) \in \Pi_M \times \Pi_N$ satisfy the property in the assertion, then so does $(\pi_1 \wedge \pi_2, \varpi_1 \wedge \varpi_2)$. The meet of $\pi_1 \wedge \pi_2$ can be constructed as follows: connect i and $i' \in \{1, \dots, M\}$ by an edge if they lie in the same block of π_1 or π_2 ; this defines a graph with vertex set $\{1, \dots, M\}$ whose connected components yield the blocks of the partition $\pi_1 \wedge \pi_2$. Denote this equivalence relation by $i \sim i'$ in $\pi_1 \wedge \pi_2$.

Now suppose $i \sim i'$ in $\pi_1 \wedge \pi_2$ and, similarly, $j \sim j'$ in $\varpi_1 \wedge \varpi_2$, so there are paths joining them, each of whose vertices lies in a block of π_1 or π_2 , and ϖ_1 or ϖ_2 , respectively. Denote these paths by

$$i = i_0 \leftrightarrow i_1 \leftrightarrow \dots \leftrightarrow i_r = i' \quad \text{and} \quad j = j_0 \leftrightarrow j_1 \leftrightarrow \dots \leftrightarrow j_s = j'.$$

We claim that $a_{ij} \in Ga_{i'j'}$. Indeed, using the above paths,

$$a_{ij} = a_{i_0j} \in Ga_{i_1j} = Ga_{i_2j} = \cdots = Ga_{i_rj} = Ga_{i'j_0} = Ga_{i'j_1} = \cdots = Ga_{i'j'},$$

and this proves the claim.

Now suppose $M = N$ and A is symmetric. Then the partition $(\pi_{\min}, \varpi_{\min})$ works for A , and $(\varpi_{\min}, \pi_{\min})$ for $A^T = A$, whence the above analysis shows $\pi_{\min} \wedge \varpi_{\min}$ works for both rows and columns of A . By minimality, it follows that $\pi_{\min} = \pi_{\min} \wedge \varpi_{\min} = \varpi_{\min}$.

Finally, suppose $R = \mathbb{C}$, $G \subset S^1$, and A is Hermitian. We claim that $(\pi_{\min}, \varpi_{\min})$ works for A as well as for $A^T = \bar{A}$; by the previous paragraph, this gives the result. To show the claim, note that if (π, ϖ) works for A and $G \subset S^1$, then (π, ϖ) works for \bar{A} , as G is closed under conjugation: if $\zeta \in G \subset S^1$ then $a_{ij} = \zeta a_{i'j'}$ if and only if $\bar{a}_{ij} = \bar{\zeta} \bar{a}_{i'j'} = \zeta^{-1} a_{i'j'}$. ■

Theorem 2.3 has a useful “symmetric” version for square matrices, in the sense that the same partition is used for both the rows and the columns. The proof is similar to that of Theorem 2.3 and is hence omitted.

Proposition 2.4. *Fix an integer $N \geq 1$, and a multiplicative subgroup G of units in a unital commutative ring R . Given $A \in R^{N \times N}$, there exists a unique minimal partition $\pi = \{I_1, \dots, I_m\} \in \Pi_N$, such that the entries of the block submatrix $A_{I_i \times I_j}$ lie in a single G -orbit, for all $i, j \in \{1, \dots, N\}$.*

The following definitions follow naturally from the previous proposition. Our focus henceforth is on complex matrices with blocks which are orbits of a fixed multiplicative subgroup G of S^1 , and primarily the case $G = \{1\}$, where the entries in any given block are identical.

Definition 2.5. Given a matrix $A \in \mathbb{C}^{N \times N}$ and a multiplicative group $G \subset S^1$, let $\pi^G(A) \in \Pi_N$ be the partition provided by Proposition 2.4 for the matrix A .

Conversely, for a given partition $\pi \in \Pi_N$, define the stratum \mathcal{S}_π^G to be

$$\mathcal{S}_\pi^G := \{A \in \mathbb{C}^{N \times N} : \pi^G(A) = \pi\}.$$

The proof of the following proposition is immediate.

Proposition 2.6. *Given $N \geq 1$ and a multiplicative subgroup G of S^1 , there is a natural stratification of the set of $N \times N$ complex matrices*

$$\mathbb{C}^{N \times N} = \bigsqcup_{\pi \in \Pi_N} \mathcal{S}_\pi^G,$$

and the stratum \mathcal{S}_π^G has closure

$$\overline{\mathcal{S}_\pi^G} = \bigsqcup_{\pi' \prec \pi} \mathcal{S}_{\pi'}^G \tag{2.1}$$

when $\mathbb{C}^{N \times N}$ is equipped with its usual topology.

Equation (2.1) may be seen as akin to the Schubert-cell decomposition of the flag variety corresponding to a semisimple Lie group.

Definition 2.7. The family of stratifications given by Proposition 2.6 will be referred to as *G-block stratifications* of the space $\mathbb{C}^{N \times N}$.

Remark 2.8. There is an important distinction between the stratification of $\mathbb{C}^{N \times N}$ considered here and that for the cone $\mathcal{P}_N(\mathbb{C})$ considered previously. In Theorem 2.1, the partition was defined to have the property that the diagonal blocks of $A \in \mathcal{P}_N(\mathbb{C})$ have rank at most one. However, for a general matrix $A \in \mathbb{C}^{N \times N}$, this extra property need not hold for $\pi^G(A)$, unless either $G = \{1\}$, or $A \in \mathcal{P}_N(\mathbb{C})$ and $G \subset S^1$. In fact, as shown in [5, Proposition 4.6], in the latter case the requirement in Theorem 2.1 that A is positive semidefinite may be relaxed by requiring A to be *3-PMP*: every principal minor of size no more than 3×3 is non-negative.

3. PMP matrices and simultaneous kernels of entrywise powers

Henceforth, we will focus on constant-block stratifications, which we call *isogenic*, and we work mainly with square complex matrices.

We begin by noting how, for a large class of Hermitian matrices, the partition $\pi^{\{1\}}(A)$ introduced in Definition 2.5 emerges naturally from the study of simultaneous kernels of entrywise powers of A .

Definition 3.1. ([5]) Given an integer $k \geq 0$, a complex Hermitian matrix is said to be *k-PMP* (principal minor positive) if every $l \times l$ principal minor is non-negative, for all $l \in \{1, \dots, k\}$.

The *k-PMP* matrices interpolate between Hermitian matrices, the case when $k = 0$, and positive semidefinite matrices, where k is maximal.

Theorem 3.2. ([5, Theorem 5.1]) *Let the Hermitian matrix $A \in \mathbb{C}^{N \times N}$ be 3-PMP, and let $\pi' = \{I'_1, \dots, I'_{m'}\}$ be any partition refined by $\pi = \pi^{\{1\}}(A) = \{I_1, \dots, I_m\}$. The following spaces are equal.*

- (1) *The simultaneous kernel of $\mathbf{1}_{N \times N}, A, \dots, A^{(N-1)}$.*
- (2) *The simultaneous kernel of $A^{\circ n}$ for all $n \geq 0$.*
- (3) *The simultaneous kernel of the block-diagonal matrices*

$$\text{diag } A_{\pi'}^{\circ n} := \bigoplus_{j=1}^{m'} A_{I'_j \times I'_j}^{\circ n}$$

for all $n \geq 0$.

- (4) *The kernel of the matrix $J_\pi := \bigoplus_{j=1}^m \mathbf{1}_{I_j \times I_j}$.*

This equality of kernels need not hold for matrices that are not 3-PMP.

Our immediate goal is to show that a similar result holds for arbitrary real powers.

Proposition 3.3. *Let the Hermitian matrix $A \in (0, \infty)^{N \times N}$ be 3-PMP, and let $\pi' = \{I'_1, \dots, I'_{m'}\}$ be any partition refined by $\pi = \pi^{\{1\}}(A) = \{I_1, \dots, I_m\}$. Fix real powers $n_1 < \dots < n_N$. The following spaces are equal.*

- (1) *The simultaneous kernel of $A^{\circ n_j}$ for $j = 1, \dots, N$.*
- (2) *The simultaneous kernel of $A^{\circ \alpha}$ for all $\alpha \in \mathbb{R}$.*
- (3) *The simultaneous kernel of the block-diagonal matrices*

$$\text{diag } A_{\pi'}^{\circ \alpha} := \bigoplus_{j=1}^{m'} A_{I'_j \times I'_j}^{\circ \alpha}$$

for all $\alpha \in \mathbb{R}$.

- (4) *The kernel of the matrix $J_\pi := \bigoplus_{j=1}^m \mathbf{1}_{I_j \times I_j}$.*

The same holds when A has non-negative entries with at least one zero entry, as long as $n_1 = 0$ and α is taken to be non-negative in (2) and (3).

These equalities of kernels need not hold for matrices that are not 3-PMP.

Remark 3.4. One subtlety in adapting the proof in [5] of Theorem 3.2 to these variants is that a key matrix identity relating $A^{\circ \alpha}$ to $A^{\circ n_1}, \dots, A^{\circ n_N}$ is no longer available in the required generality. In [4, Lemma 3.5], it is shown that if A is an $N \times N$ matrix then

$$A^{\circ M} = \sum_{j=1}^N (-1)^{N-j} D_{M,j} A^{\circ(j-1)} \quad (M \geq N),$$

where $D_{M,j}$ is a diagonal matrix composed of certain Schur polynomials evaluated at the rows of A . If one tries to generalize this identity from exponents $\{0, 1, \dots, N - 1\}$ to arbitrary real powers, as in [19], the entries of the diagonal matrices become ratios of generalized Vandermonde determinants, and such ratios are not defined for all A . As a result, the above identity does not admit a uniform generalization, so we cannot naively adapt the previous proof to show that the subspace in (2) contains that in (1). There is a similar issue when α is not greater than n_N , which cannot be resolved by modifying the arguments in [5].

In light of the preceding remark, it is heartening that all three variants above follow from an even stronger result, that holds over arbitrary subsets of \mathbb{C} and for more general functions than powers. To state this result, we introduce the following notion, over an arbitrary commutative ring.

Definition 3.5. Let X be a non-empty set and R a unital commutative ring. A set \mathcal{F} of functions from X to R has *full determinantal rank over X* if, for any k distinct points $x_1, \dots, x_k \in X$, where $k \leq |\mathcal{F}|$, there exist functions $f_1, \dots, f_k \in \mathcal{F}$ such that the determinant $\det(f_i(x_j))$ is not a zero divisor.

If R is a field, X is a finite set and \mathcal{F} has at least $|X|$ elements, then the family \mathcal{F} has full determinantal rank over X if and only if the generalized matrix $(f(x))_{x \in X, f \in \mathcal{F}}$ has full rank. If, however, the family \mathcal{F} has fewer than $|X|$ elements, then this is not the case; for example, if \mathcal{F} contains a single function f which is the indicator function for a point in X , and X contains at least two points, then the matrix $(f(x))_{x \in X}$ has full rank but zero entries, and thus \mathcal{F} does not have full determinantal rank over X .

Example 3.6. We now give several examples of such families of functions, the first three of which correspond precisely to the results above.

- (1) If X is a finite set of complex numbers with k elements then the family $\{z^{j-1} : j = 1, \dots, k\}$ has full determinantal rank over X . Thus $\{z^n : n = 0, 1, 2, \dots\}$ has full determinantal rank over any subset of \mathbb{C} .
- (2) If X is a finite set of positive real numbers with k elements then $\{x^{n_j} : j = 1, \dots, k\}$ has full determinantal rank over X for any choice of distinct real exponents n_1, \dots, n_k . Thus $\{x^\alpha : \alpha \in \mathbb{R}\}$ has full determinantal rank over any subset of $(0, \infty)$.
- (3) The family of functions $\{x^\alpha : \alpha \geq 0\}$ has full determinantal rank over $[0, \infty)$.
- (4) The exponential family $\{\exp(\alpha x) : \alpha \in \mathbb{R}\}$ has full determinantal rank over \mathbb{R} . If $\alpha_1, \dots, \alpha_k$ are distinct, and similarly for x_1, \dots, x_k , then the matrix $(\exp(\alpha_i x_j))_{i,j=1}^k$ is, up to transposition of rows and columns, a generalized Vandermonde matrix and so non-singular.
- (5) The Gaussian family $\{f_\alpha(x) := \exp(-(x - \alpha)^2) : \alpha \in \mathbb{R}\}$ has full determinantal rank over \mathbb{R} . If $\alpha_1, \dots, \alpha_k$ are distinct, and similarly for x_1, \dots, x_k , then the matrix

$$(f_{\alpha_i}(x_j))_{i,j=1}^k = \text{diag}(e^{-\alpha_1^2}, \dots, e^{-\alpha_k^2})M \text{diag}(e^{-x_1^2}, \dots, e^{-x_k^2}),$$

where $M = (\exp(2\alpha_i x_j))$ is invertible by the preceding example.

Classes of functions with the separation property of Definition 3.5 are well studied in approximation theory, under the name of *Chebyshev systems* [18].

We now provide a theorem which contains all three results described above.

Theorem 3.7. Fix a Hermitian matrix $A \in \mathbb{C}^{N \times N}$ that is 3-PMP, where $N \geq 1$. Suppose $\pi := \pi^{\{1\}}(A) = \{I_1, \dots, I_m\}$ and $\pi' = \{I'_1, \dots, I'_{m'}\}$ is any partition refined by π . Let X denote the set of entries of A and suppose \mathcal{F} is a family of complex-valued functions on X that has full determinantal rank over the entries of each row of A . The following spaces are equal.

- (1) The simultaneous kernel of $f[A]$ for all $f \in \mathcal{F}$.
- (2) The simultaneous kernel of $f[A]$ for all functions $f: X \rightarrow \mathbb{C}$.

(3) *The simultaneous kernel of the block-diagonal matrices*

$$f[\text{diag } A_{\pi'}] := \bigoplus_{j=1}^{m'} f[A_{I'_j \times I'_j}]$$

for all functions $f: X \rightarrow \mathbb{C}$.

(4) *The kernel of $J_{\pi} := \bigoplus_{j=1}^m \mathbf{1}_{I_j \times I_j}$.*

This equality of kernels need not hold for matrices that are not 3-PMP.

The proof of Theorem 3.7 relies on the following strengthening of a result obtained in the proof of [5, Theorem 5.1]; in that setting, the ring R is taken to be a field and $\mathcal{F} = \{1, x, \dots, x^{m-1}\}$.

Proposition 3.8. *Let R be a unital commutative ring, and suppose the matrix $B \in R^{m \times m}$ is such that $m \geq 1$ and*

$$b_{ii} \neq b_{ij} \quad \text{whenever } 1 \leq i < j \leq m. \tag{3.1}$$

If the family \mathcal{F} has full determinantal rank over the entries in each row of B , then $\bigcap_{f \in \mathcal{F}} \ker f[B] = \{0\}$.

Proof. We show the result by induction on m , with the case $m = 1$ being immediate. For the inductive step, we claim that if $\mathbf{u} \in \bigcap_{f \in \mathcal{F}} \ker f[B]$ then $u_1 = 0$. This reduces the problem to showing that the trailing principal $(N - 1) \times (N - 1)$ submatrix of B has the same simultaneous kernel, whence we are done by the induction hypothesis; note that if \mathcal{F} has full determinantal rank over a set X , then it has so over any subset of X .

To show the claim, let $\mathbf{r}^T = (r_1, \dots, r_m)$ be the first row of B , and apply Theorem 2.3 with $M = 1$, $N = m$ and $G = \{1\}$ to obtain a minimal partition $\varpi_{\min} = \{J_1, \dots, J_k\}$ such that \mathbf{r}^T is constant on each block. Since $r_1 \neq r_j$ for $j = 2, \dots, m$, we may take $J_1 = \{1\}$ without loss of generality. Let $\mathbf{s} \in R^k$ be the compression of \mathbf{r} obtained by deleting repeated entries, so that $s_j = r_l$ for any $l \in J_j$, with $s_1 = r_1 = b_{11}$. As \mathbf{s} has distinct entries by construction, there exist $f_1, \dots, f_k \in \mathcal{F}$ such that $\det C$ is not a zero divisor, where $C := (f_i(s_j))_{i,j=1}^k$.

If $\mathbf{u} \in \bigcap_{f \in \mathcal{F}} \ker f[B]$, then $f_i[\mathbf{r}]^T \mathbf{u} = 0$ for $i = 1, \dots, k$. Let $\mathbf{v} \in R^k$ be defined by setting $v_j := \sum_{l \in J_j} u_l$, and note that $v_1 = u_1$. It follows that $f_i[\mathbf{s}]^T \mathbf{v} = 0$ for $i = 1, \dots, k$, so that $C\mathbf{v} = 0$. By Cramer’s rule, it follows that $\det(C)\mathbf{v} = 0$ in R^k , whence $\mathbf{v} = 0$ by the hypotheses. In particular, we have that $u_1 = v_1 = 0$, as desired. ■

We can now prove the general theorem described above.

Proof of Theorem 3.7. Let V_1, \dots, V_4 be the subspaces described in parts (1) to (4) of the statement of the theorem. We will show a chain of inclusions.

Note first that $\mathbf{v} \in V_4$ if and only if $\sum_{l \in I_j} v_l = 0$ for $j = 1, \dots, m$, from which it follows that $V_4 \subset V_2 \cap V_3$. Furthermore, it is immediate that $V_2 \subset V_1$. We now claim that the inclusion $V_1 \subset V_4$ gives the result. Firstly, we then have that

$$V_4 \subset V_2 \cap V_3 \subset V_2 \subset V_1 \subset V_4$$

and secondly, this also gives the inclusion $V_3 \subset V_4$, in which case

$$V_4 \subset V_2 \cap V_3 \subset V_3 \subset V_4.$$

For the last claim, note that V_3 is the direct sum of $V'_j := \bigcap_f \ker f[A_{I'_j \times I'_j}]$ for $j = 1, \dots, m'$, where the intersection is taken over the set of all functions from X to \mathbb{C} . Each V'_j is contained in the simultaneous kernel of $f[A_{I'_j \times I'_j}]$ with $f \in \mathcal{F}$, so the inclusion $V_1 \subset V_4$ applied to each matrix $A_{I'_j \times I'_j}$ with the partition $\pi \cap I'_j$ gives the inclusion as required.

Thus, it remains to show that $V_1 \subset V_4$. We proceed as in [5, Proof of Theorem 5.1]. Let $B \in \mathbb{C}^{m \times m}$ be the *compression* of A , denoted $B = \Sigma_\pi^\downarrow(A)$, so that the (i, j) entry of B equals the value A takes on the block $I_i \times I_j$. Note that B inherits the 3-PMP property from A .

Without loss of generality, we suppose that $b_{11} \geq b_{22} \geq \dots \geq b_{mm} \geq 0$. It must hold that $b_{ii} > 0$ for $i = 1, \dots, m - 1$. If not, then $b_{jj} = 0$ for $j = i, \dots, m$, whence all the entries of B in the $\{i, \dots, m\} \times \{i, \dots, m\}$ block are zero, by the 2-PMP property, which contradicts the minimality of $\pi = \pi^{\{1\}}(A)$.

We next claim that B satisfies condition (3.1) of Proposition 3.8. If there exist $i, j \in \{1, \dots, m\}$ with $i < j$ and $b_{ij} = b_{ii}$, then

$$b_{ii}^2 \geq b_{ii}b_{jj} \geq |b_{ij}|^2 = b_{ii}^2,$$

so $b_{jj} = b_{ii}$ because $b_{ii} > 0$. Hence B is constant on the block $\{i, j\}$, which again violates the definition of π .

We can now conclude our proof that $V_1 \subset V_4$. If $\mathbf{u} \in \mathbb{C}^N$ and $\mathbf{v} \in \mathbb{C}^m$ is defined by setting $v_j := \sum_{i \in I_j} u_i$, then \mathbf{v} is the compression of $J_\pi \mathbf{u}$, whence $\mathbf{u} \in \ker J_\pi$ if and only if $\mathbf{v} = 0$. Now suppose $\mathbf{u} \in V_1 = \bigcap_{f \in \mathcal{F}} \ker f[A]$. Then $\mathbf{v} \in \bigcap_{f \in \mathcal{F}} \ker f[B] = \{0\}$, by Proposition 3.8, so $\mathbf{u} \in \ker J_\pi = V_4$ as required. This concludes the proof of the chain of inclusions.

The counterexample is as in [5]: let $N = 3k + 2$ for some $k \geq 1$, let A be the Toeplitz matrix with (i, j) entry 1 if $|i - j| \leq 1$ and 0 otherwise, and note that A is 2-PMP but not 3-PMP. It is immediate that the family $\{f_1(x) = 1, f_2(x) = x\}$ has full determinantal rank on the set $\{0, 1\}$ of entries of A and the vector $(1, -1, 0, 1, -1, 0, \dots, 1, -1)^T$ lies in $\ker f_1[A] \cap \ker f_2[A]$. However, as $\pi^{\{1\}}(A) = \{\{1\}, \dots, \{N\}\}$, so $J_\pi = \text{Id}_N$ and $\ker J_\pi = \{0\}$. ■

4. Inflation and compression

In the present section we continue to explore isogenic stratification. The proof of Theorem 3.7 used the *compression operator* Σ_π^\downarrow , which will be one of the main characters in the new act.

Throughout this section, we fix a partition $\pi = \{I_1, \dots, I_m\} \in \Pi_N$, where $N \geq 1$.

Suppose $A, B \in \overline{\mathcal{S}_\pi^{\{1\}}}$, so that these matrices are constant on the blocks defined by the partition π . Then we may write

$$A = \sum_{i,j=1}^m a_{ij} \mathbf{1}_{I_i \times I_j} \quad \text{and} \quad B = \sum_{i,j=1}^m b_{ij} \mathbf{1}_{I_i \times I_j},$$

where $\mathbf{1}_{I_i \times I_j}$ is the $N \times N$ matrix with 1 in each entry of the $I_i \times I_j$ block and 0 elsewhere. Hence

$$AB = \sum_{i,j,k=1}^m a_{ik} b_{kj} |I_k| \mathbf{1}_{I_i \times I_j} \quad \text{and} \quad A \circ B = \sum_{i,j=1}^m a_{ij} b_{ij} \mathbf{1}_{I_i \times I_j},$$

so the closure of every stratum is a subalgebra of $\mathbb{C}^{N \times N}$ for both the usual and entrywise multiplication. The isogenic stratification of the space $\mathbb{C}^{N \times N}$ is not merely into linear spaces of matrices, but into subalgebras.

Next we focus on the compression operation of a fixed stratum to a lower-dimensional space. To simplify notation, we write henceforth

$$\mathcal{S}_\pi = \mathcal{S}_\pi^{\{1\}}$$

whenever there is no danger of confusion.

Definition 4.1. For $i, j \in \{1, \dots, m\}$, let E_{ij} denote the elementary matrix with (i, j) entry equal to 1 and all other entries 0, and recall that $\mathbf{1}_{I_i \times I_j}$ is the $N \times N$ matrix with 1 in each entry of the $I_i \times I_j$ block and 0 elsewhere.

(1) Define the linear *inflation* map as the linear extension of

$$\Sigma_\pi^\uparrow: \mathbb{C}^{m \times m} \rightarrow \mathbb{C}^{N \times N}; \quad E_{ij} \mapsto \mathbf{1}_{I_i \times I_j}$$

and note that the range of Σ_π^\uparrow is $\overline{\mathcal{S}_\pi}$.

(2) Define the linear *compression* map

$$\Sigma_\pi^\downarrow: \mathbb{C}^{N \times N} \rightarrow \mathbb{C}^{m \times m}; \quad \Sigma_\pi^\downarrow(A)_{ij} := \frac{1}{|I_i| |I_j|} \sum_{p \in I_i, q \in I_j} a_{pq},$$

so that the image $B = \Sigma_\pi^\downarrow(A)$ is such that b_{ij} is the arithmetic mean of the entries in $A_{I_i \times I_j}$, for $i, j = 1, \dots, m$.

Our next result shows that we may, in the entrywise setting, compress the matrices in a given stratum and work with the resulting smaller matrices with no loss of information.

Theorem 4.2. *Let $\overline{\mathcal{S}}_\pi$ and $\mathbb{C}^{m \times m}$ be equipped with the entrywise product, so that the units for this product are $\mathbf{1}_{N \times N}$ and $\mathbf{1}_{m \times m}$, respectively. The maps*

$$\Sigma_\pi^\downarrow: \overline{\mathcal{S}}_\pi \rightarrow \mathbb{C}^{m \times m} \quad \text{and} \quad \Sigma_\pi^\uparrow: \mathbb{C}^{m \times m} \rightarrow \overline{\mathcal{S}}_\pi$$

are mutually inverse, rank-preserving isomorphisms of unital $$ -algebras. Moreover, a matrix $A \in \overline{\mathcal{S}}_\pi$ is positive semidefinite if and only if $\Sigma_\pi^\downarrow(A)$ is.*

Towards the proof of this result, we first study the inflation and compression operators. To this aim, some new terminology will be useful.

Definition 4.3.

- (1) Define the *weight* matrix $\mathcal{W}_\pi \in \mathbb{C}^{N \times m}$ to have (i, j) entry 1 if $i \in I_j$ and 0 otherwise. Let

$$D_\pi := \mathcal{W}_\pi^* \mathcal{W}_\pi = \text{diag}(|I_1|, \dots, |I_m|). \tag{4.1}$$

When the rows of \mathcal{W}_π are ordered so that the indices in I_1 are first, then the indices in I_2 , and so on, then

$$\mathcal{W}_\pi \mathcal{W}_\pi^* = \text{diag}(\mathbf{1}_{|I_1|}, \dots, \mathbf{1}_{|I_m|}) =: J_\pi.$$

- (2) For any coarsening $\pi' \prec \pi$, define the partition $\pi'_\downarrow \in \Pi_m$ so that the blocks of π'_\downarrow are made up of those indices of blocks in π to be combined to form the blocks of π' . Thus if

$$\pi = \{I_1 = \{1, 2\}, I_2 = \{3, 4\}, I_3 = \{5\}, I_4 = \{6, 7, 8\}\}$$

and

$$\pi' = \{\{1, 2\}, \{3, 4, 5\}, \{6, 7, 8\}\},$$

then $\pi'_\downarrow = \{\{1\}, \{2, 3\}, \{4\}\}$. Denote the inverse map from Π_m to Π_N by $\pi'' \mapsto \pi''_\uparrow$.

Several important properties of the operators Σ_π^\downarrow and Σ_π^\uparrow are summarized in Proposition 4.4.

Proposition 4.4.

- (1) *The map $\pi' \mapsto \pi'_\downarrow$ is a bijection between the set of all coarsenings of π in Π_N and the set Π_m .*
- (2) *For all $A \in \mathbb{C}^{m \times m}$,*

$$\Sigma_\pi^\uparrow(A) = \mathcal{W}_\pi A \mathcal{W}_\pi^* \in \overline{\mathcal{S}}_\pi. \tag{4.2}$$

Moreover, Σ_π^\uparrow is a bijection from $\mathbb{C}^{m \times m}$ onto $\overline{\mathcal{S}}_\pi$, sending the stratum $\mathcal{S}_{\pi'} \subset \mathbb{C}^{m \times m}$ to the stratum $\overline{\mathcal{S}}_{\pi'_\uparrow} \subset \overline{\mathcal{S}}_\pi$, for any $\pi' \in \Pi_m$.

(3) For all $A \in \mathbb{C}^{N \times N}$,

$$\Sigma_\pi^\downarrow(A) = D_\pi^{-1} \mathcal{W}_\pi^* A \mathcal{W}_\pi D_\pi^{-1} \in \mathbb{C}^{m \times m}. \tag{4.3}$$

Moreover, Σ_π^\downarrow restricted to $\overline{\mathcal{S}_\pi}$ is a bijection onto $\mathbb{C}^{m \times m}$, being the inverse map of Σ_π^\uparrow .

(4) The linear maps Σ_π^\uparrow and Σ_π^\downarrow are compatible with matrix multiplication in the following sense:

$$\Sigma_\pi^\uparrow(AB) = \Sigma_\pi^\uparrow(AD_\pi^{-1/2})\Sigma_\pi^\uparrow(D_\pi^{-1/2}B) \quad \text{for all } A, B \in \mathbb{C}^{m \times m} \tag{4.4}$$

and

$$\Sigma_\pi^\downarrow(AB) = \Sigma_\pi^\downarrow(A)D_\pi\Sigma_\pi^\downarrow(B) \quad \text{for all } A, B \in \overline{\mathcal{S}_\pi}. \tag{4.5}$$

Proof. Part (1) readily follows from the definitions. To see that (4.2) holds, it suffices by linearity to show that $\mathcal{W}_\pi E_{ij} \mathcal{W}_\pi^* = \mathbf{1}_{I_i \times I_j}$ for each elementary matrix E_{ij} , but this is immediate. It is also clear that Σ_π^\downarrow and Σ_π^\uparrow are mutually inverse bijections between $\overline{\mathcal{S}_\pi}$ and $\mathbb{C}^{m \times m}$. The other assertions of (2) are straightforward.

To show that (4.3) holds, it once again suffices to take A to be an arbitrary elementary matrix; the calculation is then straightforward. That Σ_π^\downarrow and Σ_π^\uparrow are inverse between $\overline{\mathcal{S}_\pi}$ and $\mathbb{C}^{m \times m}$ has already been discussed.

Finally, equation (4.4) is verified by using (4.2) and the definition of D_π . To see (4.5), note that, by (4.4),

$$\begin{aligned} \Sigma_\pi^\uparrow(\Sigma_\pi^\downarrow(A)D_\pi\Sigma_\pi^\downarrow(B)) &= \Sigma_\pi^\uparrow(\Sigma_\pi^\downarrow(A)D_\pi^{1/2}D_\pi^{1/2}\Sigma_\pi^\downarrow(B)) \\ &= \Sigma_\pi^\uparrow(\Sigma_\pi^\downarrow(A))\Sigma_\pi^\uparrow(\Sigma_\pi^\downarrow(B)) = AB, \end{aligned}$$

and this concludes the proof. ■

In other words, the map $\Sigma_\pi^\uparrow \circ \Sigma_\pi^\downarrow$ is a conditional expectation on $\mathbb{C}^{N \times N}$ corresponding to the σ -algebra generated by the blocks which define the stratum \mathcal{S}_π .

These properties of inflation and compression maps help demonstrate the $*$ -isomorphism claimed above.

Proof of Theorem 4.2. By Proposition 4.4, it suffices to show the results only for Σ_π^\uparrow . This map is linear and multiplicative for the entrywise product, by definition. Equation (4.2) shows that Σ_π^\uparrow commutes with taking the adjoint $*$, and also that Σ_π^\uparrow preserves rank, since \mathcal{W}_π has full rank. Finally, that positivity is preserved follows immediately from (4.2). ■

5. Spectral permanence

Theorem 4.2 shows that the map $\Sigma_\pi^\downarrow: \overline{\mathcal{S}_\pi} \rightarrow \mathbb{C}^{m \times m}$ is a positivity-preserving $*$ -algebra isomorphism for the entrywise product, whence the entrywise calculus is transported to a lower-dimensional space of matrices. However, it is not immediately apparent if the usual holomorphic functional calculus in $\mathbb{C}^{m \times m}$ can be transported up to the closure of each stratum. We now show how this can be accomplished with the help of different weighted inflation and compression maps.

As in the previous Section 4, we fix a partition $\pi = \{I_1, \dots, I_m\} \in \Pi_N$, where $N \geq 1$.

Definition 5.1. Define linear operators

$$\Theta_\pi^\downarrow: \mathbb{C}^{N \times N} \rightarrow \mathbb{C}^{m \times m} \quad \text{and} \quad \Theta_\pi^\uparrow: \mathbb{C}^{m \times m} \rightarrow \mathbb{C}^{N \times N}$$

by setting

$$\Theta_\pi^\downarrow(A) := D_\pi^{1/2} \Sigma_\pi^\downarrow(A) D_\pi^{1/2} = D_\pi^{-1/2} \mathcal{W}_\pi^* A \mathcal{W}_\pi D_\pi^{-1/2}$$

and

$$\Theta_\pi^\uparrow(B) := \Sigma_\pi^\uparrow(D_\pi^{-1/2} B D_\pi^{-1/2}) = \mathcal{W}_\pi D_\pi^{-1/2} B D_\pi^{-1/2} \mathcal{W}_\pi^*.$$

Theorem 5.2. *The maps Θ_π^\downarrow and Θ_π^\uparrow are mutually inverse, rank-preserving isomorphisms between the unital $*$ -algebras $\overline{\mathcal{S}_\pi}$ and $\mathbb{C}^{m \times m}$ equipped with the usual matrix multiplication. Furthermore, a matrix $A \in \overline{\mathcal{S}_\pi}$ is positive semidefinite if and only if $\Theta_\pi^\downarrow(A)$ is.*

Proof. That Θ_π^\downarrow and Θ_π^\uparrow are linear, bijective, $*$ -equivariant, and preserve rank follows from the corresponding properties of Σ_π^\downarrow and Σ_π^\uparrow from Theorem 4.2, since D_π is positive definite. Moreover, it is easily shown that Θ_π^\downarrow and Θ_π^\uparrow are mutually inverse between $\overline{\mathcal{S}_\pi}$ and $\mathbb{C}^{m \times m}$, since Σ_π^\downarrow and Σ_π^\uparrow are. Furthermore, if $A, B \in \overline{\mathcal{S}_\pi}$, then, by (4.5),

$$\Theta_\pi^\downarrow(AB) = D_\pi^{1/2} \Sigma_\pi^\downarrow(AB) D_\pi^{1/2} = D_\pi^{1/2} \Sigma_\pi^\downarrow(A) D_\pi \Sigma_\pi^\downarrow(B) D_\pi^{1/2} = \Theta_\pi^\downarrow(A) \Theta_\pi^\downarrow(B).$$

Thus the maps Θ_π^\downarrow and Θ_π^\uparrow are algebra homomorphisms for the usual matrix multiplication, and that they take the units to one another is readily verified; note that $\overline{\mathcal{S}_\pi}$ has unit

$$1_{\overline{\mathcal{S}_\pi}} = \Theta_\pi^\uparrow(1_{\mathbb{C}^{m \times m}}) = \sum_{k=1}^m |I_k|^{-1} \mathbf{1}_{I_k \times I_k}.$$

The final assertion about preserving positivity is immediate. ■

Remark 5.3. We note a couple of simple consequences of Theorem 5.2.

- (1) The matrix $\Sigma_\pi^\downarrow(A)$ is invertible if and only if $\Theta_\pi^\downarrow(A)$ is, in which case

$$\Theta_\pi^\downarrow(A)^{-1} = D_\pi^{-1/2} \Sigma_\pi^\downarrow(A)^{-1} D_\pi^{-1/2}.$$

- (2) If $A \in \overline{\mathcal{S}_\pi}$, then the Moore–Penrose pseudo-inverse is:

$$A^\dagger = \Theta_\pi^\uparrow(\Theta_\pi^\downarrow(A)^\dagger), \tag{5.1}$$

since $A = \Theta_\pi^\uparrow(\Theta_\pi^\downarrow(A))$ and Θ_π^\uparrow is a $*$ -algebra homomorphism.

We now extend the compression and inflation operators to act on vectors as well as matrices.

Definition 5.4. Define

$$\Sigma_\pi^\downarrow: \mathbb{C}^N \rightarrow \mathbb{C}^m \quad \text{and} \quad \Theta_\pi^\downarrow: \mathbb{C}^N \rightarrow \mathbb{C}^m$$

by setting

$$\Sigma_\pi^\downarrow(\mathbf{u})_j = |I_j|^{-1} \sum_{k \in I_j} u_k \quad \text{and} \quad \Theta_\pi^\downarrow(\mathbf{u})_j = |I_j|^{-1/2} \sum_{k \in I_j} u_k.$$

Similarly, define

$$\Sigma_\pi^\uparrow: \mathbb{C}^m \rightarrow \mathbb{C}^N \quad \text{and} \quad \Theta_\pi^\uparrow: \mathbb{C}^m \rightarrow \mathbb{C}^N$$

by setting

$$\Sigma_\pi^\uparrow(\mathbf{v})_j = v_k \quad \text{and} \quad \Theta_\pi^\uparrow(\mathbf{v})_j = |I_k|^{-1/2} v_k \quad (j \in I_k).$$

The following proposition summarizes basic properties of the operators Σ_π^\downarrow , Θ_π^\downarrow , Σ_π^\uparrow , and Θ_π^\uparrow acting on vectors. Its proof is omitted.

Proposition 5.5. Let $\mathbf{u} \in \mathbb{C}^N$, $\mathbf{v}, \mathbf{w} \in \mathbb{C}^m$, $A \in \overline{\mathcal{S}_\pi}$, and $B \in \mathbb{C}^{m \times m}$.

- (1) $\Theta_\pi^\downarrow(\mathbf{u}) = D_\pi^{1/2} \Sigma_\pi^\downarrow(\mathbf{u})$ and $\Theta_\pi^\uparrow(\mathbf{v}) = \Sigma_\pi^\uparrow(D_\pi^{-1/2} \mathbf{v})$.
- (2) $\Theta_\pi^\downarrow(\Theta_\pi^\uparrow(\mathbf{v})) = \mathbf{v} = \Sigma_\pi^\downarrow(\Sigma_\pi^\uparrow(\mathbf{v}))$.
- (3) $\Theta_\pi^\uparrow(\mathbf{v})^* \Theta_\pi^\uparrow(\mathbf{w}) = \mathbf{v}^* \mathbf{w}$ and $\Sigma_\pi^\uparrow(\mathbf{v})^* \Sigma_\pi^\uparrow(\mathbf{w}) = \mathbf{v}^* D_\pi \mathbf{w}$.
- (4) $\Theta_\pi^\downarrow(A\mathbf{u}) = \Theta_\pi^\downarrow(A) \Theta_\pi^\downarrow(\mathbf{u})$ and $\Sigma_\pi^\downarrow(A\mathbf{u}) = \Sigma_\pi^\downarrow(A) D_\pi \Sigma_\pi^\downarrow(\mathbf{u})$.
- (5) $\Theta_\pi^\uparrow(B\mathbf{v}) = \Theta_\pi^\uparrow(B) \Theta_\pi^\uparrow(\mathbf{v})$ and $\Sigma_\pi^\uparrow(B\mathbf{v}) = \Sigma_\pi^\uparrow(B) \Sigma_\pi^\uparrow(D_\pi^{-1} \mathbf{v})$.

Remark 5.6. We collect here some further properties of the maps Σ_π^\downarrow , Σ_π^\uparrow , Θ_π^\downarrow , and Θ_π^\uparrow .

- (1) If $A \in \overline{\mathcal{S}_\pi}$ and $\mathbf{u} \in \mathbb{C}^N$, then $A\mathbf{u}$ is constant on the blocks of the partition π , that is, $A\mathbf{u} \in \text{im } \Sigma_\pi^\uparrow = \text{im } \Theta_\pi^\uparrow$, the range of these two operators acting from \mathbb{C}^m to \mathbb{C}^N . Consequently,

$$A\mathbf{u} = \Theta_\pi^\uparrow(\Theta_\pi^\downarrow(A) \Theta_\pi^\downarrow(\mathbf{u})) = \Sigma_\pi^\uparrow(\Sigma_\pi^\downarrow(A) D_\pi \Sigma_\pi^\downarrow(\mathbf{u})).$$

(2) If $\mathbf{u}, \mathbf{v} \in \text{im } \Sigma_\pi^\uparrow = \text{im } \Theta_\pi^\uparrow$ and $A \in \overline{\mathcal{S}}_\pi$, then

$$\mathbf{u}^* \mathbf{v} = \Theta_\pi^\downarrow(\mathbf{u})^* \Theta_\pi^\downarrow(\mathbf{v}) \quad \text{and} \quad \mathbf{u}^* A \mathbf{v} = \Theta_\pi^\downarrow(\mathbf{u})^* \Theta_\pi^\downarrow(A) \Theta_\pi^\downarrow(\mathbf{v}),$$

whereas

$$\mathbf{u}^* \mathbf{v} = \Sigma_\pi^\downarrow(\mathbf{u})^* D_\pi \Sigma_\pi^\downarrow(\mathbf{v}) \quad \text{and} \quad \mathbf{u}^* A \mathbf{v} = \Sigma_\pi^\downarrow(\mathbf{u})^* D_\pi \Sigma_\pi^\downarrow(A) D_\pi \Sigma_\pi^\downarrow(\mathbf{v}).$$

Furthermore, if $\mathbf{x}, \mathbf{y} \in \mathbb{C}^m$ then

$$\Theta_\pi^\uparrow(\mathbf{x} \mathbf{y}^*) = \Theta_\pi^\uparrow(\mathbf{x}) \Theta_\pi^\uparrow(\mathbf{y})^* \quad \text{and} \quad \Sigma_\pi^\uparrow(\mathbf{x} \mathbf{y}^*) = \Sigma_\pi^\uparrow(\mathbf{x}) \Sigma_\pi^\uparrow(\mathbf{y})^*,$$

so

$$\Theta_\pi^\downarrow(\mathbf{u} \mathbf{v}^*) = \Theta_\pi^\downarrow(\mathbf{u}) \Theta_\pi^\downarrow(\mathbf{v})^* \quad \text{and} \quad \Sigma_\pi^\downarrow(\mathbf{u} \mathbf{v}^*) = \Sigma_\pi^\downarrow(\mathbf{u}) \Sigma_\pi^\downarrow(\mathbf{v})^*.$$

Our next result shows that the maps Θ_π^\uparrow and Θ_π^\downarrow preserve eigenvalues, up to the possible addition or removal of 0.

Proposition 5.7. *The following spectral permanence holds for all $A \in \overline{\mathcal{S}}_\pi$ and $B \in \mathbb{C}^{m \times m}$:*

$$\sigma(A) \setminus \{0\} = \sigma(\Theta_\pi^\downarrow(A)) \setminus \{0\} \tag{5.2}$$

$$\text{and } \sigma(B) \setminus \{0\} = \sigma(\Theta_\pi^\uparrow(B)) \setminus \{0\}. \tag{5.3}$$

In particular, $\Theta_\pi^\downarrow: \overline{\mathcal{S}}_\pi \rightarrow \mathbb{C}^{m \times m}$ and $\Theta_\pi^\uparrow: \mathbb{C}^{m \times m} \rightarrow \overline{\mathcal{S}}_\pi$ preserve the spectral radius. Furthermore, applying Θ_π^\uparrow to B adds a zero eigenvalue of geometric multiplicity $N - m$ to $\sigma(B)$, whereas applying Θ_π^\downarrow to A reduces the geometric multiplicity of the zero eigenvalue of A by $N - m$.

Proof. To prove (5.3), first let $B \mathbf{v} = \lambda \mathbf{v}$ for some $\lambda \neq 0$ and $\mathbf{v} \in \mathbb{C}^m \setminus \{0\}$. Then, by Proposition 5.5(5),

$$\Theta_\pi^\uparrow(B) \Theta_\pi^\uparrow(\mathbf{v}) = \Theta_\pi^\uparrow(B \mathbf{v}) = \lambda \Theta_\pi^\uparrow(\mathbf{v})$$

and $\Theta_\pi^\uparrow(\mathbf{v}) \neq 0$, so $\lambda \in \sigma(\Theta_\pi^\uparrow(B)) \setminus \{0\}$. Conversely, suppose $\Theta_\pi^\uparrow(B) \mathbf{u} = \lambda \mathbf{u}$ for some $\lambda \neq 0$ and $\mathbf{u} \in \mathbb{C}^N \setminus \{0\}$. Then $\mathbf{u} = \lambda^{-1} \Theta_\pi^\uparrow(B) \mathbf{u} \in \text{im } \Theta_\pi^\uparrow$, and therefore $\mathbf{u} = \Theta_\pi^\uparrow(\mathbf{v})$ for some $\mathbf{v} \in \mathbb{C}^m \setminus \{0\}$. Hence, using Proposition 5.5(5) again,

$$\Theta_\pi^\uparrow(B \mathbf{v} - \lambda \mathbf{v}) = \Theta_\pi^\uparrow(B) \Theta_\pi^\uparrow(\mathbf{v}) - \lambda \Theta_\pi^\uparrow(\mathbf{v}) = 0.$$

It follows that $B \mathbf{v} - \lambda \mathbf{v} = 0$, and this proves the first equality.

The proof of (5.2) is similar, with the help of Proposition 5.5 and that the fact that if $\mathbf{u} \in \text{im } \Theta_\pi^\uparrow$ and $\Theta_\pi^\downarrow(\mathbf{u}) = 0$ then $\mathbf{u} = 0$.

The last claim follows immediately by the rank-nullity theorem. ■

Remark 5.8. Let \mathcal{A} be a complex associative algebra with multiplicative identity $1_{\mathcal{A}}$. Then the *spectrum* of $a \in \mathcal{A}$ is defined as

$$\sigma(a; \mathcal{A}) := \{\lambda \in \mathbb{C} : \lambda 1_{\mathcal{A}} - a \text{ is not invertible in } \mathcal{A}\}.$$

In the notation of Proposition 5.7, $\sigma(A) = \sigma(A; \mathbb{C}^{N \times N})$ for any $A \in \mathbb{C}^{N \times N}$, where $\mathbb{C}^{N \times N}$ equipped with the usual matrix product.

Since $\Theta_{\pi}^{\downarrow}$ is an algebra isomorphism from $\overline{\mathcal{S}_{\pi}}$ to $\mathbb{C}^{m \times m}$, and the spectrum and the set of eigenvalues coincide in the latter, it follows from (5.3) and (5.2) that, for any $A \in \overline{\mathcal{S}_{\pi}}$,

$$\sigma(A; \overline{\mathcal{S}_{\pi}}) = \sigma(\Theta_{\pi}^{\downarrow}(A); \mathbb{C}^{m \times m}) = \sigma(\Theta_{\pi}^{\downarrow}(A)) \subset \sigma(A) \tag{5.4}$$

and

$$\sigma(A) \setminus \{0\} \subset \sigma(\Theta_{\pi}^{\downarrow}(A)) = \sigma(\Theta_{\pi}^{\downarrow}(A); \mathbb{C}^{m \times m}) = \sigma(A; \overline{\mathcal{S}_{\pi}}). \tag{5.5}$$

The matrix $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \in \mathcal{S}_{\pi_{\wedge}}$ has $\sigma(A; \overline{\mathcal{S}_{\pi_{\wedge}}}) = \{2\}$ and $\sigma(A) = \{0, 2\}$, so the inclusion in (5.4) may be strict. Similarly, the matrix $A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \in \mathcal{S}_{\pi_{\vee}}$ has $\sigma(A; \overline{\mathcal{S}_{\pi_{\vee}}}) = \sigma(A) = \{0, 1\}$, so the inclusion in (5.5) may also be strict.

The following theorem shows that the holomorphic functional calculus naturally transfers between $\mathbb{C}^{m \times m}$ and $\overline{\mathcal{S}_{\pi}}$. Its proof follows immediately from the fact that $\Theta_{\pi}^{\downarrow}$ is an algebra isomorphism.

Theorem 5.9. *Given $A \in \overline{\mathcal{S}_{\pi}}$ and $B \in \mathbb{C}^{m \times m}$, let the resolvents*

$$R(z; A) := (zJ_{\pi} - A)^{-1} \quad \text{for all } z \in \sigma(A; \overline{\mathcal{S}_{\pi}})$$

and

$$R(z; B) := (zI_{m \times m} - B)^{-1} \quad \text{for all } z \in \sigma(B; \mathbb{C}^{m \times m}) = \sigma(B).$$

If $z \in \sigma(A; \overline{\mathcal{S}_{\pi}}) = \sigma(\Theta_{\pi}^{\downarrow}(A); \mathbb{C}^{m \times m}) = \sigma(\Theta_{\pi}^{\downarrow}(A))$, then

$$\Theta_{\pi}^{\downarrow}(R(z; A)) = R(z; \Theta_{\pi}^{\downarrow}(A)).$$

Thus, the holomorphic functional calculus transfers between $\mathbb{C}^{m \times m}$ and $\overline{\mathcal{S}_{\pi}}$: if $A \in \overline{\mathcal{S}_{\pi}}$ and f is holomorphic on an open set containing $\sigma(A; \overline{\mathcal{S}_{\pi}})$, then

$$f(A) \in \overline{\mathcal{S}_{\pi}} \quad \text{and} \quad \Theta_{\pi}^{\downarrow}(f(A)) = f(\Theta_{\pi}^{\downarrow}(A)).$$

Remark 5.10. It follows from Theorem 5.2 that analogues of the general linear group GL_m , unitary group U_m , and permutation group S_m exist inside the stratum $\overline{\mathcal{S}_{\pi}}$. Furthermore, the notions of nilpotent, Hermitian, and positive semidefinite matrices are preserved in $\overline{\mathcal{S}_{\pi}}$ via Θ_{π}^{\uparrow} . Hence analogues of the Bruhat, Cholesky, and polar decompositions can also be defined on $\overline{\mathcal{S}_{\pi}}$, and all of the respective factors live inside the same stratum.

We conclude this section with some remarks on the situation for \mathcal{S}_π^G with more general $G \subset \mathbb{C}^\times$. A key feature in the definition of the compression operator Σ_π^\downarrow , and so of Θ_π^\downarrow , was the unique decomposition of a single rank-one block matrix: $\mathbf{1}_{I_i \times I_j} = \mathbf{1}_{I_i} \mathbf{1}_{I_j}^*$. When $G \neq \{1\}$, the block matrices do not possess such a decomposition. Moreover, each stratum and its closure are no longer closed under multiplication. For example, if $G = \{\pm 1\}$, $N = 2$, and $\pi = \{\{1, 2\}\}$ is the minimum partition, then

$$A = \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \in \mathcal{S}_\pi^G, \quad \text{but} \quad A^2 = \begin{pmatrix} 0 & -2 \\ 2 & 0 \end{pmatrix} \notin \mathcal{S}_\pi^G.$$

6. Ramifications

We collect in this last section several observations revealing natural links between the compression and inflation operations derived from the isogenic stratification of the matrix space and recent advances or classical examples of current interest in numerical matrix analysis.

6.1. Symmetric statistical models

We briefly discuss a related setting of statistical models and covariance matrices which exhibit symmetry with respect to a group of permutations; see [26] and the references therein. The authors fix there a subgroup \mathfrak{G} of the symmetric group S_N , and define

$$\mathcal{W}_\mathfrak{G} := \{A \in \mathbb{R}^{N \times N} : a_{ij} = a_{\sigma(i), \sigma(j)} \text{ whenever } 1 \leq i, j \leq N \text{ and } \sigma \in \mathfrak{G}\}.$$

In the framework of the present article we consider permutation groups associated to a partition $\pi = \{I_1, \dots, I_m\}$ of $\{1, \dots, N\}$, that is,

$$\mathfrak{G}_\pi := \text{Aut}(I_1) \times \dots \times \text{Aut}(I_m).$$

By contrast, in [26] the authors work with more general subgroups of S_N . Furthermore, the matrices in a given stratum $\mathcal{S}_\pi^{\{1\}}$ have diagonal blocks with equal entries, as the permutations act separately on the rows and columns of $\mathbb{C}^{N \times N}$, whereas the diagonal blocks of a square matrix in $\mathcal{W}_\mathfrak{G}$ may have different diagonal and off-diagonal entries.

Having acknowledged these differences, we now discuss the setting of [26] from the viewpoint adopted above. The first step is to establish the existence of a suitable partition associated with a given matrix.

Proposition 6.1. *Fix a unital commutative ring R and a multiplicative subgroup $G \subset R^\times$. Given a matrix $A \in R^{N \times N}$, where $N \geq 1$, there is a unique minimal partition $\varpi_{\min} = \{I_1, \dots, I_m\} \in \Pi_N$ satisfying the following two properties.*

- (1) If $i, j \in \{1, \dots, N\}$ are distinct, then all entries of the block $A_{I_i \times I_j}$ lie in a single G -orbit.
- (2) If $i \in \{1, \dots, N\}$, then all diagonal entries of $A_{I_i \times I_i}$ all lie in a single G -orbit, as do all off-diagonal entries.

Proof. As in the proof of Theorem 2.3, the key step is to show that if ϖ_1 and ϖ_2 satisfy both of the properties given above, then $\varpi_1 \vee \varpi_2$ does too. This proceeds as in that proof, and the only part which is not immediate the case of off-diagonal elements of a diagonal block, with one lying above the diagonal and the other below. However, property (2) gives that $a_{pq} \in Ga_{qp}$ if p and q are distinct and lie in the same block of ϖ_1 or ϖ_2 , so one may “cross the diagonal”. ■

Denote the partition in Proposition 6.1 by $\varpi^G(A)$ and, as above, define for each partition ϖ the set

$$\mathcal{C}_\varpi^G := \{A \in \mathbb{C}^{N \times N} : \varpi^G(A) = \varpi\}.$$

Proposition 6.2. *There is a natural stratification of $\mathbb{C}^{N \times N}$ resulting from Proposition 6.1, namely*

$$\mathbb{C}^{N \times N} = \bigsqcup_{\varpi \in \Pi_N} \mathcal{C}_\varpi^G, \quad \text{and} \quad \overline{\mathcal{C}_\varpi^G} = \bigsqcup_{\varpi' \prec \varpi} \mathcal{C}_{\varpi'}^G. \tag{6.1}$$

Furthermore, we have that $\overline{\mathcal{C}_\varpi^{\{1\}}} = \mathcal{W}_{\mathfrak{G}_\varpi}$.

Thus $\mathbb{C}^{N \times N}$ indeed admits a stratification, analogous to the situation above. However, the reason we do not proceed further along these lines is the lack of a *rank-preserving* equivalence of the stratum $\mathcal{C}_\varpi^{\{1\}}$ with any lower-dimensional space. Indeed, if $\varpi = \{I_1, \dots, I_m\}$ and $A_\varpi := \sum_{j=1}^m \mathbf{1}_{I_j \times I_j}$, then the sum of A_ϖ with any positive multiple of the $N \times N$ identity matrix is an element of $\mathcal{C}_\varpi^{\{1\}}$ with full rank.

6.2. Block correlation matrices and group kernels

A popular approach for constructing probabilistic models involving categorical data consists of grouping the input levels in such a way that correlation is constant within each group and across groups [2, 8, 17, 24]. In particular, the rapidly developing area of *group kernels* exploits this idea to obtain useful Gaussian processes for categorical variables [22, 25]. This approach yields parsimonious covariance models that can be naturally analyzed in the framework of the current paper. To elaborate, consider a categorical problem involving a potentially large number of levels N , partitioned into a typically small number of groups m of sizes n_1, \dots, n_m , so that $n_1 + \dots + n_m = N$. Assuming the

within-groups correlations and between-groups correlation are constant, the associated covariance matrix A can be written in block form

$$A = (A_{ij})_{i,j=1}^m, \tag{6.2}$$

where the diagonal blocks $A_{ii} \in \mathbb{R}^{n_i \times n_i}$ are compound symmetry matrices of the form

$$A_{ii} = \text{Id}_{n_i} + c_{ii}(\mathbf{1}_{n_i \times n_i} - \text{Id}_{n_i})$$

and the off-diagonal blocks $A_{ij} \in \mathbb{R}^{n_i \times n_j}$ are of the form

$$A_{ij} = c_{ij} \mathbf{1}_{n_i \times n_j}.$$

As shown in [24, 25], positive definiteness of A is equivalent to the positivity of the much smaller compressed matrix $\Sigma_\pi^\downarrow(A)$, where π is the partition of $\{1, \dots, N\}$ associated with the block structure of A . Earlier versions of this result may be found in [8] and [17].

Theorem 6.3. ([24, Theorem 2]) *Let $A = (A_{ij})_{i,j=1}^m$ be as above, with $0 < c_{ij} < 1$ for $i, j = 1, \dots, m$. Then A is positive definite if and only if its compression $\Sigma_\pi^\downarrow(A)$ is positive definite.*

In fact, Theorem 6.3 is a consequence of the following more general result. Recall the Loewner order: if A and B are Hermitian square complex matrix of the same size, then $A \geq B$ if and only if $A - B$ is positive semidefinite.

Theorem 6.4. ([24, Theorem 1]) *Let $B = (B_{ij})_{i,j=1}^m$ be an arbitrary Hermitian block matrix with $B_{ij} \in \mathbb{R}^{n_i \times n_j}$ for $i, j = 1, \dots, m$. Denote by π the partition of $\{1, \dots, n_1 + \dots + n_m\}$ associated with the block structure of B and let $\overline{B_{ii}} := \Sigma_\pi^\downarrow(B_{ii}) \in \mathbb{R}$ be the arithmetic mean of the entries of the block B_{ii} . If*

$$B_{ii} \geq \overline{B_{ii}} \mathbf{1}_{n_i \times n_i} \quad (i = 1, \dots, m)$$

then the following are equivalent.

- (1) *The matrix B is positive semidefinite.*
- (2) *The matrix $\Sigma_\pi^\downarrow(B)$ and the diagonal blocks B_{11}, \dots, B_{mm} are positive semidefinite.*

The result also holds when “positive semidefinite” is replaced by “positive definite” in both (1) and (2).

Note that Theorem 6.3 follows from Theorem 6.4 because

$$A_{ii} - \overline{A_{ii}} \mathbf{1}_{n_i \times n_i} = (1 - c_{ii})(\text{Id}_{n_i} - n_i^{-1} \mathbf{1}_{n_i \times n_i})$$

is positive semidefinite when $0 < c_{ii} < 1$. More generally, the same result holds if the diagonal blocks have the form

$$A_{ii} = d_i \text{Id}_{n_i} + c_{ii}(\mathbf{1}_{n_i \times n_i} - \text{Id}_{n_i}), \quad \text{where } d_i \geq c_{ii}.$$

For completeness, we provide a self-contained proof of Theorem 6.4 in the language developed above.

Proof of Theorem 6.4. The implication (1) \implies (2) is clear with the help of Theorem 4.2. Conversely, if $C := \Sigma_\pi^\downarrow(B)$ is positive semidefinite, then so is $\Sigma_\pi^\uparrow(C)$, by Theorem 4.2 again. The block-diagonal matrix

$$D = \text{diag}(B_{11} - \overline{B_{11}}\mathbf{1}_{n_1 \times n_1}, \dots, B_{mm} - \overline{B_{mm}}\mathbf{1}_{n_m \times n_m})$$

is positive semidefinite by assumption, and so $B = \Sigma_\pi^\uparrow(C) + D$ is positive semidefinite. This shows that (2) \implies (1).

For the positive-definite version, that (1) \implies (2) is again clear, with the help of (4.3) and the fact that W_π there has full rank. For the converse, suppose $C := \Sigma_\pi^\downarrow(B)$ is positive definite and note that $B = \Sigma_\pi^\uparrow(C) + D$ is the sum of positive semidefinite matrices. Thus, if \mathbf{v} is such that $\mathbf{v}^T B \mathbf{v} = 0$ then \mathbf{v} lies in the kernels of D and $\Sigma_\pi^\uparrow(C)$. Since each B_{ii} is invertible, and rank is subadditive, the kernel of each block of D is at most one dimensional. Hence, the kernel of D is spanned by vectors of the form $\Sigma_\pi^\uparrow(\mathbf{e}_i)$ for $i \in I$, where $\mathbf{e}_1, \dots, \mathbf{e}_m$ is the canonical basis of \mathbb{R}^m and $I \subset \{1, \dots, m\}$. However, by Proposition 5.5(5), we have that

$$\Sigma_\pi^\uparrow(C) \sum_{i \in I} v_i \Sigma_\pi^\uparrow(\mathbf{e}_i) = \Sigma_\pi^\uparrow\left(CD_\pi \sum_{i \in I} v_i \mathbf{e}_i\right) = \mathbf{0} \iff \sum_{i \in I} v_i \mathbf{e}_i = \mathbf{0} \quad (v_i \in \mathbb{R})$$

since C and D_π are invertible. Hence D has trivial kernel and B is positive definite. ■

An explicit expression for the eigendecomposition of the block matrices featuring in Theorem 6.3 was obtained in [8] and [17], as well as in [2] for the case where the diagonal blocks are of the form

$$A_{ii} = d_i \text{Id}_{n_i} + c_{ii}(\mathbf{1}_{n_i \times n_i} - \text{Id}_{n_i}).$$

We provide a proof of the last result using our stratification language and some results from Section 5.

Theorem 6.5. ([2, Theorem 1]) *Let $A = (A_{ij})_{i,j=1}^m$ be a real matrix with*

$$A_{ij} = \begin{cases} d_i \text{Id}_{n_i} + c_{ii}(\mathbf{1}_{n_i \times n_i} - \text{Id}_{n_i}) & \text{if } i = j, \\ c_{ij} \mathbf{1}_{n_i \times n_j} & \text{if } i \neq j. \end{cases}$$

Then $A = QDQ^T$, where Q is an orthogonal matrix,

$$D = A' \oplus \bigoplus_{j=1}^m (d_j - c_{jj}) \text{Id}_{n_j-1}$$

and $A' = (a'_{ij})_{i,j=1}^m$ with

$$a'_{ij} = \begin{cases} d_i + c_{ii}(n_i - 1) & \text{if } i = j, \\ c_{ij}\sqrt{n_i n_j} & \text{if } i \neq j. \end{cases}$$

Proof. As above, let $\overline{A_{ii}}$ denote the arithmetic mean of the entries of the block A_{ii} , so that $\overline{A_{ii}} = n_i^{-1}(d_i + (n_i - 1)c_{ii})$ for $i = 1, \dots, m$, and let $\pi = \{I_1, \dots, I_m\}$ be the partition associated with the block structure of A . Then

$$\begin{aligned} A &= \Theta_\pi^\uparrow(A') + (A_{11} - \overline{A_{11}}\mathbf{1}_{n_1 \times n_1}) \oplus \dots \oplus (A_{mm} - \overline{A_{mm}}\mathbf{1}_{n_m \times n_m}) \\ &= \Theta_\pi^\uparrow(A') + \left(\bigoplus_{j=1}^m (d_j - c_{jj})\text{Id}_{n_j}\right) \left(\bigoplus_{j=1}^m (\text{Id}_{n_j} - n_j^{-1}\mathbf{1}_{n_j \times n_j})\right). \end{aligned} \tag{6.3}$$

We claim that

$$\sigma(A) = \sigma(A') \cup \{d_1 - c_{11}, \dots, d_m - c_{mm}\}, \tag{6.4}$$

where $d_j - c_{jj}$ has multiplicity $n_j - 1$ for $j = 1, \dots, m$. If this holds then, since A and D are symmetric and have the same spectrum, there exists an orthogonal matrix Q such that $A = QDQ^T$.

It remains to show the claim (6.4), and we do so by explicitly producing an orthonormal eigenbasis. First, let V_j be the orthogonal complement of $\mathbf{1}_{I_j}$ in \mathbb{R}^{I_j} , padded by zeros to form a subspace of \mathbb{R}^N , where $N := n_1 + \dots + n_m$. A direct calculation shows that every non-zero vector in V_j is an eigenvector of A with eigenvalue $d_j - c_{jj}$. As the subspaces V_1, \dots, V_m are pairwise orthogonal, this eigenvalue has multiplicity $n_j - 1$ as required.

Next, let $\mathbf{w}_1, \dots, \mathbf{w}_m \in \mathbb{R}^m$ be an orthonormal eigenbasis for A' , with eigenvalues $\lambda_1, \dots, \lambda_m$. Then $\{\Theta_\pi^\uparrow(\mathbf{w}_1), \dots, \Theta_\pi^\uparrow(\mathbf{w}_m)\}$ is an orthonormal set, by Proposition 5.5(3), which lies in the span of $\{\mathbf{1}_{I_1}, \dots, \mathbf{1}_{I_m}\}$, by definition, so is orthogonal to V_i for $i = 1, \dots, m$. Furthermore, if $\mathbf{v} \in \mathbb{R}^m$ then Proposition 5.5(5) implies that

$$\bigoplus_{j=1}^m (\text{Id}_{n_j} - n_j^{-1}\mathbf{1}_{n_j \times n_j})\Theta_\pi^\uparrow(\mathbf{v}) = (\text{Id}_N - \Theta_\pi^\uparrow(\text{Id}_m))\Theta_\pi^\uparrow(\mathbf{v}) = \mathbf{0}_N.$$

Hence, by (6.3) and Proposition 5.5(5), we conclude that

$$A\Theta_\pi^\uparrow(\mathbf{w}_j) = \Theta_\pi^\uparrow(A')\Theta_\pi^\uparrow(\mathbf{w}_j) + \mathbf{0}_N = \Theta_\pi^\uparrow(A'\mathbf{w}_j) = \lambda_j\Theta_\pi^\uparrow(\mathbf{w}_j). \quad \blacksquare$$

In the same spirit as here, several important matrix calculations involving the matrix A , including powers, exponential, logarithm, and Gaussian log likelihood, can be performed by working with the compressed matrix A' . See [2] for more details.

6.3. Transversal matrix structures

The classical structures of Hankel and Toeplitz matrices interact with isogenic block stratification in a very rigid manner, as we explain below.

Consider first a positive semidefinite Hankel matrix $H = (c_{j+k})_{j,k=0}^N$ with real entries. It is well known (see, for example, [1]) that there exists at least one positive measure μ on the real line having c_j as power moments:

$$c_j = \int_{\mathbb{R}} x^j d\mu(x) \quad (j = 0, \dots, 2N).$$

Then, if H belongs to a stratum other than the topmost, $\mathcal{S}_{\pi_{\vee}}$, there are distinct indices j and k such that $c_{2j} = c_{j+k} = c_{2k}$. Consequently,

$$\int_{\mathbb{R}} (x^j - x^k)^2 d\mu(x) = 0,$$

so the measure μ is supported by at most three points, $-1, 0$ and 1 . It follows for $N \geq 2$ that H has rank at most 3 and extends uniquely to an infinite positive semidefinite Hankel matrix, whereas if $N = 1$ then $c_0 = c_1 = c_2$ and

$$\int_{\mathbb{R}} (1 - x)^2 d\mu(x) = 0,$$

so the measure μ is a point mass at 1.

A similar situation arises from the analysis of a positive semidefinite Toeplitz matrix $T = (s_{k-j})_{j,k=0}^N$. Here, we assume the entries to be complex and the positivity of T implies that

$$s_0 \geq 0 \quad \text{and} \quad s_{-j} = \overline{s_j} \quad (j = 0, \dots, N).$$

The solution to the truncated trigonometric moment problem guarantees a positive measure ν on $[-\pi, \pi)$ such that

$$s_j = \int_{-\pi}^{\pi} e^{j\theta} d\nu(\theta) \quad (j = -N, \dots, N).$$

If, as before, T has some non-trivial isogenic block structure, then there exists an index j satisfying

$$\int_{-\pi}^{\pi} e^{-j\theta} d\nu(\theta) = \int_{-\pi}^{\pi} d\nu(\theta) = \int_{-\pi}^{\pi} e^{j\theta} d\nu(\theta).$$

It follows that

$$\int_{-\pi}^{\pi} |1 - e^{j\theta}|^2 d\nu(\theta) = 0,$$

so the measure ν is the sum of at most j point masses, situated on the vertices of the regular polygon determined by that equation $z^j = 1$. Thus, the matrix T is degenerate and extends uniquely, by periodicity to an infinite positive semidefinite Toeplitz matrix.

6.4. Stability of semigroups

The simple observation that the spectral radius of a square matrix belonging to the closure of a certain stratum is preserved by the push-down operation Θ_π^\perp immediately resonates with stability criteria of evolution semigroups. To be more specific, consider the time-invariant linear homogeneous system of differential equations,

$$\frac{du}{dt}(t) = Au(t),$$

where A is an element of $\mathbb{C}^{N \times N}$ and $u : [0, \infty) \rightarrow \mathbb{C}^N$. As is well known, the solution

$$u(t) = \exp(tA)u(0) \tag{6.5}$$

decays exponentially as $t \rightarrow \infty$, for arbitrary initial data $u(0)$, if and only if the spectrum of A lies in the open left half-plane. In this case, the system is called asymptotically stable.

A great deal of work has been done to establish asymptotic-stability criteria for linear systems, in both the time-invariant and then time-dependent cases. A relatively early work in this area is [15], where the distance from a stable matrix to the unstable region is described definitively.

The isogenic block stratification introduced in the present article can be used in some cases to simplify the verification of stability. If the Cayley transform $X = (A - I)(A + I)^{-1}$ of A belongs to a closed stratum $\overline{\mathcal{S}_\pi}$ and the spectrum of X belongs to the open unit disk, then so does that of the compression $\Theta_\pi^\perp(X)$. Furthermore, the converse is true, which gives the following asymptotic-stability criterion.

Proposition 6.6. *Suppose the Cayley transform X of the generator A of the linear system (6.5) belongs to the stratum $\overline{\mathcal{S}_\pi}$. The system is asymptotically stable if and only if the spectral radius of the compression $\Theta_\pi^\perp(X)$ is less than 1.*

Much more involved, but well studied due to its important applications to control theory, is the case of a time-varying linear system

$$\frac{du}{dt}(t) = A(t)u(t).$$

As before, it may be beneficial to examine the compression of the Cayley transform of the family of matrices $A(t)$, but we do not enter into the details here.

One of the challenges of modern stability theory of switched dynamical systems is the computation of the joint numerical radius of a tuple of non-commuting matrices. More precisely, if A_1, A_2, \dots, A_n are complex $N \times N$ matrices, one wants to certify that

$$\lim_{m \rightarrow \infty} \max_{\sigma \in \{1, 2, \dots, n\}^m} \|A_{\sigma(1)} A_{\sigma(2)} \cdots A_{\sigma(m)}\|^{1/m} < 1.$$

While the general case of the corresponding weak inequality is known to be undecidable, sufficient conditions for the strong inequality are known and widely used; see [21] and references therein. Once again, being fortunate enough to have the matrices A_1, A_2, \dots, A_n in a high-codimension isogenic stratum may considerably simplify, via compression, the verification of such criteria.

6.5. Hierarchical matrices

With its origins in the theory of numerical approximation of integral equations, a novel chapter of applied linear algebra emerged in the last two decades having the concept of *hierarchical matrix* at its center [3, 13]. The philosophy behind this powerful new tool is to exploit, via effective numerical schemes, the sparsity and hidden redundancy in a large matrix of relative low rank. Starting with the factorization of a large matrix into a product of lower-rank rectangular matrices of smaller size, a systematic study of minimizing computational-cost algorithms for matrix multiplication, polar decomposition, Cholesky decomposition and much more were devised by Hackbusch, his disciples and an increasing number of followers. The foundational work [12] is complemented by the informative survey [14].

The matrices belonging to the closure $\overline{\mathcal{S}_\pi}$ of a high-codimension stratum defined by the partition $\pi = \{I_1, I_2, \dots, I_m\}$ of $\{1, \dots, N\}$ are hierarchical in an obvious way. Moreover, the compression $\Theta_\pi^\perp : \overline{\mathcal{S}_\pi} \rightarrow \mathbb{C}^{m \times m}$ naturally aligns with the main concepts of the hierarchical matrix theory. We indicate here a few common trends.

The isogenic structure of $A \in \overline{\mathcal{S}_\pi}$ is reflected by the factorization

$$A = \mathcal{W}_\pi B \mathcal{W}_\pi^*,$$

where B is an $m \times m$ matrix and $\mathcal{W}_\pi \in \mathbb{C}^{N \times m}$; see Proposition 4.4. Thus every element of $\overline{\mathcal{S}_\pi}$ can be decomposed with a universal left factor \mathcal{W}_π and a right factor depending on m^2 complex parameters. Moreover, the weighted compression Θ_π^\perp built on the same data (the partition π and the parameters determining each isogenic block) is a $*$ -algebra homomorphism. Specifically, a matrix $A \in \overline{\mathcal{S}_\pi}$ has the multiplicative decomposition

$$A = \mathcal{W}_\pi D_\pi^{-1} \mathcal{W}_\pi^* \Theta_\pi^\perp(A) \mathcal{W}_\pi D_\pi^{-1} \mathcal{W}_\pi^*,$$

and the correspondence $A \mapsto \Theta_\pi^\perp(A)$ preserves all matrix operations.

Proposition 6.7. *If the matrix $A \in \overline{\mathcal{S}_\pi}$, then compression $\Theta_\pi^\perp(A)$ has the same spectrum and singular values as A , with the possible exception of the value zero.*

Proof. Proposition 5.7 gives that $\sigma(A) \setminus \{0\} = \sigma(\Theta_\pi^\perp(A)) \setminus \{0\}$. Moreover, the singular values of A are the eigenvalues of its modulus $|A| = \sqrt{A^*A}$, but A^*A and $\Theta_\pi^\perp(A)^* \Theta_\pi^\perp(A)$ have the same non-zero eigenvalues. ■

Lifting the polar decomposition is slightly more subtle. Let $B = \Theta_\pi^\downarrow(A)$, where $A \in \overline{\mathcal{S}_\pi}$. The polar decomposition $B = U_B|B|$ lifts immediately to $A = V_A|A|$, but $V_A := \Theta_\pi^\uparrow(U_B)$ is only a partial isometry. There is, however, a unitary matrix $U_A \in \mathbb{C}^{N \times N}$ such that $U_A|A| = V_A|A|$. The initial and final spaces of V_A are equal and contain the range of A , and are spanned by $\{\mathbf{1}_{I_k} : k = 1, \dots, m\}$. If W is the orthogonal projection onto the orthonormal complement of this space then $U_A = V_A + W$ is as desired.

The lifting of an LU decomposition is similarly not quite immediate. As before, let B be the compression of the matrix $A \in \overline{\mathcal{S}_\pi}$ and suppose $B = LU$, where L is a lower-triangular matrix and U is upper triangular. Then the liftings $\Theta_\pi^\uparrow(L)$ and $\Theta_\pi^\uparrow(U)$ will be only block triangular, being elements of $\overline{\mathcal{S}_\pi}$. To obtain a genuine LU decomposition, one has to depart from the isogenic structure and split the products of rank-one matrices in a consistent pattern. For example, one can factor the product of $k \times n$ and $n \times m$ isogenic matrices as follows:

$$\begin{pmatrix} a & \cdots & a \\ \vdots & \ddots & \vdots \\ a & \cdots & a \end{pmatrix} \begin{pmatrix} b & \cdots & b \\ \vdots & \ddots & \vdots \\ b & \cdots & b \end{pmatrix} = \begin{pmatrix} a\sqrt{n} & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ a\sqrt{n} & 0 & \cdots & 0 \end{pmatrix} \begin{pmatrix} b\sqrt{n} & \cdots & b\sqrt{n} \\ 0 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & 0 \end{pmatrix}.$$

Finally, on the lighter side, we state the following theorem to give a demonstration of the general paradigm for results one can derive using the compression-inflation procedure described above.

Theorem 6.8. *Let A, B and C be real matrices belonging to $\overline{\mathcal{S}_\pi}$, where π is a partition of $\{1, 2, \dots, N\}$ with m elements. The classical equations of Lyapunov, Sylvester and Riccati,*

$$X + XA^T = C \quad AX + XB = C \quad \text{and} \quad AX + XA^T - XBX = C,$$

have a solution $X \in \mathbb{C}^{N \times N}$, with $X = X^T$ for the Riccati case, if and only if the compressed equations have a solution in $\mathbb{C}^{m \times m}$.

6.6. Coherent matrix organization

Complex multivariate data, such as those arising from the measurement of neuronal structures, cannot be handled without regularization and compression of the corresponding large matrices. Among the many techniques to analyze and transform such matrices, the method of *coherent matrix organization* proposed by Gavish and Coifman [10] stands out for its elegance and universality. These authors propose organizing a matrix using a natural metric which clusters the entries by proximity; the emerging tree of finer and finer partitions offers a canonical compression by averaging along strata, with tight control of the resulting approximation error.

The isogenic stratification of a matrix and the compression maps discussed above align perfectly with Gavish and Coifman's idea. Our purely theoretical setting is an extremal one, in the sense that we employ the discrete metric for clustering. However, it is notable that the isogenic structure and the associated inflation map offer a rapid matrix-completion algorithm within a prescribed stratum, in the spirit of recent advances in coherent matrix organization; see, for instance, [9, 11, 20]. We do not expand here some clear consequences of this non-accidental similarity.

Conflict of interest. On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

- [1] N. AKHIEZER, *The Classical Moment Problem and Some Related Questions in Analysis*, Hafner Publishing Co., New York, 1965.
- [2] I. ARCHAKOV and P.R. HANSEN, A canonical representation of block matrices with applications to covariance and correlation matrices, *Preprint* (2020).
- [3] M. BEBENDORF, *Hierarchical Matrices*, Springer-Verlag, Berlin, 2008.
- [4] A. BELTON, D. GUILLOT, A. KHARE and M. PUTINAR, Matrix positivity preservers in fixed dimension. I, *Adv. Math.*, **298** (2016), 325–368; doi: 10.1016/j.aim.2016.04.016.
- [5] A. BELTON, D. GUILLOT, A. KHARE and M. PUTINAR, Simultaneous kernels of matrix Hadamard powers, *Linear Algebra Appl.*, **576** (2019), 142–157; doi: 10.1016/j.laa.2018.03.035.
- [6] D. P. BLECHER, Real positive maps and conditional expectations on operator algebras, *Positivity and Its Applications* (E. Kikianty et al., eds.), Trends in Math., Birkhäuser, Cham, 2021, pp. 63–102; doi: 10.1007/978-3-030-70974-7_5.
- [7] S. BÖRM, L. GRASEDYCK and W. HACKBUSCH, Introduction to hierarchical matrices with applications, *Eng. Analysis with Boundary Elements*, **27** (2003), 405–422; doi: 10.1016/S0955-7997(02)00152-2.
- [8] J. CADIMA, F. L. CALHEIROS and I. P. PRETO, The eigenstructure of block-structured correlation matrices and its implications for principal component analysis, *J. Appl. Stat.*, **37** (2010), 577–589; doi: 10.1080/02664760902803263.
- [9] Y. CHEN, S. BHOJANAPALLI, S. SHANGAVI and R. WARD, Coherent matrix completion, *Proc. Machine Learning Research*, **32** (2014), 674–682; .
- [10] M. GAVISH and R. R. COIFMAN, Sampling, denoising and compression of matrices by coherent matrix organization, *Applied Comp. Harm. Analysis*, **33** (2012), 354–369; doi: 10.1016/j.acha.2012.02.001.
- [11] D. GROSS, Recovering low-rank matrices from few coefficients in any basis, *IEEE Trans. Information Theory*, **57** (2011), 1548–1566; doi: 10.1109/TIT.2011.2104999.
- [12] W. HACKBUSCH, A sparse matrix arithmetic based on \mathcal{H} -matrices I. Introduction to \mathcal{H} -matrices, *Computing*, **62** (1999), 89–108; doi: 10.1007/s006070050015.
- [13] W. HACKBUSCH, *Hierarchical Matrices: Algorithms and Analysis*, Springer, Heidelberg, 2015.

- [14] W. HACKBUSCH, Survey on the technique of hierarchical matrices, *Vietnam J. Math.*, **44** (2016), 71–101; doi: 10.1007/s10013-015-0168-5.
- [15] D. HINRICHSSEN and A. J. PRITCHARD, Stability radii of linear systems, *Systems Control Lett.*, **7** (1986), 1–10; doi: 10.1016/0167-6911(86)90094-0.
- [16] R. A. HORN, The theory of infinitely divisible matrices and kernels, *Trans. Amer. Math. Soc.*, **136** (1969), 269–286; doi: 10.1090/S0002-9947-1969-0264736-5.
- [17] J. HUANG and L. YANG, Correlation matrix with block structure and efficient sampling methods, *J. Comput. Finance*, **14** (2010), 81–94; doi: 10.21314/JCF.2010.216.
- [18] S. KARLIN and W. J. STUDDEN, *Tchebycheff Systems: With Applications in Analysis and Statistics*, Pure and Applied Mathematics, Vol. XV, Interscience Publishers John Wiley and Sons, New York – London – Sydney, 1966.
- [19] A. KHARE and T. TAO, On the sign patterns of entrywise positivity preservers in fixed dimension, *Amer. J. Math.*, **143** (2021), 1863–1929; doi: 10.1353/ajm.2021.0049.
- [20] G. MISHNE, R. TALMON, I. COHEN, R. COIFMAN and Y. KLUGER, Data-driven tree transforms and metrics, *IEEE Trans. Signal. Inform. Proc. Networks*, **4** (2018), 451–466; doi: 10.1109/TSIPN.2017.2743561.
- [21] P.A. PARRILO and A. JADBABAIE, Approximation of the joint spectral radius using sums of squares, *Linear Algebra Appl.*, **428** (2008), 2385–2402; doi: 10.1016/j.laa.2007.12.027.
- [22] P. Z. G. QIAN, H. WU and C. F. J. WU, Gaussian process models for computer experiments with qualitative and quantitative factors, *Technometrics*, **50** (2008), 383–396; doi: 10.1198/0040170080000000262.
- [23] A. J. ROTHMAN, E. LEVINA and J. ZHU, Generalized thresholding of large covariance matrices, *J. Amer. Statist. Assoc.*, **104** (2009), 177–186; doi: 10.1198/jasa.2009.0101.
- [24] O. ROUSTANT and Y. DEVILLE, On the validity of parametric block correlation matrices with constant within and between group correlations, *Preprint* (2017), arXiv: 1705.09793.
- [25] O. ROUSTANT, E. PADONOU, Y. DEVILLE, A. CLÉMENT, G. PERRIN, J. GIORLA and H. WYNN, Group kernels for Gaussian process metamodels with categorical inputs, *SIAM/ASA J. Uncert. Quant.*, **8** (2020), 775–806; doi: 10.1137/18M1209386.
- [26] P. SHAH and V. CHANDRASEKARAN, Group symmetry and covariance regularization, *Electron. J. Statist.*, **6** (2012), 1600–1640; doi: 10.1214/12-EJS723.
- [27] R. P. STANLEY, *Enumerative Combinatorics. Volume 1, second edition*, Cambridge University Press, Cambridge, 2012.

A. BELTON, Lancaster University, Lancaster, UK; *e-mail*: a.belton@lancaster.ac.uk
D. GUILLOT, University of Delaware, Newark, DE, USA; *e-mail*: dguillot@udel.edu
A. KHARE, Indian Institute of Science, Bangalore – 560012, India; and India and Analysis and Probability Research Group, Bangalore – 560012, India; *e-mail*: khare@iisc.ac.in

M. PUTINAR, University of California at Santa Barbara, CA, USA and Newcastle University, Newcastle upon Tyne, UK; *e-mail*: mputinar@math.ucsb.edu, mi-hai.putinar@ncl.ac.uk

Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit (<http://creativecommons.org/licenses/by/4.0/>).