

2023-07-23

# Content of nitrate and nitrite in commercial and self-made beetroot juices and the effect of storage temperature

Bescos Garcia, Raul

<https://pearl.plymouth.ac.uk/handle/10026.1/21064>

---

10.1002/fsn3.3575

Food Science and Nutrition

Wiley Open Access

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

1 **Title:** Content of nitrate and nitrite in commercial and self-made beetroot juices and the  
2 effect of storage temperature.

3 **Running title:** Nitrate and nitrite content in beetroot juice

4  
5 **Authors:**

6 Raul Bescos<sup>1\*</sup>, Mark L Rollason<sup>1</sup>, Tanisha S Davies<sup>1</sup>, Patricia Casas-Agustench<sup>1</sup>

7  
8 **Institutional affiliations:**

9 <sup>1</sup> School of Health Professions, Faculty of Health, University of Plymouth, Plymouth, PL4  
10 8AA, UK.

11  
12 **\*Corresponding author:**

13 Raul Bescos PhD

14 Lecturer in Physiology

15 Exercise, Nutrition and Health Laboratory, Drake campus (Link Building, ground floor)

16 PL4 8AA, University of Plymouth

17 United Kingdom

18 [Raul.Bescos@plymouth.ac.uk](mailto:Raul.Bescos@plymouth.ac.uk)

19 +44 0 7599 034710

20  
21 **Author contributions:** RB and MR contributed to project conception; RB, MR, TD and PC-  
22 A analysed the data. RB, MR, TD and PC-A authors contributed to data interpretation. RB  
23 and PC-A drafted the initial paper. All authors revised and approved the final manuscript.

**Abstract**

Popularity of beetroot juice (BJ) is growing due to its high inorganic nitrate content ( $\text{NO}_3^-$ ) and its potential physiological benefits. However, the content of  $\text{NO}_3^-$  is not indicated in most commercial BJs and it can be affected by seasonal changes and storage conditions. This study analysed the content of  $\text{NO}_3^-$  and nitrite ( $\text{NO}_2^-$ ) in five and two commercial and self-made BJs, respectively, that were purchased in the summer and winter period. The effect of storage temperature (20°C, 4°C and -20°C) and pH was also analysed. In non-concentrated BJs, the  $\text{NO}_3^-$  content was  $34 \pm 20\%$  ( $P = 0.075$ ) in the winter than in the summer.  $\text{NO}_3^-$  was fully degraded in self-made BJ after 3 days at 20°C. This effect was attenuated by 78% and 82% when it was kept at 4°C and -20°C, respectively. The addition of lemon juice (5%) to self-made BJ was another useful approach to avoid  $\text{NO}_3^-$  degradation for 3 days when it was kept at 20°C. Regarding  $\text{NO}_2^-$ , self-made BJ had higher concentration ( $0.097 \pm 0.01$  mg/mL) compared to commercial BJs ( $< 0.1$  mg/mL;  $P = 0.001$ ). The pH of self-made BJ was higher ( $6.3 \pm 0.1$ ) compared to commercial BJs ( $4.5 \pm 0.3$ ;  $P = 0.001$ ). These results suggest that the content of  $\text{NO}_3^-$  in non-concentrated BJs can substantially differ across the year and this is an important factor to take into account when recommending BJs to promote some of its potential physiological benefits.

**Keywords:** beet, nitrate, nitrite, nitric oxide, nitrogen.

## 42 **Introduction**

43 Beetroot is one of the main dietary sources of inorganic nitrate ( $\text{NO}_3^-$ )<sup>1</sup>, a natural ion that has  
44 been traditionally considered harmful due to the risk of formation of nitrosamines that can lead  
45 to cancer<sup>2,3</sup>. As a consequence, the European Food Safety Authority (EFSA) established an  
46 Acceptable Daily Intake (ADI) for  $\text{NO}_3^-$  of 3.7 mg/kg body mass/day that is still valid<sup>4</sup>.  
47 However, this view has substantially changed over the last decade due to new evidence  
48 suggesting that consumption of vegetables rich in  $\text{NO}_3^-$ , which can exceed the ADI levels, is  
49 safe and it can enhance nitric oxide (NO) bioavailability<sup>5-7</sup>. Increased NO bioavailability due  
50 to  $\text{NO}_3^-$  consumption has been associated with enhanced exercise capacity especially in  
51 moderate-trained individuals and some clinical populations<sup>7,8</sup>. Consequently, beetroot juice is  
52 currently listed within category A (products with the most scientific evidence to enhance  
53 exercise performance) in the Sport Supplement Framework of the Australian Institute of Sport  
54<sup>9</sup>.

55 The minimum amount of  $\text{NO}_3^-$  that can elicit improvements in exercise performance is about  
56 5 mmol (310 mg)<sup>8</sup>, however, the content of  $\text{NO}_3^-$  in most commercial beetroot juices is not  
57 indicated in the list of ingredients as it is not required by the law. This is important since the  
58 content of  $\text{NO}_3^-$  in vegetables can substantially change depending on several factors including  
59 environmental conditions, soil clay content, organic matter content, nitrogen fertilization and  
60 type of beet<sup>10</sup>. Previous research in lettuce and spinach has shown large variations in the  
61 content of  $\text{NO}_3^-$  and nitrite ( $\text{NO}_2^-$ ) across different seasons being higher in the winter than in  
62 the summer<sup>11</sup>. Similar data in beets or beetroot juice is missing, but it can be hypothesised that  
63 similar variations can also occur across the year.

64 Given the potential ergogenic effects of beetroot juice, the popularity of this product is  
65 increasing among professional and recreational athletes with an average increase in the sales

66 of beetroot of 8% per year since 2016<sup>12</sup>. However, some people dislike the taste of this product  
67<sup>13</sup>. One approach to make beetroot juice more attractive is by mixing it with other juices such  
68 as lemon and apple juice. They can also act as natural preservatives due to their antioxidant  
69 compounds (e.g ascorbic acid). However, the effect of adding lemon juice on the  $\text{NO}_3^-$  content  
70 of beetroot juice has not been reported. This study investigated whether the addition of lemon  
71 juice into self-made beetroot juice affects the content of  $\text{NO}_3^-$  and  $\text{NO}_2^-$  and acidity (pH) levels.

72 Another important issue that has not been analysed especially in commercial beetroot juices is  
73 the stability of  $\text{NO}_3^-$ . While small shots (70 ml) are easily consumed at once, large bottles (0.5  
74 – 1 L) can last longer and the storage conditions can affect the  $\text{NO}_3^-$  content. Fresh beetroot  
75 juice (from natural beets) may contain bacteria that can reduce  $\text{NO}_3^-$  into  $\text{NO}_2^-$  increasing the  
76 concentration of the second<sup>14</sup>. This is relevant because while inorganic  $\text{NO}_3^-$  is safe even at  
77 high doses,  $\text{NO}_2^-$  can cause serious harm at considerably lower levels. Thus, it is important to  
78 consider the content of  $\text{NO}_2^-$  in beetroot juice as well. In this study, we investigated the effect  
79 of 3 different storage temperatures (room: 20°C; fridge: 4°C; freezer: -20°C) on the content of  
80  $\text{NO}_3^-$  and  $\text{NO}_2^-$  in commercial and self-made beetroot juice.

81 In summary, the main goals of this study were to: 1) analyse the content of  $\text{NO}_3^-$  and  $\text{NO}_2^-$  of  
82 commercial and self-made beetroot juices at different periods of the year; 2) analyse the effect  
83 of the storage temperature on the content of  $\text{NO}_3^-$  and  $\text{NO}_2^-$  in commercial and self-made  
84 beetroot juice; 3) investigate the effect of adding lemon into self-made beetroot juice on the  
85  $\text{NO}_3^-$  and  $\text{NO}_2^-$  content and acidity (pH). According to this, the main hypotheses of this study  
86 were that: 1)  $\text{NO}_3^-$  content of commercial and self-made beetroot juice will differ across  
87 different periods of the year; 2)  $\text{NO}_3^-$  in commercial and self-made beetroot juice will be  
88 degraded more quickly at higher temperatures; 3) the addition of lemon juice into self-made  
89 beetroot juice will reduce  $\text{NO}_3^-$  degradation in self-made beetroot juice.

## 90 **Methods**

91 We analysed the  $\text{NO}_3^-$  and  $\text{NO}_2^-$  content in five commercial beetroot juices that are commonly  
92 used by professional and recreational athletes and two-self made beetroot juices (Table 1).  
93 Commercial juices and raw beets were purchased in June 2021 and February 2022 and stored  
94 for less than one week at room temperature or under refrigeration as recommended by  
95 manufacturers before they were analysed.

### 96 *Preparation of products*

97 Self-made beetroot juice (SBJ) was prepared using whole beets (*Beta Vulgaris*) from a local  
98 supermarket (Plymouth, UK). Beetroot was washed with tap water, and then with ultrapure  
99 water (Purelab OptionQ, Elga Veolia, UK). The outer skin and inedible parts were removed  
100 before being chopped into small pieces and weighed using an electronic scale (Precisa XB  
101 3200C, Switzerland). Then, beetroot was juiced using an electric juicer machine (Waring  
102 11JE65, USA). Lemon was bought in a local supermarket and juiced using a fruit juicer. Then,  
103 2.5 mL (5%) of lemon juice was mixed with 47.5 mL (95%) of fresh beetroot juice (SBJ),  
104 which is similar to the volume of lemon juice added into some commercial beetroot juices  
105 analysed in this study (Table 1). Commercial beetroot juices were opened on the first day of  
106 analyses. All beetroot juices were filtered using a Whatman® filter paper number 1 and  
107 centrifuged at 3,500 rpm for 10 min to remove solid parts.

### 108 *Analysis of nitrate ( $\text{NO}_3^-$ ) and nitrite ( $\text{NO}_2^-$ )*

109 All beetroot samples were centrifuged at 13,000 rpm at 4°C for 10 min before analysis. The  
110 content of  $\text{NO}_3^-$  and  $\text{NO}_2^-$  of each product was analysed using a dedicated high-performance  
111 liquid chromatography (HPLC) analyser (ENO-30; Eicom USA) as previously described <sup>15</sup>.  
112 Briefly,  $\text{NO}_3^-$  and  $\text{NO}_2^-$  were separated on a reverse-phase separation column packed with  
113 polystyrene polymer (NO-PAK 4.6 x 50 mm, EICOM, Amuza Inc, US), and  $\text{NO}_3^-$  was reduced

114 to  $\text{NO}_2^-$  in a reduction column packed with copper-plated cadmium filins (NO-RED EICOM,  
115 Amuza Inc, US).  $\text{NO}_2^-$  was mixed with a Griess reagent to form a purple azo dye in a reaction  
116 coil. The separation and reaction columns and the reaction coil were placed in a column oven  
117 set at  $35^\circ\text{C}$ . The absorbance of the color of the product dye at 540 nm was measured with a  
118 flow-through spectrophotometer (NOD-30, Eicom). The mobile phase (10% methanol, 0.15M  
119  $\text{NaCl}/\text{NH}_4\text{Cl}$  and 0.5 g/L 4Na-EDTA) and reactor phase (10% methanol, 1.25% HCl containing  
120 5 g/L of sulphanilamide with 0.25 g/L of N-naphthylethylenediamine) were delivered at a flow  
121 rate of 0.33 mL/min and 0.10 mL/min, respectively. A standard curve was produced by  
122 injecting 10  $\mu\text{L}$  of water with sodium  $\text{NO}_3^-$  ( $\text{NaNO}_3^-$  / 7631-99-4, Sigma Aldrich, UK) and  
123 sodium  $\text{NO}_2^-$  ( $\text{NaNO}_2^-$  / 7632-00-0, Sigma Aldrich, UK) at different concentrations (7.8  $\mu\text{M}$ ,  
124 15.6  $\mu\text{M}$ , 31.2  $\mu\text{M}$ , 62.5  $\mu\text{M}$ , 125  $\mu\text{M}$  and 250  $\mu\text{M}$ ). Beetroot samples were diluted 1:200 using  
125 a carrier solution containing 10% methanol, 0.15M  $\text{NaCl}/\text{NH}_4\text{Cl}$  and 0.5 g/L 4Na-EDTA.  
126 Samples were analysed (10  $\mu\text{L}$ ) in duplicate on the first day and single on third and seventh  
127 day given the small coefficient of variation of  $\text{NO}_3^-$  ( $2.1 \pm 1.9\%$ ) and  $\text{NO}_2^-$  ( $4.8 \pm 3.0\%$ )  
128 analyses.

### 129 *pH measurements*

130 Measurements of pH were performed using a single electrode digital pH meter (Lutron  
131 Electronic Enterprise Co Ltd., Model PH-208, Taiwan) that was calibrated following the  
132 manufacturer's instructions prior each use.

### 133 *Storage temperature*

134 The effect of different storage temperatures on the  $\text{NO}_3^-$  and  $\text{NO}_2^-$  content was only analysed in  
135 the first batch (June 2021). Eppendorf (1.5 mL) and Falcon tubes (3 mL) were filled with each  
136 product and kept at three different temperatures ( $20^\circ\text{C}$ ;  $4^\circ\text{C}$ ;  $-20^\circ\text{C}$ ) to analyse  $\text{NO}_3^-$ ,  $\text{NO}_2^-$  and  
137 pH on the first (baseline), third and seventh day using the same methods described above. All

138 the tubes were wrapped with aluminium foil to preserve the samples from light oxidation.  
139 Samples at -20°C were thawed the same day of the analysis. Then, all samples were centrifuged  
140 at 13,000 rpm at 4°C for 10 min before analysis was undertaken.

#### 141 *Statistical analyses*

142 Data are presented as mean  $\pm$  standard deviation (SD). Differences in  $\text{NO}_3^-$  and  $\text{NO}_2^-$  content  
143 and pH between different beetroot juices were compared using a one-way of analysis of  
144 variance (ANOVA). *Post hoc* analyses were performed using Tukey HSD. Data were analysed  
145 using the statistical software SPSS (version 28). The level of significance was set at  $P < 0.05$ .

146

## 147 **Results**

### 148 *Juices*

149 Raw beetroot (SBJ) in the summer (102 g) and winter (178 g) yielded 53 (52%) and 107  
150 (60%) mL of juice, respectively.

### 151 *$\text{NO}_3^-$ and $\text{NO}_2^-$ content in beetroot juices in the summer and winter*

152 As expected, concentrated beetroot juice (JW2) had the highest content of  $\text{NO}_3^-$  ( $6.3 \pm 0.2$   
153 mg/mL;  $P = 0.001$ ) compared to non-concentrated commercial (JW1:  $1.1 \pm 0.2$  mg/mL; BN:  
154  $1.1 \pm 0.1$  mg/mL; BT:  $1.6 \pm 0.2$  mg/mL; CW:  $0.8 \pm 0.1$  mg/mL) and self-made juices (SBJ:  $1.4$   
155  $\pm 0.2$  mg/mL; SBJL:  $1.3 \pm 0.2$  mg/mL) (Figure 1A). The content of  $\text{NO}_3^-$  of concentrated  
156 beetroot juice (JW2) was similar in the summer ( $6.3 \pm 0.2$  mg/mL) and winter ( $6.4 \pm 0.2$   
157 mg/mL;  $P > 0.05$ ). Non-concentrated beetroot juices (JW1, BN, BT, CW, SBJ), had in average  
158  $34 \pm 20\%$  more  $\text{NO}_3^-$  in the summer ( $1.2 \pm 0.3$  mg/mL) than in the winter ( $0.8 \pm 0.3$  mg/mL;  
159  $P = 0.075$ ) (Figure 1A). These differences were more pronounced in JW1, BN and BT juices  
160 (from  $0.7 \pm 0.1$ ,  $0.5 \pm 0.1$  mg/mL and  $0.8 \pm 0.1$  mg/mL in the winter to  $1.1 \pm 0.1$ ,  $1.1 \pm 0.1$



161 and 1.6 mg/m in the summer;  $P < 0.001$ ) than in CW and SBJ (from  $0.6 \pm 0.1$  and  $1.3 \pm 0.2$   
162 mg/mL in the winter to  $0.8 \pm 0.1$  and  $1.4 \pm 0.2$  mg/mL in the summer;  $P > 0.05$ ) (Figure 1A).

163 Using the  $\text{NO}_3^-$  results from each product, we calculated the amount of beetroot juice that was  
164 needed to achieve the minimum dose of  $\text{NO}_3^-$  to enhance exercise performance (5 mmol of  
165  $\text{NO}_3^- = 310$  mg) (Figure 2). With the exception of concentrated (JW2) and self-made juice  
166 (SBJ), an average of  $258 \pm 162$  mL more beetroot juice from the winter batches of commercial  
167 beetroot juices was needed to achieve such amount compared to the summer batches.

168 The content of  $\text{NO}_2^-$  is shown in Figure 1B. SBJ ( $0.097 \pm 0.01$  mg/mL) and SBJL ( $0.090 \pm 0.01$   
169 mg/mL) had the highest content of  $\text{NO}_2^-$  compared to the commercial beetroot juices (JW1:  
170  $0.030 \pm 0.01$  mg/mL; JW2:  $0.035 \pm 0.01$  mg/mL; BN:  $< 0.01 \pm 0.01$  mg/mL; BT:  $0.032 \pm 0.01$   
171 mg/mL; CW:  $0.023 \pm 0.01$  mg/mL;  $P = 0.001$ ) (Figure 1B). The content of  $\text{NO}_2^-$  was slightly  
172 lower in JW1 and BT juices in the winter (JW1:  $0.011 \pm 0.01$  mg/mL; BT:  $0.017 \pm 0.01$  mg/mL)  
173 than in the summer (JW1:  $0.030 \pm 0.01$  mg/mL; BT:  $0.032 \pm 0.01$  mg/mL;  $P = 0.110$ ), while  
174 JW2 had slightly higher content of  $\text{NO}_2^-$  in the winter ( $0.110 \pm 0.01$  mg/mL) than in the summer  
175 ( $0.097 \pm 0.01$  mg/mL;  $P = 0.101$ ).

#### 176 *pH of beetroot juice*

177 Results of pH are shown in Figure 1C. SBJ had the highest pH ( $6.3 \pm 0.1$ ) compared to  
178 commercial juices (mean pH from all the commercial beetroot juices =  $4.5 \pm 0.3$ ;  $P = 0.001$ )  
179 and SBJL ( $3.6 \pm 0.1$ ;  $P = 0.001$ ) (Figure 1C). Overall, the average pH of commercial juices  
180 (JW1, JW2, BN, BT, CW) was slightly lower in the winter ( $4.2 \pm 0.2$ ) than in the summer ( $4.5$   
181  $\pm 0.3$ ;  $P = 0.239$ ).

#### 182 *Effect of storage temperature on $\text{NO}_3^-$ , nitrite and pH*

183 The content of  $\text{NO}_3^-$  in juices stored at 20°C, 4°C and -20°C for 1, 3 and 7 days during the  
184 summer is shown in Figure 3.

185 A reduction of 24% (from  $6.3 \pm 0.2$  mg/mL to  $4.8 \pm 0.2$  mg/mL;  $P < 0.001$ ) and 46% (from  $6.3$   
186  $\pm 0.2$  mg/mL to  $3.4 \pm 0.2$  mg/mL,  $P < 0.001$ ) in  $\text{NO}_3^-$  was observed when concentrated beetroot  
187 juice (JW2) was kept at  $20^\circ\text{C}$  for 3 and 7 days, respectively (Figure 3A). A similar effect was  
188 observed when JW2 was kept for 3 days at  $4^\circ\text{C}$  (from  $6.3 \pm 0.2$  mg/mL to  $4.7 \pm 0.2$  mg/mL;  $P$   
189  $< 0.001$ ) and  $-20^\circ\text{C}$  (from  $6.3 \pm 0.2$  mg/mL to  $3.4 \pm 0.2$  mg/mL;  $P < 0.001$ ) (Figure 3B).

190  $\text{NO}_3^-$  was degraded in SBJ after 3 days at  $20^\circ\text{C}$  (from  $1.4 \pm 0.1$  mg/mL to  $0.04 \pm 0.01$  mg/mL;  
191  $P < 0.001$ ) (Figure 3A). This reduction was attenuated by 78% (from  $1.4 \pm 0.1$  mg/mL to  $1.1 \pm$   
192  $0.1$  mg/mL) and 82% (from  $1.4 \pm 0.1$  mg/mL to  $1.2 \pm 0.1$  mg/mL) when it was kept at  
193  $4^\circ\text{C}$  and  $-20^\circ\text{C}$  for 3 days, respectively (Figure 3B and 3C).

194 The addition of 5% lemon juice was also effective to fully attenuate the reduction of  $\text{NO}_3^-$  in  
195 SBJ for 3 days (from  $1.3 \pm 0.1$  mg/mL to  $1.3 \pm 0.1$  mg/mL) at  $20^\circ\text{C}$  (Figure 3A). Furthermore,  
196 the addition of lemon juice was useful to preserve 62% of  $\text{NO}_3^-$  in SBJ (from  $1.3 \pm 0.1$  mg/mL  
197 to  $0.8 \pm 0.1$  mg/mL) when it was kept at  $20^\circ\text{C}$  for 7 days (Figure 3A).

198 Regarding the  $\text{NO}_2^-$  content, an abrupt increase was observed in SBJ on day 3 at  $20^\circ\text{C}$  (from  
199  $(0.097 \pm 0.01$  mg/mL to  $1.5 \pm 0.2$  mg/mL;  $P < 0.001$ ) (Figure 3D). This effect was inhibited  
200 when SBJ was stored at  $4^\circ\text{C}$  for 3 (from  $0.097 \pm 0.01$  mg/mL to  $0.01 \pm 0.001$  mg/mL) and 7  
201 days (from  $0.097 \pm 0.01$  mg/mL to  $0.01 \pm 0.001$  mg/mL) and when it was stored at  $-20^\circ\text{C}$  for  
202 the same duration (3 days: from  $0.01 \pm 0.001$  mg/mL to  $0.01 \pm 0.001$  mg/mL; 7 days: from  
203  $0.01 \pm 0.001$  mg/mL to  $0.01 \pm 0.001$  mg/mL) (Figures 3E and 3F). Furthermore, the addition  
204 of lemon juice to self-made juice (SBJL) was effective to inhibit the increase in  $\text{NO}_2^-$  when it  
205 was kept at  $20^\circ\text{C}$  for 3 (from  $0.097 \pm 0.01$  mg/mL to  $0.087 \pm 0.01$  mg/mL) and 7 days (from  
206  $0.090 \pm 0.01$  mg/mL to  $0.11 \pm 0.01$  mg/mL), respectively (Figure 3D).

207 The pH of all juices, except in SBJL, remained relatively stable on day 3 and 7 at 20°C, 4°C  
208 and -20°C (Figures 3G, 3H and 3I). The pH of SBJL increased from day 1 to day 3 when it was  
209 kept at 20°C (from  $3.6 \pm 0.1$  to  $4.7 \pm 0.1$ ;  $P < 0.001$ ).

210

## 211 **Discussion**

212 The main finding of this study was that the content of  $\text{NO}_3^-$  in non-concentrated commercial  
213 beetroot juices was on average  $34 \pm 20\%$  lower in the winter than in the summer . Differences  
214 in the content of  $\text{NO}_3^-$  in concentrated commercial (JW2) ( $1.9 \pm 0.7\%$ ) and self-made beetroot  
215 juices (SBJ, SBJL) ( $5.7 \pm 2.1\%$ ) were smaller.

216  $\text{NO}_3^-$  is the main form of nitrogen used by crop to synthesise amino acids. They absorb  $\text{NO}_3^-$   
217 from the soil via transporter proteins in the root cell membrane <sup>16</sup>. Thus, the amount of  $\text{NO}_3^-$  in  
218 vegetables depends on the level of this ion in the soil, which can substantially differ across the  
219 year. For example, in the UK it has been reported that the soil is poorer in  $\text{NO}_3^-$  in the winter  
220 because wet conditions (rainfalls) can wash out  $\text{NO}_3^-$  into the groundwater, a phenomenon  
221 known as  $\text{NO}_3^-$  leaching <sup>17</sup>. For this reason, it is feasible to use additional fertilizer (nitrogen)  
222 in autumn and winter in some vulnerable areas to improve the crops yield <sup>17</sup>. Four of the  
223 commercial beetroot juices (JW1, JW2, BN, BT) analysed in this study indicated that beetroot  
224 used was organic so nitrogen fertilizers were not supposed to be used during the growth of the  
225 crop. Interestingly, all of them, except the concentrated juice (JW2), had lower content of  $\text{NO}_3^-$   
226 when they were bought and analysed in the winter, which may suggest that beetroot were grown  
227 over the summer. However, this information was not provided by the commercial companies.  
228 Light conditions, use of organic matter (animal manure), and storage conditions are also  
229 important factors to take into account as they can affect the content of  $\text{NO}_3^-$  in vegetables <sup>18-</sup>  
230 <sup>20</sup>. Commercial companies can obtain beetroot from different locations and areas given the

231 large amount of product needed to constantly supply the market, which can modify the content  
232 of  $\text{NO}_3^-$  in the final product. Furthermore, there is no regulation about labelling the content of  
233  $\text{NO}_3^-$  in commercial beetroot juice, its origin or when crop were harvested. This is relevant  
234 given the potential physiological implications of  $\text{NO}_3^-$  and the variations in the content of this  
235 ion observed in this study in some commercial products. Although individuals can always  
236 choose to consume larger-than-recommended amounts, potential disadvantages to doing so  
237 include increased cost, greater volume to ingest, higher intake of oxalate and potential side  
238 effects.

239 Only two of the commercial juices analysed in this study (JW1 and JW2) reported an estimated  
240 value of  $\text{NO}_3^-$  in the serving size (Table 1). The first juice (JW1) claimed that the  $\text{NO}_3^-$  content  
241 was on average 800 mg per litre (0.8 mg/mL). Compared to this, we found that the  $\text{NO}_3^-$  content  
242 of this product was 38% higher in the summer batch (June 2020) (1.1 mg/mL) and 14% lower  
243 in the winter batch (February 2021) (0.69 mg/mL). On the other hand, the  $\text{NO}_3^-$  content from  
244 a concentrated product from the same commercial brand (JW2) was 47% and 50% higher in  
245 summer and winter batch compared to the claimed  $\text{NO}_3^-$  content. Our results are in agreement  
246 to a previous study indicating that the  $\text{NO}_3^-$  content of the same beetroot juice was 23% higher  
247 than the claimed  $\text{NO}_3^-$  content <sup>21</sup>. However, they did not compare the content of  $\text{NO}_3^-$  of the  
248 same product across different periods of the year. Furthermore, both studies, showed that  
249 commercial concentrated beetroot shots (JW2) had nearly 5 times more  $\text{NO}_3^-$  than commercial  
250 non-concentrated and fresh beetroot juice. Concentrated beetroot shots appeared in the market  
251 a decade ago to provide the minimal dose of inorganic  $\text{NO}_3^-$  (5 mmol = 310 mg/serving) that  
252 has been suggested to enhance exercise capacity in a small volume <sup>22</sup>. Although the method  
253 to concentrate beetroot juice is not reported in the label, this process is usually performed by  
254 removing part of the water of the juice <sup>23</sup>.

255 Regarding the effect of temperature storage on the content of  $\text{NO}_3^-$  and  $\text{NO}_2^-$ , rapid degradation  
256 of  $\text{NO}_3^-$  occurred in self-made beetroot juice (SBJ) when it was kept at  $20^\circ\text{C}$  for 3 days, but  
257 this reaction was attenuated by storing it at low temperatures and by adding lemon juice  
258 (SBJL), a natural source of ascorbic acid. This is in agreement to our hypothesis suggesting  
259 that low temperatures and the addition of lemon juice can help to attenuate  $\text{NO}_3^-$  degradation  
260 in beetroot juice. Ascorbic acid is widely used in the food industry for its antioxidant and  
261 stabilising properties <sup>24</sup>. Two of the commercial juices analysed in this study (JW2 and BN)  
262 contained lemon juice, two more apple juice (JW1 and CW) and another one (CW) was fortified  
263 with ascorbic acid. Despite the addition of lemon juice into concentrated beetroot juice (JW2),  
264 we found a rapid reduction in the content of  $\text{NO}_3^-$  occurred over day 3 and 7 that was not  
265 attenuated at low temperatures. According to this, rapid consumption of concentrated beetroot  
266 juice is recommended to enhance  $\text{NO}_3^-$  intake. This is in agreement to the recommendations  
267 from the commercial companies indicating to keep the juice refrigerated and consume it within  
268 3 – 7 days once opened. The addition of lemon and apple juices can also help to enhance the  
269 organoleptic characteristics of beetroot juice for some people who dislike the taste of beetroot  
270 <sup>13</sup>.

271 The content of  $\text{NO}_2^-$  was very low ( $< 0.1 \text{ mg/mL}$ ) in all the juices at baseline, however, a rapid  
272 increase was observed in self-made beetroot juice (SBJ) on day 3 at  $20^\circ\text{C}$ . This could happen  
273 due to the activity of  $\text{NO}_3^-$  reductase enzymes or microorganisms present in beetroot as the  
274 decrease of  $\text{NO}_3^-$  was accompanied by the increase of  $\text{NO}_2^-$ . From a safety point of view, the  
275 levels of  $\text{NO}_2^-$  achieved on day 3 were quite low to cause harm in healthy individuals as doses  
276 above  $100 \text{ mg/kg}$  of body mass are required to produce serious side effects in humans <sup>25</sup>.  
277 According to our results, the consumption of over 4 litres of beetroot juice rich in  $\text{NO}_2^-$  over a  
278 relatively short period of time may be needed to reach this quantity of  $\text{NO}_2^-$ . However, a word

279 of caution is needed about beetroot juice overload among athletes thinking ‘the more the  
280 better’.

281 The pH of commercial beetroot juices was more acidic than self-made juice (SBJ), which can  
282 be related to lacto-fermentation and addition of ascorbic acid in commercial juices. Two  
283 commercial juices of this study were lacto-fermented (BN and BT), which consists of the  
284 addition of lactic acid bacteria consuming sugars to produce acid compounds and carbon  
285 dioxide by fermentation<sup>26</sup>. Three commercial juices also contained lemon juice (JW2 and BN)  
286 or vitamin C as an additive (CW). Addition of lemon juice into self-made beetroot juice (SBJL)  
287 is a useful approach to maintain the content of  $\text{NO}_3^-$  as we demonstrated in this study. On the  
288 other hand, further research is needed to investigate whether lacto-fermentation and/or addition  
289 of other juices can modify  $\text{NO}_3^-$  bioavailability.

290 This study had some limitations that are worth discussing. First, it was based on a Masters  
291 thesis that was performed during Covid-19 pandemic when students had to deal with laboratory  
292 restrictions. Bottles of five of the most consumed brands of beetroot juice in the UK were  
293 analysed in the summer (June) and winter (February) season. The batch code of each product  
294 was not recorded, but we believe that juices belonged from different batches given the time  
295 gap (8 months) between the purchase and analyses of them. Despite this limitation, our results  
296 are still interesting indicating that the content of  $\text{NO}_3^-$  can substantially differ especially in  
297 commercial non-concentrated beetroot juices. We also had limitations to increase the sample  
298 size of different beetroot juices given the duration of each chromatogram (10 min) to analyse  
299  $\text{NO}_3^-$  and  $\text{NO}_2^-$ . We could analyse a maximum of 42 samples in a day. In the winter batches  
300 (February 2021), only baseline analyses of  $\text{NO}_3^-$ ,  $\text{NO}_2^-$  and pH were performed due to time  
301 constraints. All the analyses ( $\text{NO}_3^-$ ,  $\text{NO}_2^-$  and pH) were performed in duplicate during the first  
302 day (7 samples x 3 different temperatures) to ensure the reproducibility of the results.  
303 Regarding the effect of storage temperature, only 1.5 and 3 mL of each beetroot juice were

304 taken and stored at the respective temperature prior to testing, which it may not represent of  
305 what would happen in larger volumes (e.g 500 mL) of juice.

306 In summary, this study showed that the  $\text{NO}_3^-$  content of commercial beetroot juices can  
307 substantially differ across different batches. Reduction of the content of  $\text{NO}_3^-$  in concentrated  
308 commercial beetroot juice and fresh beetroot juice occurs quickly at room temperature (20°C).  
309 Furthermore, it is possible to obtain similar quantities of  $\text{NO}_3^-$  from self-made beetroot juice  
310 compared to non-concentrated commercial beetroot juices, but it must be kept at low  
311 temperatures (4°C and -20°C) and/or mixed with lemon juice to avoid  $\text{NO}_3^-$  degradation. These  
312 findings are relevant to individuals (e.g nutritionists, athletes, coaches, etc.) and researchers  
313 interested in the physiological effects of beetroot juice supplementation. Indeed, given the  
314 possible variation in the  $\text{NO}_3^-$  content of beetroot juice, scientists looking at the physiological  
315 effect of dietary  $\text{NO}_3^-$  in beetroot juice should measure the content of  $\text{NO}_3^-$  in the supplement.

316 **Acknowledgement:** We thank the technical and funding support of the University of  
317 Plymouth.

318 **Declaration of interest:** This study was not funded by any commercial company. We also  
319 declare to not have any conflict of interest.

320 **Data availability:** Data will be available on request from the authors.

321 **Word count:** 3904

**322 References**

- 323 1. Bailey SJ, Winyard P, Vanhatalo A, et al. Dietary nitrate supplementation reduces the O<sub>2</sub> cost  
324 of low-intensity exercise and enhances tolerance to high-intensity exercise in humans. *J Appl Physiol*.  
325 2009;107(4):1144-55.
- 326 2. Zhang F-X, Miao Y, Ruan J-G, et al. Association between nitrite and nitrate intake and risk of  
327 gastric cancer: a systematic review and meta-analysis. *Med Sci Monit: Int Med J Experim Clin Res*.  
328 2019;25:1788.
- 329 3. Bryan NS, Alexander DD, Coughlin JR, Milkowski AL, Boffetta P. Ingested nitrate and nitrite  
330 and stomach cancer risk: An updated review. *Food Chem Toxicol*. 2012;50(10):3646-3665.
- 331 4. EFSA, Mortensen A, Aguilar F, et al. Re-evaluation of sodium nitrate (E 251) and potassium  
332 nitrate (E 252) as food additives. *Efsa journal*. 2017;15(6):e04787.
- 333 5. Lundberg JO, Carlström M, Weitzberg E. Metabolic Effects of Dietary Nitrate in Health and  
334 Disease. *Cell Metab*. 2018;28(1):9-22.
- 335 6. Hultström M, Amorim de Paula C, Porcelli S, et al. Commentaries on Viewpoint: Can elite  
336 athletes benefit from dietary nitrate supplementation? Commentaries on Viewpoint: Can elite athletes  
337 benefit from dietary nitrate supplementation? Commentaries on Viewpoint: Can elite athletes benefit  
338 from dietary nitrate. *J Appl Physiol*. 2015;119(6):762-769.
- 339 7. Senefeld JW, Wiggins CC, Regimbal RJ, Dominelli PB, Baker SE, Joyner MJ. Ergogenic Effect of  
340 Nitrate Supplementation: A Systematic Review and Meta-analysis. *Med Sci Sports Exerc*. 2020;  
341 52(10):2250-2261.
- 342 8. Shannon OM, Allen JD, Bescos R, et al. Dietary Inorganic Nitrate as an Ergogenic Aid: An Expert  
343 Consensus Derived via the Modified Delphi Technique. *Sports Med*. 2022; 52(10):2537-2558.
- 344



- 345 9. Australian Institute of Sport A. Supplements and sports food in high performance sport.  
346 Accessed 22/03, 2023. [https://www.ais.gov.au/\\_data/assets/pdf\\_file/0014/1000841/Position-](https://www.ais.gov.au/_data/assets/pdf_file/0014/1000841/Position-Statement-Supplements-and-Sports-Foods.pdf)  
347 [Statement-Supplements-and-Sports-Foods.pdf](https://www.ais.gov.au/_data/assets/pdf_file/0014/1000841/Position-Statement-Supplements-and-Sports-Foods.pdf)
- 348 10. Gallardo EJ, Coggan AR. What Is in Your Beet Juice? Nitrate and Nitrite Content of Beet Juice  
349 Products Marketed to Athletes. *Int J Sports Nutr Exerc Metab.* 2019;29(4):345.
- 350 11. Ashworth A, Bescos R. Dietary nitrate and blood pressure: evolution of a new nutrient? *Nutr*  
351 *Res Rev.* 2017:1-12.
- 352 12. Ysart G, Clifford R, Harrison N. Monitoring for nitrate in UK-grown lettuce and spinach. *Food*  
353 *Addit Contam.* Jul 1999;16(7):301-6.
- 354 13. Grand View Research. Sugar Beet Juice Extract Market Size. Accessed 26/09/2022,  
355 <https://www.grandviewresearch.com/industry-analysis/sugar-beet-juice-extract-market>
- 356 14. Babateen AM, Shannon OM, O'Brien GM, et al. Acceptability and Feasibility of a 13-Week Pilot  
357 Randomised Controlled Trial Testing the Effects of Incremental Doses of Beetroot Juice in Overweight  
358 and Obese Older Adults. *Nutrients.* 2021;13(3).
- 359 15. Corleto KA, Singh J, Jayaprakasha GK, Patil BS. Storage Stability of Dietary Nitrate and Phenolic  
360 Compounds in Beetroot (*Beta vulgaris*) and Arugula (*Eruca sativa*) Juices. *J Food Sci.* 2018;83(5):1237-  
361 1248.
- 362 16. Lundberg JO, Larsen FJ, Weitzberg E. Supplementation with nitrate and nitrite salts in exercise:  
363 a word of caution. *J Appl Physiol.* 2011;111(2):616-7.
- 364 17. Thomas B, Smallwood S, Cutler C, Bescos R. The oral nitrate-reducing capacity correlates with  
365 peak power output and peak oxygen uptake in healthy humans. *Nitric Oxide.* 2019;87:43-51.
- 366 18. Dechorgnat J, Nguyen CT, Armengaud P, et al. From the soil to the seeds: the long journey of  
367 nitrate in plants. *J Experim Bot.* 2010;62(4):1349-1359.
- 368 19. Gov.UK. Using nitrogen fertilisers in nitrate vulnerable zones. 26/09/2022,  
369 <https://www.gov.uk/guidance/using-nitrogen-fertilisers-in-nitrate-vulnerable-zones>

- 370 20. Santamaria P. Nitrate in vegetables: toxicity, content, intake and EC regulation. *J Sci Food*  
371 *Agric.* 2006;86(1):10-17.
- 372 21. Weightman RM, Huckle AJ, Roques SE, Ginsburg D, Dyer CJ. Factors influencing tissue nitrate  
373 concentration in field-grown wild rocket (*Diplotaxis tenuifolia*) in southern England. *Food Addit*  
374 *Contam Part A Chem Anal Control Expo Risk Assess.* 2012;29(9):1425-35.
- 375 22. Santamaria P, Elia A, Serio F, Todaro E. A survey of nitrate and oxalate content in fresh  
376 vegetables. *J Sci Food Agric.* 1999; 79(13):1882-1888.
- 377 23. Jones AM, Thompson C, Wylie LJ, Vanhatalo A. Dietary Nitrate and Physical Performance. *Ann*  
378 *Rev Nutr.* 2018;38(1):303-328.
- 379 24. Bazaria B, Kumar P. Compositional Changes in Functional Attributes of Vacuum Concentrated  
380 Beetroot Juice. *J Food Process Preserv.* 2016;40(6):1215-1222.
- 381 25. Liao M-L, Seib PA. Chemistry of L-ascorbic acid related to foods. *Food Chem.* 1988;30(4):289-  
382 312.
- 383 26. Klewicka E, Zduńczyk Z, Juśkiewicz J, Klewicki R. Effects of Lactofermented Beetroot Juice  
384 Alone or with N-nitroso-N-methylurea on Selected Metabolic Parameters, Composition of the  
385 Microbiota Adhering to the Gut Epithelium and Antioxidant Status of Rats. *Nutrients.* 2015;7(7):5905-  
386 5915.

387 **Table 1:** Beetroot juices analysed in this study.

<b>Brand</b>	<b>Code</b>	<b>Product</b>	<b>Serving size (mL)</b>	<b>Claimed nitrate content (mg/serving)</b>	<b>Characteristics</b>
<b>Juices</b>					
James White	JW1	Beet it organic juice	1,000	800	90% organic beetroot juice + 10% organic apple juice
James White (300 mg)	JW2	Beet it sport	70	300	98% organic concentrated beetroot juice + 2% lemon juice
Biona	BN	Beetroot pressed juice	500	-	Beetroot juice partially lacto fermented + lemon juice
Biotta	BT	Beetroot juice	500	-	100% organic pressed beetroot juice lacto-fermented.
Cawston	CW	Brilliant beetroot juice	1,000	-	90% pressed beetroot juice + 10% pressed apple juice + vitamin C
Fresh beetroot juice	SBJ	Fresh beetroot juice 1	50	-	100% pressed beetroot juice
Fresh beetroot juice + lemon juice	SBJL	Fresh beetroot juice	50	-	95% pressed beetroot juice + 5% pressed lemon juice

389 **Figure legends**

390 **Figure 1:** Content of nitrate ( $\text{NO}_3^-$ ) (A), nitrite ( $\text{NO}_2^-$ ) (B), and pH (C) in commercial and self-  
391 made beetroot juices in two different periods of the year. (*a* represents statistical differences  
392 between beetroot juices; *b* represents statistical differences between beetroot juice batches in  
393 the summer and winter.

394 **Figure 2:** Estimated amount of beetroot juice required to achieve 5 mmol of nitrate ( $\text{NO}_3^-$ )  
395 from commercial and self-made beetroot juices in the summer and winter.

396 **Figure 3:** Effect of storage temperature on the content of nitrate ( $\text{NO}_3^-$ ) (A-C), nitrite ( $\text{NO}_2^-$ )  
397 (D-F) and pH (G-I) of beetroot juice at baseline (day 1) and 3 and 7 days after opening the  
398 package (commercial beetroot juice) or after preparing self-made beetroot juice. (*a* represents  
399 statistical differences between day 1 and day 3; *b* represents statistical differences between day  
400 1 and day 7; *c* represents statistical differences between day 3 and day 7).