

2023-06

Terrestrial carbon isotope stratigraphy of the EoceneOligocene transition, Petrockstowe and Bovey basins, Devon, UK

Chaanda, MS

<https://pearl.plymouth.ac.uk/handle/10026.1/21172>

10.1016/j.pgeola.2023.05.003

Proceedings of the Geologists' Association

Elsevier BV

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1 **Terrestrial carbon isotope stratigraphy of the Eocene–Oligocene transition, Petrockstowe and**
2 **Bovey basins, Devon, UK**

3 Mohammed S. Chaanda^a, Stephen T. Grimes^a, Rhodri M. Jerrett^b, Mark Anderson^a, Melanie J. Leng^c
4 Meriel E. Fitzpatrick^a, Gregory D. Price^{a*}

5

6 ^aSchool of Geography, Earth and Environmental Sciences, Plymouth University, Drake Circus,
7 Plymouth, PL4 8AA, UK

8 ^bDepartment of Earth and Environmental Sciences, University of Manchester, Oxford Road,
9 Manchester, M13 9PL, UK

10 ^cBritish Geological Survey, Keyworth, Nottingham, NG12 5GG, UK.

11 ABSTRACT

12 The terrestrial sediments of the Petrockstowe and Bovey basins in Devon, UK were examined. Their
13 age is considered to be Eocene and Oligocene. The sediments (kaolinitic clays, silts, sands, gravels,
14 and lignites) from both basins were analysed for carbon isotopes of organic material, in conjunction
15 with total organic carbon and palynological analyses used to unravel the type of and provenance of
16 organic matter present. Within the Petrockstowe Basin, the lowermost interval examined shows a
17 palynological distribution dominated by phytoclasts, whilst the upper part of the core is dominated
18 by higher concentrations of palynomorphs (up to 90%) and an increase in amorphous organic
19 matter consistent (up to 37%) with a change from sand-filled fluvial channels followed by an
20 ephemeral lake or lake margin setting. Our palynological data from South John Acres Lane Quarry
21 section, Bovey Basin, show that within the lignites palynomorphs are high again (up to 95%)
22 consistent with them representing more ephemeral lakes or lake margins periodically exposed with
23 mires. Our palynological data set further allows us to determine that isotope trends are not overly
24 determined by the source of carbon in the basins. Our study suggests that the observed patterns
25 were primarily produced by variations of the isotope ratios of terrestrial atmospheric carbon
26 reservoirs. Even with our less than well constrained biostratigraphic control, the data indicate that
27 the carbon isotope excursions seen in the Eocene and Oligocene could be associated with several
28 transient carbon isotopic shifts (associated with the Palaeocene-Eocene Thermal Maximum). Our

29 findings therefore appear to lend support to the surface ocean and atmosphere behaving as
30 coupled reservoirs at this time.

31

32 Keywords: Eocene Oligocene Petrockstowe Bovey terrestrial carbon isotope, Palynological analyses

33

34 *Corresponding author.

35 E-mail address: g.price@plymouth.ac.uk (Gregory D. Price)

36

37

38 **1. Introduction**

39 The terrestrial sediments of the Petrockstowe and Bovey basins in Devon, UK offer an
40 opportunity to examine the carbon cycle during the Eocene–Oligocene transition (~33.9 million
41 years ago), an interval, that saw a climate shift from a largely ice-free greenhouse conditions to an
42 icehouse world (Miller et al., 2009; Coxall and Wilson, 2011; Hutchinson et al. 2021). At this time
43 major changes in fauna and flora record a shift toward more cold-climate-adapted species (e.g., Sun
44 et al., 2014). For this pivotal interval in Earth's climate, our understanding the role of the carbon
45 cycle, is much more limited (Coxall and Wilson, 2011; Armstrong McKay et al. 2016). Across the
46 Eocene–Oligocene boundary a positive carbon isotope($\delta^{13}\text{C}$) excursion of ~1.0‰ is typically
47 recorded in the marine record followed by a decline to pre-excursion values (~0.7‰) in the
48 Oligocene (e.g., Nilsen et al. 2003; Armstrong McKay et al. 2016). This well-documented
49 perturbation may be used to correlate marine and terrestrial sections from around the globe as
50 previous studies have shown that $\delta^{13}\text{C}$ obtained from terrestrial organic material such as wood and
51 lignites typically records a global signal (e.g., Heimhofer et al., 2003; Collinson et al. 2003; Gröcke et
52 al. 2005; Bechtel et al. 2009; Hodgson et al. 2011; Jerrett et al. 2015; Lenz et al. 2022).

53 The Petrockstowe and Bovey basins lie on the Sticklepath - Lustleigh Fault and owe their
54 origin to subsidence within this zone (Blyth 1962, Dearman 1963). About 600 m of Eocene–
55 Oligocene sediments are present in the Petrockstowe Basin (Freshney et al. 1979) and ~ 1200 m in
56 the Bovey Basin (Edwards 1976). In this study, we present new organic $\delta^{13}\text{C}$ data from terrestrial
57 Eocene–Oligocene aged sediments from these basins. Our data is used to improve age constraints
58 on the succession via comparison of our terrestrial $\delta^{13}\text{C}$ record with that of the extensively
59 described time-equivalent marine sections. The similarity and magnitude of the $\delta^{13}\text{C}$ excursions
60 between terrestrial and marine records can also be used to assess whether these archives behaved
61 as coupled reservoirs during this time.

62

63 **2. Location and Geological setting**

64 The Petrockstowe Basin, nr Newton Abbott, Devon, UK (Fig. 1), lies on the Sticklepath –
65 Lustleigh Fault (Dearman 1963; Holloway and Chadwick, 1986). The fault became active during the
66 Paleogene and most activity ceased before deposition of the upper part of the Bovey Formation
67 (Blyth 1962). Bristow and Robson (1994) proposed a structural model for the development of the
68 basin – a pull-push model – and suggested that the development was in two phases: an early,

69 transitional phase, during which much of the sedimentation occurred, and a subsequent
70 transpressional phase in which boundary thrust faults developed. Geophysical measurements,
71 confirmed by a British Geological Survey (BGS) borehole from the centre of the basin proved a basin
72 fill of 660 m of sands and silts (Freshney et al. 1979). The sediments, kaolinitic clays, silts, sands,
73 gravels, and lignite, were likely to have been derived from weathering granite under warm
74 temperate or sub-tropical conditions of the early Paleogene (Bristow 1968; Edwards 1976). The
75 succession consists of fining-upwards cycles comprising of one or more of a gravel lag, gravelly
76 sands, silty sands, and are probably representative of point bar and swale-fill deposits of a river
77 system. These interstratify with clays and silts of lacustrine origin (Freshney 1970; Freshney et al.
78 1979). Based on palynological evidence, Turner (cited by Freshney et al. 1979) suggested these
79 were deposited in a subtropical climate with palms, ferns and heathers and many plants with
80 swamp affinities.

81 The Bovey Basin is located between Newton Abbot, Kingsteignton and Bovey Tracey, Devon
82 UK (Fig. 2) and is 45 km southeast of the Petrockstowe Basin. The Bovey Basin lies southeast of the
83 Dartmoor granite and is approximately 7 km from east to west and 5 km from north to south. In the
84 northern and eastern boundaries of the basin there are sedimentary contacts between the
85 Dartmoor granite and the Upper Greensand and Aller Gravel. The bulk of the basin is filled by a
86 thick (~ 1200 m) succession of Paleogene kaolinitic clays, silty clays, silts, lignites and sands,
87 referred to as the Bovey Formation (Edwards 1976). Edwards (1976) proposed a morphological sub-
88 division of the basin into two parts, lying to the north and to the south of Newton Abbot. The part
89 between Bovey Tracey and Newton Abbot is considered as the main basin; the second part lies
90 south of Newton Abbot and is referred to as Decoy Basin (Fig. 2). Edwards and Freshney (1982)
91 proposed an informal sub-division of the Bovey Formation into 'lower', 'middle' and 'upper'. The
92 'lower' is not exposed and the 'middle' and 'upper' Bovey Formation includes 14 members, some of
93 uncertain stratigraphic position or lateral equivalents and are described in detail by Edwards and
94 Freshney (1982) and Selwood et al. (1984). Of the top 350 exposed at the surface (Edwards 1976),
95 48 m of the Abbok Clay and Sand and Southacre Clay and Lignite members of the 'middle' Bovey
96 Formation were examined in the exposed working section at the South John Acre Lane Quarry.
97 Chandler (1957) and Edwards (1976), suggest that during the Oligocene the lignites accumulated in
98 swamps with associated fluvial sands and plant debris swept in from a warm hinterland into a lake

99 basin lying on Palaeozoic strata (see also Selwood et al. 1984). The lake was surrounded by
100 marshland tree covered slopes (Chandler 1957).

101

102 **3. Age of the Petrockstowe and Bovey basins**

103 The age of the Bovey lignites has for a long time been debated (Chandler 1964). Based upon
104 the macroflora in the lignite beds of the Bovey Formation, these were originally regarded as
105 Miocene, but later assigned to the Oligocene (Chandler 1957, 1964). Likewise, Wilkinson (1979)
106 cited by Selwood et al. (1984) noted that pollen from a borehole, near Heathfield that penetrated
107 185 m of Blatchford Sand (upper Bovey Formation), 69 m of South Acre Clay and Lignite and 51 m
108 of Abbrook Clay and Sand and from below 290 m depth, Eocene indicators like *Anacolosidites* and
109 *Pompeckjoidaepollenites* were observed (Fig. 3). It is also important to note that the Blatchford
110 Sand is an obsolete unit name and has been replaced by the Woolley Grit Member. The South Acre
111 Clay and Lignite Member is therefore likely to be early to middle Oligocene in age (Selwood et al.
112 1984) and the Abbrook Clay and Sand Member would contain the Eocene–Oligocene boundary.
113 Freshney et al. (1982) also suggested that the lowermost ~ 700–800 m of the Bovey Formation
114 could probably be assigned to Eocene (see also Wilkinson et al. 1980; Wilkinson and Boulter, 1981).
115 For the Petrockstowe Basin, Turner (cited by Freshney et al. 1979) reported that pollen data
116 indicate a boundary between the Oligocene and Eocene at ~120 m depth in the BGS Borehole No.1
117 (Fig. 3).

118 **4. Methods**

119 From Petrockstowe 2 borehole cores (Petrockstowe 1A and 1B) held in the core repository
120 at the BGS, Keyworth, Nottingham, UK were logged, and sub sampled. The sampled section was 640
121 m long, and samples were collected, on average, every 6 m. Within the region of the Eocene
122 Oligocene boundary as proposed by Turner (cited by Freshney et al. 1979), as well as the early
123 Eocene of Core 1B higher resolution sampling was undertaken. In the Bovey Basin, the Abbrook
124 Clay and Sand and Southacre Clay and Lignite members of the Bovey Formation from the accessible
125 exposed working section at South John Acre Lane Quarry (Grid Reference SX 858758) were
126 sampled. The sampled section was 48 m and samples were collected, on average, every 0.6 m. It
127 was necessary to excavate the sediment surface by up to 0.5 m before sampling with a trowel to
128 ensure fresh samples. All sediment types were sampled.

129 For the determination of the carbon isotope composition of total organic carbon ($\delta^{13}\text{C}_{\text{TOC}}$),
130 samples were ground to a fine powder using an agate pestle and mortar. Powdered samples were
131 decarbonated by placing each sample in a 50 ml polypropylene centrifuge tube and treating with
132 10% HCl for 1 h until any carbonate had reacted. Samples were then rinsed with deionized water,
133 centrifuged, and rinsed again until neutrality was reached (using universal indicator paper). For
134 $\delta^{13}\text{C}_{\text{TOC}}$ analysis, samples were weighed, to achieve ~ 0.5 mg TOC, into a tin capsule and placed into
135 a Carlo Erba 1500 EA for analysis using an online VG Triple Trap Mass Spectrometer. The $\delta^{13}\text{C}_{\text{TOC}}$
136 results were calibrated against Vienna PeeDee Belemnite (V-PDB) through laboratory (BROC1) and
137 International Standards (NBS19, NBS22, CH6). Standards were evenly distributed throughout the
138 individual isotope runs to correct for daily drift. The mean standard deviation on replicate $\delta^{13}\text{C}_{\text{TOC}}$
139 analyses of laboratory standard (BROC1) and soil (SOILB) was between $\pm 0.1\%$ and 0.5% (1
140 Standard Deviation, σ) for $\delta^{13}\text{C}_{\text{TOC}}$. Replicate analyses showed an average precision of $\pm 0.1\%$. TOC
141 content for each sample was measured using a Carlo Erba 1500 elemental analyser with acetanilide
142 used as the calibration standard.

143 Palynological analyses were used to unravel the type of organic matter associated with the
144 sediments and as a means of determining the source of carbon reservoir in the basins. Samples
145 were processed using standard palynological processing (Brown, 2008)(hydrochloric acid followed
146 by hydrofluoric acid for demineralisation). Slides were studied using a Zeiss standard microscope,
147 normally using standard transmitted light. This is the first time such a method was used in both
148 Petrockstowe and Bovey basins. To achieve this, counts of >300 organic matter types from each
149 sample was made. There are several schemes to classify different components of the particulate
150 organic matter (e.g., Tyson, 1995, Aggarwal et al., 2019). Four main categories of palynological
151 matter were identified in this study (Fig. 4): (1) Non-opaque phytoclasts includes woody remains,
152 tracheid material, poorly lignified, tissue fragments derived from higher plants, yellowish-brown
153 organic remains (2) Opaque phytoclast includes palynodebris with irregular shapes and charcoal (3)
154 Palynomorphs in this study include pollen, spores and undifferentiated forms (4) amorphous
155 organic matter (AOM) and other palynodebris which appears grey, pale yellow or brown in colour,
156 partly translucent masses of variable thickness and with no cellular detail. The AOM group probably
157 originates from bacteria, phytoplankton and degraded organic aggregates. Their size varies from <5
158 to about $45\ \mu\text{m}$ in diameter.

159 5. Results

160 5.1 Total Organic Carbon (%TOC)

161 The sediments from the Petrockstowe core have highly varying wt. % TOC values ranging
162 from 0.02 to 42.7 wt. % TOC (Fig. 5). Unsurprisingly, the highest %TOC values coincide with the
163 lignitic clays and lignites. These lignitic clays and lignites are seen in the middle and upper parts of
164 the core (Core 1A). The lower part (core 1B) consists mostly of gravels, sands, and clays with very
165 low wt. % TOC contents. The wt. % TOC values from the South John Acres Lane Quarry section,
166 Bovey Basin, range from 0.1 to 61.8 % (Fig. 6). As for the Petrockstowe Basin, the highest wt. % TOC
167 values coincide with either lignitic clays or the lignites. These sediments are seen within the
168 Southacre Clay and Lignite Member whereas the underlying Abbrook Clay and Sand Member is
169 dominated by sands and silty clays with fewer lignitic clay beds.

170 5.2 Palynology

171 Within the cores of the Petrockstowe Basin, the lowermost interval shows a palynological
172 distribution dominated by phytoclasts with at certain intervals nearly 100% and low palynomorphs
173 and AOM. The upper part of the core is dominated by much higher concentrations of palynomorphs
174 and an increase in AOM (up to 37%) and low concentrations of phytoclasts.

175 With respect to South John Acres Lane Quarry section, Bovey Basin, phytoclasts are highest in the
176 Abbrook Clay and Sand Member. When phytoclasts are high (opaque phytoclasts reach 91%), the
177 palynomorphs shows lowest concentrations and vice versa. High AOM concentrations (up to 66%)
178 are seen at the base of Abbrook Clay and Sand Member and decline upwards. In the overlying
179 Southacre Clay and Lignite Member, dominated by lignites, opaque phytoclast concentrations are
180 low, palynomorphs consistently high (spore-pollen and non-opaque phytoclasts reach 95%, Fig. 6).

181 5.3 Carbon Isotopes ($\delta^{13}C_{TOC}$)

182 The $\delta^{13}C_{TOC}$ values of samples in the Petrockstowe cores range from -28.5‰ to -23.5‰
183 with a mean value of -26.5‰ . As can be seen in Figure 5, values at the base of core 1B begins with
184 a $\delta^{13}C_{TOC}$ value of $\sim -26.1\text{‰}$ at 645 m. A carbon isotope excursion with a magnitude of $\sim 2.5\text{‰}$ can
185 be seen at ~ 586 m depth with $\delta^{13}C_{TOC}$ values reaching a minimum of -28.6‰ . The entire excursion
186 occurs over a thickness of ~ 19 m, from 586 – 605 m. The data then shows a return to more positive
187 values of -26.2‰ at 585 m. Thereafter, the $\delta^{13}C_{TOC}$ values remain relatively consistent between 584
188 m and 540 m with $\delta^{13}C_{TOC}$ in the range of -27.0‰ and -26.3‰ . There is a lack of core (because of

189 poor recovery) between 513.59 m to 431.60 m. In the upper part of the Petrockstowe 1A core, the
190 $\delta^{13}\text{C}_{\text{TOC}}$ values generally vary around -26.0‰ . In the uppermost (Oligocene) part of the core the
191 most positive $\delta^{13}\text{C}_{\text{TOC}}$ values are seen.

192 At South John Acres Lane Quarry section, Bovey Basin the $\delta^{13}\text{C}_{\text{TOC}}$ values range between –
193 27.8‰ to -22.5‰ with a mean value of -26.0‰ (Fig. 6). In this succession, $\delta^{13}\text{C}_{\text{TOC}}$ values show
194 limited variability. In the uppermost (Oligocene) part of the section within the lignitic clays and
195 lignites the most positive $\delta^{13}\text{C}_{\text{TOC}}$ values (-22.5‰) are found.

196

197 **6. Discussion**

198 *6.1 Palynological Interpretation*

199 In the Petrockstowe 1A and 1B cores, close to the base of the succession the high opaque
200 phytoclast content, together with low TOC values (down to 0.1 wt. %), and low AOM and non-
201 opaque phytoclast contents may be related to local oxidation of organic matter (Figs. 5, 7) or
202 diagenesis. Opaque phytoclasts are typically derived from the oxidation of structured organic
203 matters (translucent brown wood, tracheids, cuticle, etc.) and along with a low proportion of the
204 other organic matter types has been documented in oxic swamps and river sediments (e.g., Martín-
205 Closas et al., 2005; Pieńkowski and Waksmundzka, 2009). In the upper part of the Petrockstowe
206 section (Fig. 5) palynomorphs dominate which could indicate suboxic/anoxic waters (Tyson, 1995).
207 Consequently, a restriction of water circulation rather than productivity, may serve as the
208 controlling factor for the organic rich sediment accumulation. Also, the fluctuating, but relatively
209 high percentages of palynomorphs and AOM (up to 37%), could suggest diverse source of areas of
210 the organic matter (e.g., Martín-Closas et al. 2005) with deposition within an ephemeral lake or lake
211 margin. These observations agree with the Petrockstowe 1A and 1B cores, representing a
212 succession of sand-filled fluvial channels followed by an ephemeral lake or lake margin setting (Fig.
213 5, see Freshney et al., 1979). This represents an overall deepening-up sequence.

214 In the Abbrook Clay and Sand and Southacre Clay and Lignite members, the base sees a
215 diversity in palynological types (Fig. 6) with high percentage of AOM and phytoclasts potentially
216 reflecting enhanced preservation in low energy, stagnant, oxygen depleted environment (Tyson
217 1995, Fig. 7). Only a single high opaque phytoclast level is seen, possibly associated with deposition

218 within an oxidising environment. In the overlying Southacre Clay and Lignite Member, dominated
219 by lignites, palynomorph concentrations are consistently high and as such suggests a depositional
220 environment associated with a swamp or ephemeral lake or marginal lake. These observations are
221 in agreement that the Abbrook Clay and Sand and Southacre Clay and Lignite members of the
222 Bovey Formation, represents a long-lived lake followed by sediments representing more ephemeral
223 lakes or lake margins periodically exposed with mires (see Chandler, 1964; Edwards, 1976).

224 Our palynological data set further allows us to determine whether isotope trends are carbon
225 source related. For example, within the Abbrook Clay and Sand Member, when phytoclasts are
226 high, the palynomorphs shows lowest concentrations and vice versa. High, but variable, AOM
227 concentrations are also seen. No correspondence is seen here with fluctuations in carbon isotopes,
228 suggesting organic matter associated with the sediments is not overly determining the source of
229 the carbon reservoir in the basins. Nevertheless, changes in the dominance of gymnosperms,
230 angiosperms or pteridophytes/bryophytes within the vegetation could be of importance.

231 *6.2 Carbon isotope trends*

232 Carbon isotopic ratios from terrestrial organic materials have been previously used to study
233 global carbon-isotope excursions in the Cenozoic (Collinson et al. 2003, Bechtel et al. 2009;
234 Holdgate et al. 2009; Hodgson et al. 2011; Fang et al., 2013; Jerrett et al. 2015; Garel et al. 2020;
235 Lenz et al. 2022). These studies (which use discrete plant fragments, lignites or disseminated
236 organic matter) identify reproducible patterns in atmospheric carbon isotopic compositions. There
237 are just a few terrestrially sourced high-resolution carbon isotope stratigraphies to compare our
238 Eocene and Oligocene data to (e.g., Holdgate et al. 2009; Garel et al. 2020). Nevertheless, our
239 $\delta^{13}\text{C}_{\text{TOC}}$ data are, consistent with terrestrially sourced $\delta^{13}\text{C}$ values of the Eocene (e.g., Collinson et al.
240 2003, Bechtel et al. 2009; Hodgson et al. 2011). Considering the carbon isotope excursion of -2.5‰
241 from the lower part of Petrockstowe core 1B (Fig. 5), the magnitude of this excursion falls within
242 the lower limit of that is associated with the Palaeocene-Eocene Thermal Maximum (PETM), which
243 ranges from -2.4 to -6.3‰ (see summary of McInerney and Wing 2011). This suggests it could be
244 related to this event. However, biostratigraphically there is limited data from the Petrockstowe
245 core (see Turner cited by Freshney et al. 1979). The biostratigraphic constraints allow the carbon
246 isotope excursion to also be associated with one of the other transient carbon isotopic shifts that
247 occurred after the Palaeocene-Eocene Thermal Maximum i.e., the Eocene Thermal Maximum (ETM-
248 2). For example, the magnitude of the ETM-2 carbon isotope excursion documented in the

249 continental succession of the McCullough peaks, Bighorn Basin, Wyoming, USA, using paleosol
250 carbonate is of -3.8‰ (Abels et al. 2012).

251 The presence of the Eocene – Oligocene boundary in the Petrockstowe core 1A has been
252 proposed, based on pollen data, by Turner (cited by Freshney et al. 1979). The Abbrook Clay and
253 Sand Member is also likely to contain the Eocene–Oligocene boundary (Selwood et al. 1984) but
254 because of the limited biostratigraphic data the exact positioning of the boundary is less certain.
255 The Eocene – Oligocene boundary is one of the most prominent abrupt climatic events in the
256 Cenozoic and is considered to represent the initiation of major permanent Palaeogene ice sheets
257 on Antarctica (Miller et al., 2009; Coxall and Wilson, 2011; Hutchinson et al. 2021). The glaciation of
258 Antarctica is thought to result from the tectonic opening of Southern Ocean gateways, which
259 enabled the formation of the Antarctic Circumpolar Current and the subsequent thermal isolation
260 of the Antarctic continent (e.g., Zachos et al. 2001) Modelling studies implicate low atmospheric
261 CO_2 also as an important factor (DeConto and Pollard, 2003). The carbon isotope changes across
262 this boundary are, however, less pronounced, and certainly less well documented in the terrestrial
263 system. This is perhaps due to the lack of suitable terrestrial sections to study.

264 Nevertheless, the marine records (Zachos et al. 2001; Coxall and Wilson, 2011) show a $\delta^{13}\text{C}$
265 excursion of $\sim 1.0\text{‰}$ in benthic foraminifera, peaking in the earliest Oligocene and followed by a
266 decline to $\sim 0.5\text{‰}$, 1 million years after the boundary. The Petrockstowe and Bovey $\delta^{13}\text{C}_{\text{TOC}}$ data do
267 show some correspondence with this marine record whereby for the Eocene stable but the most
268 negative carbon values are observed, whereas the most positive carbon isotope values are present
269 in the Oligocene. More positive $\delta^{13}\text{C}_{\text{TOC}}$ values have been linked to increased organic carbon burial
270 (Coxall and Wilson, 2011). Our study therefore supports the notion that the surface ocean and
271 atmosphere behaved as coupled reservoirs at this time, similar to other times in the Cenozoic
272 (Jerrett et al. 2015; Cui et al., 2021; Lenz et al. 2022), as opposed to a decoupled system (cf.
273 Holdgate et al. 2009; Fang et al., 2013), but more data is required to fully test this possibility.

274 **7. Conclusions**

275 In conclusion, and in agreement with Freshney et al., (1979), our palynological observations
276 show that the Petrockstowe 1A and 1B cores, represent a succession of sand-filled fluvial channels
277 followed by an ephemeral lake or lake margin setting. The Abbrook Clay and Sand and Southacre
278 Clay and Lignite members of the Bovey Formation, represents a long-lived lake followed by

279 sediments representing more ephemeral lakes or lake margins periodically exposed with mires (see
280 Chandler, 1964; Edwards, 1976). Our palynological data set further allows us to determine that
281 isotope trends are not overly determined by the source of carbon in the basins.

282 Our study suggests that the observed $\delta^{13}\text{C}_{\text{TOC}}$ trends in the Eocene–Oligocene of the Petrockstowe
283 and Bovey basins were primarily produced by variations of the carbon isotope ratios of terrestrial
284 atmospheric carbon reservoirs. Even with our less than well constrained biostratigraphic control,
285 the data indicate that the carbon isotope excursions seen in the Eocene and Oligocene could be
286 associated with a number of transient global carbon isotopic shifts (e.g., the PETM). Our findings
287 therefore appear to lend support to the surface ocean and atmosphere behaving as coupled
288 reservoirs at this time.

289 **Acknowledgements**

290 We would like to thank the British Geological Survey for access to cores and Sibelco UK for access
291 and information regarding South John Acres Lane Quarry. MC was funded by the Tertiary Education
292 Trust Fund (TETFund) Abuja, via the intervention of academic staff training and development
293 programme to the Federal University of Petroleum Resources, Effurun, Nigeria. This manuscript
294 benefited from the extensive and constructive reviews of two anonymous reviewers, who are duly
295 thanked.

296 **References**

- 297 Abels, H.A., Clyde, W.C., Gingerich, P.D., Hilgen, F.J., Fricke, H.C., Bowen, G.J., Lourens, L.J. 2012.
298 Terrestrial carbon isotope excursions and biotic change during Palaeogene hyperthermals.
299 Nature Geoscience, 5, 326–329.
- 300 Armstrong McKay D.I., Tyrrell, T., Wilson, P.A. 2016 Global carbon cycle perturbation across the
301 Eocene-Oligocene climate transition. Paleoclimatology, 31, 311–329.
- 302 Aggarwal N., Agrawal, S., Thakur, B. 2019. Palynofloral, palynofacies and carbon isotope of Permian
303 coal deposits from the Godavari Valley Coalfield, South India: Insights into the age,
304 palaeovegetation and palaeoclimate. International Journal of Coal Geology, 214, 103285
- 305 Aggarwal N., Mathews, R.P., Ansarie, A.H., Thakur, B. Agrawal, S., 2022. Palaeoenvironmental
306 reconstruction for the Permian (lower Gondwana) succession of the Godavari Valley

- 307 Coalfield in southern India based on a combined palynofacies, carbon isotope, and
308 biomarker study. *Journal of Palaeogeography*, 2022, 11, 123–144.
- 309 Bechtel, A., Gratzer, R., Sachsenhofer, R.F., Gusterhuber, J., Lucke, A., Puttmann, W., 2008.
310 Biomarker and carbon isotope variation in coal and fossil wood of Central Europe through
311 the Cenozoic. *Palaeogeography, Palaeoclimatology, Palaeoecology* 262, 166–175.
- 312 Blyth, F.G.H. 1962. The structure of the north-eastern tract of the Dartmoor granite. *Quarterly*
313 *Journal of the Geological Society*, 118, 435.
- 314 Bristow, C. 1968. The derivation of the Tertiary sediments in the Petrockstowe Basin, North Devon,
315 *The Proceedings of the Ussher Society*, 2, 29–35.
- 316 Bristow, C., Robson, J. 1994. Palaeogene basin development in Devon. *Transactions of the*
317 *Institution of Mining and Metallurgy. Section B. Applied Earth Science*, 103, B163–B173.
- 318 Brown C.A., 2008. *Palynological Techniques. Second Edition.* Riding J.B., Warny, S. (editors).
319 American Association of Stratigraphic Palynologists, Dallas, USA, 137 p.
- 320 Chandler, M.E.J. 1957. The Oligocene Flora of the Bovey Tracey Lake Basin, Devonshire. *British*
321 *Museum (Natural History)*, 73–123.
- 322 Chandler, M.E.J. 1964. *The Lower Tertiary Floras.* British Museum (Natural History), London, 151 pp.
- 323 Collinson, M., Hooker, J., Gröcke, D. 2003. Cobham lignite bed and penecontemporaneous
324 macrofloras of southern England: A record of vegetation and fire across the Paleocene-
325 Eocene Thermal Maximum. *Special Papers-Geological Society of America*, 333–350.
- 326 Collinson, M.E., Steart, D.C., Harrington, G., Hooker, J.J., Scott, A.C., Allen, L.O., Glasspool, I.J.,
327 Gibbson, S.J. 2009. Palynological evidence of vegetation dynamics in response to
328 palaeoenvironmental change across the onset of the Paleocene-Eocene Thermal Maximum
329 at Cobham, Southern England. *Grana*, 48, 38–66.
- 330 Coxall, H. K., Wilson, P.A., 2011, Early Oligocene glaciation and productivity in the eastern
331 equatorial Pacific: Insights into global carbon cycling, *Paleoceanography*, 26, PA2221,
332 doi:10.1029/2010PA002021.

- 333 Cui, Y., Diefendorf, A.F., Kump, L.R. Jiang, S.J. Freeman, K.H. 2021. Synchronous Marine and
334 Terrestrial Carbon Cycle Perturbation in the High Arctic During the PETM. *Paleoceanography*
335 and *Paleoclimatology* 36, Article Number e2020PA003942
- 336 DeConto, R.M., Pollard, D. 2001. Rapid Cenozoic glaciation of Antarctica induced by declining
337 atmospheric CO₂. *Nature* 421, 245–249.
- 338 Dearman, W., 1963. Wrench-faulting in Cornwall and south Devon. *Proceedings of the Geologists’*
339 *Association*, 74, 265–287.
- 340 Edwards, R.A. 1976. Tertiary sediments and structure of the Bovey Basin, south Devon. *Proceedings*
341 *of the Geologists Association*, 87, 1–26.
- 342 Fang, L., Bjerrum, C.J., Hesselbo, S.P., Kotthoff, U., McCarthy, F.M.G., Huang, B., Ditchfield, P.W.,
343 2013, Carbon-isotope stratigraphy from terrestrial organic matter through the Monterey
344 event, Miocene, New Jersey margin (IODP Expedition 313). *Geosphere*, 9, 1303–1318.
- 345 Freshney, E. 1970. Cyclical sedimentation in the Petrockstowe Basin. *Proceedings of the Ussher*
346 *Society*, 2, 179–189.
- 347 Freshney, E.C., Beer, K.E. and Wright, J.E. 1979. *Geology of the Country around Chulmleigh.*
348 *Institute of Geological Sciences* 69p.
- 349 Freshney, E.C., Edwards, R.A., Isaac, K.P., Witte, G., Wilkinson, G.C., Boulter, M.C. and Bain, J.A.
350 1982. A Tertiary basin at Dutson, near Launceston, Cornwall, England. *Proceedings of the*
351 *Geologists Association*, 93, 395–402.
- 352 Garel, S., Dupuis, C., Quesnel, F., Jacob, J., Yans, J., Magioncalda, R., Flehoc, C., Schnyder, J. 2020
353 Multiple early Eocene carbon isotope excursions associated with environmental changes in
354 the Dieppe-Hampshire Basin (NW Europe). *BSGF - Earth Sciences Bulletin*, 191, 33.
- 355 Gröcke, D.R., Price, G.D., Robinson, S.A., Baraboshkin, E., Ruffell, A.H., and Mutterlose, J., 2005, The
356 Valanginian (Early Cretaceous) positive carbon-isotope event recorded in terrestrial plants:
357 *Earth and Planetary Science Letters*, 240, 495–509.
- 358 Hacquebard, P.A., Donaldson, J.R., 1969. Carboniferous coal deposition associated with floodplain
359 and limnic environments in Nova Scotia. In: Dapples, E.C., Hopkins, M.E. (Eds.), *Environment*
360 *of Coal Deposition. Geol. Soc. America Spec. Papers* 114, 143–191.

- 361 Heimhofer, U., Hochuli, P.A., Burla, S., Andersen, N., Weissert, H., 2003. Terrestrial carbon-isotope
362 records from coastal deposits (Algarve, Portugal): a tool for chemostratigraphic correlation
363 on an intrabasinal and global scale. *Terra Nova*, 15, 8–13.
- 364 Hodgson, E., Grimes, S.T., Fitzpatrick, M.E.J., Price, G.D., Hart, M.B., Leng, M.J. 2011. Paleogene
365 carbon isotope excursions in the Bunkers Hill borehole: Hampshire Basin, UK. *Proceedings of*
366 *the Geologists Association*, 122, 460–471.
- 367 Holdgate, G.R., McGowran, B., Fromhold, T., Wagstaff, B.E. Gallagher, S.J., Wallace, M.W., Sluiter,
368 I.R.K., Whitelaw, M., 2009 Eocene–Miocene carbon-isotope and floral record from brown
369 coal seams in the Gippsland Basin of southeast Australia. *Global and Planetary Change*, 65,
370 89–103.
- 371 Holloway, S., Chadwick, R.A. 1986. The Sticklepath-Lustleigh fault zone: Tertiary sinistral
372 reactivation of a Variscan dextral strike-slip fault. *Journal of the Geological Society*, 143,
373 447–452.
- 374 Hutchinson, D.K., Coxall, H.K., Lunt, D.J., Steinthorsdottir, M de Boer, A.M., Baatsen, M., von der
375 Heydt, A., Huber, M., Kennedy-Asser, A.T., Kunzmann, L., Ladant, J.B., Lear, C.H., Moraweck,
376 K., Pearson, P.N., Piga, E., Pound, M.J., Salzmann, U., Scher, HD., Sijp, W.P., Sliwinska, K.K.,
377 Wilson, P.A., Zhang, Z.S., 2021 The Eocene-Oligocene transition: a review of marine and
378 terrestrial proxy data, models and model data comparisons. *Climate of the Past*, 17, 269–
379 315.
- 380 Jerrett, R.M., Price, G.D., Grimes, S.T., and Dawson, A.T., 2015, A paleoclimatic and
381 paleoatmospheric record from peatlands accumulating during the Cretaceous–Paleogene
382 boundary event, Western Interior Basin, Canada: *Geological Society of America Bulletin*,
383 127, 1564–1582.
- 384 Lenz, O. K., Montag, M., Wilde, V., Methner, K., Riegel, W., and Mulch, A. 2022 Early Eocene carbon
385 isotope excursions in a lignite-bearing succession at the southern edge of the proto-North
386 Sea (Schöningen, Germany). *Clim. Past*, 18, 2231–2254.
- 387 Marchionni, D.L., 1980. Petrography and depositional environments of the Liddel seam, Upper
388 Hunter Valley, New South Wales. *Int. J. Coal Geol.* 1, 36–61.

- 389 Martín-Closas, C., Permanyer, A. and Vila, M.-J. 2005. Palynofacies distribution in a lacustrine basin.
390 *Geobios*, 38, 197–210.
- 391 McInerney, F.A., Wing, S.L. 2011. The Paleocene-Eocene thermal maximum: a perturbation of
392 carbon cycle, climate, and biosphere with implications for the future. *Annual Review of*
393 *Earth and Planetary Sciences*, 39, 489–516.
- 394 Miller, K.G., Wright, J.D., Katz, M.E., Wade, B.S., Browning, J.V. and Cramer B.S. 2009, Climate
395 threshold at the Eocene-Oligocene transition: Antarctic ice sheet influence on ocean
396 circulation, in *SPE452: The Late Eocene Earth—Hothouse, Icehouse, and Impacts*, Geol. Soc.
397 of Am. Spec. Pap., 452, 80301.
- 398 Nilsen, E. B., Anderson, L. D., and Delaney, M. L. 2003. Paleoproductivity, nutrient burial, climate
399 change and the carbon cycle in the western equatorial Atlantic across the Eocene/Oligocene
400 boundary. *Paleoceanography*, 18, 1057. <https://doi.org/10.1029/2002PA000804>
- 401 Pieńkowski, G., Waksmundzka, M., 2009. Palynofacies in Lower Jurassic epicontinental deposits of
402 Poland: tool to interpret sedimentary environments. *Episodes*, 32, 21–32.
- 403 Selwood, E.B., Edwards, R., Chester, J., Hamblin, R., Henson, M., Riddolls, B. and Waters, R. 1984.
404 *Geology of the Country around Newton Abbot*. Natural Environment Research Council.
405 London: HMSO. 212 p.
- 406 Sun, J., Ni, X., Bi, S., Wu, W., Ye, J., Meng, J., Windley, B.F., 2014 Synchronous turnover of flora,
407 fauna, and climate at the Eocene–Oligocene Boundary in Asia. *Scientific Reports*, 4, 7463
- 408 Tyson, R.V. 1995. *Sedimentary Organic Matter: Organic Facies and Palynofacies*. Springer.
- 409 Wilkinson, G.C., Boulter, M.C. 1981. Oligocene pollen and spores from the western part of the
410 British Isles. *Palaeontographica Abteilung B*, 175, 27–83.
- 411 Wilkinson, G.C., Bazley, R.A.B., Boulter, M.C. 1980. The geology and palynology of the Oligocene
412 Lough Neagh Clays, Northern Ireland. *Journal of the Geological Society*, 137, 65–75.
- 413 Zachos, J.C., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms, and aberrations in
414 global climate 65 Ma to present. *Science*, 292, 686–693.
- 415
- 416

417

418 **Figure 1.** A. Map showing the southern part of the UK and location of inset. B. General geological
419 map of the southwest UK showing Petrockstowe and Bovey basins (modified from Bristow
420 and Robson 1994).

421 **Figure 2.** A. Map showing the location of Petrockstowe and Bovey basins (modified from Bristow
422 and Robson 1994). B. Geological map of the Petrockstowe Basin showing the location of
423 Borehole 1 (cores 1A and 1B) and the relative positions of the axial trough and the marginal
424 shelves with their dividing fault (modified from Freshney 1970; Freshney et al., 1979). C.
425 Geological map of the Bovey Basin showing the location of the South John Acres Lane
426 Quarry (modified from Selwood et al. 1984).



427 **Figure 3.** Summary stratigraphic logs (and correlation) of the Bovey succession with data derived
428 from (Edwards 1976) and Wilkinson (1979) cited by Selwood et al. (1984) and Petrockstowe
429 with age data from Turner (cited by Freshney et al. 1979).

430 **Figure 4.** Phytoclasts from the Petrockstowe and Bovey basins. (A) Opaque lath-shaped phytoclast,
431 sample MC95 Petrockstowe Basin; (B) Large opaque lath shaped phytoclast; sample MC3
432 Petrockstowe; (C) Multicellular fungal 'fruiting body', sample MC19 Petrockstowe; (D) Mass
433 of melanised fungal hyphae; sample MC58, Petrockstowe. (E) Multicellular fungal 'fruiting
434 body' sample MC19, Petrockstowe. (F) Well preserved, pale brown in colour, partly
435 translucent AOM sample SJAL029, Bovey Basin. (G) well preserved pale yellow AOM seen in
436 transmitted white light; sample SJAL002, Bovey Basin (H) is a cross section of plant fragment
437 sample SJAL013, Bovey Basin; (I) Phytoclast (biostructured) composed of gymnosperm
438 tracheids; sample MC76 Petrockstowe Basin.

439 **Figure 5.** TOC and $\delta^{13}\text{C}_{\text{TOC}}$ data, compared to palynological data, Petrockstowe Basin, Devon. Age
440 assignments based on Wilkinson (1979) cited by Selwood et al. (1984).

441 **Figure 6.** TOC and $\delta^{13}\text{C}_{\text{TOC}}$ data, compared to palynological data, Bovey Basin, Devon. Age
442 assignments of Turner (cited by Freshney et al. 1979).

443 **Figure 7.** Distribution of the different categories of palynological matter in the ternary diagram
444 (proposed by Hacquebard and Donaldson, 1969 and modified by Marchionni, 1980; and
445 Aggarwal et al., 2022).










Fig 3






Fig 5






Fig 6



opaque phytoclast

- Petrockstowe
- Bovey



Our m/s has not been published previously (except in the form of an abstract) and it is not under consideration for publication elsewhere. The publication is approved by all authors.

